

## THE TOTAL $\{k\}$ -DOMATIC NUMBER OF DIGRAPHS

SEYED MAHMOUD SHEIKHOESLAMI

*Department of Mathematics*  
*Azərbaycan University of Tarbiat Moallem*  
*Tarbriz, I.R. Iran*

**e-mail:** s.m.sheikholeslami@azaruniv.edu

AND

LUTZ VOLKMANN

*Lehrstuhl II für Mathematik*  
*RWTH Aachen University*  
*52056 Aachen, Germany*

**e-mail:** volkm@math2.rwth-aachen.de

### Abstract

For a positive integer  $k$ , a *total  $\{k\}$ -dominating function* of a digraph  $D$  is a function  $f$  from the vertex set  $V(D)$  to the set  $\{0, 1, 2, \dots, k\}$  such that for any vertex  $v \in V(D)$ , the condition  $\sum_{u \in N^-(v)} f(u) \geq k$  is fulfilled, where  $N^-(v)$  consists of all vertices of  $D$  from which arcs go into  $v$ . A set  $\{f_1, f_2, \dots, f_d\}$  of total  $\{k\}$ -dominating functions of  $D$  with the property that  $\sum_{i=1}^d f_i(v) \leq k$  for each  $v \in V(D)$ , is called a *total  $\{k\}$ -dominating family* (of functions) on  $D$ . The maximum number of functions in a total  $\{k\}$ -dominating family on  $D$  is the *total  $\{k\}$ -domatic number* of  $D$ , denoted by  $d_t^{\{k\}}(D)$ . Note that  $d_t^{\{1\}}(D)$  is the classic total domatic number  $d_t(D)$ . In this paper we initiate the study of the total  $\{k\}$ -domatic number in digraphs, and we present some bounds for  $d_t^{\{k\}}(D)$ . Some of our results are extensions of well-know properties of the total domatic number of digraphs and the total  $\{k\}$ -domatic number of graphs.

**Keywords:** digraph, total  $\{k\}$ -dominating function, total  $\{k\}$ -domination number, total  $\{k\}$ -domatic number.

**2010 Mathematics Subject Classification:** 05C69.

## 1. INTRODUCTION

In this paper,  $D$  is a finite and simple digraph with vertex set  $V = V(D)$  and arc set  $A = A(D)$ . The order  $|V|$  of  $D$  is denoted by  $n = n(D)$ . We write  $d_D^+(v) = d^+(v)$  for the *outdegree* of a vertex  $v$  and  $d_D^-(v) = d^-(v)$  for its *indegree*. The *minimum* and *maximum indegree* are  $\delta^-(D)$  and  $\Delta^-(D)$ . The sets  $N^+(v) = \{x | (v, x) \in A(D)\}$  and  $N^-(v) = \{x | (x, v) \in A(D)\}$  are called the *outset* and *inset* of the vertex  $v$ . If  $X \subseteq V(D)$ , then  $D[X]$  is the subdigraph induced by  $X$ . For an arc  $(x, y) \in A(D)$ , the vertex  $y$  is an *outer neighbor* of  $x$  and  $x$  is an *inner neighbor* of  $y$ . We write  $K_n^*$  for the *complete digraph* of order  $n$ . Consult [5] for the notation and terminology which are not defined here.

For a positive integer  $k$ , a *total  $\{k\}$ -dominating function* ( $T\{k\}$ DF) of a digraph  $D$  with  $\delta^-(D) \geq 1$  is a function  $f$  from the vertex set  $V(D)$  to the set  $\{0, 1, 2, \dots, k\}$  such that for any vertex  $v \in V(D)$ , the condition  $\sum_{u \in N^-(v)} f(u) \geq k$  is fulfilled. The *weight* of a  $T\{k\}$ DF  $f$  is the value  $\omega(f) = \sum_{v \in V(D)} f(v)$ . The *total  $\{k\}$ -domination number* of a digraph  $D$ , denoted by  $\gamma_t^{\{k\}}(D)$ , is the minimum weight of a  $T\{k\}$ DF of  $D$ . A  $\gamma_t^{\{k\}}(D)$ -*function* is a total  $\{k\}$ -dominating function of  $D$  with weight  $\gamma_t^{\{k\}}(D)$ . Note that  $\gamma_t^{\{1\}}(D)$  is the classical total domination number  $\gamma_t(D)$ . If  $F$  is a minimum total dominating set of a digraph  $D$  with  $\delta^-(D) \geq 1$ , then the function  $f$  from  $V(D)$  to  $\{0, 1, 2, \dots, k\}$  with  $f(v) = k$  for  $v \in F$  and  $f(x) = 0$  for  $x \in V(D) - F$  is a total  $\{k\}$ -dominating function of  $D$  and therefore

$$\gamma_t^{\{k\}}(D) \leq k|F| = k\gamma_t(D).$$

In this paper we always assume that  $D$  is a digraph with  $\delta^-(D) \geq 1$ .

A set  $\{f_1, f_2, \dots, f_d\}$  of distinct total  $\{k\}$ -dominating functions of  $D$  with the property that  $\sum_{i=1}^d f_i(v) \leq k$  for each  $v \in V(D)$ , is called a *total  $\{k\}$ -dominating family* (of functions) on  $D$ . The maximum number of functions in a total  $\{k\}$ -dominating family ( $T\{k\}$ D family) on  $D$  is the *total  $\{k\}$ -domatic number* of  $D$ , denoted by  $d_t^{\{k\}}(D)$ . The total  $\{k\}$ -domatic number is well-defined and

$$(1) \quad d_t^{\{k\}}(D) \geq 1, \text{ for all digraphs } D \text{ with } \delta^-(D) \geq 1,$$

since the set consisting of the function  $f : V(D) \rightarrow \{0, 1, 2, \dots, k\}$  defined by  $f(v) = k$  for each  $v \in V(D)$ , forms a  $T\{k\}$ D family on  $D$ . The total domatic number of a digraph was introduced by Jacob and Arumugam in [6].

Our purpose in this paper is to initiate the study of the total  $\{k\}$ -domatic number in digraphs. We first study basic properties and bounds for the total  $\{k\}$ -domatic number of a digraph. In addition, we determine the total  $\{k\}$ -domatic number of some classes of digraphs. Some of our results are extensions of well-know properties of the total domatic number of digraphs and the total  $\{k\}$ -domatic number of graphs (see, for example, [2, 3, 4, 6, 8]).

We start with the following observation.

**Observation 1.** *Let  $k$  be an integer, and let  $D$  be a digraph with  $\delta^-(D) \geq 1$ . Then  $\gamma_t^{\{k\}}(D) \geq k+1$ , with equality if and only if there exists a subset  $S \subseteq V(D)$  of size  $k+1$  such that  $D[S]$  is a complete digraph, and each vertex  $x \in V(D) - S$  has at least  $k$  inner neighbors in  $S$ .*

**Proof.** Let  $f$  be a  $\gamma_t^{\{k\}}(D)$ -function, and let  $v \in V(D)$  be an arbitrary vertex. The definition implies that  $\sum_{x \in N^-(v)} f(x) \geq k$ . If  $\sum_{x \in N^-(v)} f(x) \geq k+1$ , then  $\gamma_t^{\{k\}}(D) \geq k+1$ . If  $\sum_{x \in N^-(v)} f(x) = k$ , then let  $u \in N^-(v)$  be a vertex such that  $f(u) \geq 1$ . Since  $\sum_{x \in N^-(u)} f(x) \geq k$  and  $u \notin N^-(u)$ , we deduce that  $\omega(f) = \sum_{x \in V(D)} f(v) \geq \sum_{x \in (N^-(u) \cup \{u\})} f(x) \geq k+1$  and therefore  $\gamma_t^{\{k\}}(D) \geq k+1$ .

Assume that  $\gamma_t^{\{k\}}(D) = k+1$ . Let  $f$  be a  $\gamma_t^{\{k\}}(D)$ -function. If there exists a vertex  $v$  such that  $f(v) \geq 2$ , then we obtain the contradiction  $\sum_{x \in N^-(v)} f(x) \leq k+1-2 = k-1$ . Hence  $f(x) = 1$  or  $f(x) = 0$  for each vertex  $x \in V(D)$ . Let  $S \subseteq V(D)$  such that  $f(x) = 1$  for each  $x \in S$ . Then  $|S| = k+1$ ,  $D[S]$  is a complete digraph, and each vertex  $x \in V(D) - S$  has at least  $k$  inner neighbors in  $S$ .

Conversely, assume that there exists a subset  $S \subseteq V(D)$  of size  $k+1$  such that  $D[S]$  is a complete digraph, and each vertex  $x \in V(D) - S$  has at least  $k$  inner neighbors in  $S$ . Define the function  $f$  by  $f(x) = 1$  for  $x \in S$  and  $f(x) = 0$  for  $x \in V(D) - S$ . Then  $f$  is a total  $\{k\}$ -dominating function of  $D$  such that  $\omega(f) = k+1$ . Since  $\gamma_t^{\{k\}}(D) \geq k+1$ , we deduce that  $\gamma_t^{\{k\}}(D) = k+1$ . ■

## 2. PROPERTIES OF THE $\{k\}$ -DOMATIC NUMBER

In this section we mainly present basic properties of  $d_t^{\{k\}}(D)$  and bounds on the total  $\{k\}$ -domatic number of a digraph.

**Theorem 2.** *If  $D$  is a digraph of order  $n$ , then  $\gamma_t^{\{k\}}(D) \cdot d_t^{\{k\}}(D) \leq kn$ . Moreover, if  $\gamma_t^{\{k\}}(D) \cdot d_t^{\{k\}}(D) = kn$ , then for each  $T\{k\}D$  family  $\{f_1, f_2, \dots, f_d\}$  on  $D$  with  $d = d_t^{\{k\}}(D)$ , each function  $f_i$  is a  $\gamma_t^{\{k\}}(D)$ -function and  $\sum_{i=1}^d f_i(v) = k$  for all  $v \in V(D)$ .*

**Proof.** Let  $\{f_1, f_2, \dots, f_d\}$  be a  $T\{k\}D$  family on  $D$  such that  $d = d_t^{\{k\}}(D)$ . Then

$$\begin{aligned} d \cdot \gamma_t^{\{k\}}(D) &= \sum_{i=1}^d \gamma_t^{\{k\}}(D) \leq \sum_{i=1}^d \sum_{v \in V(D)} f_i(v) = \sum_{v \in V(D)} \sum_{i=1}^d f_i(v) \\ &\leq \sum_{v \in V(D)} k = kn. \end{aligned}$$

If  $\gamma_t^{\{k\}}(D) \cdot d_t^{\{k\}}(D) = kn$ , then the two inequalities occurring in the proof become equalities. Hence for the  $T\{k\}D$  family  $\{f_1, f_2, \dots, f_d\}$  on  $D$  and for each  $i$ ,

$\sum_{v \in V(D)} f_i(v) = \gamma_t^{\{k\}}(D)$ . Thus each function  $f_i$  is a  $\gamma_t^{\{k\}}(D)$ -function, and  $\sum_{i=1}^d f_i(v) = k$  for all  $v \in V(D)$ . ■

The special case  $k = 1$  in Theorem 2 can be found in [6].

**Corollary 3.** *Let  $k, n$  be two positive integers. If  $k + 1$  is a divisor of  $n$  and  $\frac{n}{k+1} \geq 2$ , then  $d_t^{\{k\}}(K_n^*) = \frac{kn}{k+1}$ .*

**Proof.** Applying Observation 1 and Theorem 2, we see that  $d_t^{\{k\}}(K_n^*) \leq \frac{kn}{k+1}$ .

Now we consider a partition of  $V(K_n^*)$  into  $s = \frac{n}{k+1}$  sets  $V_1, V_2, \dots, V_s$  such that  $|V_i| = k + 1$  for each  $i$ . Let  $V_i = \{v_1^i, v_2^i, \dots, v_{k+1}^i\}$  for  $1 \leq i \leq s$ . Define, for  $1 \leq i \leq s$  and  $1 \leq j \leq k$ ,

$$f_i^j(v_1^i) = \dots = f_i^j(v_j^i) = 1, f_i^j(v_{j+1}^i) = \dots = f_i^j(v_{k+1}^i) = 1 \text{ and} \\ f_i^j(x) = 0 \text{ otherwise, where the indices } i + 1 \text{ are taken modulo } s.$$

It is easy to see that  $\{f_i^j \mid 1 \leq i \leq \frac{n}{k+1}, 1 \leq j \leq k\}$  is a  $T\{k\}D$  family on  $K_n^*$ , and therefore  $d_t^{\{k\}}(K_n^*) \geq \frac{kn}{k+1}$ . Since  $k + 1$  is a divisor of  $n$ , the proof is complete. ■

A further consequence of Theorem 2 and Observation 1 now follows.

**Corollary 4.** *If  $k \geq 2$  is an integer, and  $D$  is a digraph of order  $k + 1$ , then  $d_t^{\{k\}}(D) \leq k - 1$ .*

**Proof.** Since  $\gamma_t^{\{k\}}(D) \geq k + 1$ , it follows from Theorem 2 that  $d_t^{\{k\}}(D) \leq k$ . If  $\gamma_t^{\{k\}}(D) \geq k + 2$ , then Theorem 2 implies  $d_t^{\{k\}}(D) \leq k - 1$  immediately. If  $\gamma_t^{\{k\}}(D) = k + 1$  and  $d_t^{\{k\}}(D) = k$ , then for the  $T\{k\}D$  family  $\{f_1, f_2, \dots, f_k\}$  on  $D$ , each function  $f_i$  is a  $\gamma_t^{\{k\}}(D)$ -function, and Observation 1 leads to the contradiction that  $f_1 \equiv f_2 \equiv \dots \equiv f_k$ . This completes the proof. ■

**Corollary 5.** *If  $k$  is a positive integer, and  $D$  is a digraph of order  $n$ , then  $d_t^{\{k\}}(D) \leq \frac{kn}{k+1}$ , with equality only if  $k + 1$  is a divisor of  $n$  and  $\frac{n}{k+1} \geq 2$  when  $k \geq 2$ .*

**Proof.** Since  $\gamma_t^{\{k\}}(D) \geq k + 1$ , it follows from Theorem 2 that  $d_t^{\{k\}}(D) \leq \frac{kn}{\gamma_t^{\{k\}}(D)} \leq \frac{kn}{k+1}$ , and this is the desired inequality.

Assume that  $d_t^{\{k\}}(D) = \frac{kn}{k+1}$ . Since  $(k, k + 1) = 1$ ,  $k + 1$  must be a divisor of  $n$ . If  $k \geq 2$ , then it follows from Corollary 4 that  $\frac{n}{k+1} \geq 2$ . ■

Corollary 3 demonstrates that Corollary 5 is sharp.

**Theorem 6.** *If  $D$  is a digraph of order  $n$  and  $k$  a positive integer, then*

$$\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq nk + 1.$$

**Proof.** Applying Theorem 2, we obtain  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \frac{kn}{d_t^{\{k\}}(D)} + d_t^{\{k\}}(D)$ .

Note that  $d_t^{\{k\}}(G) \geq 1$ , by inequality (1), and that Corollary 5 implies that  $d_t^{\{k\}}(D) \leq n$ . Using these inequalities, and the fact that the function  $g(x) = x + (kn)/x$  is decreasing for  $1 \leq x \leq \sqrt{kn}$  and increasing for  $\sqrt{kn} \leq x \leq n$ , we obtain  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \max\{kn + 1, \frac{kn}{n} + n\} = nk + 1$ , and this is the desired bound. ■

If  $C_n$  denotes a directed cycle on  $n$  vertices, then the function  $f : V(C_n) \rightarrow \{0, 1, \dots, k\}$  defined by  $f(x) = k$  for each  $x \in V(C_n)$  is the unique total  $\{k\}$ -dominating function of  $C_n$  and hence  $\gamma_t^{\{k\}}(C_n) = nk$  and  $d_t^{\{k\}}(C_n) = 1$ . This demonstrates that Theorem 6 is sharp.

**Theorem 7.** Let  $D$  be a digraph of order  $n \geq 3$ , and let  $k \geq 1$  be an integer. If  $d_t^{\{k\}}(D) \geq 2$ , then  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \frac{kn}{2} + 2$ .

**Proof.** Theorem 2 implies that  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \gamma_t^{\{k\}}(D) + \frac{kn}{\gamma_t^{\{k\}}(D)}$ . It follows from Observation 1 and Theorem 2 that  $k + 1 \leq \gamma_t^{\{k\}}(D) \leq kn/2$ . Using these inequalities, and the fact that the function  $g(x) = x + (kn)/x$  is decreasing for  $k + 1 \leq x \leq \sqrt{kn}$  and increasing for  $\sqrt{kn} \leq x \leq kn/2$ , we obtain

$$\gamma_t^{\{k\}}(G) + d_t^{\{k\}}(G) \leq \max\left\{k + 1 + \frac{kn}{k+1}, \frac{kn}{2} + 2\right\} = \frac{kn}{2} + 2,$$

and this is the desired bound. ■

**Theorem 8.** If  $D$  is a digraph and  $k \geq 1$  an integer, then  $d_t^{\{k\}}(D) \leq \delta^-(D)$ . Moreover, if  $d_t^{\{k\}}(D) = \delta^-(D)$ , then for each function of any  $T\{k\}D$  family  $\{f_1, f_2, \dots, f_d\}$  and for all vertices  $v$  of indegree  $\delta^-(D)$ ,  $\sum_{u \in N^-(v)} f_i(u) = k$  and  $\sum_{i=1}^d f_i(u) = k$  for every  $u \in N^-(v)$ .

**Proof.** Let  $\{f_1, f_2, \dots, f_d\}$  be a  $T\{k\}D$  family on  $D$  such that  $d = d_t^{\{k\}}(D)$ , and let  $v$  be a vertex of minimum indegree  $\delta^-(D)$ . Since  $\sum_{u \in N^-(v)} f_i(u) \geq k$  for all  $i \in \{1, 2, \dots, d\}$ , we obtain  $kd \leq \sum_{i=1}^d \sum_{u \in N^-(v)} f_i(u) = \sum_{u \in N^-(v)} \sum_{i=1}^d f_i(u) \leq \sum_{u \in N^-(v)} k = k\delta^-(D)$ , and this leads to the desired bound.

If  $d_t^{\{k\}}(D) = \delta^-(D)$ , then the two inequalities occurring in the proof become equalities, which leads to the two properties given in the statement. ■

The special case  $k = 1$  in Theorem 8 can be found in [6].

**Observation 9.** Let  $D$  be a digraph with the property that the underlying graph is connected and bipartite. If  $k \geq 1$  is an integer, then  $\gamma_t^{\{k\}}(D) \geq 2k$ .

**Proof.** Let  $f$  be a  $\gamma_t^{\{k\}}(D)$ -function, and let  $V_1$  and  $V_2$  be the partite sets of the underlying graph. If  $w_i \in V_i$ , then the definition implies that  $\sum_{x \in N^-(w_i)} f(x) \geq k$  for  $i = 1, 2$ . It follows that  $w(f) = \sum_{x \in V(D)} f(x) = \sum_{x \in V(D)-V_1} f(x) + \sum_{x \in V(D)-V_2} f(x) \geq \sum_{x \in N^-(w_2)} f(x) + \sum_{x \in N^-(w_1)} f(x) \geq 2k$ , thus  $\gamma_t^{\{k\}}(D) \geq 2k$ . ■

**Corollary 10.** *If  $K_{p,p}^*$  is the complete bipartite digraph and  $k \geq 1$  an integer, then  $d_t^{\{k\}}(K_{p,p}^*) = p$ .*

**Proof.** Theorem 2 and Observation 9 show that  $d_t^{\{k\}}(K_{p,p}^*) \leq p$ .

Now let  $\{u_1, u_2, \dots, u_p\}$  and  $\{v_1, v_2, \dots, v_p\}$  be the partite sets of the complete bipartite digraph. Define  $f_i(u_i) = f_i(v_i) = k$  and  $f_i(x) = 0$  for each vertex  $x \in V(D) - \{u_i, v_i\}$  and each  $i \in \{1, 2, \dots, p\}$ . Then we observe that  $f_i$  is a  $T\{k\}$ DF of  $K_{p,p}^*$  for each  $i \in \{1, 2, \dots, p\}$ . Therefore  $\{f_1, f_2, \dots, f_p\}$  is a  $T\{k\}$ D family on  $K_{p,p}^*$ . Consequently,  $d_t^{\{k\}}(K_{p,p}^*) \geq p$  and so  $d_t^{\{k\}}(K_{p,p}^*) = p$ . ■

Corollary 10 demonstrates that Theorem 8 is sharp.

**Theorem 11.** *Let  $k \geq 1$  be an integer and  $D$  a digraph of order  $n$  with  $\delta^-(D) \geq 1$ . If  $\delta^-(D) \mid k$ , then  $d_t^{\{k\}}(D) \geq \delta^-(D) - 1$ .*

**Proof.** If  $\delta^-(D) = 1$ , then the result is immediate.

Let  $\delta^-(D) \geq 2$  and let  $V(D) = \{v_1, v_2, \dots, v_n\}$ . Define  $f_i : V(D) \rightarrow \{0, 1, \dots, k\}$  by

$$f_i(v_j) = \begin{cases} \frac{k}{\delta^-(D)} + 1 & \text{if } j = i, \\ \frac{k}{\delta^-(D)} & \text{if } j \neq i, \end{cases} \quad \text{for every } 1 \leq i \leq \delta^-(D) - 1 \text{ and } 1 \leq j \leq n.$$

Then for each  $v \in V(D)$  and each  $1 \leq i \leq \delta^-(D) - 1$ ,

$$\sum_{u \in N^-(v)} f_i(u) \geq \sum_{u \in N^-(v)} \frac{k}{\delta^-(D)} \geq \frac{k}{\delta^-(D)} \delta^-(D) = k.$$

Hence  $f_i$  is a  $T\{k\}$ DF of  $D$  for each  $1 \leq i \leq \delta^-(D) - 1$ . Now, since  $\delta^-(D) \mid k$ , we have

$\sum_{i=1}^{\delta^-(D)-1} f_i(v) \leq \frac{k}{\delta^-(D)} (\delta^-(D) - 2) + \left( \frac{k}{\delta^-(D)} + 1 \right) = k + \left( 1 - \frac{k}{\delta^-(D)} \right) \leq k$  for each  $v \in V(D)$ . Thus  $\{f_1, f_2, \dots, f_{\delta^-(D)-1}\}$  is a  $T\{k\}$ D family on  $D$ , and the proof is complete. ■

**Theorem 12.** *Let  $k \geq 1$  be an integer and  $D$  a digraph of order  $n$ . If  $\delta^-(D) \nmid k$ , then  $d_t^{\{k\}}(D) \geq \left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor$ .*

**Proof.** Let  $V(D) = \{v_1, v_2, \dots, v_n\}$ . Define  $f_i : V(D) \rightarrow \{0, 1, \dots, k\}$  by

$$f_i(v_j) = \begin{cases} \lfloor \frac{k}{\delta^-(D)} \rfloor & \text{if } j = i, \\ \lceil \frac{k}{\delta^-(D)} \rceil & \text{if } j \neq i, \end{cases} \quad \text{for every } 1 \leq i \leq \left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor \text{ and } 1 \leq j \leq n.$$

Then for each  $v \in V(D)$  and each  $1 \leq i \leq \left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor$ ,

$$\sum_{u \in N^-(v)} f_i(u) \geq \left\lfloor \frac{k}{\delta^-(D)} \right\rfloor + \left\lceil \frac{k}{\delta^-(D)} \right\rceil (\delta^-(D) - 1) \geq \left\lceil \frac{k}{\delta^-(D)} \right\rceil \delta^-(D) - 1 \geq k.$$

Hence  $f_i$  is a  $T\{k\}$ DF of  $D$  for each  $i$ . Since  $\delta^-(D) \nmid k$ , we have

$$\sum_{i=1}^{\left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor} f_i(v) \leq \left\lceil \frac{k}{\delta^-(D)} \right\rceil \cdot \left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor \leq \left\lceil \frac{k}{\delta^-(D)} \right\rceil \cdot \frac{k}{\lceil k/\delta^-(D) \rceil} = k$$

for each  $v \in V(D)$ . Thus  $\{f_1, f_2, \dots, f_{\left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor}\}$  is a  $T\{k\}$ D family on  $D$ , and the proof is complete. ■

Using Theorems 2, 8, 11 and 12, we will improve Theorem 6 considerably for some cases.

**Corollary 13.** *Let  $k \geq 1$  be an integer, and let  $D$  be a digraph of order  $n$ . If  $\delta^-(D) > k$ , then  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq n + k$ .*

**Proof.** Since  $\delta^-(D) > k$ , it follows from Theorem 12 that

$$d_t^{\{k\}}(D) \geq \left\lfloor \frac{k}{\lceil k/\delta^-(D) \rceil} \right\rfloor = k.$$

In addition, Theorem 8 implies that  $d_t^{\{k\}}(D) \leq \delta^-(D) \leq n$ . Using these two inequalities, and the fact that the function  $g(x) = x + (kn)/x$  is decreasing for  $k \leq x \leq \sqrt{kn}$  and increasing for  $\sqrt{kn} \leq x \leq n$ , Theorem 2 leads to

$$\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \frac{kn}{d_t^{\{k\}}(D)} + d_t^{\{k\}}(D) \leq \max \left\{ \frac{kn}{k} + k, \frac{kn}{n} + n \right\} = n + k.$$

This is the desired bound, and the proof is complete. ■

**Corollary 14.** *Let  $k \geq 1$  be an integer, and let  $D$  be a digraph of order  $n$  with  $\delta^-(D) \geq 2$ . If  $\delta^-(D) \mid k$ , then  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \frac{kn}{\delta^-(D)-1} + \delta^-(D) - 1$ .*

**Proof.** Since  $\delta^-(D) \mid k$ , Theorem 11 shows that  $d_t^{\{k\}}(D) \geq \delta^-(D) - 1$ , and Theorem 8 implies that  $d_t^{\{k\}}(D) \leq \delta^-(D)$ . Using these two inequalities and Theorem 2, we obtain the desired bound as follows  $\gamma_t^{\{k\}}(D) + d_t^{\{k\}}(D) \leq \frac{kn}{d_t^{\{k\}}(D)} + d_t^{\{k\}}(D) \leq \max \left\{ \frac{kn}{\delta^-(D)-1} + \delta^-(D) - 1, \frac{kn}{\delta^-(D)} + \delta^-(D) \right\} = \frac{kn}{\delta^-(D)-1} + \delta^-(D) - 1$ . ■

Let  $D$  be a digraph. By  $D^{-1}$  we denote the digraph obtained by reversing all arcs of  $D$ . A digraph without directed cycles of length 2 is called an *oriented graph*. An oriented graph  $D$  is a *tournament* when either  $(x, y) \in A(D)$  or  $(y, x) \in A(D)$  for each pair of distinct vertices  $x, y \in V(D)$ .

**Theorem 15.** *For every oriented graph  $D$  with  $\delta^-(D) \geq 1$  and  $\delta^-(D^{-1}) \geq 1$ ,  $d_t^{\{k\}}(D) + d_t^{\{k\}}(D^{-1}) \leq n - 1$ . If  $d_t^{\{k\}}(D) + d_t^{\{k\}}(D^{-1}) = n - 1$ , then  $D$  is a regular tournament.*

**Proof.** Since  $\delta^-(D) + \delta^-(D^{-1}) \leq n - 1$ , Theorem 8 leads to

$$d_t^{\{k\}}(D) + d_t^{\{k\}}(D^{-1}) \leq \delta^-(D) + \delta^-(D^{-1}) \leq n - 1.$$

If  $D$  is not a tournament or  $D$  is a non-regular tournament, then  $\delta^-(D) + \delta^-(D^{-1}) \leq n - 2$ , and hence we deduce from Theorem 8 that

$$d_t^{\{k\}}(D) + d_t^{\{k\}}(D^{-1}) \leq \delta^-(D) + \delta^-(D^{-1}) \leq n - 2. \quad \blacksquare$$

Now we present further lower bounds on the total  $\{k\}$ -domatic number.

**Theorem 16.** *Let  $k \geq 1$  be an integer, and  $D$  a digraph with  $\delta^-(D) = \delta^- \geq 1$ .*

- (i) *If  $k < \delta^-$ , then  $d_t^{\{k\}}(D) \geq k$ .*
- (ii) *If  $k = p\delta^-$  with an integer  $p \geq 1$ , then  $d_t^{\{k\}}(D) \geq \delta^- - 1$ .*
- (iii) *If  $k = p\delta^- + r$  with integers  $p, r \geq 1$  and  $r \leq \delta^- - 1$ , then*  

$$d_t^{\{k\}}(D) \geq \left\lceil \frac{p(\delta^- - 1) + 1}{p + 1} \right\rceil.$$

**Proof.** (i) If  $k < \delta^-$ , then Theorem 12 implies immediately  $d_t^{\{k\}}(D) \geq k$ .

(ii) If  $k = p\delta^-$ , then Theorem 11 implies immediately  $d_t^{\{k\}}(D) \geq \delta^- - 1$ .

(iii) If  $k = p\delta^- + r$  with integers  $p, r \geq 1$  and  $r \leq \delta^- - 1$ , then  $\lceil \frac{k}{\delta^-} \rceil = p + 1$  and therefore we deduce from Theorem 12 that

$$d_t^{\{k\}}(D) \geq \left\lfloor \frac{k}{\lceil k/\delta^- \rceil} \right\rfloor = \left\lfloor \frac{k}{p+1} \right\rfloor = \left\lfloor \frac{p\delta^- + r}{p+1} \right\rfloor \geq \frac{p\delta^- + r}{p+1} - \frac{p}{p+1} \geq \frac{p(\delta^- - 1) + 1}{p+1}.$$

This leads to the desired bound, and the proof is complete.  $\blacksquare$

**Corollary 17.** *If  $k \geq 1$  is an integer and  $D$  a digraph with  $\delta^-(D) \geq 1$ , then*  

$$d_t^{\{k\}}(D) \geq \min \left\{ k, \frac{\delta^-(D)}{2} \right\}.$$

The *complement*  $\overline{D}$  of a digraph  $D$  is that digraph with vertex set  $V(D)$  such that for two arbitrary different vertices  $u$  and  $v$  the arc  $(u, v)$  belongs to  $\overline{D}$  if and only if  $(u, v)$  does not belong to  $D$ .



**Theorem 18.** *Let  $k \geq 1$  be an integer, and let  $D$  be an  $r$ -diregular digraph of order  $n \geq 3$  with  $1 \leq r \leq n - 2$ . Then  $d_t^{\{k\}}(D) + d_t^{\{k\}}(\overline{D}) \geq \min \{k + 1, \lceil \frac{n-1}{2} \rceil\}$ .*

**Proof.** Assume first that  $k < \delta^-(D)$ . Then it follows from Theorem 16 (i) that  $d_t^{\{k\}}(D) \geq k$  and thus  $d_t^{\{k\}}(D) + d_t^{\{k\}}(\overline{D}) \geq k + 1$ .

Assume next that  $k \geq \delta^-(D)$  and  $k < \delta^-(\overline{D})$ . Then Theorem 16 (i) implies  $d_t^{\{k\}}(\overline{D}) \geq k$  and so  $d_t^{\{k\}}(D) + d_t^{\{k\}}(\overline{D}) \geq k + 1$ .

Finally assume that  $k \geq \delta^-(D)$  and  $k \geq \delta^-(\overline{D})$ . Applying Theorem 16 (ii) and (iii), we observe that  $d_t^{\{k\}}(D) \geq \delta^-(D)/2$  and  $d_t^{\{k\}}(\overline{D}) \geq \delta^-(\overline{D})/2$ , and hence we deduce that

$$d_t^{\{k\}}(D) + d_t^{\{k\}}(\overline{D}) \geq \frac{\delta^-(D)}{2} + \frac{\delta^-(\overline{D})}{2} = \frac{n-1}{2}.$$

Combining these inequalities, we obtain the desired bound.  $\blacksquare$

**Theorem 19.** *For every digraph  $D$  of order  $n$ ,  $d_t^{\{k\}}(D) \geq \lfloor \frac{n}{n-\delta^-(D)} \rfloor$ .*

**Proof.** Let  $S$  be any subset of  $V(D)$  with  $|S| \geq n - \delta^-(D)$ . If  $v \in V(D) - S$ , then there exists at least one vertex  $u \in S$  such that  $(u, v) \in A(D)$ . Let  $S_1, S_2, \dots, S_{\lfloor \frac{n}{n-\delta^-(D)} \rfloor}$  be disjoint subsets of  $V(D)$  each of cardinality  $n - \delta^-(D)$ . Define  $f_i : V(G) \rightarrow \{0, 1, \dots, k\}$  by

$$f_i(v) = \begin{cases} k & \text{if } v \in S_i, \\ 0 & \text{otherwise,} \end{cases}$$

for each  $1 \leq i \leq \lfloor \frac{n}{n-\delta^-(D)} \rfloor$ .

Since  $|S_i| = n - \delta^-(D)$ , it is clear that  $f_i$  is a total  $\{k\}$ -dominating function of  $D$  for each  $i$ . Since also  $S_i$  are disjoint subsets of  $V(D)$ , then for every  $v \in V(D)$   $\sum_{i=1}^{\lfloor \frac{n}{n-\delta^-(D)} \rfloor} f_i(v) \leq k$ . Thus  $\{f_1, f_2, \dots, f_{\lfloor \frac{n}{n-\delta^-(D)} \rfloor}\}$  is a  $T\{k\}D$  family on  $D$ , and the proof is complete.  $\blacksquare$

The special case  $k = 1$  in Theorems 15 and 19 can be found in [6].

### 3. THE TOTAL $\{k\}$ -DOMATIC NUMBER OF GRAPHS

The *total  $\{k\}$ -dominating function* of a graph  $G$  is defined in [7] as a function  $f : V(G) \rightarrow \{0, 1, 2, \dots, k\}$  such that  $\sum_{x \in N_G(v)} f(x) \geq k$  for all  $v \in V(G)$ . The sum  $\sum_{x \in V(G)} f(x)$  is the weight  $w(f)$  of  $f$ . The minimum of weights  $w(f)$ , taken over all total  $\{k\}$ -dominating functions  $f$  on  $G$  is called the *total  $\{k\}$ -domination number* of  $G$ , denoted by  $\gamma_t^{\{k\}}(G)$ . In the special case  $k = 1$ ,  $\gamma_t^{\{k\}}(G)$  is the classical total domination number  $\gamma_t(G)$ .

A set  $\{f_1, f_2, \dots, f_d\}$  of distinct total  $\{k\}$ -dominating functions on  $G$  such that  $\sum_{i=1}^d f_i(v) \leq k$  for each  $v \in V(G)$ , is called a *total  $\{k\}$ -dominating family* on  $G$ . The maximum number of functions in a total  $\{k\}$ -dominating family on  $G$  is the *total  $\{k\}$ -domatic number* of  $G$ , denoted by  $d_t^{\{k\}}(G)$ . This parameter was introduced by Sheikholeslami and Volkmann in [8] and has been studied in [1]. In the case  $k = 1$ , we write  $d_t(G)$  instead of  $d_t^{\{1\}}(G)$  which was introduced by Cockayne, Dawes and Hedetniemi [3], and has been studied in many articles.

The *associated digraph*  $D(G)$  of a graph  $G$  is the digraph obtained from  $G$  when each edge  $e$  of  $G$  is replaced by two oppositely oriented arcs with the same ends as  $e$ . Since  $N_{D(G)}^-(v) = N_G(v)$  for each vertex  $v \in V(G) = V(D(G))$ , the following useful observation is valid.

**Observation 20.** *If  $D(G)$  is the associated digraph of a graph  $G$ , then*

$$\gamma_t^{\{k\}}(D(G)) = \gamma_t^{\{k\}}(D) \text{ and } d_t^{\{k\}}(D(G)) = d_t^{\{k\}}(D).$$

There are a lot of interesting applications of Observation 20. Using Theorems 2 and 6, we obtain the next results immediately.

**Corollary 21** [8]. *If  $k \geq 1$  is an integer and  $G$  a graph of order  $n$  without isolated vertices, then  $\gamma_t^{\{k\}}(G) \cdot d_t^{\{k\}}(G) \leq kn$ .*

The case  $k = 1$  in Corollary 21 leads to the well-known inequality  $\gamma_t(G) \cdot d_t(G) \leq n$ , given by Cockayne, Dawes and Hedetniemi [3] in 1980.

**Corollary 22** [8]. *If  $k \geq 1$  is an integer and  $G$  a graph of order  $n$  without isolated vertices, then  $\gamma_t^{\{k\}}(G) + d_t^{\{k\}}(G) \leq nk + 1$ .*

**Corollary 23** [3]. *If  $G$  is graph of order  $n$  without isolated vertices, then  $\gamma_t(G) + d_t(G) \leq n + 1$ .*

Theorem 7 and Observation 20 lead to the following bound.

**Corollary 24** [8]. *Let  $k \geq 1$  be an integer and  $G$  a graph of order  $n$  without isolated vertices. If  $d_t^{\{k\}}(G) \geq 2$ , then  $\gamma_t^{\{k\}}(G) + d_t^{\{k\}}(G) \leq \frac{kn}{2} + 2$ .*

**Corollary 25** [4]. *If  $G$  is a graph of order  $n$  without isolated vertices and if  $d_t(G) \geq 2$ , then  $\gamma_t(G) + d_t(G) \leq \frac{n}{2} + 2$ .*

Since  $\delta^-(D(G)) = \delta(G)$ , the next result follows from Observation 20 and Theorem 8.

**Corollary 26** [8]. *If  $k \geq 1$  is an integer and  $G$  a graph without isolated vertices, then  $d_t^{\{k\}}(G) \leq \delta(G)$ .*

The case  $k = 1$  in Corollary 26 can be found in [3]. Theorem 11 and Observation 20 imply the next result.

**Corollary 27** [2]. *Let  $k \geq 1$  be an integer and  $G$  a graph of order  $n$  without isolated vertices. If  $\delta(G) \mid k$ , then  $d_t^{\{k\}}(G) \geq \delta(G) - 1$ .*

Finally, the next theorem follows from Theorem 18 and Observation 20.

**Corollary 28** [1]. *For every  $\delta$ -regular graph of order  $n \geq 5$  in which neither  $G$  nor  $\overline{G}$  have isolated vertices,  $d_t^{\{k\}}(G) + d_t^{\{k\}}(\overline{G}) \geq \min\{k + 1, \lceil \frac{n-2}{2} \rceil\}$ .*

## REFERENCES

- [1] H. Aram, S.M. Sheikholeslami and L. Volkmann, *On the total  $\{k\}$ -domination and  $\{k\}$ -domatic number of a graph*, Bull. Malays. Math. Sci. Soc. (to appear).
- [2] J. Chen, X. Hou and N. Li, *The total  $\{k\}$ -domatic number of wheels and complete graphs*, J. Comb. Optim. (to appear).
- [3] E.J. Cockayne, R.M. Dawes and S.T. Hedetniemi, *Total domination in graphs*, Networks **10** (1980) 211–219.  
doi:10.1002/net.3230100304
- [4] E.J. Cockayne, T.W. Haynes, S.T. Hedetniemi, Z. Shanchao and B. Xu, *Extremal graphs for inequalities involving domination parameters*, Discrete Math. **216** (2000) 1–10.  
doi:10.1016/S0012-365X(99)00251-4
- [5] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in graphs* (New York: Marcel Dekker, Inc., 1998).
- [6] K. Jacob and S. Arumugam, *Domestic number of a digraph*, Bull. Kerala Math. Assoc. **2** (2005) 93–103.
- [7] N. Li and X. Hou, *On the total  $\{k\}$ -domination number of Cartesian products of graphs*, J. Comb. Optim. **18** (2009) 173–178.  
doi:10.1007/s10878-008-9144-2
- [8] S.M. Sheikholeslami and L. Volkmann, *The total  $\{k\}$ -domatic number of a graph*, J. Comb. Optim. **23** (2012) 252–260.  
doi:10.1007/s10878-010-9352-4

Received 31 March 2011

Revised 29 August 2011

Accepted 30 August 2011

