

## ERDŐS–KO–RADO FROM INTERSECTING SHADOWS

GYULA O.H. KATONA AND ÁKOS KISVÖLCSEY

*Alfréd Rényi Institute of Mathematics,  
Hungarian Academy of Sciences  
1053 Budapest, Reáltanoda u. 13–15, Hungary*

**e-mail:** {ohkatona,ksvlcs}@renyi.hu

### Abstract

A set system is called  $t$ -intersecting if every two members meet each other in at least  $t$  elements. Katona determined the minimum ratio of the shadow and the size of such families and showed that the Erdős–Ko–Rado theorem immediately follows from this result. The aim of this note is to reproduce the proof to obtain a slight improvement in the Kneser graph. We also give a brief overview of corresponding results.

**Keywords:** Kneser graph, coclique, intersecting family, shadow.

**2010 Mathematics Subject Classification:** 05C35, 05D05.

### 1. INTRODUCTION

Throughout the paper we will investigate subsets of an  $n$ -element underlying set  $[n] = \{1, 2, \dots, n\}$ .  $\binom{[n]}{k}$  will denote the collection of all  $k$ -element subsets of  $[n]$ . A family  $\mathcal{F}$  is said to be  $k$ -uniform if  $\mathcal{F} \subseteq \binom{[n]}{k}$ .

$\mathcal{F} \subseteq \binom{[n]}{k}$  is called *intersecting* if it does not contain disjoint sets. In general,  $\mathcal{F}$  is  *$t$ -intersecting* if  $|F_1 \cap F_2| \geq t$  for all  $F_1, F_2 \in \mathcal{F}$ .

The *Kneser graph*,  $\text{Kn}(n, k)$ , is the graph whose vertices are the  $k$ -element subsets of  $[n]$ , i.e.  $V(\text{Kn}(n, k)) = \binom{[n]}{k}$  and two vertices are connected iff the two corresponding sets are disjoint. A *coclique* in a graph is a set of vertices, such that no two vertices in the set are adjacent. An intersecting family is a coclique in the corresponding Kneser graph. The maximum size of a coclique in a graph  $G$  is denoted by  $\alpha(G)$ .

The following theorem is one of the famous results in extremal combinatorics:

**Theorem 1** (Erdős, Ko, Rado [3]). *If  $k \leq n/2$ , then*

$$\alpha(\text{Kn}(n, k)) = \binom{n-1}{k-1}.$$

Obviously, the family consisting of the  $k$ -subsets that contain 1 has size  $\binom{n-1}{k-1}$ , so only the  $\leq$  part is interesting.

Let  $\mathcal{F} \subseteq \binom{X}{k}$  be a family of  $k$ -element sets; for  $l \leq k$ , the  $l$ -shadow of  $\mathcal{F}$  is defined as  $\Delta_l \mathcal{F} = \{G : |G| = l, \text{ and there exists } F \in \mathcal{F} \text{ such that } G \subset F\}$ . It is clear that  $\mathcal{F} = \binom{[2k-t]}{k}$  is  $t$ -intersecting and  $\Delta_l \mathcal{F} = \binom{[2k-t]}{l}$ . The next theorem shows that this is the extremal case in some sense.

**Theorem 2** (Katona [5]). *Assume that  $\mathcal{F}$  is a  $k$ -uniform,  $t$ -intersecting family. Then for  $l \geq k - t$ ,*

$$\frac{|\Delta_l \mathcal{F}|}{|\mathcal{F}|} \geq \frac{\binom{2k-t}{l}}{\binom{2k-t}{k}}.$$

## 2. A GENERALIZATION OF THE EKR THEOREM

In this section we deduce a slight generalization of the EKR theorem from Theorem 2.

For a set  $A \subseteq V(\text{Kn}(n, k))$ , the neighborhood of  $A$  is denoted by  $N(A)$ . Similarly, for a given  $k$ -uniform family  $\mathcal{F}$ , let us introduce the notation  $\mathcal{N}(\mathcal{F}) = \{H \in \binom{[n]}{k} : \text{there exists } F \in \mathcal{F} \text{ such that } H \cap F = \emptyset\}$  as the “neighborhood” of  $\mathcal{F}$ .

**Theorem 3.** *If  $k \leq n/2$  and  $C$  is a coclique in the Kneser graph,  $\text{Kn}(n, k)$ , then*

$$\frac{|C|}{|C| + |N(C)|} \leq \frac{k}{n}.$$

Since  $C$  is a coclique,  $C$  and  $N(C)$  are disjoint, so  $|C| + |N(C)| \leq |V(\text{Kn}(n, k))| = \binom{n}{k}$  and the EKR theorem follows.

**Proof of Theorem 3.** To apply Theorem 2, let  $\mathcal{F}$  be the intersecting  $k$ -uniform family that corresponds to  $C$ . Let  $\mathcal{F}^c$  be the family of complements, i.e.  $\mathcal{F}^c = \{[n] \setminus F : F \in \mathcal{F}\} \subseteq \binom{[n]}{n-k}$ . For each pair  $F_1, F_2 \in \mathcal{F}$ , we have  $|F_1 \cup F_2| \leq 2k - 1$ , thus  $\mathcal{F}^c$  is  $t$ -intersecting for  $t = n - 2k + 1$ . By Theorem 2,

$$\frac{|\Delta_k \mathcal{F}^c|}{|\mathcal{F}^c|} \geq \frac{\binom{2(n-k)-(n-2k+1)}{k}}{\binom{2(n-k)-(n-2k+1)}{n-k}} = \frac{n-k}{k}.$$

$|\mathcal{F}^c| = |\mathcal{F}|$  and  $\Delta_k \mathcal{F}^c \subseteq \mathcal{N}(\mathcal{F})$ , because for every  $H \in \Delta_k \mathcal{F}^c$ ,  $H \subseteq [n] \setminus F$  for some  $F \in \mathcal{F}$  and  $H \cap F = \emptyset$ . Thus,

$$\frac{|N(C)|}{|C|} = \frac{|\mathcal{N}(\mathcal{F})|}{|\mathcal{F}|} \geq \frac{n-k}{k}$$

and we are done. ■

### 3. SIMILAR RESULTS

Let  $A \subseteq V(\text{Kn}(n, k))$ . For another slight generalization, we denote by  $I(A)$  the family of isolated points in  $A$ , that is,

$$I(A) = \{a \in A : (a, b) \notin E(\text{Kn}(n, k)) \text{ for all } b \in A\}.$$

In his paper, Borg [1] extended Daykin's proof [2] of the EKR theorem to obtain the following improvement:

**Theorem 4** (Borg). *If  $A \subseteq V(\text{Kn}(n, k))$  and  $k \leq n/2$ , then*

$$|I(A)| + \frac{k}{n} |A \setminus I(A)| \leq \binom{n-1}{k-1}.$$

It is easy to see that Theorems 3 and 4 are equivalent.

First, let  $A$  be an arbitrary subgraph of  $\text{Kn}(n, k)$ .  $C := I(A)$  is a coclique, so by Theorem 3,  $\frac{|I(A)|}{|I(A)| + |N(I(A))|} \leq \frac{k}{n}$ .

By definition,  $I(A)$ ,  $A \setminus I(A)$  and  $N(I(A))$  are disjoint, hence  $|I(A)| + |A \setminus I(A)| + |N(I(A))| \leq \binom{n}{k}$ .

These two inequalities now imply Theorem 4.

On the other hand, if  $C$  is a coclique, let  $A := V(\text{Kn}(n, k)) \setminus N(C)$ . By definition,  $C$  and  $N(C)$  are disjoint, and  $C \subseteq I(A)$ . Thus, by Theorem 4,

$$|C| + \frac{k}{n} |V(\text{Kn}(n, k)) \setminus N(C) \setminus C| \leq \binom{n-1}{k-1},$$

and Theorem 3 follows.

Remember that though the two theorems are equivalent, their proofs are quite different: while Theorem 3 is proved as a consequence of the theorem on shadows of intersecting families, Borg uses the Kruskal–Katona theorem [6, 7] to verify Theorem 4.

**Remark 5.** In [1], Borg also showed that Theorem 4 (and so Theorem 3) yields Hilton's theorem [4] for cross-intersecting sub-families of  $\binom{[n]}{k}$ .

Recently, J. Wang and H. Zhang [8, 9] investigated similar problems in general circumstances. A graph  $G = (V, E)$  is called *vertex-transitive* if its automorphism group,  $\text{Aut}(G)$ , acts transitively on  $V$ , i.e. for every  $u, v \in V$  there exists a  $\gamma \in \text{Aut}(G)$  such that  $\gamma(u) = v$ .

The following theorem is the analogue of Theorem 3 for arbitrary vertex-transitive graph.

**Theorem 6** (Zhang). *Let  $G = (V, E)$  be a vertex-transitive simple graph. If  $C \subseteq V$  is a coclique, then*

$$\frac{|C|}{|C| + |N(C)|} \leq \frac{\alpha(G)}{|V|}.$$

Note that the EKR theorem and Theorem 6 together imply Theorem 3.

#### REFERENCES

- [1] P. Borg, *A short proof of a cross-intersecting theorem of Hilton*, Discrete Math. **309** (2009) 4750–4753.  
doi:10.1016/j.disc.2008.05.051
- [2] D.E. Daykin, *Erdős–Ko–Rado from Kruskal–Katona*, J. Combin. Theory (A) **17** (1974) 254–255.  
doi:10.1016/0097-3165(74)90013-2
- [3] P. Erdős, C. Ko and R. Rado, *Intersection theorems for systems of finite sets*, Quart. J. Math., Oxford **12** (1961) 313–320.
- [4] A.J.W. Hilton, *An intersection theorem for a collection of families of subsets of a finite set*, J. London Math. Soc. (2) **15** (1977) 369–376.  
doi:10.1112/jlms/s2-15.3.369
- [5] G.O.H. Katona, *Intersection theorems for systems of finite sets*, Acta Math. Hungar. **15** (1964) 329–337.  
doi:10.1007/BF01897141
- [6] G.O.H. Katona, *A theorem of finite sets* in: Theory of Graphs, Proc. Colloq. Tihany, 1966, P. Erdős and G.O.H. Katona (Eds.) (Akadémiai Kiadó, 1968) 187–207.
- [7] J.B. Kruskal, *The number of simplices in a complex* in: Math. Optimization Techniques, R. Bellman (Ed.) (Univ. of Calif. Press, Berkeley, 1963) 251–278.
- [8] J. Wang and H.J. Zhang, *Cross-intersecting families and primitivity of symmetric systems*, J. Combin. Theory (A) **118** (2011) 455–462.  
doi:10.1016/j.jcta.2010.09.005
- [9] H.J. Zhang, *Primitivity and independent sets in direct products of vertex-transitive graphs*, J. Graph Theory **67** (2011) 218–225.  
doi:10.1002/jgt.20526

Received 28 April 2011

Revised 6 August 2011

Accepted 8 August 2011