

STRONG EQUALITY BETWEEN THE ROMAN DOMINATION AND INDEPENDENT ROMAN DOMINATION NUMBERS IN TREES

MUSTAPHA CHELLALI¹

LAMDA-RO, Department of Mathematics
University of Blida
B.P. 270, Blida, Algeria
e-mail: mchellali@hotmail.com

AND

NADER JAFARI RAD²

Department of Mathematics, Shahrood University of Technology
Shahrood, Iran
and
School of Mathematics
Institute for Research in Fundamental Sciences (IPM),
P.O. Box 19395-5746, Tehran, Iran
e-mail: n.jafarirad@gmail

Abstract

A Roman dominating function (RDF) on a graph $G = (V, E)$ is a function $f : V \rightarrow \{0, 1, 2\}$ satisfying the condition that every vertex u for which $f(u) = 0$ is adjacent to at least one vertex v for which $f(v) = 2$. The weight of an RDF is the value $f(V(G)) = \sum_{u \in V(G)} f(u)$. An RDF f in a graph G is independent if no two vertices assigned positive values are adjacent. The Roman domination number $\gamma_R(G)$ (respectively, the independent Roman domination number $i_R(G)$) is the minimum weight of an RDF (respectively, independent RDF) on G . We say that $\gamma_R(G)$ strongly equals $i_R(G)$, denoted by $\gamma_R(G) \equiv i_R(G)$, if every RDF on G of minimum weight is independent. In this paper we provide a constructive characterization of trees T with $\gamma_R(T) \equiv i_R(T)$.

Keywords: Roman domination, independent Roman domination, strong equality, trees.

2010 Mathematics Subject Classification: 05C69.

¹This research was supported by "Programmes Nationaux de Recherche: Code 8/u09/510".

²This research was in part supported by a grant from IPM (No. 90050042).

1. INTRODUCTION

We consider finite, undirected, and simple graphs G with vertex set $V = V(G)$ and edge set $E = E(G)$. The *open neighborhood* of a vertex $v \in V$ is $N(v) = N_G(v) = \{u \in V \mid uv \in E\}$ and the *closed neighborhood* of v is $N[v] = N_G[v] = N_G(v) \cup \{v\}$. If D is a subset of $V(G)$, then the subgraph induced by D in G is denoted by $G[D]$. The *degree* of v , denoted by $d_G(v)$, is the cardinality of its open neighborhood. A vertex of degree one is called a *leaf*, and its neighbor is called a *support vertex*. If v is a support vertex, then L_v will denote the set of the leaves attached at v . A support vertex v is called *strong* if $|L_v| > 1$. A tree T is a *double star* if it contains exactly two vertices that are not leaves. A double star with respectively p and q leaves attached at each support vertex is denoted by $S_{p,q}$. For a vertex v in a rooted tree T , we denote by $D(v)$ the set of all descendants of v . The *maximal subtree* at v is the subtree of T induced by $D(v) \cup \{v\}$, and is denoted by T_v .

For a graph G , let $f : V(G) \rightarrow \{0, 1, 2\}$ be a function, and let $(V_0; V_1; V_2)$ be the ordered partition of $V = V(G)$ induced by f , where $V_i = \{v \in V(G) : f(v) = i\}$ for $i = 0, 1, 2$. There is a 1 – 1 correspondence between the functions $f : V(G) \rightarrow \{0, 1, 2\}$ and the ordered partitions $(V_0; V_1; V_2)$ of $V(G)$. So we will write $f = (V_0; V_1; V_2)$.

A function $f : V(G) \rightarrow \{0, 1, 2\}$ is a *Roman dominating function* (RDF) on G if every vertex u of G for which $f(u) = 0$ is adjacent to at least one vertex v of G for which $f(v) = 2$. The weight of a Roman dominating function f on G is the value $f(V(G)) = \sum_{u \in V(G)} f(u)$. The *Roman domination number* of a graph G , denoted by $\gamma_R(G)$, is the minimum weight of a Roman dominating function on G . A function $f = (V_0; V_1; V_2)$ is called a $\gamma_R(G)$ -function or γ_R -function for G if it is a Roman dominating function on G and $f(V(G)) = \gamma_R(G)$. Roman domination has been introduced by Cockayne *et al.* [1] and has been further studied for example in [5, 6, 7].

A function $f = (V_0; V_1; V_2)$ is called an *independent Roman dominating function* (IRDF) on G if f is an RDF and no two vertices in $V_1 \cup V_2$ are adjacent. The *independent Roman domination number* $i_R(G)$ is the minimum weight of an independent Roman dominating function of G . A function $f = (V_0; V_1; V_2)$ is called an $i_R(G)$ -function or i_R -function for G if it is an IRDF on G and $f(V(G)) = i_R(G)$.

Observe that for every graph G , $\gamma_R(G) \leq i_R(G)$. Clearly if G is a graph with $\gamma_R(G) = i_R(G)$, then every $i_R(G)$ -function is also a $\gamma_R(G)$ -function. However not every $\gamma_R(G)$ -function is an $i_R(G)$ -function even when $\gamma_R(G) = i_R(G)$. For example the double star $S_{2,3}$ has two $\gamma_R(S_{2,3})$ -functions but only one $\gamma_R(S_{2,3})$ -function is an $i_R(S_{2,3})$ -function. We say that $\gamma_R(G)$ and $i_R(G)$ are *strongly equal*, denoted by $\gamma_R(G) \equiv i_R(G)$, if every $\gamma_R(G)$ -function is an $i_R(G)$ -function. Note that Haynes and Slater in [4] were the first to introduce strong equality between

two parameters. Also in [2] and [3], Haynes, Henning and Slater gave constructive characterizations of trees with strong equality between some domination parameters.

In this paper we present a constructive characterization of trees T with $\gamma_R(T) \equiv i_R(T)$. If f is an RDF on a graph G and H is a subgraph of G , then we denote the restriction of f on H by $f|_{V(H)}$.

2. TREES T WITH $\gamma_R(T) \equiv i_R(T)$

We begin by the following results that will be useful for the next.

Proposition 1 (Cockayne *et al.* [1]). *Let $f = (V_0; V_1; V_2)$ be any $\gamma_R(G)$ -function. Then*

- (i) *The subgraph induced by the vertices of V_1 has maximum degree one.*
- (ii) *No edge of G joins V_1 to V_2 .*




Figure 1. A tree in \mathcal{T} .

Proposition 2 (Jafari Rad and Volkmann [7]). *If v is a vertex in a graph G such that $\gamma_R(G - v) > \gamma_R(G)$, then $f(v) = 2$ for every $\gamma_R(G)$ -function f .*

Let \mathcal{T} be the family of trees that can be obtained from k ($k \geq 1$) disjoint stars of centers x_1, x_2, \dots, x_k , where each star has order at least three, attached by edges from their center vertices either to a single vertex or to a same leaf of a path P_2 . If T is a tree of \mathcal{T} , then let us call the vertex adjacent to the centers of stars, the special vertex of T . Note that if T belongs to \mathcal{T} , then $\gamma_R(T) \equiv i_R(T)$.

Now we present a constructive characterization of trees T with $\gamma_R(T) \equiv i_R(T)$. For this purpose, we define a family of trees as follows: Let \mathcal{F} be the collection of trees T that can be obtained from a sequence T_1, T_2, \dots, T_k ($k \geq 1$) of trees, where T_1 is a star $K_{1,t}$ with $t \geq 2$, $T = T_k$, and, if $k \geq 2$, then T_{i+1} can

be obtained recursively from T_i by one of the following operations. Also for any tree T_i of \mathcal{F} we let f_i be a $\gamma_R(T_i)$ -function.




Figure 2. The \mathcal{O}_i Operations.

- Operation \mathcal{O}_1 : Assume y is a leaf of T_i with $f_i(y) = 0$ and whose support vertex z is either strong or satisfies $\gamma_R(T_i - z) > \gamma_R(T_i)$. Then T_{i+1} is obtained from T_i by adding a new vertex x and adding the edge xy .
- Operation \mathcal{O}_2 : Assume y is a vertex of T_i . Then T_{i+1} is obtained from T_i by adding a tree $T \in \mathcal{T}$ of special vertex x and adding the edge xy with the condition that if x is a support vertex, then y satisfies $\gamma_R(T_i - y) \geq \gamma_R(T_i)$.
- Operation \mathcal{O}_3 : Assume y is a vertex of T_i assigned 0 or 1 for every $\gamma_R(T_i)$ -function. Then T_{i+1} is obtained from T_i by adding a path $P_3 = u-v-w$ and adding the edge wy .

Lemma 3. *If T_i is a tree with $\gamma_R(T_i) \equiv i_R(T_i)$ and T_{i+1} is a tree obtained from T_i by Operation \mathcal{O}_1 , then $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$.*

Proof. Since $\gamma_R(T_i) \equiv i_R(T_i)$, it is clear that every $i_R(T_i)$ -function with y assigned 0 can be extended to an IRDF for T_{i+1} by assigning 1 to x . Hence $\gamma_R(T_{i+1}) \leq i_R(T_{i+1}) \leq i_R(T_i) + 1 = \gamma_R(T_i) + 1$. Now let f be a $\gamma_R(T_{i+1})$ -function. If $f(y) = 1$, then $f(x) = 1$ and $f|_{V(T_i)}$ is an RDF for T_i . If $f(y) = 0$, then $f(x) = 1$ (else $f(x) = 2$ and we can make a change to obtain $f(x) = 1$ and

$f(y) = 1$) and $f|_{V(T_i)}$ is an RDF for T_i . In both cases, $\gamma_R(T_i) \leq \gamma_R(T_{i+1}) - 1$ and equality throughout the above chain is obtained. Now assume that $f(y) = 2$. Then $f(x) = 0$ and by Proposition 1 we may assume that $f(z) = 0$. If z has a leaf neighbor, say z' , then $f(z') = 1$ and we can change $f(z') = 1$ to $f(z') = 0$, $f(z) = 0$ to $f(z) = 2$, $f(y) = 2$ to $f(y) = 0$ and $f(x) = 0$ to $f(x) = 1$. Clearly we are in the previous situation. Hence we may assume that z is not a support vertex. Then consider the function f' on $V(T_i - z)$ defined by $f'(a) = f(a)$ if $a \in V(T_i) - \{y, z\}$, and $f'(y) = 1$. Then f' is an RDF for $T_i - z$ and so $\gamma_R(T_i - z) \leq f'(V(T_i - z)) - 1 = \gamma_R(T_{i+1}) - 1$. Now since z is a support vertex in T_i but not strong, it satisfies $\gamma_R(T_i - z) > \gamma_R(T_i)$. Then we obtain $\gamma_R(T_i) < \gamma_R(T_i - z) \leq \gamma_R(T_{i+1}) - 1$, implying that $\gamma_R(T_{i+1}) > \gamma_R(T_i) + 1$, which is impossible. Thus for the next we may assume that for any $\gamma_R(T_{i+1})$ -function y is not assigned 2.

Next we shall show that $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$. Assume to the contrary that $h = (V_0; V_1; V_2)$ is a $\gamma_R(T_{i+1})$ -function such that $V_1 \cup V_2$ is not independent. Thus there are two adjacent vertices $u, v \in V_1 \cup V_2$. Recall that $h(y) \in \{0, 1\}$. If $h(y) = 0$, then $h(x) = 1$, and so $h|_{V(T_i)} = (V_0; V_1 - \{x\}; V_2)$ is a $\gamma_R(T_i)$ -function. But $h|_{V(T_i)}$ is not independent since u, v belong to $(V_1 - \{x\}) \cup V_2$, contradicting $\gamma_R(T_i) \equiv i_R(T_i)$. If $h(y) = 1$, then $h(x) = 1$. By Proposition 1, $h(z) \neq 2$, and so $h|_{V(T_i - z)}$ is an RDF for $V(T_i - z)$. Observe that z cannot be a support vertex in T_{i+1} . Now by using the fact that z verifies $\gamma_R(T_i - z) > \gamma_R(T_i)$, we obtain $\gamma_R(T_i) < \gamma_R(T_i - z) \leq h(V(T_i - z)) \leq \gamma_R(T_{i+1}) - 1$, which is impossible. Therefore $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$. ■

Lemma 4. *If T_i is a tree with $\gamma_R(T_i) \equiv i_R(T_i)$ and T_{i+1} is a tree obtained from T_i by Operation \mathcal{O}_2 , then $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$.*

Proof. Let $T \in \mathcal{T}$ be the added tree of special vertex x . Recall that T is obtained from k ($k \geq 1$) disjoint stars of centers x_1, x_2, \dots, x_k , each of order at least three, attached by edges xx_j at x , where x may be a single vertex or belongs to a path $P_2 = x-x'$.

Clearly every $i_R(T_i)$ -function can be extended to an IRDF for T_{i+1} by assigning 2 to every x_j , 1 to x' (if x' exists), and 0 to x and every leaf of T different to x' . Hence $\gamma_R(T_{i+1}) \leq i_R(T_{i+1}) \leq i_R(T_i) + 2k + t = \gamma_R(T_i) + 2k + t$, where $t = 1$ if x' exists and $t = 0$ otherwise. Now let f be a $\gamma_R(T_{i+1})$ -function. Without loss of generality we can assume that $f(x_j) = 2$ for every j . Hence every leaf adjacent to some x_i is assigned 0. If $f(x) = 0$ and $f(x') = 1$ (if x' exists), then $f|_{V(T_i)}$ is an RDF for T_i implying that $i_R(T_i) \leq \gamma_R(T_{i+1}) - 2k - t$. Equality throughout the above inequality chain is obtained. Now if either $f(x) = 2$ and $f(x') = 0$ or $f(x) = 0$ and $f(x') = 2$, then we can change by assigning 1 to x' and y , and 0 to x . Clearly we are in the previous situation.

Assume now that $\gamma_R(T_{i+1})$ and $i_R(T_{i+1})$ are not strongly equal and let $h =$

$(V_0; V_1; V_2)$ be a $\gamma_R(T_{i+1})$ -function such that $V_1 \cup V_2$ is not independent. Let u and v be any two adjacent vertices in $V_1 \cup V_2$. If $h(x) = 0$, then clearly u, v belong to $V(T_i)$ and $h|_{V(T_i)}$ is a $\gamma_R(T_i)$ -function that is not independent, a contradiction with $\gamma_R(T_i) \equiv i_R(T_i)$. If $h(x) = 1$, then $h(x') = 1$ (if x' exists) and so $h|_{V(T_i)}$ is an RDF for T_i with weight $\gamma_R(T_{i+1}) - 2k - t - 1 < \gamma_R(T_i)$, which is impossible. Finally assume that $h(x) = 2$. We may assume that x' exists for otherwise we can decrease the weight of h by assigning 0 to x and 1 to y . Hence $h(x') = 0$ and $h(y) = 0$. Then $h|_{V(T_i-y)}$ is an RDF for $T_i - y$ and so $h(V(T_i - y)) = \gamma_R(T_{i+1}) - 2k - 2$. Now since x is a support vertex in T , y must satisfy $\gamma_R(T_i - y) \geq \gamma_R(T_i)$, implying that $\gamma_R(T_{i+1}) - 2k - 2 = h(V(T_i - y)) \geq \gamma_R(T_i - y) \geq \gamma_R(T_i)$. Therefore we have $\gamma_R(T_{i+1}) \geq \gamma_R(T_i) + 2k + 2$, a contradiction. Consequently $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$. ■

Lemma 5. *If T_i is a tree with $\gamma_R(T_i) \equiv i_R(T_i)$ and T_{i+1} is a tree obtained from T_i by Operation \mathcal{O}_3 , then $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$.*

Proof. Clearly every $i_R(T_i)$ -function can be extended to an IRDF for T_{i+1} by assigning 0 to u, w and 2 to v . Hence $\gamma_R(T_{i+1}) \leq i_R(T_{i+1}) \leq i_R(T_i) + 2 = \gamma_R(T_i) + 2$. Now let f be a $\gamma_R(T_{i+1})$ -function. If $f(v) = 2$, then $f(w) = f(u) = 0$ and $f|_{V(T_i)}$ is an RDF for T_i . Hence $\gamma_R(T_i) \leq \gamma_R(T_{i+1}) - 2$. If $f(v) = 1$, then $f(u) = 1$ and w must be assigned 0. It follows that $f|_{V(T_i)}$ is an RDF for T_i and so $\gamma_R(T_i) \leq \gamma_R(T_{i+1}) - 2$. Now assume that $f(v) = 0$. Then $f(u) = 2$ and $f(w) \notin \{1, 2\}$. It follows that $f|_{V(T_i)}$ is an RDF for T_i and so $\gamma_R(T_i) \leq \gamma_R(T_{i+1}) - 2$. For all cases, we obtain $\gamma_R(T_{i+1}) = \gamma_R(T_i) + 2$, implying that $i_R(T_{i+1}) = i_R(T_i) + 2$.

Assume now that $\gamma_R(T_{i+1})$ is not strongly equal to $i_R(T_{i+1})$ and let h be a $\gamma_R(T_{i+1})$ -function that is not independent. Thus there are two adjacent vertices a and b assigned positive values. If $h(v) = 2$, then $h(w) = h(u) = 0$ and $h|_{V(T_i)}$ is a $\gamma_R(T_i)$ -function, where $a, b \in V(T_i)$, contradicting $\gamma_R(T_i) \equiv i_R(T_i)$. If $h(v) = 1$, then $h(u) = 1$ and $h(w) = 0$. It follows that $h(y) = 2$ and $h|_{V(T_i)}$ is a $\gamma_R(T_i)$ -function for which y is assigned 2, a contradiction with the construction. Thus we assume that $h(v) = 0$. Hence $h(u) = 2$. If $h(w) = 1$, then $h|_{V(T_i)}$ is an RDF for T_i of weight $\gamma_R(T_i) - 1$, which is impossible. If $h(w) = 2$, then we change $h(w) = 2$ to $h(w) = 1$ and $h(y) = 0$ to $h(y) = 1$ and we obtain the previous situation. Thus $h(w) = 0$ implying that $h(y) = 2$. But then $h|_{V(T_i)}$ is a $\gamma_R(T_i)$ -function for which y is assigned 2, a contradiction with the construction. Therefore $\gamma_R(T_{i+1}) \equiv i_R(T_{i+1})$. ■

We now are ready to establish our main result.

Theorem 6. *Let T be a tree. Then $\gamma_R(T) \equiv i_R(T)$ if and only if $T = K_1$ or $T \in \mathcal{F}$.*

Proof. Obviously, if $T = K_1$, then $\gamma_R(T) \equiv i_R(T)$. Now suppose that $T \in \mathcal{F}$. Then there is a sequence of trees T_1, T_2, \dots, T_k ($k \geq 1$) such that T_1 is a star $K_{1,t}$

with $t \geq 2$, $T = T_k$, and, if $k \geq 2$, then T_{i+1} can be obtained recursively from T_i by an operation \mathcal{O}_1 , \mathcal{O}_2 or \mathcal{O}_3 for $i = 1, \dots, (k-1)$. We use an induction on the number of operations performed to construct T . Clearly the property is true if $k = 1$. This establishes the basis case. Assume now that $k \geq 2$ and that the result holds for all trees $T \in \mathcal{F}$ that can be constructed from a sequence of length at most $k-1$, and let $T' = T_{k-1}$. By the induction hypothesis, $\gamma_R(T') \equiv i_R(T')$. By construction T is obtained from T' by using Operation \mathcal{O}_1 , \mathcal{O}_2 or \mathcal{O}_3 . Hence by Lemmas 3, 4 and 5 it follows that $\gamma_R(T) \equiv i_R(T)$.

Conversely, let T be a tree of order n with $\gamma_R(T) \equiv i_R(T)$. Clearly if $n = 1$, then $T = K_1$. Hence we assume that T has order $n \geq 2$. We use an induction on the order n . Since a path P_2 has a $\gamma_R(P_2)$ -function that is not independent, we assume that $n \geq 3$. If $n = 3$, then $T = P_3$ which belongs to \mathcal{F} , establishing the base case. Assume that every tree T' of order $2 \leq n' < n$ with $\gamma_R(T') \equiv i_R(T')$ is in \mathcal{F} . Let T be a tree of order n with $\gamma_R(T) \equiv i_R(T)$ and let f be a $\gamma_R(T)$ -function. Since stars of order at least three belong to \mathcal{F} , we may assume that T has diameter at least three. If $\text{diam}(T) = 3$, then T is a double star $S_{1,p}$ with $p \geq 1$ and $T \in \mathcal{F}$ because it is obtained from a star $K_{1,p+1}$ by using Operation \mathcal{O}_1 . Therefore assume that $\text{diam}(T) \geq 4$.

We now root T at a leaf r of a longest path. Let u be a vertex at distance $\text{diam}(T) - 1$ from r on a longest path starting at r such that $|L_u|$ is as small as possible. Let v, w be the parents of u and v on this path, respectively. Clearly $f(u) \neq 1$, else u and its leaves belong to V_1 , contradicting $\gamma_R(T) \equiv i_R(T)$. We consider the following cases.

Case 1. $f(u) = 2$. Then $f(v) = 0$ and $f(u') = 0$ for every $u' \in L_u$.

Subcase 1.1. v is a support vertex. Then $f(v') = 1$ for every $v' \in L_v$. If v is adjacent to two leaves v' and v'' , then we can change $f(v) = 0$ to $f(v) = 2$ and $f(v') = f(v'') = 1$ to $f(v') = f(v'') = 0$. Clearly we obtain a $\gamma_R(T)$ -function for which $V_1 \cup V_2$ is not independent. Hence v is adjacent to a unique leaf v' . So $|L_v| = 1$.

Suppose that $|L_u| = 1$ and let u' be the unique leaf neighbor of u . Consider the function h on $V(T)$ defined by $h(x) = f(x)$ if $x \in V(T) - \{u, u', v, v'\}$, $h(u') = 1, h(u) = 0, h(v) = 2$ and $h(v') = 0$. Then h is a $\gamma_R(T)$ -function and $h(w) = 0$. Furthermore $d_T(v) = 3$, for otherwise every child y of v different from u is assigned 2, a contradiction. Let T' be the tree obtained from T by removing u' . Note that v is a strong support vertex in T' . Clearly $h|_{V(T')}$ is both an RDF and an IRDF for T' implying that $\gamma_R(T') \leq \gamma_R(T) - 1$ and $i_R(T') \leq i_R(T) - 1$. Since every $\gamma_R(T')$ -function can be extended to an RDF for T by assigning 1 to u' we obtain $\gamma_R(T) = \gamma_R(T') + 1$. Also $i_R(T') \leq i_R(T) - 1 = \gamma_R(T) - 1 = \gamma_R(T')$ and so $i_R(T') = \gamma_R(T')$. It follows that $i_R(T) = i_R(T') + 1$ and so $i_R(T') = \gamma_R(T')$. On the other hand, if $\gamma_R(T')$ and $i_R(T')$ are not strongly equal, then every $\gamma_R(T')$ -function for which $V_1 \cup V_2$ is not independent can be extended to a $\gamma_R(T)$ -function

by assigning 1 to u' , a contradiction with $\gamma_R(T) \equiv i_R(T)$. Therefore $\gamma_R(T') \equiv i_R(T')$ and by induction on T' , we have $T' \in \mathcal{F}$. We conclude that $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_1 .

Assume now that $|L_u| \geq 2$. By our choice of u , every child of v which is a support vertex is adjacent to at least two leaves. Hence T_v is a tree of \mathcal{T} . Let $u = u_1, u_2, \dots, u_k$ with $k \geq 1$, denote the support vertices adjacent to v in T_v , and let $T' = T - T_v$. Since $\text{diam}(T) \geq 4$, T' is nontrivial. We observe that $f|_{V(T')}$ is both an RDF and IRDF for T' implying that $\gamma_R(T') \leq \gamma_R(T) - 2k - 1$ and $i_R(T') \leq i_R(T) - 2k - 1$. Equality is obtained by the fact that every $\gamma_R(T')$ -function (resp. $i_R(T')$ -function) can be extended to an RDF (resp. an IRDF) for T by assigning 2 to every u_i , 0 to v and every leaf in T_v except v' , and 1 to v' . On the other hand, observe that if w satisfies $\gamma_R(T' - w) \leq \gamma_R(T') - 1$, then every $\gamma_R(T' - w)$ -function can be extended to a $\gamma_R(T)$ -function that is not independent by assigning 2 to v and every u_i and 0 to the remaining vertices, a contradiction with $\gamma_R(T) \equiv i_R(T)$. Thus w satisfies $\gamma_R(T' - w) \geq \gamma_R(T')$. If $\gamma_R(T')$ and $i_R(T')$ are not strongly equal, then every $\gamma_R(T')$ -function which is not independent can be extended to a $\gamma_R(T)$ -function, contradicting $\gamma_R(T) \equiv i_R(T)$. It follows that $\gamma_R(T') \equiv i_R(T')$ and by induction on T' we have $T' \in \mathcal{F}$. Therefore $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_2 .

Subcase 1.2. v is not a support vertex. We first assume that $d_T(v) \geq 3$. Then all children of v are support vertices and each one is assigned 2. If some child b of v is adjacent to only one leaf b' , then we can change $f(b) = 2$ to $f(b) = 1$ and $f(b') = 0$ to $f(b') = 1$. We then obtain a $\gamma_R(T)$ -function that is not independent, a contradiction. Thus every child of v is adjacent to at least two leaves. Let $T' = T - T_v$. Observe that T_v belongs to \mathcal{T} . Then $\gamma_R(T) \leq \gamma_R(T') + 2(d_T(v) - 1)$ since every $\gamma_R(T')$ -function can be extended to an RDF for T by assigning 2 to every support vertex in T_v . Likewise, $i_R(T) \leq i_R(T') + 2(d_T(v) - 1)$. Both equalities are obtained from the fact that $f|_{V(T')}$ is an RDF and IRDF for T' . It follows that $\gamma_R(T') = i_R(T')$. Now if T' admits a $\gamma_R(T')$ -function that is not independent, then such a function can be extended to a $\gamma_R(T)$ -function that is not independent, a contradiction with $\gamma_R(T) \equiv i_R(T)$. Thus every $\gamma_R(T')$ -function is independent, that is $\gamma_R(T') \equiv i_R(T')$. By induction on T' we have $T' \in \mathcal{F}$ and so $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_2 .

Now assume that $d_T(v) = 2$. If $|L_u| \geq 2$, then we consider $T' = T - T_v$. Observe that $T_v \in \mathcal{T}$. It is easy to see that $\gamma_R(T) = \gamma_R(T') + 2$ and $i_R(T) = i_R(T') + 2$, and so $\gamma_R(T') = i_R(T')$. Since every $\gamma_R(T')$ -function can be extended to a $\gamma_R(T)$ -function, it follows that $\gamma_R(T') \equiv i_R(T')$. By induction on T' we have $T' \in \mathcal{F}$ and so $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_2 . Now assume that $|L_u| = 1$, and let u' be the unique leaf adjacent to u . If $f(u) = 2$, then we change $f(u) = 2$ to $f(u) = 1$ and $f(u') = 0$ to $f(u') = 1$. We obtain a $\gamma_R(T)$ -function that is not independent, a contradiction. Thus $f(u) \in \{0, 1\}$ for

every $\gamma_R(T)$ -function f . Let $T' = T - T_v$. Then $\gamma_R(T') \leq \gamma_R(T) - 2$ and $i_R(T') \leq i_R(T) - 2$. Both equalities hold since every $\gamma_R(T')$ -function (respectively, $\gamma_R(T')$ -function) can be extended to an RDF (respectively, IRDF) for T by assigning 0 to u', v and 2 to u . Hence $i_R(T') = \gamma_R(T')$. Note that since $f(w) \neq 2$, w is assigned 0 or 1 for every $\gamma_R(T')$ -function. Now it is clear that $\gamma_R(T') \equiv i_R(T')$ and by induction on T' we have $T' \in \mathcal{F}$. It follows that $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_3 .

Case 2. $f(u) = 0$. Then $f(u') > 0$ for every $u' \in L_u$. It follows that $|L_u| \leq 2$, for otherwise we can decrease the weight of f by changing the assignment of u and its leaves. Now if $L_u = \{u', u''\}$, then $f(u') = f(u'') = 1$ and $f(v) = 2$. In this case we change $f(u) = 0$ to $f(u) = 2$ and $f(u') = f(u'') = 1$ to $f(u') = f(u'') = 0$. Clearly we obtain a $\gamma_R(T')$ -function that is not independent, a contradiction. Hence $|L_u| = 1$. Let u' be the leaf adjacent to u . If $f(u') = 2$, then we must have $f(v) = 0$ and so we can change $f(u') = 2$ to $f(u') = 0$ and $f(u) = 0$ to $f(u) = 2$. Hence we are in Case 1. Thus we assume that $f(u') = 1$ and so $f(v) = 2$. We consider the following subcases.

Subcase 2.1. v is a support vertex. Then $f(v') = 0$ for every $v' \in L_v$. Let T' be the tree obtained from T by removing u' . As seen in Subcase 1.1 we obtain $\gamma_R(T') \equiv i_R(T')$ and by induction on T' , $T' \in \mathcal{F}$. Since T is obtained from T' by using Operation \mathcal{O}_1 , we have $T \in \mathcal{F}$.

Subcase 2.2. v is not a support vertex but has degree at least three. Thus every child of v is a support vertex with degree two. Also every support vertex in T_v is assigned 0 and every leaf is assigned 1. Now let T' be the tree obtained from T by removing u' . It is easy to see that $\gamma_R(T) = \gamma_R(T') + 1$ and $i_R(T) = i_R(T') + 1$. Hence $\gamma_R(T') = i_R(T')$. On the other hand suppose that $\gamma_R(T' - v) \leq \gamma_R(T')$ and let f' be any $\gamma_R(T' - v)$ -function. Then u is an isolated vertex in $T' - v$ and is assigned 1. Also we may assume, without loss of generality, that every child of v different from u is assigned 2 in $T' - v$. Hence f' can be extended to a $\gamma_R(T)$ -function for T by assigning 1 to u' . But then the resulting $\gamma_R(T)$ -function is not independent, a contradiction. It follows that v satisfies $\gamma_R(T' - v) > \gamma_R(T')$ and so by Proposition 2, v is assigned 2 for every $\gamma_R(T')$ -function. Using this fact and the fact that every $\gamma_R(T')$ -function can be extended to a $\gamma_R(T)$ -function by assigning 1 to u' , we obtain $\gamma_R(T') \equiv i_R(T')$. By induction on T' we have $T' \in \mathcal{F}$ and so $T \in \mathcal{F}$ since it is obtained from T' by using Operation \mathcal{O}_1 .

Subcase 2.3. $d_T(v) = 2$. Recall that since $f(v) = 2$, we have $f(w) = 0$. Then we can make a change to obtain $f(u') = 0, f(u) = 2, f(v) = 0$ and $f(w) = 1$. Since $\gamma_R(T) \equiv i_R(T)$, no vertex of $N(w) - \{v\}$ is assigned a positive value. Now let $T' = T - T_v$. As seen in Subcase 1.2 (when $d_T(v) = 2$) w is not assigned 2 for every $\gamma_R(T')$ -function, $\gamma_R(T) = \gamma_R(T') + 2$, $i_R(T) = i_R(T') + 2$ and $\gamma_R(T') \equiv i_R(T')$.

By induction on T' we have $T' \in \mathcal{F}$ and so $T \in \mathcal{F}$ because it is obtained from T' by using Operation \mathcal{O}_3 . ■

We close with the following problem.

Problem. *Characterize other classes of graphs (or regular graphs) with strong equality between the Roman domination and the independent Roman domination numbers.*

REFERENCES

- [1] E.J. Cockayne, P.A. Dreyer Jr., S.M. Hedetniemi and S.T. Hedetniemi, *On Roman domination in graphs*, Discrete Math. **278** (2004) 11–22.
doi:10.1016/j.disc.2003.06.004
- [2] T.W. Haynes, M.A. Henning and P.J. Slater, *Strong equality of domination parameters in trees*, Discrete Math. **260** (2003) 77–87.
doi:10.1016/S0012-365X(02)00451-X
- [3] T.W. Haynes, M.A. Henning and P.J. Slater, *Strong equality of upper domination and independence in trees*, Util. Math. **59** (2001) 111–124.
- [4] T.W. Haynes and P.J. Slater, *Paired-domination in graphs*, Networks **32** (1998) 199–206.
doi:10.1002/(SICI)1097-0037(199810)32:3<199::AID-NET4>3.0.CO;2-F
- [5] M.A. Henning, *A characterization of Roman trees*, Discuss. Math. Graph Theory **22** (2002) 325–334.
doi:10.7151/dmgt.1178
- [6] M.A. Henning, *Defending the Roman Empire from multiple attacks*, Discrete Math. **271** (2003) 101–115.
doi:10.1016/S0012-365X(03)00040-2
- [7] N. Jafari Rad and L. Volkmann, *Changing and unchanging the Roman domination number of a graph*, Util. Math. **89** (2012) 79–95.

Received 8 December 2010

Revised 24 November 2011

Accepted 5 April 2012