

## FRACTIONAL $\mathcal{Q}$ -EDGE-COLORING OF GRAPHS

JÚLIUS CZAP

*Department of Applied Mathematics and Business Informatics,  
Faculty of Economics, Technical University of Košice,  
Němcovej 32, SK-040 01 Košice, Slovakia*

**e-mail:** julius.czap@tuke.sk

AND

PETER MIHÓK<sup>†</sup>

*Department of Applied Mathematics and Business Informatics,  
Faculty of Economics, Technical University of Košice,  
Němcovej 32, SK-040 01 Košice, Slovakia  
and*

*Mathematical Institute of the Slovak Academy of Sciences,  
Grešákova 6, SK-040 01 Košice, Slovakia*

### Abstract

An additive hereditary property of graphs is a class of simple graphs which is closed under unions, subgraphs and isomorphism. Let  $\mathcal{Q}$  be an additive hereditary property of graphs. A  $\mathcal{Q}$ -edge-coloring of a simple graph is an edge coloring in which the edges colored with the same color induce a subgraph of property  $\mathcal{Q}$ . In this paper we present some results on fractional  $\mathcal{Q}$ -edge-colorings. We determine the fractional  $\mathcal{Q}$ -edge chromatic number for matroidal properties of graphs.

**Keywords:** fractional coloring, graph property.

**2010 Mathematics Subject Classification:** 05C15, 05C70, 05C72.

### 1. INTRODUCTION

Our terminology and notation will be standard. The reader is referred to [1, 11] for undefined terms. All graphs considered in this paper are simple, i.e. they have no loops or multiple edges.

---

<sup>†</sup> Peter Mihók passed away on March 27, 2012.

We denote the class of all finite simple graphs by  $\mathcal{I}$ . A *graph property*  $\mathcal{Q}$  is a non-empty isomorphism-closed subclass of  $\mathcal{I}$ . We also say that a graph  $G$  has property  $\mathcal{Q}$  whenever  $G \in \mathcal{Q}$ . The fact that  $H$  is a subgraph of  $G$  is denoted by  $H \subseteq G$  and the disjoint union of two graphs  $G$  and  $H$  is denoted by  $G \cup H$ . A property  $\mathcal{Q}$  is called *additive* if  $G \cup H \in \mathcal{Q}$  whenever  $G \in \mathcal{Q}$  and  $H \in \mathcal{Q}$ . A property  $\mathcal{Q}$  is called *hereditary* if  $G \in \mathcal{Q}$  and  $H \subseteq G$  implies  $H \in \mathcal{Q}$ . The set of all additive hereditary properties will be denoted by  $\mathbb{L}$ .

We list several well-known additive hereditary properties

$$\begin{aligned}\mathcal{D}_k &= \{G \in \mathcal{I} : \text{each subgraph of } G \text{ contains a vertex of degree at most } k\}, \\ \mathcal{I}_k &= \{G \in \mathcal{I} : G \text{ does not contain } K_{k+2}\}, \\ \mathcal{J}_k &= \{G \in \mathcal{I} : \chi'(G) \leq k\}, \\ \mathcal{O}_k &= \{G \in \mathcal{I} : \text{each component of } G \text{ has at most } k+1 \text{ vertices}\}, \\ \mathcal{S}_k &= \{G \in \mathcal{I} : \Delta(G) \leq k\}, \\ \mathcal{B} &= \{G \in \mathcal{I} : G \text{ is a bipartite graph}\},\end{aligned}$$

where  $K_{k+2}$  denotes the complete graph on  $k+2$  vertices,  $\chi'(G)$  is the *edge chromatic number* (chromatic index) and  $\Delta(G)$  is the *maximum degree* of the graph  $G$ .

Generalized colorings of edges or/and vertices of graphs under restrictions given by graph properties have recently attracted much attention, see e.g. [2, 3, 4, 6, 7, 8, 10] and references therein.

By using the class of additive hereditary properties, there is the following generalization of edge coloring. Let  $\mathcal{Q} \in \mathbb{L}$  and let  $t$  be a positive integer. A *t-fold Q-edge-coloring* of a graph is an assignment of  $t$  distinct colors to each edge such that each color class induces a subgraph of property  $\mathcal{Q}$ . The smallest number  $k$  such that  $G$  admits a  $t$ -fold  $\mathcal{Q}$ -edge-coloring with  $k$  colors is the  $(t, \mathcal{Q})$ -*chromatic index* of  $G$ , denoted by  $\chi'_{t,\mathcal{Q}}(G)$ . Clearly, a 1-fold  $\mathcal{O}_1$ -edge-coloring is a usual proper edge coloring and hence  $\chi'_{1,\mathcal{O}_1}(G) = \chi'(G)$ .

Another generalization of edge coloring is fractional edge coloring. The *fractional chromatic index* of a graph  $G$  is defined in the following way:  $\chi'_f(G) = \lim_{t \rightarrow \infty} \frac{\chi'_{t,\mathcal{O}_1}(G)}{t}$ . If we replace the property  $\mathcal{O}_1$  by any other additive hereditary graph property  $\mathcal{Q}$  in the definition of the fractional chromatic index, then we obtain the *fractional Q-chromatic index* of a graph  $G$  and we denote it  $\chi'_{f,\mathcal{Q}}(G)$ .

A *hypergraph*  $\mathcal{H}$  is a pair  $(S, X)$ , where  $S$  is a finite set and  $X$  is a family of subsets of  $S$ . The elements of  $X$  are called *hyperedges*. A *t-fold covering* of a hypergraph  $\mathcal{H}$  is a collection (multiset) of hyperedges which includes every element of  $S$  at least  $t$  times. The smallest cardinality of such a multiset is called the *t-fold covering number* of  $\mathcal{H}$  and is denoted  $k_t(\mathcal{H})$ . The *fractional covering number* of  $\mathcal{H}$  is defined as  $k_f(\mathcal{H}) = \lim_{t \rightarrow \infty} \frac{k_t(\mathcal{H})}{t}$ .

For given simple graph  $G = (V, E)$  and additive hereditary property  $\mathcal{Q}$ , let  $\mathcal{H}_G = (E_G, \mathcal{Q}_G)$  denote the hypergraph whose vertex set is the edge set of  $G$  and the hyperedges are those subsets of  $E(G) = E_G$  which induce a graph of property  $\mathcal{Q}$ . Since  $\mathcal{Q}$  is hereditary, we can formulate the  $(t, \mathcal{Q})$ -chromatic index of the graph  $G$  as the  $t$ -fold covering number of the hypergraph  $\mathcal{H}_G$ . There is a natural one-to-one correspondence between the color classes of  $G$  and the hyperedges of  $\mathcal{H}_G$ . Therefore the following assertion holds.

**Claim 1.** *The fractional  $\mathcal{Q}$ -chromatic index of the graph  $G$  is equal to the fractional covering number of the hypergraph  $\mathcal{H}_G = (E_G, \mathcal{Q}_G)$ .*

A *matroid*  $\mathcal{M} = (S, I)$  is a hypergraph which satisfies the following three conditions:

1.  $\emptyset \in I$ ,
2. if  $X \in I$  and  $Y \subseteq X$ , then  $Y \in I$ ,
3. if  $X, Y \in I$  and  $|X| > |Y|$ , then there is an  $x \in X \setminus Y$  such that  $Y \cup \{x\} \in I$ .

In [12] the fractional covering number of matroids is determined. Let  $X$  be a subset of the ground set  $S$  of a matroid  $\mathcal{M}$ . The rank of  $X$ , denoted  $\rho(X)$ , is defined as the maximum cardinality of an independent subset of  $X$  (a subset of  $X$  which belongs to  $I$ ).

**Theorem 2** [12]. *If  $\mathcal{M} = (S, I)$  is a matroid, then*

$$k_f(\mathcal{M}) = \max_{X \subseteq S; X \neq \emptyset} \frac{|X|}{\rho(X)}.$$

In this paper, by combining Claim 1 and Theorem 2, we give a general formula for the fractional  $\mathcal{Q}$ -chromatic index. Afterwards, by this formula and with other results from the literature, we determine the exact values of  $\chi'_{f, \mathcal{Q}}(G)$  for so-called  $\mathcal{Q}$ -matroidal graphs.

## 2. RESULTS

Let  $G = (V, E)$  be a graph and let  $\mathcal{Q}$  be an additive hereditary property. If the hypergraph  $(E_G, \mathcal{Q}_G)$  is a matroid, then  $G$  is called  *$\mathcal{Q}$ -matroidal*. Let  $\mathcal{Q}^{\mathcal{M}}$  denote the set of all  $\mathcal{Q}$ -matroidal graphs. A property  $\mathcal{Q}$  is called *matroidal* if every graph  $G$  is  $\mathcal{Q}$ -matroidal. Schmidt [13] proved the existence of uncountably many matroidal properties.

A subset of the edge set of a graph is called  *$\mathcal{Q}$ -independent* if it induces a graph of property  $\mathcal{Q}$ . For a graph  $H$  let  $\mathcal{Q}(H)$  denote the maximum cardinality of a  $\mathcal{Q}$ -independent subset of  $E(H)$ .

**Lemma 3.** Let  $a_i, b_i > 0$  for  $i = 1, \dots, n$ . Then  $\frac{a_1 + \dots + a_n}{b_1 + \dots + b_n} \leq \max_i \left\{ \frac{a_i}{b_i} \right\}$ .

**Proof.** By induction on  $n$ . ■

**Theorem 4.** Let  $\mathcal{Q} \in \mathbb{L}$  and let  $G \in \mathcal{Q}^{\mathcal{M}}$ . Then

$$(1) \quad \chi'_{f, \mathcal{Q}}(G) = \max \frac{|E(H)|}{\mathcal{Q}(H)},$$

where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

**Proof.** Since  $G$  is  $\mathcal{Q}$ -matroidal, the hypergraph  $\mathcal{H}_G = (E_G, \mathcal{Q}_G)$  is a matroid. Claim 1 with Theorem 2 imply that

$$\chi'_{f, \mathcal{Q}}(G) = \max_{X \subseteq E_G; X \neq \emptyset} \frac{|X|}{\rho(X)} = \max \frac{|E(H)|}{\mathcal{Q}(H)},$$

where the maximum is taken over all nontrivial subgraphs  $H$  of  $G$ .

Now we show that we may restrict our attention to connected  $H$ . Suppose that the maximum on the right-hand side of (1) is achieved for a graph  $H$  with more than one component. Let  $H = H_1 \cup \dots \cup H_n$ , where  $H_i$  are the components of  $H$ . If one of these components, say  $H_j$ , is an empty graph (set of isolated vertices), then  $\frac{|E(H)|}{\mathcal{Q}(H)} = \frac{|E(H - H_j)|}{\mathcal{Q}(H - H_j)}$ . Thus we can assume that each component has at least one edge. Then  $\frac{|E(H)|}{\mathcal{Q}(H)} = \frac{|E(H_1)| + \dots + |E(H_n)|}{\mathcal{Q}(H_1) + \dots + \mathcal{Q}(H_n)} \leq \max_i \left\{ \frac{|E(H_i)|}{\mathcal{Q}(H_i)} \right\}$ . ■

We can now determine the fractional  $\mathcal{Q}$ -chromatic index for  $\mathcal{Q}$ -matroidal graphs. The following question arises: Which graphs are  $\mathcal{Q}$ -matroidal for given properties  $\mathcal{Q}$ ?

Each hereditary property  $\mathcal{Q}$  can be determined by the set of *minimal forbidden subgraphs*  $F(\mathcal{Q}) = \{G \in \mathcal{I}; G \notin \mathcal{Q} \text{ but } G \setminus \{e\} \in \mathcal{Q} \text{ for each } e \in E(G)\}$ . For example:  $F(\mathcal{O}_k) = \{H; H \text{ is a tree on } k+2 \text{ vertices}\}$ ;  $F(\mathcal{I}_k) = \{K_{k+2}\}$ . Simões-Pereira [14] proved that if  $F(\mathcal{Q})$  is finite, then  $\mathcal{Q}$  is not matroidal.

In [9] there is the following characterization of  $\mathcal{Q}$ -matroidal graphs.

**Theorem 5** [9]. A graph  $G = (V, E)$  is  $\mathcal{Q}$ -matroidal if and only if for each  $\mathcal{Q}$ -independent set  $I \subseteq E$  and for each edge  $e \in E \setminus I$  the graph  $G[I \cup \{e\}]$  induced by  $I \cup \{e\}$  contains at most one minimal forbidden subgraph of  $\mathcal{Q}$ .

By Theorem 5 each graph  $G$  which contains either at most one minimal forbidden subgraph of  $\mathcal{Q}$  or only edge-disjoint minimal forbidden subgraphs of  $\mathcal{Q}$  is  $\mathcal{Q}$ -matroidal.

**Lemma 6** [9]. The property  $\mathcal{Q}^{\mathcal{M}}$  belongs to  $\mathbb{L}$  for every  $\mathcal{Q} \in \mathbb{L}$ .

By Lemma 6 we can characterize the structure of  $\mathcal{Q}$ -matroidal graphs by describing the set of minimal forbidden subgraphs  $F(\mathcal{Q}^M)$ .

For any two given graphs  $G_1$  and  $G_2$  with a common induced subgraph  $H$  we construct the graph  $G = (G_1; H; G_2)$  by amalgamation of  $G_1$  and  $G_2$  with respect to  $H$  so that  $V(G) = V(G_1) \cup V(G_2)$ ,  $E(G) = E(G_1) \cup E(G_2)$  and  $H = (V(G_1) \cap V(G_2), E(G_1) \cap E(G_2))$ .

In the following  $P_n$  and  $C_n$  will denote the path and the cycle on  $n$  vertices, respectively.  $D_n$  will denote the complement of  $K_n$ .

**Theorem 7** [9]. *Let  $G$  be a graph and let  $k \geq 1$ . Then*

- $G \in F(\mathcal{O}_k^M)$  if and only if  $G \in T \setminus \{K_{1,k+2}; C_{k+2}\}$ , where  $T$  is the set of all trees on  $k+3$  vertices and all unicyclic graphs on  $k+2$  vertices,
- $G \in F(\mathcal{S}_k^M)$  if and only if  $G = (K_{1,k+1}; K_2 \cup D_p; K_{1,k+1})$  for some  $0 \leq p \leq k$  and  $k \geq 2$ , where  $K_2$  joins the central vertices of the stars,
- $G \in F(\mathcal{I}_k^M)$  if and only if  $G = (K_{k+2}; K_r; K_{k+2})$  for some  $2 \leq r \leq k+1$ ,
- $G \in F(\mathcal{B}^M)$  if and only if  $G = (C_{2p+1}; P_q; C_r)$  for some  $p \geq 1$ ,  $q \geq 2$  and  $r \geq 3$ .

The seminal result on fractional edge colorings is due to Edmonds [5]. For a graph  $G$  we define  $\Gamma(G) = \max \frac{2|E(H)|}{|V(H)| - 1}$ , where the maximization is over every induced subgraph  $H$  of  $G$  with  $|V(H)| \geq 3$  and  $|V(H)|$  odd.

**Theorem 8** [5]. *Let  $G$  be a graph. Then*

$$\chi'_{f, \mathcal{J}_1}(G) = \chi'_{f, \mathcal{S}_1}(G) = \chi'_{f, \mathcal{O}_1}(G) = \chi'_f(G) = \max\{\Delta(G), \Gamma(G)\}.$$

**Lemma 9.** *Every graph is  $\mathcal{D}_1$ -matroidal.*

**Proof.** Clearly,  $F(\mathcal{D}_1)$  is a set of cycles. Moreover, if we add an edge to a tree (forest) we obtain exactly (at most) one cycle. So the claim follows from Theorem 5. ■

Although all graphs are  $\mathcal{D}_1$ -matroidal, for  $k \geq 2$  the characterization of  $\mathcal{D}_k$ -matroidal graphs seems to be difficult.

**Theorem 10.** *Let  $G$  be a graph. Then*

$$\chi'_{f, \mathcal{D}_1}(G) = \max \frac{|E(H)|}{|V(H)| - 1},$$

where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

**Proof.** From Lemma 9 it follows that  $G$  is  $\mathcal{D}_1$ -matroidal. Any spanning tree of a connected graph  $H$  on  $n$  vertices has  $n - 1$  edges, therefore  $\mathcal{D}_1(H) = |V(H)| - 1$ . Theorem 4 implies  $\chi'_{f, \mathcal{D}_1}(G) = \max_{H \subseteq G} \frac{|E(H)|}{\mathcal{D}_1(H)} = \max_{H \subseteq G} \frac{|E(H)|}{|V(H)| - 1}$ . ■

**Corollary 11.** *Let  $G$  be a graph and let  $\mathcal{Q} \in \mathbb{L}$  such that  $\mathcal{D}_1 \subseteq \mathcal{Q}$ . Then*

$$\chi'_{f, \mathcal{Q}}(G) \leq \max \frac{|E(H)|}{|V(H)| - 1},$$

where the maximization is over all connected nontrivial subgraphs  $H$  of  $G$ .

**Lemma 12.** *Let  $k \geq 1$ . The graph  $G$  is  $\mathcal{I}_k$ -matroidal if and only if any two complete graphs on  $k + 2$  vertices are edge-disjoint in  $G$ .*

**Proof.** Assume that  $G$  contains two complete graphs on  $k + 2$  vertices which have  $r \geq 2$  vertices in common. These  $r$  vertices induce  $K_r$ , hence  $G$  contains  $(K_{k+2}; K_r; K_{k+2})$  as a subgraph. So  $G \notin \mathcal{I}_k^{\mathcal{M}}$  since  $(K_{k+2}; K_r; K_{k+2}) \in F(\mathcal{I}_k^{\mathcal{M}})$  (see Theorem 7).

If  $G \notin \mathcal{I}_k^{\mathcal{M}}$ , then  $G$  contains a forbidden subgraph  $(K_{k+2}; K_r; K_{k+2})$  for some  $2 \leq r \leq k + 1$ , thus it contains two complete graphs on  $k + 2$  vertices which share an edge. ■

Let  $H_{k+2}$  denote the number of complete graphs on  $k + 2$  vertices in the graph  $H$ .

**Theorem 13.** *Let  $G$  be an  $\mathcal{I}_k$ -matroidal graph,  $k \geq 1$ . Then*

$$\chi'_{f, \mathcal{I}_k}(G) = \max \frac{|E(H)|}{|E(H)| - H_{k+2}},$$

where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

**Proof.** From Theorem 4 it follows that  $\chi'_{f, \mathcal{I}_k}(G) = \max_{H \subseteq G} \frac{|E(H)|}{\mathcal{I}_k(H)}$ . So it is sufficient to show that  $\mathcal{I}_k(H) = |E(H)| - H_{k+2}$ .

Lemma 12 implies that any two complete graphs on  $k + 2$  vertices are edge-disjoint in every subgraph  $H$  of  $G$ . Hence, if we remove less than  $H_{k+2}$  edges from  $H$ , then the obtained graph still contains at least one  $K_{k+2}$ . Therefore  $\mathcal{I}_k(H) \leq |E(H)| - H_{k+2}$ .

On the other hand, if we remove one edge from each  $K_{k+2}$ , then the remaining edges form an  $\mathcal{I}_k$ -independent set, hence  $\mathcal{I}_k(H) \geq |E(H)| - H_{k+2}$ . ■

**Lemma 14.** *Let  $k \geq 2$ . The graph  $G$  is  $\mathcal{S}_k$ -matroidal if and only if no two vertices of degree at least  $k + 1$  are incident in  $G$ .*

**Proof.** Let  $uv$  be an edge of  $G$  such that its endvertices have degree at least  $k+1$ . Let  $G_1$  be a subgraph of  $G$  which contains only the edges incident with  $u$  or  $v$ . Clearly,  $G_1$  contains a subgraph  $G_2$  in which the vertices  $u$  and  $v$  are joined by an edge and they have degree  $k+1$ . Let  $p$  denote the number of common neighbors of  $u$  and  $v$  in  $G_2$ . Observe that  $G_2 = (K_{1,k+1}; K_2 \cup D_p; K_{1,k+1})$ , consequently  $G_2 \in F(\mathcal{S}_k^{\mathcal{M}})$ . So  $G$  cannot be  $\mathcal{S}_k$ -matroidal.

If  $G \notin \mathcal{S}_k^{\mathcal{M}}$ , then it contains a minimal forbidden subgraph  $(K_{1,k+1}; K_2 \cup D_p; K_{1,k+1})$  for some  $0 \leq p \leq k$ . The central vertices of these stars are joined by an edge and they have degree  $k+1$ . ■

**Theorem 15.** *Let  $G$  be an  $\mathcal{S}_k$ -matroidal graph,  $k \geq 2$ . Then*

$$\chi'_{f, \mathcal{S}_k}(G) = \max \frac{|E(H)|}{|E(H)| - \sum_{\substack{v \in V(H) \\ \deg_H(v) \geq k+1}} (\deg_H(v) - k)},$$

where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

**Proof.** Let  $H$  be a subgraph of  $G$ . If for every vertex  $v$  of  $H$  of degree at least  $k+1$  we remove  $\deg_H(v) - k$  edges incident with it, then we obtain a graph whose maximum degree is at most  $k$ . Therefore

$$\mathcal{S}_k(H) \geq |E(H)| - \sum_{\substack{v \in V(H) \\ \deg_H(v) \geq k+1}} (\deg_H(v) - k).$$

The opposite inequality follows from the fact that no two vertices of degree at least  $k+1$  are incident in  $G$ , thus neither in  $H \subseteq G$  (see Lemma 14). Therefore the claim follows from Theorem 4. ■

**Lemma 16.** *The graph  $G$  is  $\mathcal{B}$ -matroidal if and only if no odd cycle of  $G$  shares an edge with any other cycle of  $G$ .*

**Proof.**  $G \notin \mathcal{B}^{\mathcal{M}}$  if and only if  $G$  contains a minimal forbidden subgraph  $(C_{2p+1}; P_q; C_r)$  for some  $p \geq 1$ ,  $q \geq 2$  and  $r \geq 3$ . Equivalently,  $G$  contains an odd cycle which shares an edge with an other cycle. ■

**Corollary 17.** *If  $G \in \mathcal{B}^{\mathcal{M}}$ , then the odd cycles of  $G$  are edge-disjoint.*

Let  $oc(G)$  denote the number of odd cycles in the graph  $G$ .

**Theorem 18.** *Let  $G$  be a  $\mathcal{B}$ -matroidal graph. Then*

$$\chi'_{f, \mathcal{B}}(G) = \max \frac{|E(H)|}{|E(H)| - oc(H)},$$

where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

**Proof.** Let  $H$  be a subgraph of  $G$ . If we remove one edge from every odd cycle of  $H$ , then the remaining edges induce a bipartite graph, hence  $\mathcal{B}(H) \geq |E(H)| - oc(H)$ .

The odd cycles in  $H$  are edge-disjoint (see Corollary 17), thus we must remove at least  $oc(H)$  edges from  $E(H)$  to obtain a  $\mathcal{B}$ -independent set. Therefore  $\mathcal{B}(H) \leq |E(H)| - oc(H)$ .

Consequently,  $\mathcal{B}(H) = |E(H)| - oc(H)$  and hence the assertion follows from Theorem 4. ■

**Lemma 19.** *Let  $k \geq 1$ . The graph  $G$  is  $\mathcal{O}_k$ -matroidal if and only if  $G$  either belongs to  $\mathcal{O}_k$  or it is isomorphic to  $K_{1,p}$ ,  $p \geq k+1$ , to  $C_{k+2}$  or to a tree on  $k+2$  vertices.*

**Proof.**  $G$  is  $\mathcal{O}_k$ -matroidal if and only if it does not contain any subgraph from  $F(\mathcal{O}_k^{\mathcal{M}})$ . So the claim follows from Theorem 7. ■

Clearly, if  $G \in \mathcal{O}_k$ , then its fractional  $\mathcal{O}_k$ -edge chromatic number equals one. If  $G \in \mathcal{O}_k^{\mathcal{M}} \setminus \mathcal{O}_k$ , then it has  $k+2$  vertices or it is a star on at least  $k+3$  vertices.

**Theorem 20.** *Let  $G \in \mathcal{O}_k^{\mathcal{M}} \setminus \mathcal{O}_k$  and let  $|V(G)| = k+2$ ,  $k \geq 2$ . Then*

$$\chi'_{f, \mathcal{O}_k}(G) = \frac{|E(G)|}{|E(G)| - \lambda(G)},$$

where  $\lambda(G)$  is the edge-connectivity of  $G$ .

**Proof.** Let  $H$  be a connected subgraph of  $G$ . If  $E(H)$  is not  $\mathcal{O}_k$ -independent, then either  $|E(H)| = k+2$  or  $|E(H)| = k+1$ . In the first case  $H = C_{k+2}$ , hence  $\mathcal{O}_k(H) = |E(H)| - 2$ . In the second case  $H$  is a tree, therefore  $\mathcal{O}_k(H) = |E(H)| - 1$ . Thus the claim follows from Theorem 4. ■

**Theorem 21.** *Let  $G \in \mathcal{O}_k^{\mathcal{M}} \setminus \mathcal{O}_k$  and let  $|V(G)| = k+i$ ,  $k \geq 2$ ,  $i \geq 3$ . Then*

$$\chi'_{f, \mathcal{O}_k}(G) = \frac{|E(G)|}{|E(G)| - i + 1} = \frac{k+i-1}{k}.$$

**Proof.** It follows from Theorem 4 and from the fact that  $G$  is a star. ■

### 3. EXAMPLES

**Example 22.** *Let  $K_{2,3}$  denote the complete bipartite graph on  $2+3$  vertices. We will show that  $\chi'_{f, \mathcal{S}_2}(K_{2,3}) = \frac{3}{2}$ .*



Solution 1.

From Lemma 14 it follows that  $K_{2,3} \in \mathcal{S}_2^{\mathcal{M}}$ . From Theorem 15 we have  $\chi'_{f, \mathcal{S}_2}(K_{2,3}) = \max \frac{|E(H)|}{|E(H)| - \sum_{\substack{v \in V(H) \\ \deg_H(v)=3}} 1}$ , where the maximum is taken over all connected nontrivial subgraphs  $H$  of  $G$ .

If  $H$  is a connected subgraph of  $G$ , then either  $H \in \mathcal{S}_2$  or it is a graph from Figure 1. So  $\chi'_{f, \mathcal{S}_2}(K_{2,3}) = \max\{1, \frac{3}{2}, \frac{4}{3}, \frac{5}{4}\} = \frac{3}{2}$ .




Figure 1. Connected subgraphs of  $K_{2,3}$  which are not in  $\mathcal{S}_2$ .

Solution 2.

Fractional  $\mathcal{Q}$ -edge-colorings may be viewed in several ways. We present an equivalent definition. Let  $r, s$  be positive integers with  $r \geq s$ . An  $(r, s)$ -fractional  $\mathcal{Q}$ -edge-coloring of  $G$  is an assignment of  $s$ -element subsets of  $\{1, \dots, r\}$  to the edges of  $G$  such that each color class induces a graph of property  $\mathcal{Q}$ . Then the fractional  $\mathcal{Q}$ -edge chromatic number of  $G$  is defined as

$$\chi'_{f, \mathcal{Q}}(G) = \inf \left\{ \frac{r}{s} : G \text{ has an } (r, s)\text{-fractional } \mathcal{Q}\text{-edge-coloring} \right\}.$$

Note that in this definition we can replace the infimum by the minimum.

For each  $(r, s)$ -fractional  $\mathcal{S}_2$ -edge-coloring of  $K_{2,3}$  and for each color  $i \in \{1, \dots, r\}$  the following holds: at most four edges are colored with sets containing the color  $i$ . On the other hand, every edge is assigned with an  $s$ -element color set. This implies that  $4r \geq 6s$ , hence  $\chi'_{f, \mathcal{S}_2}(K_{2,3}) \geq \frac{3}{2}$ .

To prove the inequality  $\chi'_{f, \mathcal{S}_2}(K_{2,3}) \leq \frac{3}{2}$  we construct a  $(3, 2)$ -fractional  $\mathcal{S}_2$ -edge-coloring of  $K_{2,3}$ , see Figure 2.




Figure 2. A  $(3, 2)$ -fractional  $\mathcal{S}_2$ -edge-coloring of the graph  $K_{2,3}$ .

The following results immediately follows from Theorems 13, 15 and 18.

**Example 23.** If  $k \geq 1$ , then  $\chi'_{f, \mathcal{I}_k}(K_{k+2}) = \frac{\binom{k+2}{2}}{\binom{k+2}{2} - 1}$ ,  $\chi'_{f, \mathcal{S}_k}(K_{1,k+1}) = \frac{k+1}{k}$   
and  $\chi'_{f, \mathcal{B}}(C_{2k+1}) = \frac{2k+1}{2k}$ .

### Acknowledgment

The authors would like to thank anonymous referees for many helpful comments and constructive suggestions.

### REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, *Graph Theory* (Springer, 2008).  
doi:10.1007/978-1-84628-970-5
- [2] M. Borowiecki, A. Kemnitz, M. Marangio and P. Mihók, *Generalized total colorings of graphs*, Discuss. Math. Graph Theory **31** (2011) 209–222.  
doi:10.7151/dmgt.1540
- [3] I. Broere, S. Dorfling and E. Jonck, *Generalized chromatic numbers and additive hereditary properties of graphs*, Discuss. Math. Graph Theory **22** (2002) 259–270.  
doi:10.7151/dmgt.1174
- [4] M.J. Dorfling and S. Dorfling, *Generalized edge-chromatic numbers and additive hereditary properties of graphs*, Discuss. Math. Graph Theory **22** (2002) 349–359.  
doi:10.7151/dmgt.1180
- [5] J. Edmonds, *Maximum matching and a polyhedron with 0,1-vertices*, J. Res. Nat. Bur. Standards **69B** (1965) 125–130.
- [6] G. Karafová, *Generalized fractional total coloring of complete graphs*, Discuss. Math. Graph Theory, accepted.
- [7] A. Kemnitz, M. Marangio, P. Mihók, J. Oravcová and R. Soták, *Generalized fractional and circular total coloring of graphs*, preprint.
- [8] K. Kilakos and B. Reed, *Fractionally colouring total graphs*, Combinatorica **13** (1993) 435–440.  
doi:10.1007/BF01303515
- [9] P. Mihók, *On graphs matroidal with respect to additive hereditary properties*, Graphs, Hypergraphs and Matroids II, Zielona Góra (1987) 53–64.
- [10] P. Mihók, Zs. Tuza and M. Voigt, *Fractional  $\mathcal{P}$ -colourings and  $\mathcal{P}$ -choice-ratio*, Tatra Mt. Math. Publ. **18** (1999) 69–77.
- [11] J.G. Oxley, *Matroid Theory* (Oxford University Press, Oxford, 1992).
- [12] E.R. Scheinerman and D.H. Ullman, *Fractional Graph Theory* (John Wiley & Sons, 1997).

- [13] R. Schmidt, *On the existence of uncountably many matroidal families*, Discrete Math. **27** (1979) 93–97.  
doi:10.1016/0012-365X(79)90072-4
- [14] J.M.S. Simões-Pereira, *On matroids on edge sets of graphs with connected subgraphs as circuits*, Proc. Amer. Math. Soc. **38** (1973) 503–506.  
doi:10.2307/2038939

Received 3 November 2011

Revised 29 May 2012

Accepted 29 May 2012

