

NOTE

SMALLEST REGULAR GRAPHS OF GIVEN DEGREE AND DIAMETER

MARTIN KNOR¹

Slovak University of Technology in Bratislava
Faculty of Civil Engineering, Department of Mathematics
Radlinského 11, 813 68 Bratislava

e-mail: knor@math.sk

Abstract

In this note we present a sharp lower bound on the number of vertices in a regular graph of given degree and diameter.

Keywords: regular graph, degree/diameter problem, extremal graph.

2010 Mathematics Subject Classification: 05C35, 05C12.

1. INTRODUCTION

The *degree/diameter* problem consists in determination of the largest order $N(d, k)$ of a graph with (maximum) degree d and diameter k . An upper bound for $N(d, k)$ is the *Moore bound* $M(d, k) = 1 + d + d(d - 1) + \cdots + d(d - 1)^{k-1}$ and graphs achieving this bound are called *Moore graphs*. As shown in [1, 3, 5], Moore graphs exist only when $d = 2$ or $k = 1$ or when $k = 2$ and the degree is either 3 or 7 or possibly 57. For all other pairs (d, k) we have $N(d, k) \leq M(d, k) - 2$, see [2, 4]. Recently, there are plenty of papers dealing with the degree/diameter problem, some of them constructing “large” graphs of given degree and diameter, which increases the lower bound for $N(d, k)$ for special pairs (d, k) , other decreasing $N(d, k)$ for special classes of graphs. For a nice survey see [7].

In this note we consider the inverse of degree/diameter problem. Since usually the degree/diameter problem is formulated for regular graphs (although some authors require only that d is the maximum degree), we ask what is the minimum

¹The author acknowledges partial support by Slovak research grants VEGA 1/0781/11 and APVV-0223-10.

order $n(d, k)$ of a regular graph of degree d and diameter k . In this note we answer this question completely.

We start with some notation. Let G be a graph, $G = (V(G), E(G))$. For two of its vertices, say x and y , by $\text{dist}_G(x, y)$ we denote their distance in G . By $N_i(x)$ we denote the set of vertices that are at distance i from x . As usual, $N_1(x)$ is often abbreviated to $N(x)$. The longest distance in G is the *diameter* $\text{diam}(G)$. The complete graph on n vertices is denoted by K_n and the discrete graph on n vertices (the complement of K_n) is denoted by D_n . If G is a graph, then by $G^{(-1)}$ (and $G^{(-2)}$) we denote a graph obtained from G by removing all the edges of one 1-factor (one 2-factor).

If G and H are graphs, then $G + H$ denotes the *join* of G and H , that is, a graph obtained from the disjoint union of G and H by adding all edges xy , where $x \in V(G)$ and $y \in V(H)$. The *sequential join* of graphs G_1, G_2, \dots, G_r is denoted by $G_1 + G_2 + \dots + G_r$ and is defined by

$$G_1 + G_2 + \dots + G_r = (G_1 + G_2) \cup (G_2 + G_3) \cup \dots \cup (G_{r-1} + G_r).$$

Thus, one can obtain $G_1 + G_2 + \dots + G_r$ from the disjoint union $G_1 \cup G_2 \cup \dots \cup G_r$ by adding all edges xy where $x \in V(G_i)$ and $y \in V(G_{i+1})$ for $i = 1, 2, \dots, r-1$. To simplify the expressions, instead of

$$\dots + \underbrace{G + G + \dots + G}_{k \text{ times}} + \dots \quad \text{we write} \quad \dots + (G)_k + \dots$$

Finally, denote by $G \div H$ a graph obtained from the disjoint union of G and H by adding all edges of one 1-factor, every edge of which joins a vertex of G with a vertex of H . Obviously, $G \div H$ is defined only if $|V(G)| = |V(H)|$. Analogously as in the case of join, by $G_1 \div G_2 \div \dots \div G_r$ we denote the graph $(G_1 \div G_2) \cup (G_2 \div G_3) \cup \dots \cup (G_{r-1} \div G_r)$. We can form also more complicated expressions using both $+$ and \div . In such a way, $K_1 + D_2 \div D_2 \div K_2$ is a cycle of length 7; see Figure 1.




Figure 1. The graph $K_1 + D_2 \div D_2 \div K_2$.

2. RESULTS

For small diameters we have the following statement.

Proposition 1. *Let $d \geq 2$. We have*

- (i) $n(d, 1) = d + 1$;
- (ii) *if d is even, then $n(d, 2) = d + 2$;*
- (iii) *if d is odd, then $n(d, 2) = d + 3$;*
- (iv) $n(d, 3) = 2d + 2$.

Proof. The case $k = 1$ is obvious since K_{d+1} is the unique graph of diameter 1 and degree d .

Let $k = 2$. Let G be a d -regular graph of diameter 2, and let $x, y \in V(G)$ such that $\text{dist}_G(x, y) = 2$. Then $\{x\} \cup N(x) = N_0(x) \cup N_1(x)$, which gives $|N_0(x)| + |N_1(x)| = d + 1$. Since $y \in N_2(x)$, we have $|V(G)| = |N_0(x)| + |N_1(x)| + |N_2(x)| \geq d + 2$, which gives $n(d, 2) \geq d + 2$. However, if d is odd then $|V(G)|$ cannot be odd and so $n(d, 2) \geq d + 3$ in this case. If d is even then $K_{d+2}^{(-1)}$ is a d -regular graph of diameter 2 on $d + 2$ vertices, which shows $n(d, 2) \leq d + 2$; while if d is odd then $K_{d+3}^{(-2)}$ is a d -regular graph of diameter 2 on $d + 3$ vertices, which shows $n(d, 2) \leq d + 3$.

Finally, let $k = 3$. Analogously as above, let G be a d -regular graph of diameter 3, and let $x, y \in V(G)$ such that $\text{dist}_G(x, y) = 3$. Then $\{x\} \cup N(x) = N_0(x) \cup N_1(x)$, which gives $|N_0(x)| + |N_1(x)| = d + 1$, and $\{y\} \cup N(y) \subseteq N_2(x) \cup N_3(x)$, which gives $|N_2(x)| + |N_3(x)| \geq d + 1$. Thus, $|V(G)| = |N_0(x)| + |N_1(x)| + |N_2(x)| + |N_3(x)| \geq 2d + 2$, and so $n(d, 3) \geq 2d + 2$. On the other hand, denote by $K_{n,n}$ a complete bipartite graph on $2n$ vertices in which the two partite sets have n vertices each. Then $K_{d+1, d+1}^{(-1)}$ is a d -regular graph of diameter 3 on $2d + 2$ vertices, which shows $n(d, 3) \leq 2d + 2$. ■

Now we turn our attention to larger diameters. Since there are only two 2-regular graphs of diameter k , namely the cycle on $2k$ vertices and the cycle on $2k + 1$ vertices, we have the following trivial observation.

Proposition 2. *If $k \geq 4$, then $n(2, k) = 2k$.*

For larger degrees we have a slightly different bound.

Theorem 3. *Let $k = 3j + t$, where $k \geq 4$ and $0 \leq t \leq 2$, and let $d \geq 3$. Then $n(d, k) = (d + 1)(j + 1) + t + \delta$, where $\delta = 1$ if either d is odd and $t = 1$ or d is even and $t = 2$. Otherwise $\delta = 0$.*

Proof. First we prove a lower bound for $n(d, k)$. Let G be a regular graph of degree d and diameter k and let $x, y \in V(G)$ such that $\text{dist}_G(x, y) = k$. Denote $n_i = |N_i(x)|$. Since $x \in N_0(x)$, we have $\{x\} \cup N(x) \subseteq N_0(x) \cup N_1(x)$. Thus, $n_0 + n_1 \geq d + 1$. Analogously $n_{k-1} + n_k \geq d + 1$ since $y \in N_k(x)$. Further, for every i , $1 \leq i \leq j-1$, we have $n_{3i-1} + n_{3i} + n_{3i+1} \geq d + 1$ since for $z_i \in N_{3i}(x)$ it holds $\{z_i\} \cup N(z_i) \subseteq N_{3i-1}(x) \cup N_{3i}(x) \cup N_{3i+1}(x)$. Finally, if $t \geq 1$ then $n_{k-1-\ell} \geq 1$ where $1 \leq \ell \leq t$. Summing up all these inequalities we get

$$|V(G)| = \sum_{i=0}^k n_i \geq (d+1)(j+1) + t.$$

If $t = 2$ then we use $n_{k-3} \geq 1$ and $n_{k-2} \geq 1$. But if d is even then G cannot have a bridge, and so $n_{k-3} + n_{k-2} \geq 3$. Thus, we get $|V(G)| = \sum_{i=0}^k n_i \geq (d+1)(j+1) + t + 1$ in this case.

Similarly, if $t = 1$ and d is odd then $(d+1)(j+1) + t$ is an odd number. But a regular graph of odd degree cannot have an odd number of vertices, and so $|V(G)| = \sum_{i=0}^k n_i \geq (d+1)(j+1) + t + 1$ also in this case.

To prove the upper bound we construct extremal graphs, that is, regular graphs of degree d and diameter k on $n(d, k)$ vertices. First we define an extremal graph G for odd d . The case $k = 4$ is treated separately. If $d = 3$ then one extremal graph G is on Figure 2. For $d \geq 5$ we set $G = K_2 + K_{d-1}^{(-2)} + D_2 \div D_2 + K_{d-1}$.




Figure 2. An extremal graph for $d = 3$ and $k = 4$.

Recall that $k = 3j + t$. To cover the remaining diameters, that is, 5, 6, 7, ..., in the next we assume $j \geq 1$ if $t = 2$, and $j \geq 2$ if $t = 0$ or $t = 1$:

$$G = K_2 + K_{d-1}^{(-1)} + (K_1 + K_1 + K_{d-1})_{j-1} + K_1 + K_1 + K_{d-1}^{(-1)} + K_2, \text{ if } t = 2;$$

$$G = K_2 + K_{d-1}^{(-1)} + (K_1 + K_1 + K_{d-1})_{j-2} + K_1 + K_1 + K_{d-1} \div K_{d-1}^{(-1)} + K_2, \text{ if } t = 0;$$

$$G = K_2 + K_{d-1}^{(-1)} + (K_1 + K_1 + K_{d-1})_{j-2} + K_1 + K_1 + K_{d-1}^{(-1)} + D_2 \div D_2 + K_{d-1}, \text{ if } t = 1.$$

Now we define an extremal graph G for even d . To cover all possible diameters, that is, 4, 5, 6, ..., in the next we assume $j \geq 1$ if $t = 1$ or $t = 2$, and $j \geq 2$ if $t = 0$:

$$G = K_3 + K_{d-2}^{(-1)} + (K_1 + K_2 + K_{d-2})_{j-1} + K_1 + D_2 + K_{d-1}, \text{ if } t = 1;$$

$$G = K_3 + K_{d-2}^{(-1)} + (K_1 + K_2 + K_{d-2})_{j-1} + K_1 + K_2 + K_{d-2}^{(-2)} + K_3, \text{ if } t = 2;$$

$G = K_3 + K_{d-2}^{(-1)} + (K_1 + K_2 + K_{d-2})_{j-2} + K_1 + K_2 + K_{d-2} \div K_{d-2}^{(-1)} + K_3$, if $t = 0$.

Observe that in all these graphs, whenever we removed a 1-factor out of K_q , then the number of vertices q was even. Obviously, in each case G has diameter k and it is a matter of routine to check that G is a regular graph of degree d . (For example, a vertex in the last copy of $K_{d-2}^{(-1)}$ in the last graph is joined to 1 vertex of K_{d-2} , $d-4$ vertices of $K_{d-2}^{(-1)}$ and to 3 vertices of K_3 , so its degree is $1 + d - 4 + 3 = d$.) Also, in each of these cases the number of vertices of G attains the bound of the theorem. To verify this statement it suffices to check the number of vertices for the smallest admissible values of j since in each case in the brackets we have exactly $d + 1$ vertices. ■

By Proposition 2, if $d = 2$ then $n(d, k) = dk$. However, for higher degrees we get $n(d, k) \sim \frac{1}{3}dk$. Denote by $n_{\text{VT}}(d, k)$ the minimum number of vertices in a vertex-transitive d -regular graph with diameter k . As shown in [6], for $k \geq 4$ and “large” d we have $n_{\text{VT}}(d, k) \sim \frac{2}{3}dk$, and so $n_{\text{VT}}(d, k) \doteq 2n(d, k)$ in this case. On the other hand, since the extremal graphs constructed in the proof of Proposition 1 are vertex-transitive, we have $n_{\text{VT}}(d, k) = n(d, k)$ when $k \leq 3$.

REFERENCES

- [1] E. Bannai and T. Ito, *On finite Moore graphs*, J. Fac. Sci. Tokyo Univ. **20** (1973) 191–208.
- [2] E. Bannai and T. Ito, *Regular graphs with excess one*, Discrete Math. **37** (1981) 147–158.
doi:10.1016/0012-365X(81)90215-6
- [3] R.M. Damerell, *On Moore graphs*, Proc. Cambridge Philos. Soc. **74** (1973) 227–236.
doi:10.1017/S0305004100048015
- [4] P. Erdős, S. Fajtlowicz and A.J. Hoffman, *Maximum degree in graphs of diameter 2*, Networks **10** (1980) 87–90.
doi:10.1002/net.3230100109
- [5] A.J. Hoffman and R.R. Singleton, *On Moore graphs with diameter 2 and 3*, IBM J. Res. Develop. **4** (1960) 497–504.
doi:10.1147/rd.45.0497
- [6] M. Knor and J. Širáň, *Smallest vertex-transitive graphs of given degree and diameter*, J. Graph Theory, (to appear).
- [7] M. Miller and J. Širáň, *Moore graphs and beyond: A survey of the degree-diameter problem*, Electron. J. Combin., Dynamic survey No. D14 (2005), 61pp.

Received 7 March 2012

Revised 17 September 2012

Accepted 2 October 2012