

THE DISTANCE ROMAN DOMINATION NUMBERS OF GRAPHS

HAMIDEH ARAM, SEPIDEH NOROUZIAN,

SEYED MAHMOUD SHEIKHOESLAMI

Department of Mathematics
Azərbaycan Şahid Mədanı University
Tabriz, I.R. Iran

e-mail: s.m.sheikholeslami@azaruniv.edu

AND

LUTZ VOLKMANN

Lehrstuhl II für Mathematik
RWTH Aachen University
52056 Aachen, Germany

e-mail: volkm@math2.rwth-aachen.de

Abstract

Let k be a positive integer, and let G be a simple graph with vertex set $V(G)$. A k -distance Roman dominating function on G is a labeling $f : V(G) \rightarrow \{0, 1, 2\}$ such that for every vertex with label 0, there is a vertex with label 2 at distance at most k from each other. The *weight* of a k -distance Roman dominating function f is the value $\omega(f) = \sum_{v \in V} f(v)$. The k -distance Roman domination number of a graph G , denoted by $\gamma_R^k(G)$, equals the minimum weight of a k -distance Roman dominating function on G . Note that the 1-distance Roman domination number $\gamma_R^1(G)$ is the usual Roman domination number $\gamma_R(G)$. In this paper, we investigate properties of the k -distance Roman domination number. In particular, we prove that for any connected graph G of order $n \geq k + 2$, $\gamma_R^k(G) \leq 4n/(2k + 3)$ and we characterize all graphs that achieve this bound. Some of our results extend these ones given by Cockayne *et al.* in 2004 and Chambers *et al.* in 2009 for the Roman domination number.

Keywords: k -distance Roman dominating function, k -distance Roman domination number, Roman dominating function, Roman domination number.

2010 Mathematics Subject Classification: 05C69.

1. TERMINOLOGY AND INTRODUCTION

In this paper, G is a simple graph with vertex set $V = V(G)$ and edge set $E = E(G)$. Denote by K_n the *complete graph*, by C_n the *cycle* and by P_n the *path* of order n , respectively. Given two graphs G_1 and G_2 such that $V(G_1) \cap V(G_2) = \emptyset$, the *disjoint union* is the graph $G_1 \cup G_2$ with vertex set $V(G_1) \cup V(G_2)$ and edge set $E(G_1) \cup E(G_2)$. Let k be a positive integer. For two vertices x and y , let $d(x, y)$ denote the distance between x and y in G . The *girth* $g(G)$ of a graph G is the length of its shortest cycle. For a vertex $v \in V(G)$, the *open k -neighborhood* $N_{k,G}(v)$ is the set $\{u \in V(G) \mid u \neq v \text{ and } d(u, v) \leq k\}$ and the *closed k -neighborhood* $N_{k,G}[v]$ is the set $N_{k,G}(v) \cup \{v\}$. The *open k -neighborhood* $N_{k,G}(S)$ of a set $S \subseteq V$ is the set $\bigcup_{v \in S} N_{k,G}(v)$, and the *closed k -neighborhood* $N_{k,G}[S]$ of S is the set $N_{k,G}(S) \cup S$. The *k -degree* of a vertex v is defined as $\deg_{k,G}(v) = |N_{k,G}(v)|$. The minimum and maximum k -degree of a graph G are denoted by $\delta_k(G)$ and $\Delta_k(G)$, respectively. If $\delta_k(G) = \Delta_k(G)$, then the graph G is called *distance- k -regular*. The *k -th power* G^k of a graph G is the graph with vertex set $V(G)$ where two different vertices u and v are adjacent if and only if the distance $d(u, v)$ is at most k in G . Now we observe that $N_{k,G}(v) = N_{1,G^k}(v) = N_{G^k}(v)$, $N_{k,G}[v] = N_{1,G^k}[v] = N_{G^k}[v]$, $\deg_{k,G}(v) = \deg_{1,G^k}(v) = \deg_{G^k}(v)$, $\delta_k(G) = \delta_1(G^k) = \delta(G^k)$ and $\Delta_k(G) = \Delta_1(G^k) = \Delta(G^k)$. Consult [6, 10] for the notation and terminology which are not defined here.

Let $k \geq 1$ be an integer. A set $D \subseteq V(G)$ is a *k -distance dominating set* of G if every vertex in $V(G) - D$ is within distance k of at least one vertex in D . The *k -distance domination number* $\gamma^k(G)$ of G is the minimum cardinality among all k -distance dominating sets of G .

A *k -distance Roman dominating function* (k DRDF) on a graph $G = (V, E)$ is a function $f : V \rightarrow \{0, 1, 2\}$ satisfying the condition that for every vertex v for which $f(v) = 0$, there is a vertex u for which $f(u) = 2$ and $d(u, v) \leq k$. The *weight* of a k DRDF f is the value $\omega(f) = \sum_{v \in V} f(v)$. The *k -distance Roman domination number* of a graph G , denoted by $\gamma_R^k(G)$, equals the minimum weight of a k DRDF on G . A $\gamma_R^k(G)$ -*function* is a k -distance Roman dominating function of G with weight $\gamma_R^k(G)$. A k -distance Roman dominating function $f : V \rightarrow \{0, 1, 2\}$ can be represented by the ordered partition (V_0, V_1, V_2) (or (V_0^f, V_1^f, V_2^f) to refer f) of V , where $V_i = \{v \in V \mid f(v) = i\}$. In this representation, its weight is $\omega(f) = |V_1| + 2|V_2|$. Since $V_1^f \cup V_2^f$ is a k -distance dominating set when f is a k DRDF, and since placing weight 2 at the vertices of a k -distance dominating set yields a k DRDF, we have

$$(1) \quad \gamma^k(G) \leq \gamma_R^k(G) \leq 2\gamma^k(G).$$

Note that the 1-distance Roman domination number $\gamma_R^1(G)$ is the usual *Roman domination number* $\gamma_R(G)$. The definition of the Roman dominating function was

given multiplicity by Steward [9] and ReVelle and Rosing [8]. Cockayne *et al.* [3] as well as Chambers *et al.* [2] have given a lot of results on Roman domination.

Our purpose in this paper is to initiate the study of the k -distance Roman domination number and establish some bounds for the k -distance Roman domination number of a graph. Some of our results extend many well-known results.

2. SOME BASIC RESULTS

We start with some known results on the classical Roman domination number.

Theorem A [4]. *For any graph G of order n and maximum degree $\Delta \geq 1$,*

$$\gamma_R(G) \geq \frac{2n}{\Delta + 1}.$$

Theorem B [3]. *For any graph G of order n and minimum degree δ ,*

$$\gamma_R(G) \leq \frac{2 + \ln((1 + \delta)/2)}{\delta + 1}n.$$

Theorem C [2]. *For any tree T of order $n \geq 3$, $\gamma_R(T) \leq 4n/5$.*

Theorem D [2]. *If G is a graph of order $n \geq 3$, then*

$$\gamma_R(G) + \gamma_R(\overline{G}) \leq n + 3.$$

Furthermore, equality holds only when G or $\overline{G}$ is C_5 or $\frac{n}{2}K_2$.

The next two observations are straightforward to verify.

Observation 1. *Let $f = (V_0, V_1, V_2)$ be any γ_R^k -function of a graph G . Then*

- (a) $\Delta_k(G[V_1]) \leq 1$.
- (b) *If $w \in V_1$, then $N_{k,G}(w) \cap V_2 = \emptyset$.*
- (c) *If $u \in V_0$, then $|V_1 \cap N_{k,G}(u)| \leq 2$.*
- (d) V_2 *is a γ^k -set of the induced subgraph $G[V_0 \cup V_2]$.*
- (e) *Let $H = G[V_0 \cup V_2]$. Then each vertex $v \in V_2$ with $N_{k,G}(v) \cap V_2 \neq \emptyset$ has at least two private neighbors relative to V_2 in the graph H .*

Observation 2. *Let $k \geq 1$ be an integer, and let G be a graph of order $n \geq 2$. If $\text{diam}(G) \leq k$, then $\gamma_R^k(G) = \gamma_R(K_n) = 2$.*

Observation 3. *If $k \geq 1$ is an integer and G is a graph of order n with $\Delta_k(G) \geq 1$, then*

$$\gamma_R^k(G) \geq \frac{2n}{\Delta_k(G) + 1}.$$

Proof. Using the facts $\gamma_R^k(G) = \gamma_R(G^k)$, $\Delta_k(G) = \Delta(G^k)$ and Theorem A, we obtain

$$\gamma_R^k(G) = \gamma_R(G^k) \geq \frac{2n}{\Delta(G^k) + 1} = \frac{2n}{\Delta_k(G) + 1}.$$

■

Applying Theorem B, we obtain analogously the next result.

Observation 4. For any graph G of order n ,

$$\gamma_R^k(G) \leq \frac{2 + \ln((1 + \delta_k(G))/2)}{\delta_k(G) + 1} n.$$

Observation 5. If $k \geq 1$ is an integer and G is a graph of order n with $\Delta_k(G) \geq 1$, then

$$\gamma_R^k(G) \leq n - \Delta_k(G) + 1.$$

Proof. Let v be a vertex of G such that $\deg_{k,G}(v) = \Delta_k(G)$. Then $f = (N_{k,G}(v), V(G) - N_{k,G}[v], \{v\})$ is a k DRDF on G with weight $n - \Delta_k(G) + 1$ and therefore $\gamma_R^k(G) \leq n - \Delta_k(G) + 1$. ■

Let $k \geq 1$ be an integer, and let H be a graph with $\Delta_k(H) = n(H) - 1 \geq 2$. Now let $G = rK_1 \cup sK_2 \cup H$ for two integers $r, s \geq 0$. Then $\Delta_k(G) = \Delta_k(H)$ and

$$\gamma_R^k(G) = r + 2s + 2 = n(G) - \Delta_k(G) + 1.$$

This family of graphs demonstrates that the upper bound in Observation 5 is sharp.

Observation 6. Let $k \geq 1$ be an integer, and let G be a graph of order $n \geq 2$. Then $\gamma_R^k(G) = 2$ if and only if $n = 2$ or $n \geq 3$ and $\Delta_k(G) = n - 1$.

Proof. Assume first that $n = 2$ or $n \geq 3$ and $\Delta_k(G) = n - 1$. If $n = 2$, then $\gamma_R^k(G) = 2$. If $n \geq 3$ and $\Delta_k(G) = n - 1$, then Observation 5 implies that

$$2 \leq \gamma_R^k(G) \leq n - \Delta_k(G) + 1 = 2$$

and therefore $\gamma_R^k(G) = 2$.

Conversely, assume that $\gamma_R^k(G) = 2$. If $\Delta_k(G) = 0$, then it follows that $n = 2$. If $\Delta_k(G) \geq 1$, then we deduce from Observation 3 that

$$2 = \gamma_R^k(G) \geq \frac{2n}{\Delta_k(G) + 1}$$

and hence $\Delta_k(G) + 1 \geq n$. This leads to $\Delta_k(G) = n - 1$, and the proof is complete. ■

Observation 7. Let $k \geq 1$ be an integer, and let G be a graph of order n . Then $\gamma_R^k(G) = n$ if and only if $G = rK_1 \cup sK_2$ for some integers $r, s \geq 0$.

Proof. If $G = rK_1 \cup sK_2$ for some integers $r, s \geq 0$, then obviously $\gamma_R^k(G) = n$.

Conversely, assume that $\gamma_R^k(G) = n$. If $\Delta_k(G) \geq 2$, then Observation 5 leads to the contradiction $\gamma_R^k(G) \leq n - 1$. Thus $\Delta_k(G) \leq 1$ and so $G = rK_1 \cup sK_2$ for some integers $r, s \geq 0$. ■

Observation 8. Let $k \geq 1$ be an integer, and let G be a graph of order $n \geq 4$. Then $\gamma_R^k(G) = 3$ if and only if $\Delta_k(G) = n - 2$.

Proof. Assume first that $\Delta_k(G) = n - 2$. Observation 6 implies that $\gamma_R^k(G) \geq 3$. Since we deduce from Observation 5 that $\gamma_R^k(G) \leq n - \Delta_k(G) + 1 = 3$, we obtain $\gamma_R^k(G) = 3$.

Conversely, assume that $\gamma_R^k(G) = 3$. By Observation 6, we have $\Delta_k(G) \leq n - 2$. Let now $f = (V_0, V_1, V_2)$ be a $\gamma_R^k(G)$ -function. We deduce from the assumption $n \geq 4$ that $|V_1| = |V_2| = 1$. Let $V_2 = \{v\}$ and $V_1 = \{w\}$. Since $\gamma_R^k(G) = 3$, it is obvious that $N_{k,G}[v] = V(G) - \{w\}$ and thus $\Delta_k(G) \geq n - 2$. This yields $\Delta_k(G) = n - 2$, and the proof is complete. ■

Observation 9. Let $k \geq 2$ be an integer, and let G be a graph of order $n \geq 3$. Then $\gamma_R^k(G) = n - 1$ if and only if $G = K_3 \cup rK_1 \cup sK_2$ or $G = P_3 \cup rK_1 \cup sK_2$ for some integers $r, s \geq 0$.

Proof. If $G = K_3 \cup rK_1 \cup sK_2$ or $G = P_3 \cup rK_1 \cup sK_2$ for some integers $r, s \geq 0$, then obviously $\gamma_R^k(G) = n - 1$.

Conversely, assume that $\gamma_R^k(G) = n - 1$. If $\Delta_k(G) \geq 3$, then Observation 5 implies the contradiction $\gamma_R^k(G) \leq n - \Delta_k(G) + 1 \leq n - 2$. Therefore $\Delta_k(G) \leq 2$. If $\Delta_k(G) \leq 1$, then we deduce from Observation 7 the contradiction $\gamma_R^k(G) = n$. Consequently, $\Delta_k(G) = 2$. If G contains at least two components H_1 and H_2 with $\Delta_k(H_1) = \Delta_k(H_2) = 2$, then $\gamma_R^k(G) \leq n - 2$, a contradiction. Hence G has exactly one component H with $\Delta_k(H) = 2$, and the remaining components are isolated vertices or isomorphic to K_2 . If $|V(H)| \geq 4$, then the assumption $k \geq 2$ shows that $\Delta_k(G) = \Delta_k(H) \geq 3$, a contradiction. Hence $|V(H)| = 3$ and so $G = K_3 \cup rK_1 \cup sK_2$ or $G = P_3 \cup rK_1 \cup sK_2$ for some integers $r, s \geq 0$. ■

The proof of the next result is similar to that of Observation 9 and is therefore omitted.

Observation 10. Let G be a graph of order $n \geq 3$. Then $\gamma_R(G) = n - 1$ if and only if $G = H \cup rK_1 \cup sK_2$ for some integers $r, s \geq 0$, where $H \in \{C_3, C_4, C_5, P_3, P_4, P_5\}$.

Observation 11. Let $k \geq 3$ be an integer, and let G be a graph of order $n \geq 2$. Then $\gamma_R^k(G) = 2$ or $\gamma_R^k(\overline{G}) = 2$.

Proof. If $\text{diam}(G) \leq 3$, then it follows from Observation 2 that $\gamma_R^k(G) = 2$. If $\text{diam}(G) \geq 4$, then a result of Bondy and Murty [1] (page 14) implies that $\text{diam}(\overline{G}) \leq 2$. Applying again Observation 2, we see that $\gamma_R^k(\overline{G}) = 2$. ■

Observation 12. Let G be a graph of order $n \geq 2$. If $\text{diam}(G) \neq 3$, then $\gamma_R^2(G) = 2$ or $\gamma_R^2(\overline{G}) = 2$.

Proof. If $\text{diam}(G) \leq 2$, then it follows from Observation 2 that $\gamma_R^2(G) = 2$. If $\text{diam}(G) \geq 3$, then the assumption $\text{diam}(G) \neq 3$ implies that $\text{diam}(G) \geq 4$. As above, we deduce that $\text{diam}(\overline{G}) \leq 2$, and Observation 2 leads to $\gamma_R^2(\overline{G}) = 2$. ■

Observation 13. Let $k \geq 1$ be an integer, and let G be a graph of order $n \geq 2$. Then $\gamma_R^k(G) = 2\gamma^k(G)$ if and only if G has a $\gamma_R^k(G)$ -function $f = (V_0, V_1, V_2)$ with $|V_1| = 0$.

Proof. Assume first that $\gamma_R^k(G) = 2\gamma^k(G)$. Let S be a k -distance dominating set of G such that $|S| = \gamma^k(G)$. Then $f = (V(G) - S, \emptyset, S) = (V_0, V_1, V_2)$ is a k DRDF on G such that

$$\omega(f) = 2|S| = 2\gamma^k(G) = \gamma_R^k(G)$$

and therefore f is a $\gamma_R^k(G)$ -function with $|V_1| = 0$.

Conversely, let $f = (V_0, V_1, V_2)$ be a $\gamma_R^k(G)$ -function with $|V_1| = 0$ and thus $\gamma_R^k(G) = 2|V_2|$. Then V_2 is also k -distance dominating set of G , and hence we deduce that $2\gamma^k(G) \leq 2|V_2| = \gamma_R^k(G)$. Applying the second inequality in (1), we obtain the identity $\gamma_R^k(G) = 2\gamma^k(G)$, and the proof is complete. ■

The special case $k = 1$ of Observation 13 can be found in the article [3].

Next we will prove a Nordhaus-Gaddum inequality.

Theorem 14. Let $k \geq 2$ be an integer, and let G be a graph of order $n \geq 3$. Then

$$\gamma_R^k(G) + \gamma_R^k(\overline{G}) \leq n + 2.$$

Furthermore, equality holds in the bound if and only if G or $\overline{G}$ is isomorphic to $rK_1 \cup sK_2$ for two integers $r, s \geq 0$.

Proof. If neither G nor $\overline{G}$ is isomorphic to C_5 or to $\frac{n}{2}K_2$, then it follows from Theorem D that

$$\gamma_R^k(G) + \gamma_R^k(\overline{G}) \leq \gamma_R(G) + \gamma_R(\overline{G}) \leq n + 2.$$

If $G = C_5$ or $\overline{G} = C_5$, then $\gamma_R^k(G) + \gamma_R^k(\overline{G}) = 4 < 7 = n + 2$, and if $G = \frac{n}{2}K_2$ or $\overline{G} = \frac{n}{2}K_2$, then $\gamma_R^k(G) + \gamma_R^k(\overline{G}) = n + 2$, and the desired Nordhaus-Gaddum bound is proved.

If G or $\overline{G}$ is isomorphic to $rK_1 \cup sK_2$ for two integers $r, s \geq 0$, then obviously $\gamma_R^k(G) + \gamma_R^k(\overline{G}) = n + 2$.

Next assume that $\gamma_R^k(G) + \gamma_R^k(\overline{G}) = n + 2$. We distinguish two cases.

Case 1. Assume that $k \geq 3$. If $\text{diam}(G) \leq 3$, then $\gamma_R^k(G) = 2$ and therefore $\gamma_R^k(\overline{G}) = n$. According to Observation 7, we observe that $\overline{G} = rK_1 \cup sK_2$ for two integers $r, s \geq 0$. If $\text{diam}(G) \geq 4$, then $\text{diam}(\overline{G}) \leq 2$. It follows that $\gamma_R^k(\overline{G}) = 2$ and thus $\gamma_R^k(G) = n$. Applying again Observation 7, we see that $G = rK_1 \cup sK_2$ for two integers $r, s \geq 0$.

Case 2. Assume that $k = 2$. If $\text{diam}(G) \leq 2$, then $\gamma_R^2(G) = 2$ and therefore $\gamma_R^2(\overline{G}) = n$. According to Observation 7, we observe that $\overline{G} = rK_1 \cup sK_2$ for two integers $r, s \geq 0$. If $\text{diam}(G) \geq 4$, then $\text{diam}(\overline{G}) \leq 2$. It follows that $\gamma_R^2(\overline{G}) = 2$ and thus $\gamma_R^2(G) = n$, and so $G = rK_1 \cup sK_2$ for two integers $r, s \geq 0$. If $\text{diam}(\overline{G}) \leq 2$ or $\text{diam}(\overline{G}) \geq 4$, then we obtain analogously that G or $\overline{G}$ is isomorphic to $rK_1 \cup sK_2$ for two integers $r, s \geq 0$.

There remains the case that $\text{diam}(G) = \text{diam}(\overline{G}) = 3$. Let x and y be two vertices of G such that $d(x, y) = \text{diam}(G) = 3$. Obviously, $f = (V(G) - \{x, y\}, \emptyset, \{x, y\})$ is a 2DRDF on $\overline{G}$, since there is no vertex in G adjacent to both x and y . Therefore $\gamma_R^2(\overline{G}) \leq 4$. Analogously, we obtain $\gamma_R^2(G) \leq 4$ and hence

$$\gamma_R^2(G) + \gamma_R^2(\overline{G}) \leq 8 < n + 2$$

when $n \geq 7$.

Finally, assume that $4 \leq n \leq 6$. If $4 \leq n \leq 5$, then $\Delta_2(G) = \Delta_2(\overline{G}) = n - 1$ and hence $\gamma_R^2(G) = \gamma_R^2(\overline{G}) = 2$ and consequently

$$\gamma_R^2(G) + \gamma_R^2(\overline{G}) = 4 < n + 2.$$

If $n = 6$, then $\Delta_2(G) = \Delta_2(\overline{G}) \geq 4$ and hence $\gamma_R^2(G) \leq 3$ and $\gamma_R^2(\overline{G}) \leq 3$. It follows that

$$\gamma_R^2(G) + \gamma_R^2(\overline{G}) \leq 6 < n + 2,$$

and the proof is complete. ■

3. BOUNDS ON THE k -DISTANCE ROMAN DOMINATION NUMBER

Theorem 15. *If $k \geq 1$ is an integer and G a connected graph of order n with $n - \Delta(G) - k \geq 0$, then*

$$\gamma_R^k(G) \leq n - \Delta(G) - k + 2.$$

Proof. Let v be a vertex of G such that $\deg_G(v) = \Delta(G)$. If $d(u, v) \leq k$ for each $u \in V(G)$, then obviously $\gamma_R^k(G) = 2$ and we are done. If $d(w, v) > k$ for some

$w \in V(G)$, then choose a vertex u in G such that $d(u, v) = k + 1$. Let P be a shortest (u, v) -path. Then clearly $d(v, z) \leq k$ for each $z \in (V(P) - \{u\}) \cup N_G(v)$ and hence $f = ((V(P) - \{u, v\}) \cup N_G(v), V(G) - ((V(P) - \{u\}) \cup N_G(v)), \{v\})$ is a k DRDF on G with weight $n - \Delta(G) - k + 2$ and therefore $\gamma_R^k(G) \leq n - \Delta(G) - k + 2$. ■

The special case $k = 1$ of Theorem 15 can be found in [2].

Theorem 16. *If $k \geq 1$ is an integer and G a graph of order n with $\Delta = \Delta(G) \geq 3$, then*

$$\gamma_R^k(G) \geq \frac{2n(\Delta - 2)}{\Delta(\Delta - 1)^k - 2}.$$

Proof. Each vertex $v \in V(G)$ dominates at most Δ vertices at distance 1 from v , at most $\Delta(\Delta - 1)$ vertices at distance 2 from v , at most $\Delta(\Delta - 1)^2$ vertices of at distance 3 from v , and so on. Thus

$$\Delta_k(G) \leq \Delta + \Delta(\Delta - 1) + \Delta(\Delta - 1)^2 + \cdots + \Delta(\Delta - 1)^{k-1} = \Delta \cdot \frac{(\Delta - 1)^k - 1}{\Delta - 2}.$$

Applying Observation 3, we obtain the desired lower bound as follows

$$\gamma_R^k(G) \geq \frac{2n}{\Delta_k(G) + 1} \geq \frac{2n}{\Delta \cdot \frac{(\Delta - 1)^k - 1}{\Delta - 2} + 1} = \frac{2n(\Delta - 2)}{\Delta(\Delta - 1)^k - 2}.$$

■

In the case that $\Delta(G) = 2$, the proof of Theorem 16 leads to the next lower bound, and Proposition 19 below shows that this bound is sharp.

Theorem 17. *If $k \geq 1$ is an integer and G a graph of order n with $\Delta(G) = 2$, then*

$$\gamma_R^k(G) \geq \frac{2n}{2k + 1}.$$

Theorem 18. *If $k \geq 1$ is an integer and G a connected graph of order $n \geq 2$, then*

$$\gamma_R^k(G) \geq \left\lceil \frac{\text{diam}(G) + 2}{k + 1} \right\rceil.$$

Proof. The statement is obviously true for K_2 . Let G be a connected graph of order $n \geq 3$ and $f = (V_0^f, V_1^f, V_2^f)$ be a $\gamma_R^k(G)$ -function. Suppose that $P = v_1 v_2 \cdots v_{\text{diam}(G)+1}$ is a diametral path in G . This diametral path includes at most two edges from the induced subgraph $G[N[v]]$ for each $v \in V_1^f$ and at most $2k$ edges from the induced subgraph $G[N_{k,G}[v]]$ for each $v \in V_2^f$. Let $E' = \{v_i v_{i+1} \mid 1 \leq i \leq \text{diam}(G)\} \cap (\bigcup_{v \in V_1^f} E(G[N[v]]) \cup \bigcup_{v \in V_2^f} E(G[N_{k,G}[v]]))$.

Then the diametral path contains at most $|V_2^f| - 1$ edges not in E' , joining the neighborhoods at distance k of the vertices of V_2^f . Since G is a connected graph of order at least 3, $V_2^f \neq \emptyset$. Hence,

$$\begin{aligned} \text{diam}(G) &\leq 2k|V_2^f| + 2|V_1^f| + (|V_2^f| - 1) \\ &= (2k + 2)|V_2^f| + (k + 1)|V_1^f| - |V_2^f| - 1 - |V_1^f| \\ &\leq (k + 1)\gamma_R^k(G) - 2, \end{aligned}$$

and the result follows. $\blacksquare$

The next proposition is straightforward to verify.

Proposition 19. *For $n \geq 3$,*

$$\gamma_R^k(C_n) = \begin{cases} \frac{2n}{2k+1} & n \equiv 0 \pmod{2k+1}, \\ 2\lfloor \frac{n}{2k+1} \rfloor + 1 & n \equiv 1 \pmod{2k+1}, \\ 2\lfloor \frac{n}{2k+1} \rfloor + 2 & \text{otherwise.} \end{cases}$$

Theorem 20 [7]. *For a graph G of order n with $g(G) \geq 3$, $\gamma_R(G) \geq \left\lceil \frac{2g(G)}{3} \right\rceil$.*

Theorem 21. *If $k \geq 1$ is an integer and G a connected graph of order $n \geq 2$ and $\infty > g(G) \geq 2k + 1$, then*

$$\gamma_R^k(G) \geq \left\lceil \frac{2g(G)}{2k+1} \right\rceil.$$

Proof. By Theorem 20 we may assume that $k \geq 2$. First note that if G is an n -cycle then the result follows from Proposition 19. Now, let C be a cycle of length $g(G)$ in G . Since $k \geq 2$, $g(G) \geq 5$. Then a vertex not in $V(C)$, can dominate at most one vertex of C for otherwise we obtain a cycle of length less than $g(G)$ which is a contradiction. On the other hand, each vertex in $V(C)$ dominates at most $2k + 1$ vertex of $V(C)$. Let $f = (V_0, V_1, V_2)$ be a $\gamma_R^k(G)$ -function. Then obviously, $\gamma_R^k(G) = |V_1| + 2|V_2| \geq |V_1| + 2\frac{g(G)}{2k+1} \geq \frac{2g(G)}{2k+1}$. This leads to the desired bound, and the proof is complete. $\blacksquare$

The special case $k = 1$ of Theorems 18 and 21 can be found in [7].

4. CONNECTED GRAPHS

Let $k \geq 1$ be an integer. For n -vertex graphs, always $\gamma_R^k(G) \leq n$, with equality when $G = \overline{K_n}$. In this section we prove that $\gamma_R^k(G) \leq 4n/(2k + 3)$ when G is a connected n -vertex graph. Since deleting an edge cannot decrease $\gamma_R^k(G)$, it suffices to prove the bound for trees.

A *leaf* of a graph G is a vertex of degree 1, while a *support vertex* of G is a vertex adjacent to a leaf. For a vertex v in a rooted tree T , let $D(v)$ denote the set of descendants of v and $D[v] = D(v) \cup \{v\}$. The *maximal subtree* at v is the subtree of T induced by $D[v]$, and is denoted by T_v .

Theorem 22. *If $k \geq 1$ is an integer and T is a tree of order $n \geq k + 2$, then $\gamma_R^k(T) \leq 4n/(2k + 3)$.*

Proof. By Theorem C we may assume that $k \geq 2$. The proof is by induction on n . The base step handles trees with few vertices or diameter $2k$ and $2k + 1$. If $k + 2 \leq n \leq 2k + 1$ or $\text{diam}(T) \leq 2k$, then T has a k -distance dominating vertex, and $\gamma_R^k(T) = 2 < 4n/(2k + 3)$. If $\text{diam}(T) = 2k + 1$, then T has a k -distance dominating set of size 2, which yields $\gamma_R^k(T) \leq 4$. This is sufficiently small for trees with at least $2k + 4$ vertices. Let $P = v_1 v_2 \cdots v_{2k+2}$ be a longest path in T . For $n \in \{2k + 2, 2k + 3\}$ and $\text{diam}(T) = 2k + 1$, we may assume, without loss of generality, that $\deg(v_{2k+1}) = 2$. Then the function $f : V(G) \rightarrow \{0, 1, 2\}$ defined by $f(v_{k+1}) = 2, f(v_{2k+2}) = 1$ and $f(x) = 0$ otherwise, is a k DRDF on G and hence $\gamma_R^k(T) \leq 3$, which is small enough.

Hence we may assume that $\text{diam}(T) \geq 2k + 2$. For a subtree T' with n' vertices, where $n' \geq k + 2$, the induction hypothesis yields a k DRDF f' of T' with weight at most $\frac{4}{2k+3}n'$. We find a subtree T' such that adding a bit more weight to f' will yield a small enough k DRDF f for T . Let $P = v_1 v_2 \cdots v_r v_{r+1} \cdots v_{r+k+1}$ be a longest path in T chosen to maximize the $\sum_{j=1}^k \deg_T(v_{r+j})$ and let T be rooted in v_1 . We consider three cases.

Case 1. $\sum_{j=1}^k \deg_T(v_{r+j}) > 2k$. Let $T' = T - T_{v_{r+1}}$. Since $\text{diam}(T) \geq 2k + 2$, we have $n' \geq k + 2$. Define f on $V(T)$ by letting $f(x) = f'(x)$ except for $f(v_{r+1}) = 2$ and $f(x) = 0$ for each $x \in V(T_{v_{r+1}}) - \{v_{r+1}\}$. Note that f is a k DRDF for T and that

$$w(f) = w(f') + 2 \leq \frac{4(n - k - 2)}{2k + 3} + 2 < \frac{4n}{2k + 3}.$$

Case 2. $\sum_{j=1}^k \deg_T(v_{r+j}) = 2k$ and $\deg(v_r) = 2$. Let $T' = T - T_{v_r}$. If $n' = k + 1$, then T is a path on $2k + 3$ vertices and has a k DRDF of weight 4. Otherwise, the induction hypothesis applies. Define f on $V(T)$ by letting $f(x) = f'(x)$ except for $f(v_{r+1}) = 2$ and $f(v_r) = f(v_{r+2}) = \cdots = f(v_{r+k+1}) = 0$. Again f is a k DRDF, and the computation $w(f) < \frac{4n}{2k+3}$ is the same as in Case 1.

Case 3. $\sum_{j=1}^k \deg_T(v_{r+j}) = 2k$ and $\deg(v_r) > 2$. Consider two subcases.

Subcase 3.1. $d(v_r, u) \leq k + 1$ for each $u \in V(T)$. Let $S = \{u_1, \dots, u_t\}$ be the set of vertices in distance $k + 1$ from v_r . Obviously $v_1, v_{r+k+1} \in S$ and so $t \geq 2$.

On the other hand, it is clear that $n \geq t(k+1) + 1$. Define $f : V(G) \rightarrow \{0, 1, 2\}$ by $f(v_r) = 2, f(u_1) = \dots = f(u_t) = 1$ and $f(x) = 0$ otherwise. Clearly f is a k DRDF on G and hence

$$\gamma_R^k(T) \leq t + 2 \leq \frac{4(t(k+1) + 1)}{2k+3} \leq \frac{4n}{2k+3},$$

with equality if and only if $t = 2$ and $n = 2(k+1) + 1$, and this if and only if $T = P_{2k+3}$.

Subcase 3.2. $d(v_r, u) \geq k+2$ for some $u \in V(T)$. It is clear that $d(v_1, v_r) \geq k+2$. By the choice of P , $d(v_r, w) \leq k+1$ for each $w \in V(T_{v_r})$. Let T_1 and T_2 be the connected components of $T - v_{r-1}v_r$. Let $f_i = (V_0^{f_i}, V_1^{f_i}, V_2^{f_i})$ be a $\gamma_R^k(T_i)$ -function for $i = 1, 2$. Obviously $f = (V_0^{f_1} \cup V_0^{f_2}, V_1^{f_1} \cup V_1^{f_2}, V_2^{f_1} \cup V_2^{f_2})$ is a k DRDF of T . By induction hypothesis we obtain

$$(2) \quad \gamma_R^k(T) \leq \gamma_R^k(T_1) + \gamma_R^k(T_2) = \omega(f_1) + \omega(f_2) \leq \frac{4|V(T_1)|}{2k+3} + \frac{4|V(T_2)|}{2k+3} = \frac{4n}{2k+3}.$$

This completes the proof. ■

Let k be a positive integer and let $F_{m,k}$ consist of the disjoint union of m copies of P_{2k+3} plus a path through the central vertices of these copies, as illustrated in Figure 1 for $k = 2$. If $v_1v_2 \dots v_{2k+3}$ is an induced path in a graph, then a k DRDF must put total weight at least 4 on $\{v_1, v_2, \dots, v_{2k+3}\}$. In F_m , there are m disjoint induced paths on $2k+3$ vertices, so $\gamma_R^k(T) \geq 4|V(T)|/(2k+3)$ for each $T \in F_m$. Such copies of P_{2k+3} can be assembled in many ways, and this allows us to characterize the trees achieving equality in Theorem 22.




Figure 1. A member of $F_{5,2}$.

Theorem 23. *If $k \geq 1$ is an integer and T is an n -vertex tree, then $\gamma_R^k(T) = 4n/(2k+3)$ if and only if $V(T)$ can be partitioned into sets inducing P_{2k+3} such that the subgraph induced by the central vertices of these paths is connected.*

Proof. We have observed that if an induced subgraph H of G is isomorphic to P_{2k+3} , and its noncentral vertices have no neighbors outside H in G , then every k DRDF of G puts weight at least 4 on $V(H)$. Thus in any tree with such a vertex partition, weight at least 4 is needed on every set in the partition.

To show that equality requires this structure, we examine the proof of Theorem 22 more closely. The proof is by induction on n . In the base cases and Cases 1 and 2, we produce a k DRDF with weight less than $4n/(2k+3)$. In Case 3 and Subcase 3.1 with diameter $2k+2$, equality requires $T = P_{2k+3}$.

Define T_1, T_2 as in the inductive part of Case 3. The bound holds with equality only if $\gamma_R^k(T_1) = \frac{4|V(T_1)|}{2k+3}$ and $\gamma_R^k(T_2) = \frac{4|V(T_2)|}{2k+3}$. It follows from $\gamma_R^k(T_2) = \frac{4|V(T_2)|}{2k+3}$ and the first paragraph in the proof of Theorem 22 that $\text{diam}(T_2) = 2k+2$. Since $d(v_r, u) \leq k+1$ for each $u \in T_2$, from the proof of Subcase 3.1 we deduce that $T_2 = P_{2k+3}$ with central vertex v_r . By induction hypothesis, $V(T)$ can be partitioned into sets inducing P_{2k+3} such that the subgraph induced by the central vertices of these paths is connected. Suppose $\{u_1, \dots, u_{2k+3}\}$ is the partition set inducing $P_{2k+3} = u_1 u_2 \cdots u_{2k+3}$ containing v_{r-1} . We claim that $v_{r-1} = u_{k+2}$. Otherwise, we may assume, without loss of generality that, $v_{r-1} \in \{u_{k+3}, \dots, u_{2k+3}\}$. Define $f : V(G) \rightarrow \{0, 1, 2\}$ by $f(v_r) = f(u_{k+1}) = 2, f(u_{2k+3}) = 1$ and let f assign 2 to all other central vertices and 1 to all other leaves. It is easy to see that f is a k DRDF of T with weight less than $4n/(2k+3)$ which is a contradiction. Thus v_{r-1} is the central vertex of the path $P_{2k+3} = u_1 u_2 \cdots u_{2k+3}$ and the proof is complete. ■

Theorem 24. *If k is a positive integer and G is a connected n -vertex graph with $n \geq k+2$, then*

$$\gamma_R^k(G) \leq 4n/(2k+3).$$

Moreover, the equality holds if and only if G is C_{2k+3} or obtained from $\frac{n}{2k+3} P_{2k+3}$ by adding a connected subgraph on the set of centers of the components of $\frac{n}{2k+3} P_{2k+3}$.

Proof. If G has the specified form, then as remarked earlier every k DRDF puts weight at least 4 on the vertex set of each copy of P_{2k+3} .

Now suppose that $\gamma_R^k(G) = \frac{4n}{2k+3}$. Since adding edges cannot increase $\gamma_R^k(G)$, every spanning tree of G has the form specified in Theorem 22. Given a spanning tree T , let $S_1, \dots, S_k$ be the $(2k+3)$ -sets in the special partition of $V(T)$. The assignment of weight 4 that guards S_i can be chosen independently of any other S_j . If any edge of G joins vertices of S_i and S_j that are not the centers of the paths they induce, then a k DRDF with weight less than $\frac{4n}{2k+3}$ can be built as in the proof of Theorem 23. ■

The special case $k = 1$ of Theorems 22, 23 and 24 can be found in [2]. As an application of Theorem 24, we prove the next result.

Corollary 25. *Let $f = (V_0, V_1, V_2)$ be any $\gamma_R^k(G)$ -function of a connected graph G of order $n \geq 3$. Then*

- (1) $1 \leq |V_2| \leq \frac{2n}{2k+3}$ and a graph G admits a $\gamma_R^k(G)$ -function such that $|V_2| = \frac{2n}{2k+3}$ if and only if G is C_{2k+3} or is obtained from $\frac{n}{2k+3}P_{2k+3}$ by adding a connected subgraph on the set of centers of the components of $\frac{n}{2k+3}P_{2k+3}$.
- (2) $0 \leq |V_1| \leq \frac{4n}{2k+3} - 2$.
- (3) $n - \frac{4n}{2k+3} + 1 \leq |V_0| \leq n - 1$.

Proof. (1) If $V_2 = \emptyset$, then $V_1 = V(G)$ and $V_0 = \emptyset$. The RDF $(\emptyset, V(G), \emptyset)$ is not minimum since $|V_1| + 2|V_2| > \frac{4n}{2k+3}$. Hence $|V_2| \geq 1$. On the other hand, $|V_2| \leq \frac{2n}{2k+3} - |V_1|/2 \leq \frac{2n}{2k+3}$.

Let G admit a $\gamma_R^k(G)$ -function $f = (V_0, V_1, V_2)$ such that $|V_2| = \frac{2n}{2k+3}$. Then by Theorem 24, $\frac{4n}{2k+3} \leq |V_1| + 2|V_2| = \gamma_R^k(G) \leq \frac{4n}{2k+3}$ and the result follows from Theorem 24 again.

Conversely, let G be C_{2k+3} or obtained from $\frac{n}{2k+3}P_{2k+3}$ by adding a connected subgraph on the set of centers of the components of $\frac{n}{2k+3}P_{2k+3}$. If $G = C_{2k+3}$, then assign 2 to two vertices at distance 2 and 0 to the other vertices. If G is obtained from $\frac{n}{2k+3}P_{2k+3}$ by adding a connected subgraph on the set of centers of the components of $\frac{n}{2k+3}P_{2k+3}$, then assign 2 to the neighbors of centers of the components P_{2k+3} and 0 to the other vertices. Obviously f is a $\gamma_R^k(G)$ -function with the desired property.

- (2) Since $|V_2| \geq 1$, $|V_1| \leq \frac{4n}{2k+3} - 2|V_2| \leq \frac{4n}{2k+3} - 2$.

(3) The upper bound comes from $|V_0| \leq n - |V_2| \leq n - 1$. For the lower bound, adding side by side $2|V_0| + 2|V_1| + 2|V_2| = 2n$, $-|V_1| - 2|V_2| \geq \frac{-4n}{2k+3}$ and $|V_1| \leq \frac{4n}{2k+3} - 2$ gives $2|V_0| \geq 2n - \frac{8n}{2k+3} + 2$. Therefore $|V_0| \geq n - \frac{4n}{2k+3} + 1$. ■

The special case $k = 1$ of Corollary 25 can be found in [5].

REFERENCES

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications (The Macmillan Press Ltd. London and Basingstoke, 1976).
- [2] E.W. Chambers, B. Kinnersley, N. Prince and D.B. West, *Extremal problems for Roman domination*, SIAM J. Discrete Math. **23** (2009) 1575–1586. doi:10.1137/070699688
- [3] E.J. Cockayne, P.M. Dreyer Jr., S.M. Hedetniemi and S.T. Hedetniemi, *On Roman domination in graphs*, Discrete Math. **278** (2004) 11–22. doi:10.1016/j.disc.2003.06.004
- [4] E.J. Cockayne, P.J.P. Grobler, W.R. Gründlingh, J. Munganga, and J.H. van Vuuren, *Protection of a graph*, Util. Math. **67** (2005) 19–32.
- [5] O. Favaron, H. Karami and S.M. Sheikholeslami, *On the Roman domination number in graphs*, Discrete Math. **309** (2009) 3447–3451. doi:10.1016/j.disc.2008.09.043

- [6] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, Inc. NewYork, 1998).
- [7] B.P. Mobaraky and S.M. Sheikholeslami, *Bounds on Roman domination numbers of a graph*, Mat. Vesnik **60** (2008) 247–253.
- [8] C.S. ReVelle and K.E. Rosing, *Defendens imperium romanum: a classical problem in military strategy*, Amer. Math. Monthly **107** (2000) 585–594.
doi:10.2307/2589113
- [9] I. Stewart, *Defend the Roman Empire*, Sci. Amer. **281** (1999) 136–139.
- [10] D.B. West, *Introduction to Graph Theory* (Prentice-Hall, Inc, 2000).

Received 20 December 2011

Revised 4 September 2012

Accepted 5 September 2012