

## ALMOST-RAINBOW EDGE-COLORINGS OF SOME SMALL SUBGRAPHS

ELLIOT KROP

*Department of Mathematics, Clayton State University*  
*2000 Clayton State Boulevard, Morrow, GA 30260 USA*

**e-mail:** ElliotKrop@clayton.edu

AND

IRINA KROP

*DePaul University*  
*1 E. Jackson, Chicago, IL 60604 USA*

**e-mail:** irina.krop@gmail.com

### Abstract

Let  $f(n, p, q)$  be the minimum number of colors necessary to color the edges of  $K_n$  so that every  $K_p$  is at least  $q$ -colored. We improve current bounds on these nearly “anti-Ramsey” numbers, first studied by Erdős and Gyárfás. We show that  $f(n, 5, 9) \geq \frac{7}{4}n - 3$ , slightly improving the bound of Axenovich. We make small improvements on bounds of Erdős and Gyárfás by showing  $\frac{5}{6}n + 1 \leq f(n, 4, 5)$  and for all even  $n \not\equiv 1 \pmod{3}$ ,  $f(n, 4, 5) \leq n - 1$ . For a complete bipartite graph  $G = K_{n,n}$ , we show an  $n$ -color construction to color the edges of  $G$  so that every  $C_4 \subseteq G$  is colored by at least three colors. This improves the best known upper bound of Axenovich, Füredi, and Mubayi.

**Keywords:** Ramsey theory, generalized Ramsey theory, rainbow-coloring, edge-coloring, Erdős problem.

**2010 Mathematics Subject Classification:** 05A15, 05C38, 05C55.

## 1. INTRODUCTION

### 1.1. Definitions

For basic graph theoretic notation and definition see Diestel [3]. All graphs  $G$  are undirected with the vertex set  $V$  and edge set  $E$ . We use  $|G|$  for  $|V|$  and  $\|G\|$  for

$|E|$ .  $K_n$  denotes the complete graph on  $n$  vertices and  $K_{n,m}$  the bipartite graph with  $n$  vertices and  $m$  vertices in the first and second part, respectively. For any edge  $(u, v)$ , let  $\mathcal{C}(u, v)$  be the color on that edge, and for any vertex  $v$ , let  $\mathcal{C}(v)$  be the set of colors on the edges incident to  $v$ . We say that an edge-coloring is *proper* if every pair of incident edges are of different colors. If vertices  $u, v$  are adjacent, we write  $u \sim v$ .

## 1.2. Coloring edges

Given a graph  $G$  of order  $n$  and integers  $p, q$  so that  $2 \leq p \leq n$  and  $1 \leq q \leq \binom{p}{2}$ , call an edge-coloring  $(p, q)$  if every  $K_p \subseteq K_n$  receives at least  $q$  colors on its edges. Let  $f(n, p, q)$  be the minimum colors in a  $(p, q)$  coloring of  $K_n$ . This generalization of classical Ramsey functions was first mentioned by Erdős in [4] and later studied by Erdős and Gyárfás in [5]. Further, define  $\phi(n, p, q)$  to be the minimum colors in a proper  $(p, q)$  coloring of  $K_n$ .

Extending the definition, for any graph  $G$ , call an edge coloring  $(H, q)$  if every subgraph  $H \subseteq G$  receives at least  $q$  colors on its edges. Let  $f(G, H, q)$  be the minimum colors in an  $(H, q)$  coloring of the edges of  $G$ . We say that a coloring of  $H$  is *almost-rainbow* if  $q = \|H\| - 1$ , that is, one color is repeated once.

For an extended survey regarding bounds on rainbow colorings, see [7].

Using the Local Lemma, the authors in [5] were able to produce bounds for  $f(n, p, q)$ , with several difficult cases unresolved. Among those were  $f(n, 4, 3)$ ,  $f(n, 4, 4)$ ,  $f(n, 4, 5)$ , and  $f(n, 5, 9)$ . In these cases they showed that  $f(n, 4, 3) \leq c\sqrt{n}$ ,  $c\sqrt{n} \leq f(n, 4, 4) \leq cn^{\frac{2}{3}}$ ,  $\frac{5n-1}{6} \leq f(n, 4, 5) \leq n$ , and  $\frac{4}{3}n \leq f(n, 5, 9) \leq cn^{\frac{3}{2}}$ . The authors further mentioned that in this branch of generalized Ramsey theory, finding the orders of magnitude of  $f(n, 4, 4)$  and  $f(n, 5, 9)$  are “the most interesting open problems, at least to show that the latter is non-linear”. The authors then stated the linearity of said function as Problem 1.

As for  $f(n, 4, 5)$ , the authors showed that  $\frac{5(n-1)}{6} \leq f(n, 4, 5)$  with an upper bound of  $n$  for odd  $n$  and  $n - 1$  for even  $n$  if  $n - 1$  is prime.

In [9], Mubayi showed that

$$f(n, 4, 3) \leq e^{O(\sqrt{\log n})}$$

and in [8] Kostochka and Mubayi showed that for some constant  $c$ ,

$$f(n, 4, 3) \geq \frac{c \log n}{\log \log \log n}.$$

Fox and Sudakov in [6], further improved the lower bound to  $\frac{\log n}{4000}$ .

As for the other case, in [1], Axenovich showed that for some constant  $c$ ,

$$\frac{1 + \sqrt{5}}{2}n \leq f(n, 5, 9) \leq 2n^{1 + \frac{c}{\sqrt{\log n}}}.$$

In that same paper, she remarked that Tóth had communicated to her that the lower bound can be improved to  $2n - 6$ , however, the result has remained unpublished for over ten years.

In Section 2, we show

$$f(n, 5, 9) \geq \frac{7}{4}n - 3.$$

In Section 3, we make minimal improvements in the work of [5], showing  $\frac{5}{6}(n - 1) + 1 \leq f(n, 4, 5) \leq n - 1$  for even  $n$  not congruent to one mod three.

In [2], the authors showed that  $f(K_{n,n}, C_4, 3) \geq \frac{2}{3}n$ ,  $f(K_{n,n}, C_4, 3) \leq n$  for odd  $n \geq 5$ , and  $f(K_{n,n}, C_4, 3) \leq n + 1$  for even  $n \geq 5$ .

In Section 4, we show

$$f(K_{n,n}, C_4, 3) \leq n, \text{ for all } n \geq 3.$$

We believe that this upper-bound is the best possible.

## 2. ALMOST-RAINBOW FIVE-CLIQUES

### 2.1. The main tool

Let  $f(G)$  be the minimum number of colors needed to color the edges of  $G$  so that every path or cycle with four edges is at least three-colored.

Let  $\phi(G)$  be defined as  $f(G)$  above, except replace “color” by “properly color”. By arguments from [1] it is easy to see that  $f(n, 5, 9) \leq \phi(n, 5, 9) = \phi(K_n)$ .

**Lemma 1.**  $\phi(K_{2,n}) = \lceil \frac{3}{2}n \rceil$ .

**Proof.** Suppose the edges of  $G = K_{2,n}$  are properly colored so that every path of length four receives at least three colors. Call the vertices in the first part of  $G$ ,  $u$  and  $v$ . Choose a color  $a \in \mathcal{C}(u) \cap \mathcal{C}(v)$  so that for some vertices  $x, y$  in the second part of  $G$ ,  $a = \mathcal{C}(u, x) = \mathcal{C}(v, y)$ . Note that there exist colors  $b, c$  so that  $b = \mathcal{C}(u, y)$ ,  $c = \mathcal{C}(v, x)$ , and  $b, c \in (\mathcal{C}(u) \cup \mathcal{C}(v)) \setminus (\mathcal{C}(u) \cap \mathcal{C}(v))$ . Since there are two colors for every one in  $\mathcal{C}(u) \cap \mathcal{C}(v)$ , we can say that

$$(1) \quad |\mathcal{C}(u) \cap \mathcal{C}(v)| \leq \left\lfloor \frac{1}{2} |(\mathcal{C}(u) \cup \mathcal{C}(v)) \setminus (\mathcal{C}(u) \cap \mathcal{C}(v))| \right\rfloor.$$

Applying this inequality to the principle of inclusion-exclusion, we write

$$|\mathcal{C}(u) \cup \mathcal{C}(v)| = |\mathcal{C}(u)| + |\mathcal{C}(v)| - |\mathcal{C}(u) \cap \mathcal{C}(v)| \geq 2n - \frac{1}{3} |\mathcal{C}(u) \cup \mathcal{C}(v)|.$$

Solving for the union we get

$$(2) \quad |\mathcal{C}(u) \cup \mathcal{C}(v)| \geq \frac{3}{2}n.$$

For the upper bound, we construct an edge-coloring of  $G = K_{2,n}$  with  $\lceil \frac{3}{2}n \rceil$  colors. Label the vertices of the first part of  $G$ ,  $u, v$  and the second part  $\{v_1, v_2, \dots, v_n\}$ . Let  $r = \lceil \frac{n}{2} \rceil$ . Color the edges  $(v_1, u), (v_2, u), \dots, (v_r, u)$  by the colors  $1, \dots, r$ . If  $n$  is even, color the edges  $(v_n, v), (v_{n-1}, v), \dots, (v_{n-r+1}, v)$  from the set of colors  $\{1, \dots, r\}$ . If  $n$  is odd, color the edges  $(v_n, v), (v_{n-1}, v), \dots, (v_{n-r+2}, v)$  by some of the colors from the set  $\{1, \dots, r\}$ . Color the remaining edges distinctly by all the colors not previously used. Let  $i$  and  $j$  be such that  $\mathcal{C}(u, v_i) = \mathcal{C}(v, v_j)$ . Notice that for any  $k \in \{1, \dots, n\}$ ,  $\{\mathcal{C}(u, v_i), \mathcal{C}(u, v_j), \mathcal{C}(v, v_i), \mathcal{C}(v, v_k)\}$  are pairwise distinct. Hence every 4-path receives at least three colors. ■

## 2.2. A small improvement

**Theorem 2.**  $f(n, 5, 9) \geq \frac{7}{4}n - 3$ .

**Proof.** Consider a  $(5, 9)$  edge-coloring of  $G = K_n$  using  $s$  colors. Using the argument of Axenovich [1], we first assume that the coloring is not proper, so there exist incident edges  $(v_1, v_2)$  and  $(v_1, v_3)$  of the same color. For the coloring to remain  $(5, 9)$ , all edges of  $G \setminus \{(v_1, v_2), (v_1, v_3)\}$  incident to  $\{v_1, v_2, v_3\}$  must be of different colors and not  $\mathcal{C}(v_1, v_2)$  or  $\mathcal{C}(v_2, v_3)$ . Therefore,  $s \geq 3n - 7 \geq \frac{7}{4}n - 3$  for  $n \geq 5$ .

Next we assume the coloring is proper. By the pigeonhole principle there exists a color, call it  $a$ , used on at least  $\frac{\binom{n}{2}}{s}$  edges. Let  $A$  be the set of vertices adjacent to edges colored  $a$  and choose vertices  $u, v \in A$  so that  $c(u, v) = a$ .

We say that an edge is *in*  $A$  if both vertices adjacent to that edge are in  $A$ . Notice that the number of colors on the edges in  $A$  adjacent to  $u \geq 2\frac{\binom{n}{2}}{s} - 1$ , the same for  $v$ , and  $c(u, v)$  is counted both times. Let  $H$  be the complete bipartite graph with vertices  $\{u, v\}$  in the first part and the vertices of  $G \setminus A$  in the second part. Let the edge coloring of  $H$  be induced by the edge coloring of  $G$ . For any  $x \in A$  and  $y \in G$ ,  $\mathcal{C}(u, x) \neq \mathcal{C}(v, y)$ , else we produce a two-colored four-edge path. The same reasoning holds for  $y \in A$  and  $x \in G$ . This implies that the colors on the edges of  $H$  are distinct from the colors previously counted. Hence we apply Lemma 1 to  $H$  to obtain

$$(3) \quad s \geq 2\frac{\binom{n}{2}}{s} - 1 + 2\frac{\binom{n}{2}}{s} - 1 - 1 + \frac{3}{2} \left( n - 2\frac{\binom{n}{2}}{s} \right).$$


Solving for  $s$  we obtain the result. ■

## 3. ALMOST-RAINBOW FOUR-CLIQUE

We obtain a marginal improvement for the lower bound on  $f(n, 4, 5)$  and extend the even case of the upper bound from [5] to all complete graphs with orders not congruent to one modulo three.

**Theorem 3.** (i)  $\frac{5}{6}(n-1) + 1 \leq f(n, 4, 5)$ .  
(ii)  $f(n, 4, 5) \leq n-1$  for even  $n \not\equiv 1 \pmod{3}$ .

**Proof.** Given a  $(4, 5)$  coloring of the edges of  $G = K_n$ , for a fixed vertex  $u$ , let  $P_u$  denote the set of edges incident to  $u$ , whose colors are repeated on other edges incident to  $u$ . Let  $S_u$  denote the set of edges with non-repeated colors, incident to  $u$ . Let  $T_u$  denote the set of edges incident to edges from  $P_u$  of the same color.



Notice that

1.  $\mathcal{C}(P_u) \cap \mathcal{C}(S_u) = \emptyset$  by definition.
2.  $\mathcal{C}(P_u) \cap \mathcal{C}(T_u) = \emptyset$  else we obtain an induced four-colored  $K_4$  on the edges  $p \in P_u$  and  $t \in T_u$  that share the same color and the edges  $p_1, p_2 \in P_u$  that share the same color and are incident to  $t$  ( $p$  may be equal to  $p_1$ , depending on the coloring).
3.  $\mathcal{C}(S_u) \cap \mathcal{C}(T_u) = \emptyset$  else we obtain an induced four colored  $K_4$  on the the edge  $s \in S_u$  and  $t \in T_u$  of the same color and the two edges of  $P_u$  with the same color, which are incident to  $t$ .
4. For any vertex  $v$  distinct from  $u$ , if  $(u, v) \in P_u$  so that  $\mathcal{C}(u, v) = \mathcal{C}(u, w)$  for some  $w$ , then  $(u, v) \notin P_v$  and  $(v, w) \notin P_v$ .
5. For any vertex  $v$  distinct from  $u$ ,  $T_u \cap T_v = \emptyset$ .

Notice that

$$2 \sum_{u \in V(G)} |T_u| = \sum_{u \in V(G)} |P_u|,$$

so that

$$\sum_u |T_u| + \sum_u |P_u| = 3 \sum_u |T_u| = 3 \frac{1}{n} \sum_u |T_u| \times n \leq \binom{n}{2}$$

by the above claim 5, and we obtain

$$\frac{1}{n} \sum_u |T_u| \leq \frac{n-1}{6}.$$

By the pigeonhole principle, choose a vertex  $u$  so that  $|T_u| \leq \frac{n-1}{6}$ . Notice that  $n-1 = \deg u = |S_u| + |P_u| \leq |S_u| + \frac{n-1}{3}$ , so that

$$|S_u| \geq \frac{2}{3}(n-1).$$

Summing up the colors of edges incident to  $u$  we get

$$|\mathcal{C}(u)| = |S_u| + \frac{1}{2}|P_u| \geq \frac{2}{3}(n-1) + \frac{1}{6}(n-1) = \frac{5}{6}(n-1).$$

However,  $\mathcal{C}(T_u)$  must be nonempty and distinct from the colors counted above, hence

$$|\mathcal{C}(u)| \geq \frac{5}{6}(n-1) + 1.$$

For the upper bound we color the edges of  $K_n$  by a classical proper coloring (see [10] for example) and show that such a coloring is  $(4, 5)$ .

For odd  $n$ , we  $n$ -color the edges of  $K_n$  by drawing the vertices in the form of a regular  $n$ -gon and coloring the consecutive edges around the boundary in order with colors 1 to  $n$ . Next we color every edge parallel to a boundary edge by the same color as that boundary edge. Call the resulting labeled graph  $G_n$ . Notice that every  $K_4 \subseteq G_n$  with a pair of parallel edges is a non-rectangular trapezoid. Hence the coloring is  $(4, 5)$ .

For even  $n$ , choose a  $K_{n-1}$  subgraph and color it as above, obtaining  $G_{n-1}$ . Next construct the graph  $w \times G_{n-1}$ , joining the above graph to a vertex  $w$ . Since for any vertex  $u$  of  $G_{n-1}$ , there are only  $n-2$  incident edges, some color is missing. Apply this color to the edge  $(u, w)$  and continue likewise for all vertices of  $G_{n-1}$ . Call the resulting labeled graph  $G_n^*$ .

For vertices  $x, y, z \in G_n^*$  with so that  $(x, y)$  and  $(y, z)$  are boundary edges, we say that  $y$  is *opposite* an edge  $e$  if the line bisecting angle  $uvw$  is the perpendicular bisector of  $e$ . Notice that the edges opposite to  $y$  share the same color, which is not used on any edge incident to  $y$ . By the above observation,  $G_{n-1} \subseteq G_n^*$  is  $(4, 5)$ -colored, hence it is enough to show that for  $w$  as chosen above in the definition of  $G_n^*$  and any other distinct vertices  $x, y, z$  of  $G_n^*$ , the induced subgraph receives

at most one repeated color. Choose any vertex  $v \in G_n^*$ . For  $i = 1, \dots, n-2$  label the vertices with counterclockwise distance  $i$  from  $v$ ,  $u_i$ , where arithmetic of label indices is performed modulo  $n-1$ . Notice that the only edges that share the color  $\mathcal{C}(w, v)$  are  $(u_1, u_{-1}), (u_2, u_{-2}), \dots, (u_{n-2}, u_{-(n-2)})$ . For  $i = 1, \dots, \frac{n-2}{2}$ , if  $\mathcal{C}(u_i, w) = \mathcal{C}(u_{-i}, v)$ , then for any edge  $e$  opposite  $u_i$ ,  $\mathcal{C}(e) = \mathcal{C}(u_{-i}, v)$ . However, this means that

$$\begin{aligned} \mathcal{C}(u_{i-1}, u_{i+1}) = \mathcal{C}(e) = \mathcal{C}(u_{-i}, v) &\Leftrightarrow vu_{2k} = vu_{-k} \\ &\Leftrightarrow 3k \equiv 0 \pmod{(n-1)} \Leftrightarrow n \equiv 1 \pmod{3}. \end{aligned} \quad \blacksquare$$

#### 4. ALMOST-RAINBOW FOUR-CYCLES

We show the improved upper bound for the bipartite problem, when the two parts of  $G$  are of equal size.

**Theorem 4.**

$$f(K_{n,n}, G_4, 3) \leq n, \text{ for all } n \geq 3.$$

##### 4.1. The coloring

We will explore the matrix

$$G = \begin{pmatrix} 1 & 2 & 3 & \dots & r & \dots & c+1 & \dots & n \\ 3 & 1 & 2 & \dots & r-1 & \dots & c & \dots & n-1 \\ v_3 & n-1 & 1 & \dots & r-2 & \dots & c-1 & \dots & n-2 \\ \vdots & \vdots & \dots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ v_{n+1-r} & r+1 & r+2 & \dots & 1 & \dots & r+c & \dots & r \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ v_{n-1} & 3 & 4 & \dots & r+1 & \dots & c+2 & \dots & 2 \\ n-2 & u_2 & u_3 & \dots & u_r & \dots & u_{c+1} & \dots & 1 \end{pmatrix}.$$

The values of  $v_i$  and  $u_i$  will be defined shortly.

Let permutation  $\sigma$  be the  $n-1$  cycle  $(1 \ 2 \ \dots \ n-1)$ . That is,  $\sigma$  sends  $i$  to  $i+1 \pmod{n-1}$ . For a natural number  $m$  we shall write  $m \pmod{n-1}$  for its representative in  $\{1, 2, \dots, n-1\}$ . For each  $r$  we defined  $\sigma^{(r)}$  by the rule  $\sigma^{(r)}(c) \equiv r+c \pmod{n-1}$ . Let us start with the matrix

$$C = \begin{pmatrix} 2 & 3 & \dots & c+1 & \dots & n \\ \sigma^0(1) & \sigma^0(2) & \dots & \sigma^0(c) & \dots & \sigma^0(n-1) \\ \sigma^{n-2}(1) & \sigma^{n-2}(2) & \dots & \sigma^{n-2}(c) & \dots & \sigma^{n-2}(n-1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \sigma^r(1) & \sigma^r(2) & \dots & \sigma^r(c) & \dots & \sigma^r(n-1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \sigma^2(1) & \sigma^2(2) & \dots & \sigma^2(c) & \dots & \sigma^2(n-1) \end{pmatrix}.$$

We define the matrix  $G$  by adding the first column  $V = \{v_1, \dots, v_{n-1}, vu\}$  and the last row  $U = \{vu, u_2, \dots, u_n\}$  to the matrix  $C$ .

$$G = \begin{pmatrix} v_1 & 2 & 3 & \dots & c+1 & \dots & n \\ v_2 & \sigma^0(1) & \sigma^0(2) & \dots & \sigma^0(c) & \dots & \sigma^0(n-1) \\ v_3 & \sigma^{n-2}(1) & \sigma^{n-2}(2) & \dots & \sigma^{n-2}(c) & \dots & \sigma^{n-2}(n-1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ v_{n+1-r} & \sigma^r(1) & \sigma^r(2) & \dots & \sigma^r(c) & \dots & \sigma^r(n-1) \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ v_{n-1} & \sigma^2(1) & \sigma^2(2) & \dots & \sigma^2(c) & \dots & \sigma^2(n-1) \\ vu & u_2 & u_3 & \dots & u_{c+1} & \dots & u_n \end{pmatrix}.$$

The entries of  $G$  will be defined as follows: for every 4-tuple  $(i, j; l, m)$  with  $1 \leq i < j \leq n$  and  $1 \leq l < m \leq n$  the  $(2 \times 2)$  matrix

$$G(i, j; l, m) = \begin{pmatrix} a_{il} & a_{im} \\ a_{jl} & a_{jm} \end{pmatrix}.$$

We consider the colorings for the edges  $V$  and  $U$  in three types of even  $n \pmod{6}$ .

**Type 1:** Matrix  $G_1 = G$  for  $n \equiv 2 \pmod{6}$ ;  $[n = 2 + 6k, k \geq 1]$

$$a_{i,1} = \begin{cases} 1, & i = 1, \\ 3, & i = 2, \\ n, & 3 \leq i \leq \frac{n}{2} + 1, \\ 2(i-1) - n, & \frac{n}{2} + 2 \leq i \leq n-1, \\ n-2, & i = n. \end{cases}$$

$$a_{n,l} = \begin{cases} n-2l, & 1 \leq l \leq \frac{n}{2} - 1, \\ n, & \frac{n}{2} \leq l \leq n-2, \\ n-1, & l = n-1, \\ 1, & l = n. \end{cases}$$

**Type 2:** Matrix  $G_2 = G$  for  $n \equiv 6 \pmod{6}$ ;  $[n = 6 + 6k, k \geq 1]$



We define  $Y$  as  $\frac{n}{2} - 2$  for even  $k$ , and as  $\frac{n}{2} + 1$  for odd  $k$ .

$$a_{i,1} = \begin{cases} 1, & i = 1, \\ 3, & i = 2, \\ n, & 3 \leq i \leq \frac{n}{2} + 1, \\ Y, & i = \frac{n}{2} + 2, \\ 2(i-2) - n, & \frac{n}{2} + 3 \leq i \leq n-1, \\ n-2, & i = n. \end{cases}$$

$$a_{n,l} = \begin{cases} n-2, & l = 1, \\ n-2(l+1), & 2 \leq l \leq \frac{n}{2} - 2, \\ Y, & l = \frac{n}{2} - 1, \\ n, & \frac{n}{2} \leq l \leq n-2, \\ n-1, & l = n-1, \\ 1, & l = n. \end{cases}$$

Exception for  $n = 6$ ;  $[k = 0]$  the first row  $V = \{1, 5, 6, 6, 4\}$ , the last column  $U = \{3, 6, 6, 6, 5, 1\}$ .

**Type 3:** Matrix  $G_3 = G$  for  $n \equiv 4 \pmod{6}$ ;  $[n = 4 + 6k, k \geq 4]$

The regularity starts with  $n > 22$ .

$$a_{i,1} = \begin{cases} 1, & i = 1, \\ 3, & i = 2, \\ n, & 3 \leq i \leq \frac{n}{2} + 1, \\ n-9, & i = \frac{n}{2} + 2, \\ 2(i-2) - n, & \frac{n}{2} + 3 \leq i \leq \frac{5n+4}{6}, \\ 2(i-1) - n, & \frac{5n+10}{6} \leq i \leq n-1, \\ n-2, & i = n. \end{cases}$$

$$a_{n,l} = \begin{cases} n-2l, & 1 \leq l \leq \frac{n-4}{6}, \\ n-2(l+1), & \frac{n+2}{6} \leq l \leq \frac{n}{2} - 2, \\ n-9, & l = \frac{n}{2} - 1, \\ n, & \frac{n}{2} \leq l \leq n-2, \\ n-1, & l = n-1, \\ 1, & l = n. \end{cases}$$

**Exceptions:**

For  $n = 10$  we replace  $(n-9)$  with  $(n-8)$ .

For  $n = 16$  we replace  $(n-9)$  with  $(n-11)$ .

For  $n = 22$  we replace  $(n-9)$  with  $(n-5)$  and the definitions:

$$a_{i,1} = \begin{cases} 2(i-2) - n, & \frac{n}{2} + 3 \leq i \leq \frac{5n-2}{6}, \\ 2(i-1) - n, & \frac{5n+4}{6} \leq i \leq n-1. \end{cases}$$

$$a_{n,l} = \begin{cases} n-2l, & 1 \leq l \leq \frac{n-10}{6}, \\ n-2(l+1), & \frac{n-4}{6} \leq l \leq \frac{n}{2} - 2. \end{cases}$$

#### 4.2. Sketch of proof

First, we show that every 4-cycle defined in the *basic coloring* (matrix entries  $a_{ij}$  where  $1 < i \leq n, 1 \leq j < n$ ) is *almost rainbow*. That is, given  $i < j$  and  $l < m$  we show that  $a_{i,l}, a_{j,l}, a_{i,m}, a_{j,m}$  contains at least three distinct elements in the *basic coloring*.

**Step 1.** We start with the matrix  $C$  and look at two occurrences, which are identical for each of the types of even  $n \pmod 6$  specified above.

*Case 1.* We take the submatrix of  $G(i, j; l, m)$  with  $2 \leq l < m \leq n, 2 \leq i < j < n$ , and let  $s = (n + 1) - i, t = (n + 1) - j$ . A typical  $(2 \times 2)$  submatrix in this case has the form:

$$\begin{pmatrix} \sigma^s(l-1) & \sigma^s(m-1) \\ \sigma^t(l-1) & \sigma^t(m-1) \end{pmatrix}.$$

We wish to show there are three distinct elements:  $\sigma^s(l-1) \neq \sigma^t(l-1)$ ,  $\sigma^t(l-1) \neq \sigma^t(m-1)$ ,  $\sigma^s(l-1) \neq \sigma^t(m-1)$ .

Suppose  $\sigma^s(l-1) \equiv \sigma^t(l-1) \Rightarrow s \equiv t \pmod{n-1}$ , which is a contradiction.

Suppose  $\sigma^t(l-1) \equiv \sigma^t(m-1) \Rightarrow l \equiv m \pmod{n-1}$ , which is a contradiction.

Suppose  $\sigma^s(l-1) \equiv \sigma^t(m-1) \Rightarrow s + l \equiv t + m \pmod{n-1}$ , and assume there are three distinct elements:  $\sigma^t(l-1) \neq \sigma^t(m-1)$ ,  $\sigma^s(m-1) \neq \sigma^t(m-1)$ ,  $\sigma^s(m-1) \neq \sigma^t(l-1)$ . Follow the argument above the first two inequalities are correct. Suppose  $\sigma^s(m-1) \equiv \sigma^t(l-1) \Rightarrow s + m \equiv t + l \pmod{n-1}$ . Subtracting equations  $s + l \equiv t + m$  and  $s + m \equiv t + l \Rightarrow l \equiv m \pmod{n-1}$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{\sigma^s(l-1), \sigma^t(l-1), \sigma^t(m-1)\}$  or  $\{\sigma^t(l-1), \sigma^t(m-1), \sigma^s(m-1)\}$ .

*Case 2.* We take the submatrix of  $G(i, j; l, m)$  with  $2 \leq l < m \leq n, i = 1, 1 < j < n$ , and let  $r = (n + 1) - j$ . A typical  $(2 \times 2)$  submatrix has the form:

$$\begin{pmatrix} l & m \\ \sigma^r(l-1) & \sigma^r(m-1) \end{pmatrix}.$$

We wish to show there are three distinct elements:  $l \neq m, m \neq \sigma^r(m-1), l \neq \sigma^r(m-1)$ . Suppose  $m \equiv \sigma^r(m-1) \Rightarrow r \equiv 1 \pmod{n-1}$ , which is a contradiction. Suppose  $l \equiv \sigma^r(m-1) \Rightarrow l \equiv r + m - 1 \pmod{n-1}$ , and assume there are three distinct elements:  $\sigma^r(l-1) \neq \sigma^r(m-1), m \neq \sigma^r(m-1), m \neq \sigma^r(l-1)$ . The first two inequalities are correct. Suppose  $m \equiv \sigma^r(l-1) \Rightarrow m \equiv r + l - 1 \pmod{n-1}$ . Subtracting equations  $l \equiv r + m - 1$  and  $m \equiv r + l - 1 \Rightarrow r = 1 \pmod{n-1}$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{l, m, \sigma^r(m-1)\}$  or  $\{\sigma^r(l-1), \sigma^r(m-1), m\}$ .

**Step 2.** For matrix  $G(i, j; l, m)$  with  $i = 1, j = n$  and  $2 \leq l < m \leq n$  we look at five cases and consider every matrix type defined above of even  $n \pmod 6$ .

*Case 1.* We take  $G(i, j; l, m)$  with  $2 \leq l < m \leq \frac{n}{2} - 1$ ,  $i = 1$ ,  $j = n$ .

*Subcase 1.1.* Consider  $G_1$ ,

$$\begin{pmatrix} l & m \\ n-2l & n-2m \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n-2l$ ,  $n-2l \neq n-2m$ ,  $l \neq n-2m$ . Suppose  $l = n-2l \Rightarrow 3l = n$  and since  $n = 2 + 6k$  this is a contradiction. Suppose  $n-2l = n-2m \Rightarrow l = m$ , which is a contradiction. Suppose  $l = n-2m$  and we wish to show there are three distinct elements:  $l \neq m$ ,  $l \neq n-2l$ ,  $m \neq n-2l$ . As shown above the first inequality is correct. Suppose  $m = n-2l$  and since  $l = n-2m \Rightarrow l = m$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{l, n-2l, n-2m\}$  or  $\{l, m, n-2l\}$ .

*Subcase 1.2.* Consider  $G_2$ .

1.  $G_2$  with  $2 \leq l < m \leq \frac{n}{2} - 2$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n-2(l+1) & n-2(m+1) \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n-2(m+1)$ ,  $l \neq n-2(l+1)$ ,  $n-2(l+1) \neq n-2(m+1)$ . Suppose  $l = n-2l-2 \Rightarrow 3l = n-2$  and since  $n = 6 + 6k$  this is a contradiction. Suppose  $n-2l = n-2m \Rightarrow l = m$ , which is a contradiction. Suppose  $l = n-2(m+1)$  and we wish to show there are three distinct elements:  $l \neq m$ ,  $l \neq n-2(l+1)$ ,  $m \neq n-2(l+1)$ . As shown above the first inequality is correct. Suppose  $m = n-2(l+1)$  and since  $l = n-2(m+1) \Rightarrow l = m$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{l, n-2(m+1), n-2(l+1)\}$  or  $\{l, m, n-2(l+1)\}$ .

2.  $G_2$  with  $2 \leq l \leq \frac{n}{2} - 2$ ,  $m = \frac{n}{2} - 1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n-2(l+1) & Y \end{pmatrix}.$$

If  $K$  is *even*  $\Rightarrow m = \frac{n}{2} - 1$ ,  $Y = \frac{n}{2} - 2 \Rightarrow Y = m - 1$ .

We wish to show there are three distinct entries:  $l \neq m$ ,  $m \neq m-1$ ,  $l \neq m-1$ . Assume  $l = m-1$  and we wish to show there are three distinct entries:  $m \neq m-1$ ,  $m \neq n-2(l+1)$ ,  $m-1 \neq n-2(l+1)$ . Suppose  $m = n-2(l+1)$  and since  $l = m-1$  and  $m = \frac{n}{2} - 1 \Rightarrow n = 6$ , which is a contradiction. Suppose  $m-1 = n-2(l+1)$  and since  $m = \frac{n}{2} - 1$  and  $l = m-1 \Rightarrow n = 8$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{l, m, Y\}$  or  $\{m, Y, n-2(l+1)\}$ .

If  $K$  is *odd*  $\Rightarrow m = \frac{n}{2} - 1$ ,  $Y = \frac{n}{2} + 1 \Rightarrow Y = m + 2$ . Three distinct elements are  $\{l, m, Y\}$ .

*Subcase 1.3.* Consider  $G_3$ .

1.  $G_3$  with  $i = 1$ ,  $j = n$  and  $(2 \leq l < m \leq \frac{n-4}{6}$  or  $\frac{n+2}{6} \leq l < m \leq \frac{n}{2} - 2)$ . The argument is similar to above one with  $G_1$ . One of the following two sets has three distinct elements:  $\{l, n - 2l, n - 2m\}$  or  $\{l, m, n - 2l\}$ .

2.  $G_3$  with  $2 \leq l \leq \frac{n-4}{6}$ ,  $\frac{n+2}{6} \leq m \leq \frac{n}{2} - 2$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n - 2l & n - 2(m + 1) \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n - 2l$ ,  $n - 2l \neq n - 2(m + 1)$ ,  $l \neq n - 2(m + 1)$ . Suppose  $l = n - 2l \Rightarrow 3l = n$  and since  $n = 4 + 6k$  this is a contradiction. Suppose  $n - 2l = n - 2(m + 1) \Rightarrow l = m + 1$ , which is a contradiction. Suppose  $l = n - 2(m + 1)$  and we wish to show there are three distinct elements:  $l \neq m$ ,  $l \neq n - 2l$ ,  $m \neq n - 2l$ . The first two inequalities are correct. Suppose  $m = n - 2l$  and since  $l = n - 2(m + 1) \Rightarrow m = l - 2$ , which is a contradiction. One of the following two sets has three distinct elements:  $\{l, n - 2l, n - 2(m + 1)\}$  or  $\{l, m, n - 2l\}$ .

3.  $G_3$  with  $2 \leq l \leq \frac{n-4}{6}$ ,  $m = \frac{n}{2} - 1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n - 2l & n - 9 \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq m$ ,  $m \neq n - 9$ ,  $l \neq n - 9$ . Suppose  $l = n - 9$  and since  $l < \frac{n-4}{6} \Rightarrow n - 9 < \frac{n-4}{6} \Rightarrow n < 10$ , which is a contradiction. Suppose  $m = n - 9 \Rightarrow \frac{n}{2} - 1 = n - 9 \Rightarrow n = 16$ , which is a contradiction. There are three distinct elements  $\{l, m, n - 9\}$ .

4.  $G_3$  with  $\frac{n+2}{6} \leq l \leq \frac{n}{2} - 2$ ,  $m = \frac{n}{2} - 1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n - 2(l + 1) & n - 9 \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq m$ ,  $m \neq n - 9$ ,  $l \neq n - 9$ . Suppose  $l = n - 9$  and since  $l < \frac{n}{2} - 2 \Rightarrow n - 9 < \frac{n}{2} - 2 \Rightarrow n < 14$ , which is a contradiction. Suppose  $m = n - 9 \Rightarrow \frac{n}{2} - 1 = n - 9 \Rightarrow n = 16$ , which is a contradiction. There are three distinct elements  $\{l, m, n - 9\}$ .

*Case 2.* For the submatrix  $G(i, j; l, m)$  with  $i = 1$ ,  $j = n$  and  $(\frac{n}{2} \leq l < m \leq n - 2$  or  $2 \leq l \leq \frac{n}{2} - 1$ ,  $\frac{n}{2} \leq m \leq n - 2)$  three distinct elements are  $\{l, m, n\}$ .

*Case 3.* We take the submatrix  $G(i, j; l, m)$  with  $2 \leq l \leq \frac{n}{2} - 1$ ,  $m = n - 1$ ,  $i = 1$ ,  $j = n$ .

*Subcase 3.1.* Consider  $G_1$ ,

$$\begin{pmatrix} l & m \\ n-2l & n-1 \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n-1$ ,  $n-2l \neq n-1$ ,  $l \neq n-2l$ . Suppose  $n-2l = n-1 \Rightarrow l = \frac{1}{2}$ , which is a contradiction. Suppose  $l = n-2l \Rightarrow 3l = n$  and since  $n = 2 + 6k$  this is a contradiction. There are three distinct elements  $\{l, n-1, n-2l\}$ .

*Subcase 3.2.* Consider  $G_2$ .

1.  $G_2$  with  $2 \leq l \leq \frac{n}{2} - 2$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n-2(l+1) & n-1 \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n-1$ ,  $n-2(l+1) \neq n-1$ ,  $l \neq n-2(l+1)$ . Suppose  $n-2(l+1) = n-1 \Rightarrow l = -\frac{1}{2}$ , which is a contradiction. Suppose  $l = n-2(l+1) \Rightarrow 3l = n-2$  and since  $n = 6 + 6k$  this is a contradiction. There are three distinct elements  $\{l, n-1, n-2(l+1)\}$ .

2.  $G_2$  with  $l = \frac{n}{2} - 1$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ Y & n-1 \end{pmatrix}.$$

If  $K$  is *even*  $\Rightarrow Y = \frac{n}{2} - 2$ . If  $n-1 = Y \Rightarrow n-1 = \frac{n}{2} - 2 \Rightarrow n = -2$ , which is a contradiction. Three distinct entries are  $\{l, Y, n-1\}$ .

If  $K$  is *odd*  $\Rightarrow Y = \frac{n}{2} + 1$ , and three distinct entries are  $\{l, Y, n-1\}$ .

*Subcase 3.3.* Consider  $G_3$ .

1. For  $G_3$  with  $l \leq \frac{n-4}{6}$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$  three distinct elements are  $\{l, n-1, n-2l\}$  (similar to  $G_1$ .)

2.  $G_3$  with  $\frac{n+2}{6} \leq l < n-1$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$ ,

$$\begin{pmatrix} l & m \\ n-2(l+1) & n-1 \end{pmatrix}.$$

We wish to show there are three distinct entries:  $l \neq n-1$ ,  $n-2(l+1) \neq n-1$ ,  $l \neq n-2(l+1)$ . Suppose  $n-2(l+1) = n-1 \Rightarrow l = -\frac{1}{2}$ , which is a contradiction. Suppose  $l = n-2(l+1) \Rightarrow 3l = n-2$  and since  $n = 4 + 6k$  this is a contradiction. Three distinct elements are  $\{l, n-1, n-2(l+1)\}$ .

3. For  $G_3$  with  $l = \frac{n}{2} - 1$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$  three distinct entries are  $\{l, n-9, n-1\}$ .

*Case 4.* For the submatrix  $G(i, j; l, m)$  with  $\frac{n}{2} \leq l \leq n-2$ ,  $m = n-1$ ,  $i = 1$ ,  $j = n$  three distinct elements are  $\{l, n, n-1\}$ .

*Case 5.* For the submatrix  $G(i, j; l, m)$  with  $l = n - 1$ ,  $m = n$ ,  $i = 1$ ,  $j = n$  three distinct elements are  $\{n - 1, n, 1\}$ .

The argument for other steps is similar. To see the details please view the appendix to this article on ArXiv at <http://arxiv.org/> or contact the first author.

#### REFERENCES

- [1] M. Axenovich, *A generalized Ramsey problem*, Discrete Math. **222** (2000) 247–249.  
doi:10.1016/S0012-365X(00)00052-2
- [2] M. Axenovich, Z. Füredi and D. Mubayi, *On generalized Ramsey theory: the bipartite case*, J. Combin. Theory (B) **79** (2000) 66–86.  
doi:10.1006/jctb.1999.1948
- [3] R. Diestel, *Graph Theory*, Third Edition (Springer-Verlag, Heidelberg, Graduate Texts in Mathematics, Volume 173, 2005).
- [4] P. Erdős, *Solved and unsolved problems in combinatorics and combinatorial number theory*, Congr. Numer. **32** (1981) 49–62.
- [5] P. Erdős and A. Gyárfás, *A variant of the classical Ramsey problem*, Combinatorica **17** (1997) 459–467.  
doi:10.1007/BF01195000
- [6] J. Fox and B. Sudakov, *Ramsey-type problem for an almost monochromatic  $K_4$* , SIAM J. Discrete Math. **23** (2008) 155–162.  
doi:10.1137/070706628
- [7] S. Fujita, C. Magnant and K. Ozeki, *Rainbow generalizations of Ramsey theory: A survey*, Graphs Combin. **26** (2010) 1–30.  
doi:10.1007/s00373-010-0891-3
- [8] A. Kostochka and D. Mubayi, *When is an almost monochromatic  $K_4$  guaranteed?*, Combin. Probab. Comput. **17** (2008) 823–830.  
doi:10.1017/S0963548308009413
- [9] D. Mubayi, *Edge-coloring cliques with three colors on all four cliques*, Combinatorica **18** (1998) 293–296.  
doi:10.1007/PL00009822
- [10] R. Wilson, *Graph Theory*, Fourth Edition (Prentice Hall, Pearson Education Limited, 1996).

Received 6 December 2012

Revised 21 September 2012

Accepted 21 September 2012