

MAXIMUM HYPERGRAPHS WITHOUT REGULAR SUBGRAPHS

JAEHOON KIM¹

*Department of Mathematics, University of Illinois
Urbana, IL, 61801, USA*

e-mail: kim805@illinois.edu

AND

ALEXANDR V. KOSTOCHKA²

*University of Illinois at Urbana–Champaign,
Urbana, IL 61801, USA
Sobolev Institute of Mathematics
Novosibirsk 630090, Russia*

e-mail: kostochk@math.uiuc.edu

Abstract

We show that an n -vertex hypergraph with no r -regular subgraphs has at most $2^{n-1} + r - 2$ edges. We conjecture that if $n > r$, then every n -vertex hypergraph with no r -regular subgraphs having the maximum number of edges contains a full star, that is, 2^{n-1} distinct edges containing a given vertex. We prove this conjecture for $n \geq 425$. The condition that $n > r$ cannot be weakened.

Keywords: hypergraphs, set system, subgraph, regular graph.

2010 Mathematics Subject Classification: 05C65.

¹Research of this author is partially supported by the Arnold O. Beckman Research Award of the University of Illinois at Urbana-Champaign.

²Research of this author is supported in part by NSF grant DMS-0965587 and by the Ministry of education and science of the Russian Federation (Contract no. 14.740.11.0868).

1. INTRODUCTION

A natural question in graph theory is: What are the graphs not containing r -regular subgraphs? For $r \in \{1, 2\}$, the answer is easy, but for $r \geq 3$ it is not. It was a breakthrough when Tashkinov [7] proved the conjecture by Berge that every 4-regular graph contains a 3-regular subgraph. The questions on existence of r -regular subgraphs in regular or near-regular graphs were also considered in [1, 8]. Let $F(r, n)$ denote the maximum number of edges an n -vertex graph with no r -regular subgraphs have. For $r \geq 3$, it is not fully resolved how big $F(r, n)$ is. Pyber [4] showed that for every fixed r , $F(r, n) = O(n \ln n)$. On the other hand, Pyber, Rödl and Szemerédi [5] proved that $F(3, n) \geq cn \ln \ln n$.

Similar questions are also natural for hypergraphs. We view a hypergraph as a family \mathcal{F} of its edges, so $|\mathcal{F}|$ is the number of edges of \mathcal{F} . An edge e of \mathcal{F} is a k -edge if $|e| = k$. Note that we do not consider empty set as an edge. If, for some k , every edge of \mathcal{F} is a k -edge, then \mathcal{F} is k -uniform. A hypergraph \mathcal{F} is r -free if it has no r -regular sub(hyper)graphs. Mubayi and Verstraëte [3] proved that for every even integer $k \geq 4$, there exists n_k such that for each $n \geq n_k$, each n -vertex k -uniform 2-free hypergraph \mathcal{F} has at most $\binom{n-1}{k-1}$ edges, and equality holds if and only if \mathcal{F} is a full k -star, that is, \mathcal{F} consists of all $\binom{n-1}{k-1}$ edges of size k containing a given vertex. They also proved the following simpler result for non-uniform hypergraphs.

Theorem 1.1 [3]. *For $n \geq 3$, every n -vertex 2-free hypergraph \mathcal{F} satisfies $|\mathcal{F}| \leq 2^{n-1}$, and equality holds if and only if \mathcal{F} is a full star, that is, \mathcal{F} consists of 2^{n-1} distinct edges containing a given vertex.*

Our first result is the following (simple) generalization of Theorem 1.1.

Theorem 1.2. *If $2 \leq r \leq 2^{n-1}$, then the maximum number of edges in an n -vertex r -free hypergraph is $2^{n-1} + r - 2$.*

Many examples of n -vertex r -free hypergraphs with $2^{n-1} + r - 2$ edges are formed by a full star with $r - 2$ other edges. If $r \geq n$, then some extremal examples do not contain full stars. For $r = 2$, Theorem 1.1 says that if $n \geq 3$, then the only n -vertex 2-free hypergraph with 2^{n-1} edges is a full star. We conjecture the following.

Conjecture 1.3. *Let \mathcal{F} be an n -vertex r -free hypergraph with $|\mathcal{F}| = 2^{n-1} + r - 2$. If $n > r$ and $r \geq 2$, then \mathcal{F} contains a full star.*

The main results of this paper are the following.

Theorem 1.4. *Suppose \mathcal{F} is an n -vertex r -free hypergraph with $|\mathcal{F}| = 2^{n-1} + r - 2$. If $n \geq r + 2 \lceil \log r \rceil + 1$, then \mathcal{F} contains a full star.*

Theorem 1.5. *Suppose \mathcal{F} is an n -vertex r -free hypergraph with $|\mathcal{F}| = 2^{n-1} + r - 2$. If $n > r$ and $n \geq 425$, then \mathcal{F} contains a full star.*

In the next section we prove Theorem 1.2 and derive simple properties of dense r -free hypergraphs. In Section 3 we show that dense r -free hypergraphs have no small transversals. In Section 4 we prove Theorem 1.4. In the last two sections we prove Theorem 1.5.

2. PRELIMINARIES

Proof of Theorem 1.2. Let \mathcal{F} be an n -vertex r -free hypergraph with ground set N . Consider all 2^{n-1} pairs $\{A, N - A\}$ of subsets of N . In at most $r - 1$ pairs of sets both sets are edges in \mathcal{F} , otherwise we get an r -regular subgraph of \mathcal{F} with vertex set N . If there are exactly $r - 1$ such pairs, N cannot be an edge in \mathcal{F} , since N together with those $r - 1$ pairs would form an r -regular subgraph of \mathcal{F} . Thus $|\mathcal{F}| \leq 2^{n-1} + r - 2$. If $2 \leq r \leq 2^{n-1}$, then equality can be achieved.

Let $N = [n]$ and

$$\mathcal{F} = \{e : 1 \in e\} \cup \{r - 2 \text{ smallest nonempty distinct subsets of } [n] - \{1\}\}.$$

Suppose that \mathcal{F} has an r -regular subgraph \mathcal{G} . Let C_1, C_2, \dots, C_r be the edges of \mathcal{G} that contain 1, and D_1, D_2, \dots, D_s be the remaining edges of \mathcal{G} . Let $C = \bigcup_{i=1}^r C_i$ and for $i \in [r]$ let $C'_i = C - C_i$. Since all edges are distinct, $\sum_{i=1}^r |C'_i| = r|C| - \sum_{i=1}^r |C_i| = \sum_{j=1}^s |D_j|$ should hold. The left-hand side is the sum of cardinalities of at least $r - 1$ nonempty distinct sets (possibly one C'_i is empty) not containing 1, and the right-hand side is the sum of cardinalities of at most $r - 2$ smallest distinct sets not containing 1, and so, the right-hand side is less than the left-hand side. This contradiction shows that H has no r -regular subgraphs. ■

Let N be a finite set, and $n = |N|$. Let $3 \leq r < n$. A hypergraph \mathcal{F} is (N, r) -*strange* if \mathcal{F} is an r -free hypergraph with $V(\mathcal{F}) = N$ and $|\mathcal{F}| = 2^{n-1} + r - 2$ such that \mathcal{F} does not contain a *full star*, i.e., 2^{n-1} sets containing a given element.

For a set $A \subseteq N$, \bar{A} is the *complement* of A to N , i.e., $\bar{A} = N - A$. A *full pair* in \mathcal{F} is a pair $\{A, \bar{A}\}$ such that both A and \bar{A} are in \mathcal{F} . We let the set N by itself form a full pair.

In order to prove Theorems 1.4 and 1.5, we derive some properties of (N, r) -strange hypergraphs. If \mathcal{F} is (N, r) -strange, then it contains at most $r - 1$ full pairs, and so, since $|\mathcal{F}| = 2^{n-1} + r - 2$,

- (1) it contains exactly $r - 1$ full pairs.

Moreover,

- (2) for each $A \subset N$ with $N \neq A \neq \emptyset$, either $A \in \mathcal{F}$ or $\bar{A} \in \mathcal{F}$.

Furthermore, the following statements hold for each (N, r) -strange hypergraph \mathcal{F} .

Lemma 2.1. *If $A, B \in \mathcal{F}$, $A \cap B = \emptyset$ and both A and B are not in full pairs, then $A \cup B \in \mathcal{F}$.*

Proof. If $A \cup B \notin \mathcal{F}$, then $\overline{A \cup B} \in \mathcal{F}$ by (2). Thus $A, B, \overline{A \cup B}$ with $r - 1$ full pairs form an r -regular subfamily of \mathcal{F} , a contradiction. ■

Lemma 2.2. *If $A \in \mathcal{F}$ and B and C are disjoint nonempty subsets of A such that $A = B \cup C$, then at least one of B and C is in \mathcal{F} .*

Proof. Suppose that $A = B \cup C$ is a partition of A into nonempty sets and $B, C \notin \mathcal{F}$. Then by (2), \overline{B} and \overline{C} are in \mathcal{F} but not in full pairs. Thus the sets A, \overline{B} and \overline{C} together with $r - 2$ full pairs different from $\{A, \overline{A}\}$ form an r -regular subgraph of \mathcal{F} , a contradiction. ■

Corollary 2.3. *Every edge A of \mathcal{F} contains an element x_A such that $\{x_A\} \in \mathcal{F}$. In particular, the union S of 1-edges of \mathcal{F} intersects each edge of \mathcal{F} .*

Lemma 2.4. *Let A and B be edges of \mathcal{F} such that $A \cap B \neq \emptyset$. If at least one of A and B is not in a full pair, then either $A \cap B$ or $A \cup B$ is in \mathcal{F} .*

Proof. Suppose that $A \cap B, A \cup B \notin \mathcal{F}$. Then $\overline{A \cap B}$ is in \mathcal{F} , and $\overline{A \cup B}$ is either empty or in \mathcal{F} . In both cases, the sets $A, B, \overline{A \cap B}$, and $\overline{A \cup B}$ cover every element of N exactly twice. Adding $r - 2$ full pairs containing neither A nor B will give an r -regular subgraph of \mathcal{F} . ■

3. SIZES OF TRANSVERSALS OF (N, r) -STRANGE HYPERGRAPHS

A set $A \subset V(H)$ is a *transversal* of a hypergraph H if every edge of H intersects A .

Let S be a minimum transversal of a hypergraph \mathcal{F} . Then S contains all 1-edges of \mathcal{F} . If \mathcal{F} is (N, r) -strange, then by Corollary 2.3, S contains no other vertices. Thus S is exactly the union of 1-edges of \mathcal{F} . It has several useful properties.

The goal of this section is to prove the following fact. Throughout the paper, k denote $\lceil \log_2 r \rceil$.

Theorem 3.1. *Let $3 \leq r < n$ and N be a finite set with $|N| = n$. If S is the smallest transversal of an (N, r) -strange hypergraph \mathcal{F} , then $|S| \geq n - 3k - 2$.*

Let S be the smallest transversal of an (N, r) -strange hypergraph \mathcal{F} .

Lemma 3.2. *If a nonempty $S' \subset S$ is not in \mathcal{F} , then every $S' \subseteq B \subseteq N - (S - S')$ is not in \mathcal{F} , and hence every $S - S' \subseteq A \subseteq N - S'$ is in \mathcal{F} .*

Proof. Suppose that such B is in \mathcal{F} . By Lemma 2.2, either S' or $B - S'$ is in \mathcal{F} . But $(B - S') \cap S = \emptyset$, and we know that S' is not in \mathcal{F} , a contradiction. ■

From now on, in this section, we will assume that

$$(3) \quad |S| \leq n - 2k - 2.$$

Note that to prove Theorem 3.1, we could make the stronger assumption that $|S| \leq n - 3k - 3$, but we plan to use these lemmas also in the next section.

For $S' \subseteq S$ and $M \subseteq N - S$, we say that M *belongs to* S' if $S' \cup M \in \mathcal{F}$. A nonempty proper subset S' of S is *firm* if some $M \subset N - S$ with $|M| \geq 1 + k$ belongs to S' . In particular, S is firm by the following reason. For a set $A \subset N - S$ with $|A| = k + 1$, one of $A \cup S$ and $N - S - A$ is in \mathcal{F} by (2). Since S is a transversal, $N - S - A$ is not in \mathcal{F} . Thus $S \cup A \in \mathcal{F}$, so A belongs to S and S is firm.

Lemma 3.3. *Let $S' \subseteq S$ and $M \subseteq N - S$. If M belongs to S' , then every $M' \subset M$ belongs to S' .*

Proof. Since $M \cup S' \in \mathcal{F}$, by Lemma 2.2, either $S' \cup M' \in \mathcal{F}$ or $M - M' \in \mathcal{F}$. But the latter does not hold, since $M \cap S = \emptyset$. This proves the lemma. ■

Lemma 3.4. *For every partition $S = S' \cup S''$ of S into nonempty subsets, exactly one of S' and S'' is firm.*

Proof. Assume first that neither of S' and S'' is firm. Let M be a subset of $N - S$ with $|M| = 1 + k$. Since S' is not firm, $S' \cup M \notin \mathcal{F}$. Then $N - (S' \cup M) \in \mathcal{F}$, and $N - (S' \cup M) = S'' \cup (N - S - M)$. So by (3), $|N - S - M| \geq 2k + 2 - (1 + k)$, and thus S'' is firm. Assume now that both S' and S'' are firm. If a set $M \subset N - S$ with $|M| \geq k + 1$ belongs to both S' and S'' , then we will find an r -regular subgraph \mathcal{H} of \mathcal{F} .

Since $2^{|M|} \geq r$, there are at least r subsets of M . Call them A_1, A_2, \dots, A_r . Let $\mathcal{H} = \{A_i \cup S' : 1 \leq i \leq r\} \cup \{(M - A_i) \cup S'' : 1 \leq i \leq r\}$, it is a subgraph of \mathcal{F} by Lemma 3.3. By construction, \mathcal{H} is r -regular, a contradiction.

If a set $M \subset N - S$ with $k \leq |M| \leq k + 2$ belongs to neither S' nor S'' , then $N - S - M$ belongs to both, and again \mathcal{F} has an r -regular subgraph. Thus each $M \subset N - S$ with $|M| = k + 1$ belongs to exactly one of S' and S'' . Let $\mathcal{R}_{S'}$ (respectively, $\mathcal{R}_{S''}$) denote the family of $M \subset N - S$ with $|M| = k + 1$ that belongs to S' (respectively, to S''). By our assumption, both $\mathcal{R}_{S'}$ and $\mathcal{R}_{S''}$ are nonempty. Then there exist $M' \in \mathcal{R}_{S'}$ and $M'' \in \mathcal{R}_{S''}$ with $|M' \cap M''| = k$. Thus $M' \cap M''$ belongs to both S' and S'' , and so \mathcal{F} has an r -regular subgraph, a contradiction. ■

Corollary 3.5. *If S' is a firm subset of S , then every $M \subset N - S$ with $1 \leq |M| \leq n - s - (k + 1)$ belongs to S' .*

Corollary 3.6. *Every two firm subsets of S intersect each other.*

Proof. Suppose that S_1 and S_2 are two disjoint firm subsets of S . Let $M \subset N - S$ with $|M| = k + 1$. By Corollary 3.5, M belongs to both S_1 and S_2 . Then as in the proof of Lemma 3.4, \mathcal{F} has an r -regular subgraph with vertex set $S_1 \cup S_2 \cup M$, a contradiction. ■

Lemma 3.7. $s \leq r - 1$

Proof. Suppose $s \geq r$.

Case 1. $s \geq r + k$. Since the number of 1-edges in full pairs is at most $r - 1$, we can choose $k + 1 (\leq s - (r - 1))$ 1-edges of \mathcal{F} that are not in full pairs. Let S' be the union of these edges. If some $A \subseteq S'$ is not in \mathcal{F} , then \overline{A} and the 1-edges contained in A cover N once, and together with the $r - 1$ full pairs (that exist by (1)) we obtain an r -regular subgraph of \mathcal{F} covering N , a contradiction. Thus all nonempty subsets of S' are in \mathcal{F} , and the number of nonempty proper subsets of S' is at least $2^{k+1} - 2 \geq 2r - 2$. We can pair them up so that they are partitions of S' . At least $r - 1$ of such pairs exist, so together with S' they form an r -regular subgraph of \mathcal{F} , a contradiction.

Case 2. $r \leq s \leq r + k - 1$. By (3), $n - s \geq k + 1$. If there are $v_1, v_2 \in S$ such that $S - v_1, S - v_2 \notin \mathcal{F}$, then by Lemma 3.2, every $B \subseteq N - S$ satisfies $B + v_1 \in \mathcal{F}$ and $B + v_2 \in \mathcal{F}$. Since there are at least $2^{n-s} \geq r$ possible sets for B , we can find r pairs of sets $v_1 + B, v_2 + (N - S - B)$, and they will form an r -regular subgraph of \mathcal{F} on $(N - S) + v_1 + v_2$.

Thus for some $r - 1$ vertices $v_1, \dots, v_{r-1} \in S$, the sets $S - v_i$ are in \mathcal{F} . Then the family $\{v_1, \dots, v_{r-1}, S - v_1, \dots, S - v_{r-1}, S\}$ covers every $v \in S$ exactly r times, a contradiction. ■

Lemma 3.8. *No 1-edge of \mathcal{F} is firm.*

Proof. Let $S = \{v_1, v_2, \dots, v_s\}$. Suppose that $S_1 := \{v_1\}$ is firm. Then by Corollary 3.6, no subset of $S - v_1$ is firm. Hence by Lemma 3.4, the firm subsets of S are exactly the sets containing v_1 .

Since not every subset of N containing v_1 is in \mathcal{F} and $s \leq r - 1$, there are at least $r - 1 - (s - 1) = r - s$ edges W_1, \dots, W_{r-s} that are in \mathcal{F} , not 1-edges and do not contain v_1 . For $j = 1, \dots, r - s$, let $M_j = W_j - S$. Let $M = \bigcup_{j=1}^{r-s} M_j$. Choose W_1, \dots, W_{r-s} so that to minimize $|M|$.

Case 1. $|M| \leq n - s - k - 1$. Denote by \mathcal{F}' the family $\{S \cup M, \{v_2\}, \dots, \{v_s\}, S \cup M - v_2, \dots, S \cup M - v_s, W_1, \dots, W_{r-s}, S \cup M - W_1, \dots, S \cup M - W_{r-s}\}$.

Since \mathcal{F}' forms an r -regular hypergraph, \mathcal{F}' is not a subgraph of \mathcal{F} . But since $\{v_1\}$ is firm, by the choice of W_j and Corollary 3.5, every member of \mathcal{F}' is in \mathcal{F} , a contradiction. This proves Case 1.

Let $t = \max\{|A - S| : A \in \mathcal{F} \text{ and } v_1 \notin A\}$ and let $A_0 \in \mathcal{F}$ be such that $v_1 \notin A_0$ and $|A_0 - S| = t$.

Case 2. $t \geq k$. Let M_0 be any k -element subset of $A_0 - S$ and $W_0 = M_0 \cup (A_0 \cap S)$. Since $A_0 \in \mathcal{F}$ and $S \cap (A_0 - W_0) = \emptyset$, $W_0 \in \mathcal{F}$. Since $2^k \geq r$, M_0 contains some r distinct subsets M'_1, \dots, M'_r . Let $W'_i = M'_i \cup (A_0 \cap S)$ for $i = 1, \dots, r$. Since $(M_0 - M'_i) \cap S = \emptyset$, each of W'_i is in \mathcal{F} . Moreover, since $|S| \leq n - 2k - 2$, $|M_0| = k$, and $\{v_1\}$ is firm, for every $1 \leq i \leq r$, the set $(S \cup M_0) - W'_i$ contains v_1 and has at most $n - s - (k + 1)$ vertices in $N - S$. This means that $(S \cup M_0) - W'_i$ is also in \mathcal{F} . So, the family $\{W'_1, \dots, W'_r, (S \cup M_0) - W'_1, \dots, (S \cup M_0) - W'_r\}$ forms an r -regular hypergraph, a contradiction.

Case 3. $\log_2(r - s) \leq t \leq k - 1$. Let $M_0 = A_0 - S$. In our case, $2^{|M_0|} = 2^t \geq r - s$. Let M'_1, \dots, M'_{r-s} be any distinct subsets of M_0 , and for $i = 1, \dots, r - s$, let $W'_i = M'_i \cup (A_0 \cap S)$. Similarly to Case 2, since $(M_0 - M'_i) \cap S = \emptyset$, each of W'_i is in \mathcal{F} . Moreover, since $|S| \leq n - 2k - 2$, $|M_0| \leq k - 1$, and $\{v_1\}$ is firm, for every $1 \leq i \leq r$, the set $(S \cup M_0) - W'_i$ is also in \mathcal{F} . By the same reason, for every $2 \leq j \leq s$, the set $(S \cup M_0) - v_j$ is in \mathcal{F} . So, the family $\{W'_1, \dots, W'_{r-s}, (S \cup M_0) - W'_1, \dots, (S \cup M_0) - W'_{r-s}, \{v_2\}, \dots, \{v_s\}, S \cup M_0 - v_2, \dots, S \cup M_0 - v_s, S \cup M_0\}$ forms an r -regular hypergraph, a contradiction.

Case 4. $|M| \geq n - k - s$ and $t < \min\{k - 1, \log_2(r - s)\}$. Let $M_0 = A_0 - S$. We claim that $|M| \leq r - s - 2^t + t + 1$. To prove the claim, we show a way to choose W_1, \dots, W_{r-s} so that

$$(4) \quad \left| \bigcup_{j=1}^i W_j - S \right| \leq i - 2^t + t + 1,$$

for every $2^t - 1 \leq i \leq r - s$. The sets W_1, \dots, W_{2^t-1} are all the sets of the form $A_0 - X$ where $X \subseteq M_0$, $X \neq M_0$. So, for $i = 2^t - 1$, (4) holds. Suppose that for some $2^t - 1 \leq i_0 \leq r - s - 1$, we have found W_1, \dots, W_{i_0} satisfying (4) for $i = i_0$. Let \mathcal{C} be the family of members of \mathcal{F} not containing v_1 that are distinct from W_1, \dots, W_{i_0} . Since $i_j \leq r - s - 1$, there is $C \in \mathcal{C} \neq \emptyset$. Let $C' = C - S - \bigcup_{j=1}^{i_0} W_j$. If $C' = \emptyset$, then we let $C := W_{i_0+1}$ and (4) holds for $i = i_0 + 1$. Suppose $x \in C'$. Since $(C' - x) \cap S = \emptyset$, the set $C - C' + x$ is in \mathcal{F} , does not contain v_1 , and is distinct from W_1, \dots, W_{i_0} . So, letting $W_{i_0+1} = C - C' + x$ we again have that (4) holds for $i = i_0 + 1$. This proves the claim.

Let \mathcal{F}' be the family defined in Case 1. Since it is r -regular, some $W' \in \mathcal{F}'$ is not in \mathcal{F} . Then $\overline{S \cup M - W'} \in \mathcal{F}$ by (2). By the definition of t , $|N - (S \cup M)| \leq |\overline{S \cup M - W'} - S| \leq t$. Thus by (4), $n = |N| = |M| + |S| + |N - (M \cup S)| \leq (r - s - 2^t + t + 1) + s + t = r - 2^t + 2t + 1$.

If $t \geq 3$, we get $n \leq r - 1$, a contradiction.

If $t = 2$, we get $n \leq r + 1$, $|M| = r - s - 1$ and $|A_0 - S| = 2$. Then $(A_0 - S) \subset M$ with $|M| = r - s - 1$ and $|\overline{M \cup S - W'} - S| \geq 2$. Thus there are distinct A_0, A_1 with $|A_i - S| = 2$, $(A_0 - S) \cap (A_1 - S) = \emptyset$ and $v_1 \notin A_i$. Let $\{v_1, v_2, \dots, v_{r-s-6}\} \subset M - A_0 - A_1$. For $x = 1, 2, \dots, r - s - 6$, let $v_x \in M_{j_x}$. By Lemma 3.3, $W_{j_x} - M_{j_x} + v_x \in \mathcal{F}$ for every $x_1, \dots, r - s - 6$. These edges with $A \cup (A_i \cap S)$ for nonempty $A \subseteq A_i - S$ yield that $|M| \leq r - s - 2$, a contradiction.

Thus $t = 1$. Then $n = r + 1$, $|M| = r - s$ and every edge not containing v_1 is either a 2-edge or a 1-edge. And $\overline{S \cup M - W'}$ is also a 2-edge, so W' is a 1-edge. Let $W' = \{v_2\}$.

Since $|M| = r - s$ and $\overline{S \cup M - v_2} = \{v_2\} \cup (N - M - S)$, for each $w \in N - S$ there is exactly one 2-edge not containing v_1 containing w . Since $|N - S| = r - s + 1$ and the number of 1-edges in $S - v_1$ is $s - 1$, \mathcal{F} has exactly $r - s + 1 + s - 1 = r$ edges not containing v_1 .

So we have exactly two sets containing v_1 that are not in \mathcal{F} . Call them D_1, D_2 . We have $D_1 = S \cup M - v_2$. Since $|N - S| = n - s \geq 2k + 2 \geq 4$ and every vertex in $N - S$ is contained in exactly one 2-edge not containing v_1 , there are at least 4 different ways to choose W_i s to get minimum $|M|$. Each way gives different M , let two of them be M and M' . Then, by the above logic, $D_2 = S \cup M' - v_3$ for some v_3 , and $|M \cap M'| \geq 2$. Thus $D_1 \cap D_2 - S$ is not empty. Let $w \in D_1 \cap D_2 - S$. Then there are $2^{n-1} - 2$ edges in \mathcal{F} containing both w and v_1 , and exactly one edge containing w but not v_1 . Thus $\mathcal{F} - w = \{E \in \mathcal{F} : w \notin E\}$ is an $(n - 1)$ -vertex hypergraph with $2^{n-1} + r - 1$ edges. By Theorem 1.2, $\mathcal{F} - w$ contains an r -regular subgraph, a contradiction. ■

Lemma 3.9. $s \leq r - 2$.

Proof. Suppose that $s \geq r - 1$. Then by Lemma 3.7, $s = r - 1$. By Lemma 3.8, $S - v_i$ is firm (and so is in \mathcal{F}) for every $i = 1, \dots, s$. Then the $2r - 1$ sets

$$\{v_1\}, \dots, \{v_{r-1}\}, S - v_1, \dots, S - v_{r-1}, S$$

form an r -regular subgraph of \mathcal{F} , a contradiction. ■

Lemma 3.10. Let $k \geq 2$ and let B be a set with $|B| \geq 2k + 2$. Then there are at least $2^{k-2} + 1$ partitions $(B_{i,1}, B_{i,2}, B_{i,3})$ of B such that $|B_{i,1}| = \lceil (k+1)/2 \rceil$, $|B_{i,2}| = \lceil (k+1)/2 \rceil$, $|B_{i,3}| \geq k + 1$ and all $3 \lceil 2^{k-2} \rceil + 3$ parts of these partitions are distinct.

Proof. We will choose B_1 of size $\lceil (k+1)/2 \rceil$ and B_2 of size $\lceil (k+1)/2 \rceil$, so that $|B_3| = |B| - 2 \lceil k/2 \rceil \geq k + 1$.

For $k \geq 4$, we have

$$(5) \quad \binom{2k+2-\lceil \frac{k+1}{2} \rceil}{\lceil \frac{k+1}{2} \rceil} = \frac{(2k+2-\lceil \frac{k+1}{2} \rceil)(2k+1-\lceil \frac{k+1}{2} \rceil)(2k-\lceil \frac{k+1}{2} \rceil)}{\lceil \frac{k+1}{2} \rceil(2k+2-2\lceil \frac{k+1}{2} \rceil)(2k+1-2\lceil \frac{k+1}{2} \rceil)} \\ \binom{2k-2-\lceil \frac{k-2+1}{2} \rceil}{\lceil \frac{k-2+1}{2} \rceil} \geq 4 \binom{2(k-2)+2-\lceil \frac{k-1}{2} \rceil}{\lceil \frac{k-1}{2} \rceil}.$$

So, for even k ,

$$\binom{2k+2-\lceil (k+1)/2 \rceil}{\lceil (k+1)/2 \rceil} \geq 4 \binom{2(k-2)+2-\lceil (k-1)/2 \rceil}{\lceil (k-1)/2 \rceil} \\ \geq \dots \geq 4^{\frac{k-2}{2}} \binom{4}{2} = 6 \cdot 2^{k-2}.$$

For odd k ,

$$\binom{2k+2-\lceil (k+1)/2 \rceil}{\lceil (k+1)/2 \rceil} \geq 4 \binom{2(k-2)+2-\lceil (k-1)/2 \rceil}{\lceil (k-1)/2 \rceil} \geq \dots \geq 4^{\frac{k-3}{2}} \binom{6}{2} \\ = 15 \cdot 2^{k-3} \geq 6 \cdot 2^{k-2}.$$

First for each $i = 1, \dots, 2^{k-2} + 1$, choose a set $B_{i,1}$ of size $\lceil k/2 \rceil$ so that all chosen sets are distinct. Then one by one for each $i = 1, \dots, 2^{k-2} + 1$, choose a set $B_{i,2}$ of size $\lceil k/2 \rceil$ so that

- (a) $B_{i,2}$ is distinct from all $2^{k-2} + 1$ sets $B_{i',1}$ and previously chosen $B_{i',2}$, and
- (b) $B_{i,1} \cup B_{i,2}$ is distinct from all already chosen $B_{i',1} \cup B_{i',2}$.

Even at the last step (step $2^{k-2} + 1$), the number of forbidden sets is at most $3 \cdot 2^{k-2} + 1 < 6 \cdot 2^{k-2}$. So, by (5), we finish the construction. ■

Corollary 3.11. *Let B be a set with $|B| \geq 3k + 3$. Then there are at least $2^{k-2} + 1$ partitions $(B_{i,1}, B_{i,2}, B_{i,3})$ of B such that all $3\lceil 2^{k-2} \rceil + 3$ parts of these partitions are distinct, and each $B_{i,j}$ has size at least $k + 1$.*

Proof. Let $A \subseteq B$ and $|A| = k + 1$. Let $B' = B - A$. Partition A into $A_1 \cup A_2 \cup A_3$ with $|A_1| = |A_2| = k + 1 - \lceil \frac{k+1}{2} \rceil$. By Lemma 3.10, there are at least $2^{k-2} + 1$ partitions $(B'_{i,1}, B'_{i,2}, B'_{i,3})$ of B' such that all parts of these partitions are distinct, and $|B'_{i,1}| = |B'_{i,2}| = \lceil \frac{k+1}{2} \rceil, |B'_{i,3}| \geq k + 1$. Take $B_{i,j} = B'_{i,j} \cup A_i$. Then partitions $(B_{i,1}, B_{i,2}, B_{i,3})$ satisfy all the conditions. ■

Proof of Theorem 3.1. Suppose $s \leq n - 3k - 3$. Then all lemmas in this section hold, since $s \leq n - 3k - 3 \leq n - 2k - 2$.

Let S' be a smallest firm subset of S . Note that S' is not a 1-edge. Partition S' into nonempty subsets S_1 and S_2 . By the minimality of S' , sets S_1 and S_2 are not firm, and so $S - S_1$ and $S - S_2$ are firm.

Let $B := N - S$. Then $|B| = n - s \geq 3k + 3$. So, by Corollary 3.11, there are $K := \lceil 2^k/6 \rceil + 1$ partitions

$$(B_{1,1}, B_{1,2}, B_{1,3}), (B_{2,1}, B_{2,2}, B_{2,3}), \dots, (B_{K,1}, B_{K,2}, B_{K,3})$$

of B such that all $B_{i,j}$ are distinct and $|B_{i,j}| \geq k + 1$. For every $i \in \{1, \dots, K\}$ and every $j \in \{1, 2, 3\}$, the three sets $S' \cup (B - B_{i,j})$, $S_1 \cup (B - B_{i,j+1})$, and $S_2 \cup (B - B_{i,j+2})$ (where j counts modulo 3) are in \mathcal{F} (by Corollary 3.5 and the fact that $|B - B_{i,j}| \leq n - s - (k + 1)$) and cover every vertex in N exactly twice. Using such triples for $i = 1, \dots, K$ and $j = 1, 2, 3$, we cover every vertex exactly $6K \geq 2^k \geq r$ times and every set appears at most once. If $r < 6K$ and is even, then we use not all triples.

If r is odd, then we pick a full pair $(A, N - A)$. There are at most two triples $(S' \cup (B - B_{i,j}), S_1 \cup (B - B_{i,j+1}), S_2 \cup (B - B_{i,j+2}))$ containing A or $N - A$. Then we cover the set N once by the set A and $N - A$ and $r - 1$ times with $\frac{r-1}{2} \leq 3K - 2$ triples $(S' \cup (B - B_{i,j}), S_1 \cup (B - B_{i,j+1}), S_2 \cup (B - B_{i,j+2}))$ containing neither A nor $N - A$. This contradicts the choice of \mathcal{F} . Therefore $|S| \geq n - 3k - 2$. ■

4. PROOF OF THEOREM 1.4

If the theorem does not hold, then for some $3 \leq r < n$, $k = \lceil \log_2 r \rceil$ with $n \geq r + 2k + 1$, and for some n -vertex set N , there exists an (N, r) -strange hypergraph \mathcal{F} . Let S be the union of 1-edges in \mathcal{F} . By Lemma 3.9, $|S| \leq r - 2 \leq n - 2k - 3$.

Let \mathcal{S}_{nf} denote the family of non-firm subsets of S . For every $S' \in \mathcal{S}_{nf}$, let

$$\mathcal{F}_{S'} := \{W \in \mathcal{F} : W \cap S = S'\}.$$

Furthermore, let

$$\mathcal{F}_{nf} := \bigcup_{S' \in \mathcal{S}_{nf}} \mathcal{F}_{S'}.$$

Lemma 4.1. *Let $M := \bigcup_{W \in \mathcal{F}_{nf}} W - S$. Then*

- (a) $|M| \leq r - s - 2$;
- (b) $|\mathcal{F}_{nf}| \leq r - 2$.

Proof. Assume that (a) does not hold and that w_1, \dots, w_{r-s-1} are in M . Let $M' := \{w_1, \dots, w_{r-s-1}\}$. For $j = 1, \dots, r - s - 1$, let W_j be a member of \mathcal{F}_{nf} such that $w_j \in W_j$, and let $S_j = W_j \cap S$. By Lemma 3.3, $W'_j := S_j + w_j$ is in \mathcal{F} for every $j = 1, \dots, r - s - 1$. Since each S_j and 1-edges are non-firm, $S - S_j$ and $S - v_j$ are firm. Also $|N - S - M'| = n - s - (r - s - 1) = n - r + 1 \geq 2k + 2$, thus by Corollary 3.5, every set of the form $S \cup M' - S_j - w_j$ or of the form $S \cup M' - v_i$ is in \mathcal{F} . So, every member of the family $\{S \cup M', \{v_1\}, \dots, \{v_s\}, S \cup M' - \{v_1\}, \dots, S \cup M' - \{v_s\}, W'_1, \dots, W'_{r-s-1}, S \cup M' - W'_1, \dots, S \cup M' - W'_{r-s-1}\}$

is in \mathcal{F} . Moreover, together they cover every vertex in $S \cup M'$ exactly r times. This proves (a).

Suppose now that W_1, \dots, W_{r-1} are in \mathcal{F}_{nf} . Since $|M| \leq r - s - 2$, every member of the family $\{S \cup M, W_1, \dots, W_{r-1}, S \cup M - W_1, \dots, S \cup M - W_{r-1}\}$ is in \mathcal{F} . Moreover, together they cover every vertex in $S \cup M$ exactly r times. This proves (b). ■

Remark 4.2. Since no member of \mathcal{F}_{nf} contains any element in $N - S - M$, for every $w \in N - M - S$, every subset of $N - S - w$ belongs to every firm $S' \subset S$.

Let S' be a smallest firm subset of S . By Lemma 3.8 S' is not an 1-edge. Choose a partition $S' = S_1 \cup S_2$ of S' into nonempty subsets. By the minimality of S' , sets S_1 and S_2 are not firm, and so $S - S_1$ and $S - S_2$ are firm.

Fix any element $z \in N - S - M$ and let $B := N - S - z$. Since $s \leq r - 2$, $|B| \geq n - (r - 2) - 1 \geq 2k + 2$. So, by Lemma 3.10, there are $K := \lceil 2^k/6 \rceil + 1$ partitions $(B_{1,1}, B_{1,2}, B_{1,3}), (B_{2,1}, B_{2,2}, B_{2,3}), \dots, (B_{K,1}, B_{K,2}, B_{K,3})$ of B such that all $B_{i,j}$ are distinct. For every $i \in \{1, \dots, K\}$ and every $j \in \{1, 2, 3\}$, the three sets $S' \cup (B - B_{i,j})$, $S_1 \cup (B - B_{i,j+1})$, and $S_2 \cup (B - B_{i,j+2})$ (where j counts modulo 3) are in \mathcal{F} (by Remark 4.2) and cover every vertex in $N - z$ exactly twice. Using such triples for $i = 1, \dots, K$ and $j = 1, 2, 3$, we cover every vertex exactly $6K \geq 2^k \geq r$ times and every set appears at most once. If $r < 6K$ and is even, then we use not all triples. If r is odd, then we pick a full pair $(A, N - A)$. Then we cover the set N once by the set A and $N - A$ and $r - 1$ times with the triples $(S' \cup (B - B_{i,j}), S_1 \cup (B - B_{i,j+1}), S_2 \cup (B - B_{i,j+2}))$ for $\frac{r-1}{2} (\leq 3K - 2)$ triples containing neither A nor $N - A$.

5. SIZE OF ALMOST \mathcal{F} -FREE SUBSETS

A set A is *almost \mathcal{F} -free* if every $B \in \mathcal{F}$ such that $B \subseteq A$ has size 1.

The aim of this section is to prove the following theorem.

Theorem 5.1. *If $n \geq 425$, then $|T| \leq n - 15k - 6$ for each almost \mathcal{F} -free $T \subseteq N$.*

Observe that for $n \geq 425$,

$$(6) \quad n - 15k - 6 \geq \frac{n}{2} > 0 \text{ and } n > (4k + 4)(\lceil \log(k) \rceil + 6) + 2k + 6.$$

We need some notation and lemmas. Let T be a maximum almost \mathcal{F} -free set, and $Q = N - T$. Assume that $|Q| < 15k + 6$, i.e., $|T| > n - 15k - 6$. For $Q' \subseteq Q$ and $T' \subseteq T$, we say that T' *belongs to* Q' if $Q' \cup T' \in \mathcal{F}$. A nonempty subset Q' of Q is *solid* if some $T' \subset T'$ with $|T'| \geq 3 + k$ belongs to Q' .

To show that Q is solid, let $B \subset T$ with $|B| = 2$. Since T is almost \mathcal{F} -free, $B \notin \mathcal{F}$. Then $N - B = (T - B) \cup Q \in \mathcal{F}$. By (6), $|T - B| \geq n/2 - |B| = n/2 - 2 \geq k + 3$, and so Q is solid.

Lemma 5.2. *Let $Q' \subseteq Q$ and $T' \subseteq T$. If T' belongs to Q' , then every $T'' \subset T'$ with $|T''| \leq |T'| - 2$ belongs to Q' .*

Proof. Since $T' \cup Q' \in \mathcal{F}$, by Lemma 2.2, either $Q' \cup T'' \in \mathcal{F}$ or $T - T'' \in \mathcal{F}$. But the latter does not hold, since T is almost \mathcal{F} -free. This proves the lemma. ■

Lemma 5.3. *For every partition $Q = Q' \cup Q''$ of Q into nonempty subsets, exactly one of Q' and Q'' is solid.*

Proof. Assume first that Q' is not solid. By (6), there exists a set $M \subset T$ with $|M| = 3 + k$. Since Q' is not solid, $Q' \cup M \notin \mathcal{F}$. Then $N - (Q' \cup M) \in \mathcal{F}$, and $N - (Q' \cup M) = Q'' \cup (T - M)$. So, since $|T - M| \geq n - 15k - 6 - (3 + k) \geq k + 3$, Q'' is solid.

Assume now that both Q' and Q'' are solid. We will show that if a set $M \subset T$ with $|M| \geq k + 3$ belongs to both Q' and Q'' , then \mathcal{F} has an r -regular subgraph with vertex set $Q \cup M$.

If $a \in M$, then the number of distinct subsets A_1, A_2, \dots, A_r of M containing a with $2 \leq |A_i| \leq |M| - 2$ is at least

$$2^{|M|-1} - (|M| + 1) = 2^{k+2} - k - 4 = 4r - k - 4 \geq r.$$

Note that $r \geq 2$, and $M - A_i \neq A_j$, since $a \in A_j$ and $a \notin M - A_j$. Let $\mathcal{H} = \{A_i \cup Q' : 1 \leq i \leq r\} \cup \{(M - A_i) \cup Q'' : 1 \leq i \leq r\}$. By construction, \mathcal{H} is r -regular, a contradiction.

If a set $M \subset T$ with $|M| = k + 4$ belongs to neither of Q' and Q'' , then $T - M$ belongs to both, and again \mathcal{F} has an r -regular subgraph. Thus each $M \subset T$ with $|M| = k + 4$ belongs to exactly one of Q' and Q'' . Let $\mathcal{R}_{Q'}$ (respectively, $\mathcal{R}_{Q''}$) denote the family of $M \subset T$ with $|M| = k + 4$ that belong to Q' (respectively, to Q''). By our assumption, both $\mathcal{R}_{Q'}$ and $\mathcal{R}_{Q''}$ are nonempty. Then there exist $M' \in \mathcal{R}_{Q'}$ and $M'' \in \mathcal{R}_{Q''}$ with $|M' \cap M''| = k + 3$. By Lemma 5.2, $M' \cap M''$ belongs to both Q' and Q'' , and so \mathcal{F} has an r -regular subgraph, a contradiction. ■

Corollary 5.4. *If Q' is a solid subset of Q , then every $M \subset T$ with $k + 3 \leq |M| \leq |T| - (k + 3)$ belongs to S' .*

Lemma 5.5. *The number of 1-edges not in full pairs of \mathcal{F} is at most k .*

Proof. Assume that there are $k + 1$ distinct 1-edges $\{a_1\}, \{a_2\}, \dots, \{a_{k+1}\}$ not in full pairs. If some nonempty $B \subset A = \{a_1, a_2, \dots, a_{k+1}\}$ is not in \mathcal{F} , then $\overline{B} \in \mathcal{F}$ by (2). Then \overline{B} together with 1-edges contained in B cover N once and none of these is in a full pair. These sets together with $r - 1$ full pairs cover N exactly r times, a contradiction. Thus every nonempty subset of A is in \mathcal{F} .

There are 2^k distinct nonempty subsets of A containing a_1 , call them B_1, B_2, \dots, B_{2^k} . Then all nonempty sets among $B_1, B_2, \dots, B_r, A - B_1, A - B_2, \dots, A - B_r$ are in \mathcal{F} , and they form an r -regular subgraph of \mathcal{F} , a contradiction. Therefore the number of 1-edges not in full pairs of \mathcal{F} is at most k . ■

Lemma 5.6. *The number of 1-edges in full pairs in \mathcal{F} is at least $n - 4k - 2$. Thus at most $8k - 2$ elements in full pairs are neither 1-edges nor $(n - 1)$ -edges.*

Proof. By Theorem 3.1, $|S| \geq n - 3k - 2$, so the number of 1-edges is at least $n - 3k - 2$. If fewer than $n - 4k - 2$ of them are in full pairs, then we get $k + 1$ distinct 1-edges a_1, a_2, \dots, a_{k+1} not in full pairs, a contradiction to Lemma 5.5. ■

Lemma 5.7. *For each $a \in Q$, there is $A \in \mathcal{F}$ with $2 \leq |A| \leq 3$ such that $\{a\} = A \cap Q$.*

Proof. Since T is a maximum almost \mathcal{F} -free set, $T \cup \{a\}$ is not almost \mathcal{F} -free. So, there is $B \subset T \cup \{a\}$ such that $B \in \mathcal{F}$ and $|B| \geq 2$. Take a smallest such B .

If $|B| = b \geq 4$, then there is $B' \subset B$ with $|B'| = b - 2 > 1$ and $B' \subset T$. Then $B' \notin \mathcal{F}$, and by Lemma 2.2, $B - B' \in \mathcal{F}$, so $A = B - B'$ is what we need. ■

Lemma 5.8. *The set Q contains at least one solid 1-edge.*

Proof. Let B be a smallest solid set in Q . Suppose $|B| \geq 2$. Then there are disjoint nonempty $B'_1, B'_2 \subset B$ with $B'_1 \cup B'_2 = B$. By Lemma 5.3, $B_1 = Q - B'_1$ and $B_2 = Q - B'_2$ are solid.

By (6), $T \geq n - 15k - 6 \geq 3k + 9$. Let $K := \lceil 2^k/6 \rceil + 1$. Similarly to the proofs of Lemma 3.10 and Corollary 3.11, for each $i = 1, 2, \dots, K$ there are partitions $(T_{i,1}, T_{i,2}, T_{i,3})$ of T such that all $T_{i,j}$ are distinct and $|T_{i,j}| \geq k + 3$ for all $i = 1, 2, \dots, K$ and $j = 1, 2, 3$.

For every $i \in \{1, \dots, K\}$ and every $j \in \{1, 2, 3\}$, the three sets $B \cup (T - T_{i,j})$, $B_1 \cup (T - T_{i,j+1})$, and $B_2 \cup (T - T_{i,j+2})$ (where j counts modulo 3) are in \mathcal{F} (by Corollary 5.4, and the fact that $|T - T_{i,j}| \leq |T| - (k + 3)$) and cover every vertex in N exactly twice. Using such triples for $i = 1, \dots, K$ and $j = 1, 2, 3$, we cover every vertex exactly $6K \geq 2^k \geq r$ times and every set appears at most once. If $r < 6K$ and is even, then we use not all triples.

If r is odd, then we pick a full pair $(A, N - A)$. There are at most two triples $(B \cup (T - T_{i,j}), B_1 \cup (T - T_{i,j+1}), B_2 \cup (T - T_{i,j+2}))$ containing A or $N - A$. Then we cover the set N once by the sets A and $N - A$ and $r - 1$ times by $\frac{r-1}{2} \leq 3K - 2$ triples $(B \cup (T - T_{i,j}), B_1 \cup (T - T_{i,j+1}), B_2 \cup (T - T_{i,j+2}))$ containing neither A nor $N - A$. This contradicts the choice of \mathcal{F} . ■

Lemma 5.9. $|Q| < 4k + 4$.

Proof. Suppose $|Q| \geq 4k+4$. By Lemma 5.8, Q contains a solid 1-edge $\{a\}$. Let $Q-a = \{b_1, b_2, \dots, b_{4k+3}, \dots, b_{|Q|-1}\}$. By Lemma 5.7, for each $i = 1, 2, \dots, 4k+3$, we can find B_i with $2 \leq |B_i| \leq 3$ such that $B_i \cap Q = \{b_i\}$. Let $L := N - a - \bigcup_{i=1}^{4k+3} B_i$. By definition, $|\bigcup_{i=1}^{4k+3} B_i| \leq 12k+9$. Since $n \geq 13k+13$, $|L| \geq 13k+3-1-(12k+9) = k+3$. Let $L' \subseteq L$ with $|L'| = k+3$. Let $M = N - L'$. Then \mathcal{F} contains at least $n - 4k - 5$ edges $\{a_1\}, \{a_2\}, \dots, \{a_{n-4k-5}\}$ such that all $M - a_i$ are also in \mathcal{F} , since $a \in M - a_i$ and $k+3 \leq |M - a_i| \leq |T| - k - 3$. Recall that for each $i = 1, \dots, 4k+3$, $B_i \in \mathcal{F}$ and $M - B_i \in \mathcal{F}$. Since $r \leq n-1$, the edges $\{a_1\}, \dots, \{a_{r+1-4k-5}\}, M - a_1, \dots, M - a_{r+1-4k-5}, B_1, \dots, B_{4k+3}, M - B_1, \dots, M - B_{4k+3}, M$ form an r -regular subgraph of \mathcal{F} , a contradiction. ■

Lemma 5.10. *If $\{a\}$ is a solid 1-edge and $B \in \mathcal{F}$ with $a \notin B$, then $|B \cap T| < \lceil \log k \rceil + 5$.*

Proof. If there is a set B with $a \notin B$ and $|B \cap T| \geq \lceil \log k \rceil + 5$, then by Lemma 5.2, we can find $B_1, B_2, \dots, B_{8k} \in \mathcal{F}$ such that $B_i \subset B$, $B \cap Q = B_i \cap Q$, since $2^{\lceil \log k \rceil + 4} - (\lceil \log k \rceil + 5) \geq 8k$. Let $X \subset N - (B \cup Q)$ with $|X| = k+3$ and let $M = N - X$. Since at least $n - 3k - 2 - (k+3) = n - 4k - 5$ of 1-edges $\{a_i\}$ are in M , the sets $B_1, B_2, \dots, B_{4k+4}, M - B_1, \dots, M - B_{4k+4}, \{a_1\}, \{a_2\}, \dots, \{a_{r-4k-5}\}, M - a_1, M - a_2, \dots, M - a_{r-4k-5}, M$ form an r -regular subgraph of \mathcal{F} , a contradiction. ■

Lemma 5.11. *There are at most $4k+3$ sets $A_i \in \mathcal{F}$ such that no A_i is a 1-edge and no solid 1-edge a is contained in A_i .*

Proof. Suppose that there are $4k+4$ such sets $A_1, A_2, \dots, A_{4k+4}$. Then by Lemma 5.10,

$$\left| T \cap \bigcup_{i=1}^{4k+4} A_i \right| \leq (4k+4)(\lceil \log k \rceil + 5) \leq |T| - k - 3.$$

Thus, as in the proof of Lemma 5.10, we can find an r -regular subgraph of \mathcal{F} by using A_i instead of B_i . ■

Lemma 5.12. *If $\{a\}$ is a solid 1-edge, then there is at most one $D \notin \mathcal{F}$ with $a \in D$.*

Proof. Suppose $D_1, D_2 \notin \mathcal{F}$ with $a \in D_1 \cap D_2$. By Lemma 5.10, $|D_i \cap T| \geq |T| - k + 3$ for $i = 1, 2$. So, $|D_1 \cap D_2 \cap T| \geq |T| - 2k - 6$.

By Lemmas 5.10 and 5.11, at least $|T| - (4k+4)(\log k + 6)$ elements in T are covered only by 1-edges and sets containing a .

By (6), $|T| - (4k+4)(\log k + 6) - 2k - 6 > 0$. So there is $c \in D_1 \cap D_2$ such that c is not covered by any edge of size at least 2 not containing a . Since \mathcal{F} is (n, r) -strange, Then at most $2^{n-1} + 1 - 2 = 2^{n-1} - 1$ edges of \mathcal{F} contain c . Thus

the family $\mathcal{F}_c = \{A \in \mathcal{F} : c \in A\}$ has at least $2^{n-2} + r - 1$ edges on $n - 1$ vertices, and by Theorem 1.2 we get an r -regular subgraph of \mathcal{F}' which is also a subgraph of \mathcal{F} , a contradiction. ■

Proof of Theorem 5.1. By Lemma 5.8, \mathcal{F} has a solid 1-edge $\{a\}$. By Lemma 5.12, there is at most one set $D \notin \mathcal{F}$ with $a \in D$. Since \mathcal{F} is (n, r) -strange, such D exists and exactly $r - 1$ edges of \mathcal{F} do not contain a , call them B_1, B_2, \dots, B_{r-1} .

Case 1. $\bigcup_{i=1}^{r-1} B_i = N - a$. Let l be the minimum integer such that we can renumber B_1, \dots, B_{r-1} so that $\bigcup_{i=1}^l B_i = N - a$. Let $\mathcal{B} = \{B_{l+1}, B_{l+2}, \dots, B_{r-1}\}$. Let $C_1 = B_1, C_2 = B_2 - B_1, C_3 = B_3 - B_2 - B_1, \dots, C_l = B_l - B_1 - B_2 - \dots - B_{l-1}$. By the minimality of l , $C_i \neq \emptyset$ for every $i = 1, \dots, l$. By construction, $\{C_1, \dots, C_l\}$ is a partition of $N - a$.

For every $i = 1, \dots, l$, there are $2^{|C_i|} - 2$ ways to choose a nonempty proper subset A of C_i . By Lemma 2.2, for each proper subset A of C_i , one of A and $B_i - A$ is in \mathcal{F} , and hence it is in \mathcal{B} . It follows that \mathcal{B} contains at least $\frac{1}{2}(2^{|C_i|} - 2) = 2^{|C_i|-1} - 1 \geq |C_i| - 1$ sets B such that (i) $0 < |B \cap C_i| < |C_i|$ and (ii) $B \cap C_j = \emptyset$ for all $i + 1 \leq j \leq l$. Since all C_i s are disjoint, we conclude that $|\mathcal{B}| \geq \sum_{i=1}^l (|C_i| - 1) = n - 1 - l$. Together with B_1, B_2, \dots, B_l , we have at least $n - 1$ members of \mathcal{F} not containing a . This contradicts the fact that \mathcal{F} has only $r - 1 \leq n - 2$ sets not containing a .

Case 2. There is $y \in N - a - \bigcup_{i=1}^{r-1} B_i$. Since $N - D \in \mathcal{F}$ and $a \notin N - D$, $y \notin N - D$. So, $y \in D$. Thus y belongs to at most $2^{n-2} - 1$ members of \mathcal{F} containing a and to none not containing a . So, the family $\mathcal{F}' = \mathcal{F} - y$ has at least $2^{n-1} + r - 2 - (2^{n-2} - 1) = 2^{n-2} + r - 1$ members. By Theorem 1.2, \mathcal{F}' has an r -regular subgraph, which is also a subgraph of \mathcal{F} , a contradiction. ■

6. PROOF OF THEOREM 1.5

Suppose \mathcal{F} is (n, r) -strange hypergraph on N . By Theorem 5.1,

(7) every $S \subseteq N$ with $|S| \geq n - 15k - 5$ contains some $A \in \mathcal{F}$ with $|A| \geq 2$.

Let B_1, B_2, \dots, B_l be the 1-edges not in full pairs. Let $N_1 = N - B_1 - B_2 - \dots - B_l$. By Lemma 5.5, $|N_1| \geq n - k$. So, by (7), N_1 contains some $B_{l+1} \in \mathcal{F}$ with $|B_{l+1}| \geq 2$. Then by Lemma 2.2, we can choose such B_{l+1} with $2 \leq |B_{l+1}| \leq 3$. Let $N_2 = N_1 - B_{l+1}$. Since $|N_2| \geq (n - k) - 3$, again by (7) and Lemma 2.2, N_2 contains some $B_{l+2} \in \mathcal{F}$ with $2 \leq |B_{l+2}| \leq 3$. Similarly, we find B_{l+3}, \dots, B_{5k+2} . Since at least $n - 4k - 2$ of 1-edges are in full pairs, by Lemma 5.6, at most $4k + 1$ full pairs have no 1-edges. Among the at most $8k + 2$ sets in these full pairs, at most $4k + 1$ of the sets are in $\{B_1, B_2, \dots, B_{5k+2}\}$, since $|B_i| \leq 3$ and $n \geq 425$. Thus some $k + 1$ sets among $B_1, B_2, \dots, B_{5k+2}$ are not in full pairs. Call them

A_1, A_2, \dots, A_{k+1} . Then for any $I \subset [k+1]$, $A_I = \bigcup_{i \in I} A_i$ is in \mathcal{F} , otherwise $\overline{A_I}$ and $\{A_j : j \in I\}$ together with $r-1$ full pairs yield an r -regular subgraph of \mathcal{F} . Therefore \mathcal{F} contains $2^{k+1-1} \geq r$ different pairs of edges of the kind $A_I, A_{[k+1]-I}$. They form an r -regular subgraph of \mathcal{F} covering $A_{[k+1]}$, a contradiction.

REFERENCES

- [1] N. Alon, S. Friedland, and G. Kalai, *Regular subgraphs of almost regular graphs*, J. Combin. Theory (B) **37** (1984) 79–91.
doi:10.1016/0095-8956(84)90047-9
- [2] M. Kano, *Regular subgraphs of a regular graph*, Annals of the New York Academy of Sciences **576** (1989) 281–284.
doi:10.1111/j.1749-6632.1989.tb16409.x
- [3] D. Mubayi and J. Verstraëte, *Two-regular subgraphs of hypergraphs*, J. Combin. Theory (B) **99** (2009) 643–655.
doi:10.1016/j.jctb.2008.10.005
- [4] L. Pyber, *Regular subgraphs of dense graphs*, Combinatorica **5** (1985) 347–349.
doi:10.1007/BF02579250
- [5] L. Pyber, V. Rödl, and E. Szemerédi, *Dense graphs without 3-regular subgraphs*, J. Combin. Theory (B) **63** (1995) 41–54.
doi:10.1006/jctb.1995.1004
- [6] V. Rödl and B. Wysocka, *Note on regular subgraphs*, J. Graph Theory **24** (1997) 139–154.
doi:10.1002/(SICI)1097-0118(199702)24:2<139::AID-JGT2>3.0.CO;2-R
- [7] V.A. Tashkinov, *3-regular subgraphs of 4-regular graphs*, Mat. Zametki **36** (1984) 239–259..
- [8] V.A. Tashkinov, *Regular parts of regular pseudographs*, Mat. Zametki **43** (1988) 263–275.

Received 20 June 2012

Revised 30 January 2013

Accepted 31 January 2013