

## PACKING TREES INTO $n$ -CHROMATIC GRAPHS

ANDRÁS GYÁRFÁS

*Alfréd Rényi Institute of Mathematics*

**e-mail:** gyarf. andras@mta.renyi.hu

### Abstract

We show that if a sequence of trees  $T_1, T_2, \dots, T_{n-1}$  can be packed into  $K_n$  then they can be also packed into any  $n$ -chromatic graph.

**Keywords:** tree packing.

**2010 Mathematics Subject Classification:** 05C70.

Let  $T_i$  denote a tree with  $i$  edges. The author's tree packing conjecture [3] states that  $K_n$  has an edge disjoint decomposition into any given sequence  $T_1, T_2, \dots, T_{n-1}$  of trees. Gerbner, Keszegh and Palmer extended the conjecture by replacing  $K_n$  with an arbitrary  $n$ -chromatic graph (Conjecture 2 in [1]). Here we show that the extended conjecture follows from the original one. We need the following, perhaps folklore result (Theorem 1 in [4]) and for convenience we include its simple proof.

**Lemma 1.** *Let  $G$  be a  $k$ -chromatic graph with a proper  $k$ -coloring with  $k$  distinct colors. Suppose  $T$  is a tree on  $k$  vertices and each vertex of  $T$  is labeled with a different color from the same set of  $k$  colors. Then  $G$  contains a subtree that is label-isomorphic to  $T$  (labeled exactly the same way as  $T$ ).*

**Proof.** The proof is by induction, the base step is trivial for  $k = 1$ . Let  $S_t$  denote the vertices of color  $t$  in a proper  $k$ -coloring of a  $k$ -chromatic graph  $G$  with a  $k$ -element color set  $C$ . Select a leaf vertex  $P$  in a tree  $T$  labeled with the  $k$  colors of  $C$ , suppose its label is  $i$  and assume  $P$  is adjacent in  $T$  with vertex  $Q$  labeled with  $j$ . Let  $A$  denote the set of vertices in  $S_j$  adjacent to at least one vertex of  $S_i$ . Observe that  $A$  is nonempty, otherwise  $G$  would be  $(k - 1)$ -chromatic. The subgraph  $G^* \subset G$  obtained by removing  $S_i$  and  $S_j - A$  from  $V(G)$  is  $(k - 1)$ -chromatic, since the removed vertices form an independent set. Also,  $G^*$  is colored with the color set  $C - i$ . By induction,  $G^*$  contains a label-isomorphic copy of the tree  $T - P$ , its vertex with color  $j$  is in  $A$ , thus adjacent to a vertex in  $S_i$ , extending  $T - P$  to a label-isomorphic copy of  $T$ . ■

**Theorem 2.** *Suppose that  $K_n$  has an edge disjoint decomposition into a given sequence  $T_1, T_2, \dots, T_{n-1}$  of trees and  $G$  is an  $n$ -chromatic graph. Then  $G$  contains edge disjoint copies of  $T_1, T_2, \dots, T_{n-1}$ .*

**Proof.** Let  $S_1, S_2, \dots, S_n$  be a partition of  $V(G)$  into independent sets where  $G$  is an  $n$ -chromatic graph and color all vertices of  $S_i$  with color  $i$ . By assumption the complete graph on vertex set  $V = \{1, 2, \dots, n\}$  can be decomposed into  $T_1, T_2, \dots, T_{n-1}$ . Let  $G_i$  be the subgraph of  $G$  induced by

$$\bigcup_{j \in V(T_i)} S_j.$$

The graph  $G_i$  is obviously  $(i+1)$ -chromatic since it has  $i+1$  color classes and a proper coloring of  $G_i$  with at most  $i$  colors could be obviously extended to a proper coloring of  $G$  with at most  $n-1$  colors, leading to a contradiction. Applying Lemma 1 to  $G_i$ , we find a copy  $F_i$  of  $T_i$  labeled exactly the same way as  $T_i$  is labeled in  $K_n$ . Repeating this for  $i = 1, 2, \dots, n-1$ , we obtain edge disjoint copies of  $F_1, F_2, \dots, F_{n-1}$  in  $G$ , in fact they are not only edge disjoint but the union of their edge sets takes exactly one edge from each bipartite graph  $\{[S_i, S_j] : 1 \leq i < j \leq n\}$ . ■

Theorem 2 allows to transfer known tree-packing results from  $K_n$  to  $n$ -chromatic graphs. In particular, since any sequence of trees containing only paths and stars are known to be packable to  $K_n$  ([3, 5]), we get the following, conjectured in [2].

**Corollary 3.** Any sequence  $T_1, T_2, \dots, T_{n-1}$  of stars and paths is packable into any  $n$ -chromatic graph.

#### REFERENCES

- [1] D. Gerbner, B. Keszegh and C. Palmer, *Generalizations of the tree packing conjecture*, Discuss. Math. Graph Theory **32** (2012) 569–582.  
doi:10.7151/dmgt.1628
- [2] D. Gerbner, B. Keszegh and C. Palmer, *Red-blue alternating paths*, Third Emléktábla Workshop, p.25.  
<http://www.renyi.hu/emlektab/index/booklet.html>
- [3] A. Gyárfás and J. Lehel, *Packing trees of different order into  $K_n$* , Combinatorics, Proc. Fifth Hungarian Coll. Keszthely, 1976, Vol II. North Holland. Colloq. Math. Soc. J. Bolyai **18** 463–469.
- [4] A. Gyárfás, E. Szemerédi and Zs. Tuza, *Induced subtrees in graphs of large chromatic number*, Discrete Math. **30** (1980) 235–244.  
doi:10.1016/0012-365X(80)90230-7

- [5] S. Zaks and C.L. Liu, *Decomposition of graphs into trees*, Proceedings of 8-th South-eastern Conference on Combinatorics, Graph Theory and Computing, Louisiana State Univ., Baton Rouge, La. Util. Math., Congr. Numer. **XIX** (1977) 643–654.

Received 13 June 2013

Accepted 26 June 2013