

5-STARS OF LOW WEIGHT IN NORMAL PLANE MAPS WITH MINIMUM DEGREE 5

OLEG V. BORODIN¹

*Institute of Mathematics Siberian Branch
Russian Academy of Sciences
and
Novosibirsk State University
Novosibirsk, 630090, Russia*

e-mail: brdnoleg@math.nsc.ru

ANNA O. IVANOVA²

*Institute of Mathematics of Ammosov
North-Eastern Federal University
Yakutsk, 677891, Russia*

e-mail: shmgnanna@mail.ru

AND

TOMMY R. JENSEN³

*Kyungpook National University
Republic of Korea*

e-mail: tjensen@knu.ac.kr

Abstract

It is known that there are normal plane maps M_5 with minimum degree 5 such that the minimum degree-sum $w(S_5)$ of 5-stars at 5-vertices is arbitrarily large. In 1940, Lebesgue showed that if an M_5 has no 4-stars of cyclic type $(5, 6, 6, 5)$ centered at 5-vertices, then $w(S_5) \leq 68$. We improve this bound of 68 to 55 and give a construction of a $(5, 6, 6, 5)$ -free M_5 with $w(S_5) = 48$.

Keywords: graph, plane map, vertex degree, weight, light subgraph.

2010 Mathematics Subject Classification: 05C10.

¹This work was supported by grant 12-01-00631 of the Russian Foundation for Basic Research.

²The author was supported by grants 12-01-00448 and 12-01-98510 of the Russian Foundation for Basic Research.

³This research was supported by the Kyungpook National University Research Fund, 2009.

1. INTRODUCTION

A normal plane map (NPM for short) is a plane pseudograph in which loops and multiple edges are allowed, but the degree of each vertex and face is at least three. The degree of a vertex v is denoted by $d(v)$. A k -vertex is a vertex v with $d(v) = k$. A k^+ -vertex (k^- -vertex) is one of degree at least k (at most k). An NPM with minimum degree δ at least 5 is denoted by M_5 . The *weight* of a subgraph of an NPM is the sum of degrees of its vertices. A k -star $S_k(v)$ is *minor* if its center v has degree (in the NPM) at most 5. All stars considered in this note are minor. By $w(S_k)$ we denote the minimum weight of minor k -stars in a given NPM.

In 1904, Wernicke [15] proved that every M_5 has a 5-vertex adjacent to a 6^- -vertex. This result was strengthened by Franklin [8] in 1922 to the existence of a 5-vertex with two 6^- -neighbors. In 1940, Lebesgue [14, p. 36] gave an approximate description of the neighborhoods of 5-vertices in M_5 s. In particular, this description implies the results in [15, 8] and shows that there is a 5-vertex with three 8^- -neighbors.

For M_5 s, the bounds $w(S_1) \leq 11$ (Wernicke [15]) and $w(S_2) \leq 17$ (Franklin [8]) are tight. It was proved by Lebesgue [14, p. 36] that $w(S_3) \leq 24$, which was improved in 1996 by Jendrol' and Madaras [11] to the tight bound $w(S_3) \leq 23$. Furthermore, Jendrol' and Madaras [11] gave a precise description of minor 3-stars in M_5 s.

For arbitrary NPMs, the following results concerning $(d - 2)$ -stars at d -vertices, $d \leq 5$, are known. Van den Heuvel and McGuinness [10] proved (in particular) that there is a vertex v such that either $w(S_1(v)) \leq 14$ with $d(v) = 3$, or $w(S_2(v)) \leq 22$ with $d(v) = 4$, or $w(S_3(v)) \leq 29$ with $d(v) = 5$. Balogh *et al.* [1] proved that there is a 5^- -vertex adjacent to at most two 11^+ -vertices. Harant and Jendrol' [9] strengthened these results by proving (in particular) that either $w(S_1(v)) \leq 13$ with $d(v) = 3$, or $w(S_2(v)) \leq 19$ with $d(v) = 4$, or $w(S_3(v)) \leq 23$ with $d(v) = 5$. Recently, we obtained a precise description of $(d - 2)$ -stars in NPMs [6].

For M_5 s, Lebesgue [14, p. 36] proved $w(S_4) \leq 31$, which was improved by Borodin and Woodall [3] to the tight bound $w(S_4) \leq 30$. Note that $w(S_3) \leq 23$ easily implies $w(S_2) \leq 17$ and immediately follows from $w(S_4) \leq 30$ (in both cases, it suffices to delete a vertex of maximum degree from a minor star of the minimum weight). Recently, we obtained a precise description of 4-stars in M_5 s [7].

For arbitrary NPMs, the problem of describing $(d - 1)$ -stars at d -vertices, $d \leq 5$, called *pre-complete stars*, appears difficult. As follows from the double n -pyramid, the minimum weight $w(S_{d-1})$ of pre-complete stars in NPMs can be arbitrarily large. Even when $w(S_{d-1})$ is restricted by appropriate conditions,

the tight upper bounds on it are unknown. Borodin *et al.* [4, 5] proved (in particular) that if a planar graph with $\delta \geq 3$ has no edge joining two 4^- -vertices, then there is a star $S_{d-1}(v)$ with $w(S_{d-1}(v)) \leq 38 + d(v)$, where $d(v) \leq 5$ (see [5, Theorem 2.A]). Jendrol' and Madaras [12] proved that if the weight of every edge in a planar graph with $\delta \geq 3$ is at least 9, then there is a pre-complete star in which every vertex has degree at most 20, where the bound of 20 is best possible.

The more general problem of describing d -stars at d -vertices, $d \leq 5$, called *complete stars*, at the moment seems intractable for arbitrary NPMs and difficult even for M_5 s. In this note we make a modest contribution for the case of M_5 s.

The following well-known construction shows that $w(S_5)$ is unbounded in M_5 s. Take three concentric n -cycles $C^i = v_1^i \dots v_n^i$, where n is large and $1 \leq i \leq 3$, and join C^2 with C^1 by edges $v_j^2 v_j^1$ and $v_j^2 v_{j+1}^1$ whenever $1 \leq j \leq n$ (addition modulo n). The same is done with C^2 and C^3 . Finally, join all vertices of C^1 to a new n -vertex, and do the same with C^3 .

Definition. A 5-vertex v surrounded by vertices v_1, \dots, v_5 in a cyclic order is a $(5, 6, 6, 5)$ -vertex if there is a k , $k \leq 5$, such that $d(v_{k+1}) = d(v_{k+4}) = 5$, $d(v_{k+2}) \leq 6$, and $d(v_{k+3}) \leq 6$ (addition modulo 5).

Clearly, each 5-vertex in the M_5 constructed above is a $(5, 6, 6, 5)$ -vertex and, moreover, it has two 5-neighbors and two 6-neighbors. Lebesgue [14, p. 36] proved that if an M_5 has no $(5, 6, 6, 5)$ -vertices, then $w(S_5) \leq 68$.

The purpose of this note is to improve the bound of 68 to 55 (Theorem 1) and give a construction of a $(5, 6, 6, 5)$ -free M_5 with $w(S_5) = 48$ (see Figure 1). We first truncate all vertices of the dodecahedron, and then join the mid-points of the edges of each triangle and put a 2-vertex on each edge not incident with a triangle. Finally, we insert a 20-wheel inside every 20-face as shown in Figure 1. As a result, every 5-vertex has neighbors of degrees 5, 6, 7, 5, and 20 in this order.

Theorem 1. *If a normal plane map M_5 with minimum degree 5 does not contain $(5, 6, 6, 5)$ -vertices, then M_5 contains a 5-star of weight at most 55 with a 5-vertex as its center.*

Problem 2. Find best possible version of Theorem 1.

2. PROOF OF THEOREM 1

It suffices to prove the theorem for triangulations, since adding a diagonal edge into a non-triangular face of a normal plane map with $\delta = 5$ cannot create a new minor 5-star, nor can it reduce the weight of any existing minor 5-star. So




Figure 1. An M_5 without $(5, 6, 6, 5)$ -vertices such that $w(S_5) = 48$.

suppose that a triangulation T , with its sets of vertices, edges, and faces denoted by V , E , and F , respectively, is a counterexample to Theorem 1. Euler's formula $|V| - |E| + |F| = 2$ for T implies

$$(1) \quad \sum_{v \in V} (d(v) - 6) = -12.$$

Assign an *initial charge* $\mu(v) = d(v) - 6$ to each $v \in V$, so that only 5-vertices have negative initial charge. Also, put $\mu(f) = 0$ for each $f \in F$. Using the properties of our T as a counterexample, we define a local redistribution of charges, preserving their sum, such that the final charge $\mu_3(x)$ is non-negative for all $x \in V \cup F$. This will contradict the fact that the sum of the final charges is, by (1), equal to -12 . The technique of discharging is often used in solving structural and coloring problems on plane graphs.

Definition. For integer n , $n \geq 6$, put $\xi(n) = \frac{n-6}{n}$. For any 6^+ -vertex v , put $\psi(v) = \xi(d(v))$.

In what follows, it is convenient to use the upwards convexity of the increasing function $\xi(n)$ for integer $n \geq 6$, which is checked easily.

Lemma 3. For any integers p and q , where $6 \leq p < q$, we have

$$\xi(p) + \xi(q) \leq \xi(p+1) + \xi(q-1).$$

The *final charge* $\mu_3(v)$ of vertex v is defined by applying the rules R1–R3 as follows.

R1. Each 6^+ -vertex v sends $\psi(v)$ to each incident 3-face.

R2. Let $f = xyz$ be a face such that $d(x) = 5$ and $d(z) \geq 6$. Then x receives from f the following charge:

- (a) $\frac{\psi(z)}{2}$ if $d(y) = 5$, or
- (b) $\psi(y) + \psi(z)$ if $d(y) \geq 6$.

The charge of x , where $x \in V \cup F$, after applying R1 and R2 is denoted by $\mu_2(x)$.

Definition. A 5-vertex v surrounded by vertices v_1, \dots, v_5 in a cyclic order is *bad* if $d(v_1) = d(v_2) = d(v_4) = 5$ and $d(v_3) = 7$.

Note that each bad 5-vertex v satisfies $d(v_5) \geq 29$ since $w(S_5) \geq 56$ by assumption.

R3. Suppose v is bad, and let the neighbors of v_2 and v_1 in the cyclic order be v, v_3, x, y, v_1 and v, v_2, y, z, v_5 , respectively (see Figure 2).

(a) If $d(x) = 5$ (which means that v_2 is also bad), then v_1 gives $\frac{1}{14}$ to each of v and v_2 .

(b) If $d(x) \geq 6$ and v_1 is also bad (that is $d(y) = 7$, and $d(z) = 5$), then v_2 gives $\frac{1}{14}$ to each of v and v_1 .

(c) If $d(x) \geq 6$ but v_1 is not bad, then v_2 gives $\frac{1}{14}$ only to v .


Clearly, $\mu_3(v) = \mu_2(v) = 0$ for each 6^+ -vertex v , and $\mu_3(f) = \mu_2(f) \geq 0$ for each face f . It remains to show that every 5-vertex receives at least 1 in total by R1–R3. If v is a bad 5-vertex, then $\mu_2(v) \geq 5 - 6 + \xi(7) + \xi(28) = -\frac{1}{14}$, which implies that $\mu_3(v) \geq 0$.

Our next goal is to show that every non-bad 5-vertex v satisfies $\mu_2(v) \geq 0$, and then we will complete the proof of Theorem 1 by checking that in fact $\mu_3(v) \geq 0$.

Remark 4. Suppose a 5-vertex v is adjacent to a 6^+ -vertex w ; then v receives from w by R1 and R2 one of the charges $\psi(w)$, $\frac{3\psi(w)}{2}$, or $2\psi(w)$ depending on the number of 5-vertices in the two 3-faces incident with edge vw : three, two, or one, respectively.

Definition. The *type* of a 5-vertex v is the vector (d_1, \dots, d_5) of the degrees of the neighbors of v in the non-decreasing order. (So, d_1 is the smallest degree among the neighbors of v , d_2 is the second smallest, and so on.)

Lemma 5. If v is a non-bad 5-vertex, then $\mu_2(v) \geq 0$.


$$(2) \quad \frac{3}{2}\xi(27) = \frac{3}{2} \cdot \frac{21}{27} = \frac{7}{6} > \frac{8}{7} > 1.$$

Now our proof splits into four cases.

Case 2. $5 = d_1 < d_2$. By Remark 4, $\mu_{1,2}^+(v) \geq \frac{3}{2}(\xi(d_2) + \cdots + \xi(d_5))$, which is smallest when $d_2 = \cdots = d_4 = 6$ and $d_5 = 28$. Thus $\mu_{1,2}^+(v) \geq \frac{3}{2}\xi(28) > 1$ by (2).

Case 3. $5 = d_1 = d_2 < d_3$. If the two 5-neighbors of v form a 3-face with v , then $\mu_{1,2}^+(v) \geq \frac{3}{2}(\xi(d_3) + \xi(d_4) + \xi(d_5))$, which is smallest when $d_3 = d_4 = 6$ and $d_5 = 29$. Thus $\mu_{1,2}^+(v) \geq \frac{3}{2}\xi(29) > 1$ by (2).

So assume that the two 5-neighbors of v do not form a 3-face with v . By Remark 4, $\mu_{1,2}^+(v) \geq \frac{3}{2}(\xi(a) + \xi(b)) + \xi(c)$, where a, b, c is a permutation of d_3, d_4, d_5 such that the a -vertex and b -vertex form a 3-face with v . Recall that v is not a $(5, 6, 6, 5)$ -vertex by assumption. So if $d_3 = d_4 = 6$ then $c = 6$ and $\max\{a, b\} \geq 29$; thus $\mu_{1,2}^+(v) \geq \frac{3}{2}\xi(29) > 1$ by (2). Otherwise, $d_4 \geq 7$ and $\mu_{1,2}^+(v) \geq \frac{3}{2}\xi(7) + \xi(28) = \frac{3}{2} \cdot \frac{1}{7} + \frac{22}{28} = 1$.

Case 4. $5 = d_1 = d_2 = d_3 < d_4$. Then $d_4 + d_5 \geq 36$. If $6 \leq d_4 \leq 7$, then the d_4 -vertex and the d_5 -vertex form a 3-face with v , since v is neither bad nor a $(5, 6, 6, 5)$ -vertex by assumption; thus $\mu_{1,2}^+(v) \geq \frac{3}{2}\xi(d_5) \geq \frac{3}{2}\xi(29) > 1$ by (2). If $d_4 \geq 8$, then $\mu_{1,2}^+(v) \geq \xi(8) + \xi(28) = \frac{29}{28} > 1$. ■

Lemma 6. *If v is a non-bad 5-vertex, then $\mu_3(v) \geq 0$.*

Proof. If v gives nothing to bad 5-vertices by R3, then $\mu_3(v) = \mu_2(v) \geq 0$ by Lemma 5. So we may assume that v gives either $\frac{1}{14}$ or $\frac{1}{7}$ in total to bad 5-vertices by R3. It suffices to prove that $\mu_{1,2}^+(v) \geq \frac{8}{7}$.

We may assume that v is either the vertex v_1 in R3(a) or the vertex v_2 in R3(b) or R3(c). In both cases, $d(x) + d(y) \geq 34$ since $w(S_5(v_2)) \geq 56$. In the first case, $d(x) = 5$ and $d(y) \geq 29$ (see Figure 2), and so $\mu_{1,2}^+(v) \geq 2\xi(29) > \frac{8}{7}$ by (2). In the second case, $\mu_{1,2}^+(v) \geq \frac{3}{2}(\xi(6) + \xi(28)) = \frac{3}{2}\xi(28) > \frac{8}{7}$ by (2). ■

Thus we have proved $\mu_3(x) \geq 0$ for every $x \in V \cup F$, which contradicts (1) and completes the proof of Theorem 1.

Acknowledgements

The authors are greatly indebted to the anonymous referees for suggestions on improving the presentation.

REFERENCES

- [1] J. Balogh, M. Kochol, A. Pluhár and X. Yu, *Covering planar graphs with forests*, J. Combin. Theory (B) **94** (2005) 147–158.
doi:10.1016/j.jctb.2004.12.002
- [2] O.V. Borodin, *Solution of Kotzig's and Grünbaum's problems on the separability of a cycle in a planar graph*, Mat. Zametki **46(5)** (1989) 9–12 (in Russian).
- [3] O.V. Borodin and D.R. Woodall, *Short cycles of low weight in normal plane maps with minimum degree 5*, Discuss. Math. Graph Theory **18** (1998) 159–164.
doi:10.7151/dmgt.1071

- [4] O.V. Borodin, H.J. Broersma, A.N. Glebov and J. Van den Heuvel, *The structure of plane triangulations in terms of clusters and stars*, Diskretn. Anal. Issled. Oper. Ser. 1 **8(2)** (2001) 15–39 (in Russian).
- [5] O.V. Borodin, H.J. Broersma, A.N. Glebov and J. Van den Heuvel, *Minimal degrees and chromatic numbers of squares of planar graphs*, Diskretn. Anal. Issled. Oper. Ser. 1 **8(4)** (2001) 9–33 (in Russian).
- [6] O.V. Borodin and A.O. Ivanova, *Describing $(d - 2)$ -stars at d -vertices, $d \leq 5$, in normal plane maps*, Discrete Math. **313** (2013) 1700–1709.
doi:10.1016/j.disc.2013.04.026
- [7] O.V. Borodin and A.O. Ivanova, *Describing 4-stars at 5-vertices in normal plane maps with minimum degree 5*, Discrete Math. **313** (2013) 1710–1714.
doi:10.1016/j.disc.2013.04.025
- [8] P. Franklin, *The four colour problem*, Amer. J. Math. **44** (1922) 225–236.
doi:10.2307/2370527
- [9] J. Harant and S. Jendrol', *On the existence of specific stars in planar graphs*, Graphs Combin. **23** (2007) 529–543.
doi:10.1007/s00373-007-0747-7
- [10] J. Van den Heuvel and S. McGuinness, *Coloring the square of a planar graph*, J. Graph Theory **42** (2003) 110–124.
doi:10.1002/jgt.10077
- [11] S. Jendrol' and T. Madaras, *On light subgraphs in plane graphs of minimal degree five*, Discuss. Math. Graph Theory **16** (1996) 207–217.
doi:10.7151/dmgt.1035
- [12] S. Jendrol' and T. Madaras, *Note on an existence of small degree vertices with at most one big degree neighbour in planar graphs*, Tatra Mt. Math. Publ. **30** (2005) 149–153.
- [13] A. Kotzig, *From the theory of eulerian polyhedra*, Mat. Čas. **13** (1963) 20–34 (in Russian).
- [14] H. Lebesgue, *Quelques conséquences simples de la formule d'Euler*, J. Math. Pures Appl. **19** (1940) 27–43.
- [15] P. Wernicke, *Über den kartographischen Vierfarbensatz*, Math. Ann. **58** (1904) 413–426.
doi:10.1007/BF01444968

Received 21 January 2013

Revised 10 June 2013

Accepted 10 June 2013