Discussiones Mathematicae Graph Theory 34 (2014) 801–810 doi:10.7151/dmgt.1762

AN IMPLICIT WEIGHTED DEGREE CONDITION FOR HEAVY CYCLES¹

JUNQING CAI

School of Management, Qufu Normal University Rizhao, 276826, China e-mail: caijq09@163.com

Hao Li^2

Institute for Interdisciplinary Research Jianghan University, Wuhan, 430019, China LRI, UMR 8623, CNRS and Université de Paris-Sud 11 F-91405 Orsay, France

e-mail: li@lri.fr

AND

WANTAO NING

Department of Mathematics and Statistic, Xidian University Xi'an, Shaanxi 710071, China

e-mail: ningwt05@163.com

Abstract

For a vertex v in a weighted graph G, $id^w(v)$ denotes the implicit weighted degree of v. In this paper, we obtain the following result: Let G be a 2-connected weighted graph which satisfies the following conditions: (a) The implicit weighted degree sum of any three independent vertices is at least t; (b) w(xz) = w(yz) for every vertex $z \in N(x) \cap N(y)$ with $xy \notin E(G)$; (c) In every triangle T of G, either all edges of T have different weights or all edges of T have the same weight. Then G contains either a hamiltonian cycle or a cycle of weight at least 2t/3. This generalizes the result of Zhang et al. [9].

Keywords: weighted graph, hamiltonian cycles, heavy cycles, implicit degree, implicit weighted degree.

2010 Mathematics Subject Classification: 05C38.

¹This paper is partially supposed by the Scientific Research Foundation for Doctors in Qufu Normal University (2012015) and NNSF of China (11326218).

²Corresponding author.

1. INTRODUCTION

In this paper, we consider only finite, undirected and simple graphs. Notation and terminology not defined here can be found in [2]. Let G = (V(G), E(G))be a graph with vertex set V(G) and edge set E(G). Let $H \subseteq G$. For a vertex $u \in V(G), N_H(u) = \{v \in V(H) : uv \in E(G)\}$ and $d_H(u) = |N_H(u)|$. If H = G, we always use N(u) and d(u) in place of $N_G(u)$ and $d_G(u)$ respectively. $N_2(v) = \{u \in V(G) : d(u, v) = 2\}$, where d(u, v) denotes the distance between vertices u and v in G

Based on the traditional definition of degree, Zhu, Li and Deng [10] introduced the concept of implicit degrees.

Definition [10]. Let v be a vertex of a graph G and k = d(v) - 1. Set $M_2 = \max\{d(u) : u \in N_2(v)\}$ and $m_2 = \min\{d(u) : u \in N_2(v)\}$. Suppose $d_1 \leq d_2 \leq d_3 \leq \cdots \leq d_k \leq d_{k+1} \leq \cdots$ is the degree sequence of vertices in $N(v) \cup N_2(v)$. If $N_2(v) \neq \emptyset$ and $d(v) \geq 2$, define

$$d^{*}(v) = \begin{cases} m_{2}, & \text{if } d_{k} < m_{2}, \\ d_{k+1}, & \text{if } d_{k+1} > M_{2}, \\ d_{k}, & \text{otherwise}, \end{cases}$$

then the *implicit degree* of v is defined as $id(v) = \max\{d(v), d^*(v)\}$. If $N_2(v) = \emptyset$ or $d(v) \leq 1$, then id(v) = d(v).

Clearly, $id(v) \ge d(v)$ for each vertex v from the definition of implicit degree.

For a graph G, if we assign a non-negative number w(e) to every edge e, then G is called a weighted graph and w(e) is the weight of e. Clearly, an unweighted graph can be regarded as a weighted graph in which each edge is assigned a weight 1. The weight of a subgraph H of G and the weighted degree of a vertex v in G are defined as

$$w(H) = \sum_{e \in E(H)} w(e)$$
 and $d^w(v) = \sum_{u \in N(v)} w(uv)$, respectively.

Based on the idea of the definition of implicit degree, Li [7] gave the definition of implicit weighted degrees as follows.

Definition [7]. Let v be a vertex of a weighted graph G and k = d(v) - 1. Set $m_2^w = \min\{d^w(u) : u \in N_2(v)\}$ and $M_2^w = \max\{d^w(u) : u \in N_2(v)\}$. Suppose $d_1^w \leq d_2^w \leq \cdots \leq d_{k+1}^w \leq \cdots$ is the weighted degree sequence of vertices in $N(v) \cup N_2(v)$. If $N_2(v) \neq \emptyset$ and $d(v) \geq 2$, define

$$d^{w*}(v) = \begin{cases} m_2^w, & \text{if } d_k^w < m_2^w, \\ d_{k+1}^w, & \text{if } d_{k+1}^w > M_2^w, \\ d_k^w, & \text{otherwise,} \end{cases}$$

then the implicit weighted degree of v is defined as $id^w(v) = \max\{d^w(v), d^{w*}(v)\}$. If $N_2(v) = \emptyset$ or $d(v) \leq 1$, then $id^w(v) = d^w(v)$.

Clearly, $id^w(v) \ge d^w(v)$ for every vertex v from the above definition.

Let $\alpha(G)$ be the independent number of a graph G. For a positive integer $k \leq \alpha(G)$, we define $\sigma_k(G) = \min\{\sum_{j=1}^k d(x_j) : x_1, x_2, \ldots, x_k \text{ are } k \text{ independent vertices in } G\}$ and $\sigma_k^*(G) = \min\{\sum_{j=1}^k id(x_j) : x_1, x_2, \ldots, x_k \text{ are } k \text{ independent vertices in } G\}$. For a weighted graph G, let $\sigma_k^w(G) = \min\{\sum_{j=1}^k d^w(x_j) : x_1, x_2, \ldots, x_k \text{ are } k \text{ independent vertices in } G\}$ and $\sigma_k^{w*}(G) = \min\{\sum_{j=1}^k id^w(x_j) : x_1, x_2, \ldots, x_k \text{ are } k \text{ independent vertices in } G\}$ and $\sigma_k^{w*}(G) = \min\{\sum_{j=1}^k id^w(x_j) : x_1, x_2, \ldots, x_k \text{ are } k \text{ independent vertices in } G\}$. If $k > \alpha(G)$, then they are all equal to $+\infty$.

A graph G is called hamiltonian if it has a hamiltonian cycle, i.e., a cycle that contains all vertices of G. There are many results about the existence of hamiltonian cycles and long cycles in graphs in terms of the degree sum of independent vertices. The following two theorems are famous.

Theorem 1 [4]. If G is a 2-connected graph with $\sigma_1(G) \ge c/2$, then G contains either a hamiltonian cycle or a cycle of length at least c.

Theorem 2 [8]. If G is a 2-connected graph with $\sigma_2(G) \ge c$, then G contains either a hamiltonian cycle or a cycle of length at least c.

After studying $\sigma_1(G)$ and $\sigma_2(G)$ conditions, Fournier and Fraisse [6] generalized them into degree conditions on more independent vertices, i.e., the $\sigma_k(G)$.

Theorem 3 [6]. If G is a k-connected graph $(k \ge 2)$ with $\sigma_{k+1}(G) \ge c$, then G contains either a hamiltonian cycle or a cycle of length at least 2c/(k+1).

Zhang, Li and Broersma [9] gave a result about heavy cycles on weighted degree condition. The result extended Theorem 3 in the case k = 2 by adding two extra conditions.

- (C1) w(xz) = w(yz) for every vertex $z \in N(x) \cap N(y)$ with $xy \notin E(G)$;
- (C2) For every triangle T of G, either all edges of T have different weights or all edges of T have the same weight.

Theorem 4 [9]. Suppose G is a 2-connected weighted graph satisfying conditions (C1) and (C2). If $\sigma_3^w(G) \ge t$, then G contains either a hamiltonian cycle or a cycle of weight at least 2t/3.

Motivated by the result of Theorem 4, we obtain the following result.

Theorem 5. Let G be a 2-connected weighted graph satisfying conditions (C1) and (C2). If $\sigma_3^{w*}(G) \ge t$, then G contains either a hamiltonian cycle or a cycle of weight at least 2t/3.



Figure 1. Example 1.

We give the proof of Theorem 5 in the next section. Here we give a graph G with $\sigma_3^{w*}(G) > \sigma_3^w(G)$. It is said that we can get a heavier cycle by using Theorem 5 than using Theorem 4.

Example. Let G be a graph shown in Figure 1, where K_m $(m \ge 2)$ is a complete graph and x, z are adjacent to every vertex of each $K_m, N(y) = \{x, z\}$. We assign weight 2 to each edge of G. It is easy to verify that $\sigma_3^w(G) = 4m + 8$. And by the definition of implicit weighted degree, we get that $\sigma_3^{w*}(G) = 6(m + 1)$.

2. Proof of Theorem 5

Our proof of Theorem 5 is based on the following three lemmas.

Lemma 6 [5]. Let G be a connected weighted graph satisfying conditions (C1) and (C2). Then either

- (a) all edges of G have the same weight, or
- (b) G is a complete multipartite graph.

Lemma 7 [3]. Let G be a k-connected graph with at least three vertices. If $k \ge \alpha(G)$, then G is hamiltonian.

Lemma 8. Let G be a 2-connected graph such that $\sigma_3^*(G) \ge c$, then G contains either a hamiltonian cycle or a cycle of length at least 2c/3.

We give the proof of Lemma 8 in the next section.

Proof of Theorem 5. Let G be a weighted graph satisfying the conditions of Theorem 5. If $\alpha(G) \leq 2$, then G is hamiltonian by Lemma 7.

Hence we assume $\alpha(G) \geq 3$. Then by Lemma 6, either all edges of G have the same weight or G is a complete multipartite graph.

First we suppose all edges of G have the same weight m. When m = 0, there is nothing to say. Suppose $m \neq 0$. By the definitions of implicit degree and implicit weighted degree, we have $id^w(v) = m(id(v))$ for each $v \in V(G)$. Hence, $\sigma_3^*(G) = \sigma_3^{w*}(G)/m \geq t/m$. Then, G contains either a hamiltonian cycle or a cycle C of length at least 2t/3m by Lemma 8. If G is not hamiltonian , then $w(C) = m \times |E(C)| \geq m \times (2t/3m) = 2t/3$.

Now, we assume that G is a complete multipartite graph and G is not hamiltonian. Let |V(G)| = n and V_1, V_2, \ldots, V_l be a partition of V(G).

Claim 2.1 [5]. If $x, y \in V_i$, then w(xz) = w(yz) for every $z \in V(G) \setminus V_i$. In particular, $d^w(x) = d^w(y)$.

Claim 2.2 [5]. If G is not hamiltonian, then $|V_i| > n/2$ for some i with $1 \le i \le l$.

We can assume, without loss of generality, that $|V_1| > n/2$ by Claim 2.2. Let $p = |V_1|$ and q = n - p. Since G is 2-connected, $2 \le q < p$. And let $V_1 = \{v_1, v_2, \ldots, v_p\}$ and $V(G) \setminus V_1 = \{u_1, u_2, \ldots, u_q\}$.

Claim 2.3. $id^w(x) = id^w(y)$ for any two distinct vertices $x, y \in V_i$.

Proof. Since $x, y \in V_i$, $N(x) = N(y) = V(G) \setminus V_i$ and $N_2(x) \setminus \{y\} = N_2(y) \setminus \{x\} = V_i \setminus \{x, y\}$. By Claim 2.1, we know $d^w(x) = d^w(y)$ and the weighted degree sequences of vertices in $N(x) \cup N_2(x)$ and those of vertices in $N(y) \cup N_2(y)$ are the same. Then by the definition of implicit weighted degree, we can get that $id^w(x) = id^w(y)$.

Claim 2.4. $id^w(v) \ge t/3$ for each $v \in V_1$.

Proof. Since p > 2, there are three vertices v_1, v_2 and v_3 in V_1 and $\{v_1, v_2, v_3\}$ is an independent set. Hence $id^w(v_1) + id^w(v_2) + id^w(v_3) \ge t$. We assume, without loss of generality, that $id^w(v_1) \ge t/3$. Then $id^w(v) = id^w(v_1) \ge t/3$ for each $v \in V_1$ by Claim 2.3.

Claim 2.5. $id^w(v) = d^w(v)$ for any $v \in V_1$.

Proof. Suppose there is some $v \in V_1$ such that $d^w(v) < id^w(v)$. Since G is a complete multipartite graph, $N_2(v) = V_1 \setminus \{v\}$ and $N(v) = V(G) \setminus V_1$. Since $|V_1| > n/2, |N_2(v)| \ge |N(v)|$. Let $k = d(v) - 1, M_2^w = \max\{d^w(u) : u \in N_2(v)\}$ and $m_2^w = \min\{d^w(u) : u \in N_2(v)\}$. Suppose $d_1^w \le d_2^w \le \cdots \le d_{k+1}^w \le \cdots$ is the weighted degree sequence of vertices of $N(v) \cup N_2(v)$. By Claim 2.1 and the definition of $id^w(v)$, we have $M_2^w = m_2^w, d_1^w \le d_2^w \le \cdots \le d_{k+1}^w < id^w(v)$ and $id^w(v) \ne m_2^w$. Which is contrary to the definition of $id^w(v)$. So $id^w(v) = d^w(v)$ for any $v \in V_1$. By Claims 2.4 and 2.5, we can get that $d^w(v) \ge t/3$ for each $v \in V_1$. Now, we consider the cycle $C = v_1 u_1 v_2 u_2 \cdots v_q u_q v_1$. Then Claim 2.1 implies

$$w(C) = \sum_{i=1}^{q} w(v_i u_i) + \sum_{i=1}^{q-1} w(v_{i+1} u_i) + w(v_1 u_q)$$

= $\sum_{i=1}^{q} w(v_1 u_i) + \sum_{i=1}^{q-1} w(v_1 u_i) + w(v_1 u_q)$
= $2 \sum_{i=1}^{q} w(v_1 u_i) = 2d^w(v_1).$

Hence, $w(C) \ge 2t/3$. This completes the proof of Theorem 5.

3. Proof of Lemma 8

Let $P = x_1 x_2 \cdots x_p$ be a path of a graph G, for any $I \subseteq V(P)$, define $I^- = \{x_i : x_{i+1} \in I\}$ and $I^+ = \{x_i : x_{i-1} \in I\}$. We use $L_P(x_1)$ to denote the vertex x_i such that $x_1 x_i \in E(G)$ and $x_1 x_j \notin E(G)$ for any j > i on P. To prove Lemma 8 we need the following two lemmas.

Lemma 9 [1]. Let G be a 2-connected non-hamiltonian graph and $P = x_1 x_2 \cdots x_p$ be a longest path of G. Then G contains a cycle of length at least $d(x_1) + d(x_p)$.

Lemma 10 [10]. Let G be a 2-connected graph and $P = x_1x_2\cdots x_p$ be a longest path of G. If $d(x_1) < id(x_1)$ and $x_1x_p \notin E(G)$, then either

- (1) there is some $x_j \in N_P(x_1)^-$ such that $d(x_j) \ge id(x_1)$; or
- (2) $N_P(x_1)^- = N_P(x_1) \cup \{x_1\} \setminus \{L_P(x_1)\}$ and $id(x_1) = \min\{d(v) : v \in N_2(x_1)\}.$

Proof of Lemma 8. Let G be a 2-connected graph satisfying the conditions of Lemma 8 and suppose G is not hamiltonian. Then by Lemma 9, it suffices to prove that there exists a longest path $P = v_1 v_2 \cdots v_p$ in G such that $d(v_1) + d(v_p) \ge 2c/3$.

Suppose to the contrary that $d(v_1) + d(v_p) < 2c/3$ for any longest path $P = v_1v_2\cdots v_p$. Choose a longest path $P = v_1v_2\cdots v_p$ such that $d(v_1) + d(v_p)$ is as large as possible. Then $N(v_1) \cup N(v_p) \subseteq V(P)$ and there is no cycle of length p in G. We can assume, without loss of generality, that $d(v_1) < c/3$. Since G is 2-connected, $|N_P(v_1)| \ge 2$ and $|N_P(v_p)| \ge 2$. Let $k = \max\{i : v_1v_i \in E(G)\}$ and $l = \min\{j : v_pv_j \in E(G)\}$. Then $3 \le k \le p-1$ and $2 \le l \le p-2$.

Case 1. $N(v_1) = \{v_2, v_3, \dots, v_k\}.$

Since $G - v_k$ is connected, there exists an edge $v_r v_s \in E(G)$ with r < k < s. Choose such an edge $v_r v_s$ such that s is as large as possible. By the choice of P, we have $N(v_r) \subseteq V(P)$. Let

$$P_1 = v_r v_{r-1} \cdots v_1 v_{r+1} v_{r+2} \cdots v_p.$$

806

Since G has no cycle of length $p, v_r v_p \notin E(G)$. Next, we are going to look for a special vertex $v_j \in V(P)$ such that $\{v_r, v_j, v_p\}$ is an independent set of G.

Subcase 1.1. There is some j with $r+1 \leq j \leq s-1$ such that $v_r v_{j+1} \in E(G)$ and $v_r v_j \notin E(G)$. We consider the longest path $P_2 = v_j v_{j-1} \cdots v_{r+1} v_1 v_2 \cdots v_r v_{j+1}$ $v_{j+2} \cdots v_p$. Since G has no cycle of length p, we have $v_j v_p \notin E(G)$. Then we have found the vertex v_j .

Subcase 1.2. $v_r v_i \in E(G)$ for each i with $r+1 \leq i \leq s$. By the choice of P, $r \geq 4$ and there must exist some j with $2 \leq j \leq r-2$ such that $v_r v_j \notin E(G)$. (Since, otherwise, P_1 is a longest path with $d(v_r) + d(v_p) > d(v_1) + d(v_p)$, a contradiction.) We consider the longest path $P_3 = v_j v_{j-1} \cdots v_1 v_{j+1} v_{j+2} \cdots v_p$. Then $v_j v_p \notin E(G)$. So we find the vertex v_j .

Claim 3.1. $id(v_r) \le d(v_1)$ and $id(v_j) \le d(v_1)$.

Proof. Suppose $id(v_r) > d(v_1)$. We consider the longest path P_1 defined as before. If $d(v_r) = id(v_r)$, then $d(v_r) + d(v_p) > d(v_1) + d(v_p)$, a contradiction.

Suppose $d(v_r) < id(v_r)$. For convenience, we set $P_1 = y_1y_2\cdots y_p$, where $y_1 = v_r, y_2 = v_{r-1}, \ldots, y_p = v_p$. Since $v_rv_s \in E(G)$ and $v_rv_j \notin E(G)$, we have $N_{P_1}(v_r)^- \neq N_{P_1}(v_r) \cup \{v_r\} \setminus \{L_{P_1}(v_r)\}$. Hence, by Lemma 10, there exists some $y_a \in N_{P_1}(y_1)^-$ such that $d(y_a) \geq id(y_1)$. Then $y_ay_{a-1}\cdots y_1y_{a+1}y_{a+2}\cdots y_p$ is a longest path of G with $d(y_a) + d(y_p) \geq id(y_1) + d(y_p) = id(v_r) + d(v_p) > d(v_1) + d(v_p)$, which is contrary to the choice of P. Hence $id(v_r) \leq d(v_1)$.

Suppose $id(v_j) > d(v_1)$. If v_j is got in Subcase 1.1, we consider P_2 . If v_j is got in Subcase 1.2, we consider P_3 (where P_2 and P_3 are defined as before). No matter what cases happen, we have $d(v_r, v_j) = 2$. By the choice of P, we get that $d(v_r) \le d(v_1)$. So $id(v_j) \ne \min\{d(u) : u \in N_2(v_j)\}$. By similar arguments as above, we get contradictions. So $id(v_j) \le d(v_1)$.

Since $\{v_r, v_j, v_p\}$ is an independent set of G, we have $id(v_r) + id(v_j) + id(v_p) \ge c$. So $id(v_p) \ge c - id(v_r) - id(v_j) \ge c - 2d(v_1)$ by Claim 3.1. If $d(v_p) = id(v_p)$, then $d(v_1) + d(v_p) = d(v_1) + id(v_p) \ge c - d(v_1) > 2c/3$, a contradiction. Hence we can assume $d(v_p) < id(v_p)$.

Claim 3.2. $N(v_p) = \{v_l, v_{l+1}, \dots, v_{p-1}\}$ and $d(v_i) < id(v_p)$ for any $v_i \in N_P(v_p)^+$.

Proof. Suppose to the contrary that there exists some $v_i \in N_P(v_p)^+$ such that $d(v_i) \geq id(v_p)$. Then $v_1v_2\cdots v_{i-1}v_pv_{p-1}\cdots v_i$ is a longest path of G different from P with $d(v_1) + d(v_i) \geq d(v_1) + id(v_p) > d(v_1) + d(v_p)$, which is contrary to the choice of P. Therefore, Claim 3.2 holds by Lemma 10.

By Claim 3.2, we can check that $l \geq s$, otherwise, $v_1v_2\cdots v_rv_sv_{s+1}\cdots v_pv_{s-1}$ $v_{s-2}\cdots v_{r+1}v_1$ is cycle of length p, a contradiction. Since $G-v_l$ is connected, there exists an edge $v_{r'}v_{s'} \in E(G)$ such that s' < l < r'. Choose such an edge $v_{r'}v_{s'}$ such that s' is as small as possible. We can get that $v_rv_{r'} \notin E(G)$ and $v_jv_{r'} \notin E(G)$ for $v_rv_{r-1}\cdots v_1v_{r+1}v_{r+2}\cdots v_{r'-1}v_pv_{p-1}\cdots v_{r'}$ and $v_jv_{j-1}\cdots v_{r+1}v_1v_2\cdots v_rv_{j+1}v_{j+2}\cdots v_{r'-1}v_pv_{p-1}\cdots v_{r'}$ (v_j is got in Subcase 1.1) or $v_jv_{j-1}\cdots v_1v_{j+1}v_{j+2}\cdots v_{r'-1}v_pv_{p-1}\cdots v_{r'}$ (v_j is got in Subcase 1.2) are longest paths of G. So $\{v_r, v_j, v_{r'}\}$ is an independent set of G. Hence, $id(v_r) + id(v_j) + id(v_{r'}) \geq c$. Then, by Claim 3.1, $id(v_{r'}) \geq c - id(v_r) - id(v_j) \geq c - 2d(v_1)$.

Considering the following longest path

$$P' = v_{r'}v_{r'+1}\cdots v_p v_{r'-1}v_{r'-2}\cdots v_{s'+1}v_{s'}\cdots v_1,$$

we can claim $d(v_{r'}) < id(v_{r'})$. If not, $d(v_{r'}) + d(v_1) \ge c - 2d(v_1) = c - d(v_1) > 2c/3$, a contradiction.

We can get that $v_{r'}$ is not adjacent to some vertex $v_d \in N(v_p)$. If not, we have $d(v_{r'}) > d(v_p)$. Then, for the longest path P', $d(v_{r'}) + d(v_1) > d(v_p) + d(v_1)$, which is contrary to the choice of P. Now $v_{r'}v_d \notin E(G)$ and $v_{r'}v_{s'} \in E(G)$, which imply that $N_{P'}(v_{r'})^- \neq N_{P'}(v_{r'}) \cup \{v_{r'}\} \setminus \{L_{P'}(v_{r'})\}$. For convenience, let $P' = x_1x_2\cdots x_p$ such that $x_1 = v_{r'}, x_2 = v_{r'+1}, \ldots, x_p = v_1$. By Lemma 10, there exists some vertex $x_b \in N_{P'}(x_1)^-$ such that $d(x_b) \geq id(x_1)$. Then $x_bx_{b-1}\cdots x_1x_{b+1}x_{b+2}\cdots x_p$ is a longest path of G with $d(x_b) + d(x_p) \geq id(x_1) + d(x_p) = id(v_{r'}) + d(v_1) \geq c - d(v_1) > 2c/3$, which is contrary to the choice of P.

Case 2. $N(v_1) \neq \{v_2, v_3, \dots, v_k\}.$

Choose $v_j \notin N(v_1)$ with j < k such that j is as large as possible. Then $v_1v_i \in E(G)$ for each i with $j < i \leq k$. Then $\{v_1, v_j, v_p\}$ is an independent set of G. If $d(v_1) < id(v_1)$, then there exists some $v_d \in N_P(v_1)^-$ such that $d(v_d) \ge id(v_1) > d(v_1)$ by Lemma 10. Then $v_dv_{d-1}\cdots v_1v_{d+1}v_{d+2}\cdots v_p$ is a longest path of G with $d(v_d) + d(v_p) > d(v_1) + d(v_p)$, a contradiction.

Next we assume $id(v_1) = d(v_1)$. By similar arguments as in Claim 3.1, we have $id(v_j) \leq d(v_1)$. If $d(v_p) = id(v_p)$, then $id(v_1) + id(v_j) + id(v_p) \geq c$, which implies $id(v_p) \geq c - id(v_1) - id(v_j) \geq c - 2d(v_1)$. Hence $d(v_1) + d(v_p) = d(v_1) + id(v_p) \geq c - d(v_1) > 2c/3$, a contradiction.

Suppose $d(v_p) < id(v_p)$. If $N(v_p) \neq \{v_l, v_{l+1}, \ldots, v_{p-1}\}$, then there is a vertex $v_f \in N_P(v_p)^+$ such that $d(v_f) \ge id(v_p)$ by Lemma 10. So $v_f v_{f+1} \cdots v_p v_{f-1} v_{f-2} \cdots v_1$ is a longest path with $d(v_1) + d(v_f) \ge d(v_1) + id(v_p) > d(v_1) + d(v_p)$, which is contrary to choice of P.

Hence $N(v_p) = \{v_l, v_{l+1}, \ldots, v_{p-1}\}$. By the 2-connectivity of G, we know that there exists an edge $v_r v_s \in E(G)$ with r < l < s. We can easily verify that $\{v_1, v_j, v_s\}$ is an independent set of G. Then $id(v_1) + id(v_j) + id(v_s) \ge c$. Therefore, $id(v_s) \ge c - id(v_1) - id(v_j) \ge c - 2d(v_1)$.

Now for the longest path $P_5 = v_s v_{s+1} \cdots v_p v_{s-1} v_{s-2} \cdots v_1$, we claim that there exists a vertex $v_i \in N(v_p)$ such that $v_i v_s \notin E(G)$. Otherwise, $d(v_s) > d(v_p)$ and hence $d(v_1) + d(v_s) > d(v_1) + d(v_p)$, which is contrary to the choice of P.

Since $v_i v_s \notin E(G)$ and $v_s v_r \in E(G)$, $N_{P_5}(v_s)^- \neq N_{P_5}(v_s) \cup \{v_s\} \setminus \{L_{P_5}(v_s)\}$. For convenience, we let $P_5 = z_1 z_2 \cdots z_p$ with $z_1 = v_s, z_2 = v_{s+1}, \ldots, z_p = v_1$. By Lemma 10, there is a vertex $z_g \in N_{P_5}(z_1)^-$ such that $d(z_g) \geq id(z_1)$. Then $z_g z_{g-1} \cdots z_1 z_{g+1} z_{g+2} \cdots z_p$ is a longest path of G with $d(z_g) + d(z_p) \geq id(z_1) + d(z_p) = id(v_s) + d(v_1) \geq (c - 2d(v_1)) + d(v_1) > 2c/3$, a contradiction.

Now we complete the proof of Lemma 8.

Acknowledgements

The authors are very grateful to anonymous referee whose helpful comments and suggestions have led to a substantially improvement of the paper.

References

- J.A. Bondy, Large cycles in graphs, Discrete Math. 1 (1971) 121–132. doi:10.1016/0012-365X(71)90019-7
- [2] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications (Macmillan-London, Elsevier-New York, 1976).
- [3] V. Chvátal and P. Erdős, A note on hamiltonian circuits, Discrete Math. 2 (1972) 111–113. doi:10.1016/0012-365X(72)90079-9
- [4] G.A. Dirac, Some theorems on abstract graphs, Proc. Lond. Math. Soc. 2 (1952) 69-81.
- [5] H. Enomoto, J. Fujisawa and K. Ota, $A \sigma_k$ type condition for heavy cycles in weighted graphs, Ars Combin. **76** (2005) 225–232.
- [6] I. Fournier and P. Fraisse, On a conjecture of Bondy, J. Combin. Theory (B) 39 (1985) 17–26.
 doi:10.16/0095-8956(85)90035-8
- [7] P. Li, Implicit weighted degree condition for heavy paths in weighted graphs, J. Shandong Univ. (Nat. Sci.) 18 (2003) 11–13.
- [8] L. Pósa, On the circuits of finite graphs, Magyar Tud. Akad. Mat. Kutató Int. Közl 8 (1963) 355–361.
- [9] S. Zhang, X. Li and H. Broersma, A σ₃ type condition for heavy cycles in weighted graphs, Discuss. Math. Graph Theory **21** (2001) 159–166. doi:10.7151/dmgt.1140
- [10] Y. Zhu, H. Li and X. Deng, *Implicit-degrees and circumferences*, Graphs Combin. 5 (1989) 283–290. doi:10.1007/BF01788680

J. CAI, H. LI AND W. NING

Received 31 October 2011 Revised 18 November 2013 Accepted 18 November 2013