Discussiones Mathematicae Graph Theory 35 (2015) 261–270 doi:10.7151/dmgt.1797

# THE *k*-RAINBOW BONDAGE NUMBER OF A DIGRAPH

JAFAR AMJADI, NEGAR MOHAMMADI

## Seyed Mahmoud Sheikholeslami

Department of Mathematics Azarbaijan Shahid Madani University Tabriz, I.R. Iran

e-mail: {j-amjadi;s.m.sheikholeslami}@azaruniv.edu

AND

LUTZ VOLKMANN

Lehrstuhl II für Mathematik RWTH Aachen University 52056 Aachen, Germany

e-mail: volkm@math2.rwth-aachen.de

#### Abstract

Let D = (V, A) be a finite and simple digraph. A k-rainbow dominating function (kRDF) of a digraph D is a function f from the vertex set V to the set of all subsets of the set  $\{1, 2, \ldots, k\}$  such that for any vertex  $v \in V$ with  $f(v) = \emptyset$  the condition  $\bigcup_{u \in N^-(v)} f(u) = \{1, 2, \ldots, k\}$  is fulfilled, where  $N^-(v)$  is the set of in-neighbors of v. The weight of a kRDF f is the value  $\omega(f) = \sum_{v \in V} |f(v)|$ . The k-rainbow domination number of a digraph D, denoted by  $\gamma_{rk}(D)$ , is the minimum weight of a kRDF of D. The k-rainbow bondage number  $b_{rk}(D)$  of a digraph D with maximum in-degree at least two, is the minimum cardinality of all sets  $A' \subseteq A$  for which  $\gamma_{rk}(D-A') > \gamma_{rk}(D)$ . In this paper, we establish some bounds for the k-rainbow bondage number and determine the k-rainbow bondage number of several classes of digraphs.

**Keywords:** *k*-rainbow dominating function, *k*-rainbow domination number, *k*-rainbow bondage number, digraph.

2010 Mathematics Subject Classification: 05C69.

#### 1. INTRODUCTION

Let D be a finite simple digraph with vertex set V(D) = V and arc set A(D) = A. A digraph without directed cycles of length 2 is an *oriented graph*. The order n = n(D) of a digraph D is the number of its vertices. We write  $\deg_D^+(v) =$  $\deg^+(v)$  for the *outdegree* of a vertex v and  $\deg^-_D(v) = \deg^-(v)$  for its *indegree*. The minimum and maximum indegree and minimum and maximum outdegree of D are denoted by  $\delta^- = \delta^-(D), \ \Delta^- = \Delta^-(D), \ \delta^+ = \delta^+(D)$  and  $\Delta^+ = \Delta^+(D), \ \delta^+ = \delta^+(D), \ \delta^+ = \delta^+(D), \ \delta^- = \delta^-(D), \ \delta^- = \delta^$ respectively. If (u, v) is an arc of D, then we also write  $u \to v$ , and we say that v is an *out-neighbor* of u and u is an *in-neighbor* of v. For a vertex v of a digraph D, we denote the set of in-neighbors and out-neighbors of v by  $N^{-}(v) = N_{D}^{-}(v)$ and  $N^+(v) = N_D^+(v)$ , respectively. If  $X \subseteq V(D)$ , then D[X] is the subdigraph induced by X. If  $X \subseteq V(D)$  and  $v \in V(D)$ , then E(X, v) is the set of arcs from X to v. The underlying graph G[D] of a digraph D is the graph obtained by replacing each arc uv by an edge uv. Note that G[D] has two parallel edges uvwhen D contains the arcs (u, v) and (v, u). A digraph D is called *connected*, if the underlying graph G[D] is connected. For the notation and terminology not defined here, we refer the reader to [11].

Let k be a positive integer. A k-rainbow dominating function (kRDF) of a digraph D is a function f from the vertex set V(D) to the set of all subsets of the set  $\{1, 2, \ldots, k\}$  such that for any vertex  $v \in V(D)$  with  $f(v) = \emptyset$  the condition  $\bigcup_{u \in N^-(v)} f(u) = \{1, 2, \ldots, k\}$  is fulfilled. The weight of a kRDF f is the value  $\omega(f) = \sum_{v \in V} |f(v)|$ . The k-rainbow domination number of a digraph D, denoted by  $\gamma_{rk}(D)$ , is the minimum weight of a kRDF of D. A  $\gamma_{rk}(D)$ function is a k-rainbow dominating function of D with weight  $\gamma_{rk}(D)$ . Note that  $\gamma_{r1}(D)$  is the classical domination number  $\gamma(D)$ . The k-rainbow domination numbers in digraphs were investigated by Amjadi et al. in [1]. A 2-rainbow dominating function (briefly, rainbow dominating function)  $f: V \longrightarrow \mathcal{P}(\{1,2\})$ can be represented by the ordered partition  $(V_0, V_1, V_2, V_{1,2})$  (or  $(V_0^f, V_1^f, V_2^f, V_{1,2}^f)$ to refer f) of V, where  $V_0 = \{v \in V \mid f(v) = \emptyset\}$ ,  $V_1 = \{v \in V \mid f(v) = \{1\}\}$ ,  $V_2 = \{v \in V \mid f(v) = \{2\}\}$  and  $V_{1,2} = \{v \in V \mid f(v) = \{1,2\}\}$ . In this representation, its weight is  $\omega(f) = |V_1| + |V_2| + 2|V_{1,2}|$ .

**Proposition A** [1]. Let D be a digraph of order n. Then  $\gamma_{r2}(D) < n$  if and only if  $\Delta^+(D) \ge 2$  or  $\Delta^-(D) \ge 2$ .

**Proposition B** [1]. Let  $k \ge 1$  be an integer. If D is a digraph of order n, then

$$\min\{k, n\} \le \gamma_{rk}(D) \le n.$$

**Proposition C** [1]. Let D be a digraph of order  $n \ge 2$ . Then  $\gamma_{r2}(D) = 2$  if and only if n = 2 or  $n \ge 3$  and  $\Delta^+(D) = n - 1$  or there exist two different vertices u and v such that  $V(D) - \{u, v\} \subseteq N^+(u)$  and  $V(D) - \{u, v\} \subseteq N^+(v)$ . **Proposition D** [1]. Let  $k \ge 1$  be an integer. If D is a digraph of order n, then

$$\gamma_{rk}(D) \le n - \Delta^+(D) + k - 1.$$

The definition of the k-rainbow dominating function for undirected graphs was introduced by Brešar, Henning, and Rall [2] and has been studied by several authors (see for example [3, 4, 5, 9, 10, 12, 13]).

Following the ideas in [7], we initiate the study of k-rainbow bondage number on digraphs D. The k-rainbow bondage number  $b_{rk}(D)$  of a digraph D is the cardinality of a smallest set of arcs  $A' \subseteq A(D)$  for which  $\gamma_{rk}(D - A') > \gamma_{rk}(D)$ . An edge set B with  $\gamma_{rk}(D - B) > \gamma_{rk}(D)$  is called the k-rainbow bondage set. A  $b_{rk}(D)$ -set is a k-rainbow bondage set of D of size  $b_{rk}(D)$ . If B is a  $b_{rk}(D)$ -set, then clearly

(1) 
$$\gamma_{rk}(D-B) = \gamma_{rk}(D) + 1.$$

By Proposition A, we note that if D is a digraph with  $\Delta^+(D) \leq 1$  and  $\Delta^-(D) \leq 1$ , then  $\gamma_{r2}(D) = n$  and hence if  $A' \subseteq A(D)$ , then  $\gamma_{r2}(D - A') = \gamma_{r2}(D)$ . Therefore the 2-rainbow bondage number is only defined for a digraph with maximum indegree or maximum out-degree at least two.

The definition of the k-rainbow bondage number for undirected graphs was given by Dehgardi, Sheikholeslami and Volkmann [6].

The purpose of this paper is to establish some bounds for the k-rainbow bondage number of a digraph.

**Observation 1.** Let D be a digraph of order n with  $\gamma_{rk}(D) < n$ . Assume that H is a spanning subdigraph of D with  $\gamma_{rk}(H) = \gamma_{rk}(D)$ . If K = A(D) - A(H), then  $b_{rk}(H) \leq b_{rk}(D) \leq b_{rk}(H) + |K|$ .

**Proof.** Let  $F \subseteq A(D)$  be a  $b_{rk}(D)$ -set. It follows that  $\gamma_{rk}(H-F) \ge \gamma_{rk}(D-F) > \gamma_{rk}(D) = \gamma_{rk}(H)$  and hence  $b_{rk}(H) \le |F| = b_{rk}(D)$ .

Now let  $F' \subseteq A(H)$  be a  $b_{rk}(H)$ -set. We deduce that  $\gamma_{rk}(D - (K \cup F')) = \gamma_{rk}(H - F') > \gamma_{rk}(H) = \gamma_{rk}(D)$  and thus  $b_{rk}(D) \leq b_{rk}(H) + |K|$ .

**Observation 2.** If a digraph D has a vertex v such that every  $\gamma_{rk}(D)$ -function assigns a set of size at least 2 to v, then  $b_{rk}(D) \leq \deg^+(v) \leq \Delta^+$ .

**Proof.** Assume that  $A_v^+$  is the set of arcs in D with tail v and let f be a  $\gamma_{rk}(D - A_v^+)$ -function. Since  $N_{D-A_v^+}^+(v) = \emptyset$ , we deduce that  $|f(v)| \leq 1$  and hence f is not a  $\gamma_{rk}(D)$ -function. Thus  $\gamma_{rk}(D - A_v^+) > \gamma_{rk}(D)$ , and the proof is complete.

**Theorem 3.** Let k be a positive integer and let D be a digraph of order  $n \ge k+1$ . If the underlying graph of D is connected, then

$$b_{rk}(D) \le (\gamma_{rk}(D) - k + 1)\Delta(G[D]).$$

**Proof.** By Proposition B,  $\gamma_{rk}(D) \geq k$ . We proceed by induction on  $\gamma_{rk}(D)$ . If  $\gamma_{rk}(D) = k$ , then let u be a vertex in D, and let  $A_u$  denote the set of arcs incident with u. Since  $n \geq k + 1$ , we deduce from Proposition B that  $\gamma_{rk}(D - A_u) = 1 + \gamma_{rk}(D - u) \geq k + 1 > \gamma_{rk}(D)$ . This implies that  $b_{rk}(D) \leq |A_u| = \deg_{G[D]}(u)$  and hence  $b_{rk}(D) \leq \Delta(G[D])$ .

Now assume that the statement is true for any digraph of order  $n \geq k+1$ with k-rainbow domination number  $k \leq \gamma_{rk}(D) \leq s$ . Assume that D is a digraph of order  $n \geq k+1$  with  $\gamma_{rk}(D) = s+1$ . Suppose to the contrary that  $b_{rk}(D) >$  $(\gamma_{rk}(D) - k + 1)\Delta(G[D]) > \Delta(G[D])$ . Let u be an arbitrary vertex of D, and let  $A_u$  denote the set of arcs incident with u. Then  $\gamma_{rk}(D) = \gamma_{rk}(D - A_u)$ , because  $\deg_{G[D]}(u) < b_{rk}(D)$ . Let f be a  $\gamma_{rk}(D - A_u)$ -function. Obviously, |f(u)| = 1 and the function f restricted to D - u is a  $\gamma_{rk}(D - u)$ -function. This yields  $\gamma_{rk}(D - u) = \gamma_{rk}(D) - 1$ . It follows from Observation 1 that  $b_{rk}(D) \leq$  $b_{rk}(D - u) + \deg_{G[D]}(u)$ , and by the induction hypothesis we obtain

$$b_{rk}(D) \leq b_{rk}(D-u) + \deg_{G[D]}(u)$$
  

$$\leq (s-k+1)\Delta(G[D-u]) + \deg_{G[D]}(u)$$
  

$$\leq (s-k+1)\Delta(G[D]) + \Delta(G[D])$$
  

$$= ((s+1)-k+1)\Delta(G[D]) = (\gamma_{rk}(D)-k+1)\Delta(G[D]).$$

This contradiction completes the proof.

## 2. Upper Bounds on the 2-Rainbow Bondage Number

In this section we mainly present bounds on the 2-rainbow bondage number of a digraph.

**Theorem 4.** If D is a digraph, and xyz a path of length 2 in G[D] such that  $(y, x), (y, z) \in A(D)$ , then

(2) 
$$b_{r2}(D) \le \deg_{G[D]}(x) + \deg_{G[D]}(y) + \deg_{G[D]}(z) - 3 - |N^{-}(x) \cap N^{-}(y)|.$$

Moreover, if x and z are adjacent in G[D], then

(3) 
$$b_{r2}(D) \le \deg_{G[D]}(x) + \deg_{G[D]}(y) + \deg_{G[D]}(z) - 4 - |N^{-}(x) \cap N^{-}(y)|.$$

**Proof.** Let  $A_1$  be the set of arcs incident with x, y or z with the exception of (y, z) and all arcs going from  $N^-(x) \cap N^-(y)$  to y. Then

$$|A_1| \le \deg_{G[D]}(x) + \deg_{G[D]}(y) + \deg_{G[D]}(z) - 3 - |N^-(x) \cap N^-(y)|$$

and

$$|A_1| \le \deg_{G[D]}(x) + \deg_{G[D]}(y) + \deg_{G[D]}(z) - 4 - |N^-(x) \cap N^-(y)|$$

when x and z are adjacent. Now let  $D_1 = D - A_1$ . Obviously in  $D_1$ , the vertex x is isolated, z is a vertex with indegree 1, y is an in-neighbor of z, and all inneighbors of y in  $D_1$ , if any, are contained in  $N^-(x)$ . Let  $f = (V_0, V_1, V_2, V_{1,2})$  be a  $\gamma_{r2}(D_1)$ -function. Then |f(x)| = 1 and  $|f(z)| \leq 1$ .

If  $f(z) = \emptyset$ , then  $f(y) = \{1, 2\}$  and therefore  $(V_0 \cup \{x\}, V_1 - \{x\}, V_2 - \{x\}, V_{1,2})$  is a 2RDF on D of weight less than  $\omega(f)$ , and consequently (2) as well as (3) are proved.

Now assume that |f(z)| = 1. If |f(y)| = 1, then  $(V_0 \cup \{z\}, V_1 - \{y, z\}, V_2 - \{y, z\}, V_{1,2} \cup \{y\})$  is also a  $\gamma_{r2}(D_1)$ -function, and we are in the situation discussed in the previous case. However, if  $f(y) = \emptyset$ , then there exists a vertex  $w \in N^-(x) \cap N^-(y)$  such that  $f(w) = \{1, 2\}$  or there exist two vertices  $w_1, w_2 \in N^-(x) \cap N^-(y)$  such that  $f(w_1) = \{1\}$  and  $f(w_2) = \{2\}$ . Since  $w, w_1$  and  $w_2$  are in-neighbors of x in D,  $(V_0 \cup \{x\}, V_1 - \{x\}, V_2 - \{x\}, V_{1,2})$  is a 2RDF on D of weight less than f, and the proof is complete.

**Theorem 5.** If D is a digraph, and xyz a path of length 2 in G[D] such that  $(y, x), (y, z) \in A(D)$ , then

(4) 
$$b_{r2}(D) \le \deg_{G[D]}(x) + \deg^{-}(y) + \deg_{G[D]}(z) - |N^{-}(x) \cap N^{-}(y) \cap N^{-}(z)|.$$

Moreover, if x and z are adjacent in G[D], then

(5)  $b_{r2}(D) \le \deg_{G[D]}(x) + \deg^{-}(y) + \deg_{G[D]}(z) - 1 - |N^{-}(x) \cap N^{-}(y) \cap N^{-}(z)|.$ 

**Proof.** Let F be the set of arcs incident with x or z and all arcs terminating in y except the arcs  $w \to y$  for which the arcs  $w \to x$  and  $w \to z$  also occur in D. Then

$$|F| \le \deg_{G[D]}(x) + \deg^{-}(y) + \deg_{G[D]}(z) - |N^{-}(x) \cap N^{-}(y) \cap N^{-}(z)|$$

and

$$|F| \le \deg_{G[D]}(x) + \deg^{-}(y) + \deg_{G[D]}(z) - 1 - |N^{-}(x) \cap N^{-}(y) \cap N^{-}(z)|$$

when x and z are adjacent. Let now D' = D - F. In D', the vertices x, z are isolated, and all in-neighbors of y in D', if any, are contained in  $N^{-}(x) \cap N^{-}(z)$ . Let  $f = (V_0, V_1, V_2, V_{1,2})$  be a  $\gamma_{r2}(D')$ -function. Then |f(x)| = |f(z)| = 1 and we may assume, without loss of generality, that  $f(x) = f(z) = \{1\}$ .

If  $f(y) = \{1, 2\}$ , then  $(V_0 \cup \{x, z\}, V_1 - \{x, z\}, V_2, V_{1,2})$  is a 2RDF on D of weight less than  $\omega(f)$ , and therefore (4) and (5) are proved.

If |f(y)| = 1, then  $(V_0 \cup \{x, z\}, V_1 - \{x, y, z\}, V_2 - \{y\}, V_{1,2} \cup \{y\})$  is a 2RDF on D of weight less than  $\omega(f)$ , and the desired bounds are proved.

However, if  $f(y) = \emptyset$ , then there exists a vertex  $w \in N^-(x) \cap N^-(y) \cap N^-(z)$ such that  $f(w) = \{1, 2\}$  or there exist two vertices  $w_1, w_2 \in N^-(x) \cap N^-(y) \cap$  266 J. Amjadi, N. Mohammadi, S.M. Sheikholeslami and L. Volkmann

 $N^-(z)$  such that  $f(w_1) = \{1\}$  and  $f(w_2) = \{2\}$ . Since  $w, w_1$  and  $w_2$  are inneighbors of x and z in D,  $(V_0 \cup \{x, z\}, V_1 - \{x, z\}, V_2, V_{1,2})$  is a 2RDF on D of weight less than f, and the proof is complete.

**Corollary 6.** If D is a digraph with  $\delta^+(D) \ge 2$ , then  $b_{r_2}(D) \le 2\Delta(G[D]) + \delta^-(D)$ .

**Proof.** Let  $y \in V(D)$  be a vertex with deg<sup>-</sup> $(y) = \delta^{-}(D)$ . Since  $\delta^{+}(D) \geq 2$ , there exist two different vertices  $x, z \in N^{+}(y)$ . Thus G[D] contains a path xyz such that  $(y, x), (y, z) \in A(D)$ . Now the result follows from Theorem 5.

Since  $\sum_{v \in V(D)} \deg^+(v) = \sum_{v \in V(D)} \deg^-(v)$  and  $\sum_{v \in V(D)} (\deg^+(v) + \deg^-(v)) \le n\Delta(G[D])$ , we have  $\delta^-(D) \le \frac{1}{2}\Delta(G[D])$ . Now, Corollary 6 leads to the next result.

**Corollary 7.** If D is a digraph with  $\delta^+(D) \ge 2$ , then  $b_{r2}(D) \le \frac{5}{2}\Delta(G[D])$ .

For every graph G, the expression  $\deg_a(G) = \sum_{v \in V(G)} \deg(v)/|V(G)|$  is called the *average degree* of G.

**Lemma 8.** For any digraph D with  $\delta^{-}(D) \geq 1$ , there exists a pair of vertices, say u and v, that are either adjacent or at distance two in G[D] with a common in-neighbor in D, with the property that

$$\deg_{G[D]}(u) + \deg_{G[D]}(v) \le 2\deg_a(G[D]).$$

**Proof.** Suppose that the lemma is false, and let D be a connected digraph where the result does not hold. Let the vertices of degree less than or equal to  $\deg_a(G[D])$  in G[D] be  $S = \{u_1, u_2, \ldots, u_m\}$  and the vertices of degree greater than  $\deg_a(G[D])$  be  $T = \{v_1, v_2, \ldots, v_n\}$ .

Observe that no pair of vertices of S can be joined by an arc. Hence, each  $u_i \in S$  has only vertices in T as in-neighbors or out-neighbors. Also note that each  $v_j$  has at most one out-neighbor in S, for otherwise if there were two, they would contradict our assumption.

Now we proceed to sum the degrees of all vertices in the underlying graph G[D] as follows. For each  $u_i \in S$  we consider an in-neighbor  $v_j \in T$  of  $u_i$  and take  $\deg_{G[D]}(u_i) + \deg_{G[D]}(v_j)$ . By assumption, we observe that  $\deg_{G[D]}(u_i) + \deg_{G[D]}(v_j) > 2 \deg_a(G[D])$ . Furthermore, by the above remarks, these inneighbors in T must be distinct. After adding m such pairs (to exhaust S), the degree of any remaining members of T are included. But the total sum of the degrees is greater than  $|V(G[D])| \deg_a(G[D])$  which is impossible. This completes the proof.

Next we present an upper bound on the size of a digraph with given rainbow domination number and rainbow bondage number.

**Theorem 9.** Let D be a digraph of order n with  $\delta^{-}(D) \geq 1$ ,  $\delta^{+}(D) \geq 2$ and rainbow bondage number  $b_{r2}(D)$ . If  $\deg_a(G[D])$  is the average degree of the underlying graph of D, then  $b_{r2}(D) \leq 2 \deg_a(G[D]) + \Delta(G[D]) - 3$  and  $|A(D)| \geq (n/4)(b_{r2}(D) - \Delta(G[D]) + 3).$ 

**Proof.** Let D be a digraph satisfying the assumptions of the theorem. By Lemma 8, there is at least one pair of vertices, say u and v, that are either adjacent or at distance 2 from each other with a common in-neighbor, and with the property that  $\deg_{G[D]}(u) + \deg_{G[D]}(v) \le 2 \deg_a(G[D])$ . Since  $\delta^+(D) \ge 2$ , there is a path uvw in G[D] such that  $(v, u), (v, w) \in A(D)$ , a path vuw in G[D] such that  $(u, v), (u, w) \in A(D)$ , or a path uwv in G[D] such that  $(w, u), (w, v) \in A(D)$ . Since these cases are symmetrical, we only consider the first. Applying Theorem 4, we obtain

$$b_{r2}(D) \leq \deg_{G[D]}(u) + \deg_{G[D]}(v) + \deg_{G[D]}(w) - 3$$
$$\leq 2 \deg_a(G[D]) + \Delta(G[D]) - 3.$$

Since  $2|E(G[D])| = n \deg_a(G[D])$ , we have

$$4|E(G[D])| = 2n \deg_a(G[D]) \ge n(b_{r_2}(D) - \Delta(G[D]) + 3).$$

Hence

$$|A(D)| = |E(G[D])| \ge (n/4)(b_{r2}(D) - \Delta(G[D]) + 3).$$

### 3. Some Classes of Digraphs

In this section we investigate complete digraphs, complete bipartite digraphs and transitive tournaments.

**Lemma 10.** If  $K_{p,q}^*$  is the complete bipartite digraph such that  $q \ge p \ge 2k$ , then  $\gamma_{rk}(K_{p,q}^*) = 2k$ .

**Proof.** Let  $X = \{x_1, x_2, \ldots, x_p\}$  and  $Y = \{y_1, y_2, \ldots, y_q\}$  be the partite sets of  $K_{p,q}^*$ . It is easy to see that the function f defined by  $f(x_i) = f(y_i) = \{i\}$  for  $1 \le i \le k$  and  $f(x) = \emptyset$  otherwise, is a k-rainbow dominating function of  $K_{p,q}^*$  of weight 2k and hence  $\gamma_{rk}(K_{p,q}^*) \le 2k$ .

Let now f be a  $\gamma_{rk}(K_{p,q}^*)$ -function. If  $f(x_i) \neq \emptyset$  for each i, then  $\gamma_{rk}(K_{p,q}^*) = \omega(f) \ge 2k$ . So assume  $f(x_i) = \emptyset$  for some i, say i = 1. Similarly, we may assume  $f(y_1) = \emptyset$ . This implies that  $\bigcup_{i=1}^p f(x_i) = \bigcup_{i=1}^q f(y_i) = \{1, 2, \dots, k\}$ . Hence  $\gamma_{rk}(K_{p,q}^*) = \omega(f) \ge 2k$  and the proof is complete.

**Theorem 11.** Let  $k \geq 2$  be an integer and let  $K_{p,q}^*$  be the complete bipartite digraph such that  $2k + 1 \leq p \leq q$ . Then  $p + 1 \leq b_{rk}(K_{p,q}^*) \leq 2p$ .

268 J. Amjadi, N. Mohammadi, S.M. Sheikholeslami and L. Volkmann

**Proof.** Let  $X = \{x_1, x_2, \ldots, x_p\}$  and  $Y = \{y_1, y_2, \ldots, y_q\}$  be the partite sets of  $K_{p,q}^*$ . If B is an arc set of  $K_{p,q}^*$ , then define  $D = K_{p,q}^* - B$ . If D contains a vertex  $x \in X$  and a vertex  $y \in Y$  such that  $\deg_D^+(x) = q$  and  $\deg_D^+(y) = p$ , then it follows from Lemma 10 that  $2k = \gamma_{rk}(K_{p,q}^*) \leq \gamma_{rk}(D) \leq 2k$  and therefore  $\gamma_{rk}(D) = 2k$ . Hence  $b_{rk}(K_{p,q}^*) \geq p$ . Now let |B| = p and  $D = K_{p,q}^* - B$  such that, without loss of generality,  $\deg_D^+(x) \neq q$  for all  $x \in X$ . Then  $B = \{x_1y_{i_1}, x_2y_{i_2}, \ldots, x_py_{i_p}\}$  with  $y_{i_j} \in Y$  for  $1 \leq j \leq p$ . Assume that  $y_{i_1} = y_1$ . Define the function f by  $f(x_1) = f(y_1) = \{1, 2, \ldots, k\}$  and  $f(u) = \emptyset$  for  $u \in V(K_{p,q}^*) - \{x_1, y_1\}$ . It is easy to see that f is a k-rainbow dominating function of D of weight 2k. Lemma 10 implies that  $2k = \gamma_{rk}(K_{p,q}^*) \leq \gamma_{rk}(D) \leq 2k$  and thus  $\gamma_{rk}(D) = 2k$ . Consequently,  $b_{rk}(K_{p,q}^*) \geq p + 1$ .

Let now  $B_1$  be the set of all arcs incident with the vertex  $y_1$ , and let  $H = K_{p,q}^* - B_1$ . Then  $y_1$  is an isolated vertex in H and thus  $\gamma_{rk}(H) = \gamma_{rk}(K_{p,q-1}^*) + 1$ . Since  $q \ge p \ge 2k + 1$ , Lemma 10 shows that  $\gamma_{rk}(K_{p,q-1}^*) = 2k$  and thus  $\gamma_{rk}(H) = 2k + 1$ . Since  $|B_1| = 2p$ , it follows that  $b_{rk}(K_{p,q}^*) \le 2p$ , and the proof is complete.

**Conjecture 12.** For integers  $k \ge 2$  and  $q \ge p \ge 2k + 1$ ,  $b_{rk}(K_{p,q}^*) = 2p$ .

**Theorem 13.** Let  $k \ge 2$  be an integer. If  $K_n^*$  is the complete digraph of order  $n \ge k+1$ , then  $n \le b_{rk}(K_n^*) \le n+k-1$ .

**Proof.** According to Propositions B and D, we deduce that  $\gamma_{rk}(K_n^*) = k$ . If B is an arc set of  $K_n^*$ , then define  $D = K_n^* - B$ . If D contains a vertex x such that  $\deg_D^+(x) = n - 1$ , then it follows from Propositions B and D that  $\gamma_{rk}(D) = k$ . This implies that  $b_{rk}(K_n^*) \geq n$ .

Now let  $\{x_1, x_2, \ldots, x_n\}$  be the vertex set of the complete digraph  $K_n^*$ . Define the arc sets  $B_1 = \{x_1x_n, x_2x_n, \ldots, x_{n-1}x_n\}$  and  $B_2 = \{x_nx_1, x_nx_2, \ldots, x_nx_k\}$ , and let  $D = K_n^* - (B_1 \cup B_2)$ . Then it is easy to see that  $b_{rk}(D) = b_{rk}(K_{n-1}^*) + 1 = k + 1$ . Since  $\gamma_{rk}(K_n^*) = k$ , we obtain  $b_{rk}(K_n^*) \leq |B_1| + |B_2| = n - 1 + k$ , and this is the desired upper bound.

**Theorem 14.** If  $K_n^*$  is the complete digraph of order  $n \ge 3$ , then  $b_{rk}(D) = b_{rk}(K_{n-1}^*) + 1 = k + 1$ .

**Proof.** By Theorem 13, we have  $b_{r2}(K_n^*) \ge n$ .

Now let  $\{x_1, x_2, \ldots, x_n\}$  be the vertex set of  $K_n^*$ . We define the arc set B of  $K_n^*$  by  $B = \{x_1x_2, x_2x_3, \ldots, x_{n-1}x_n, x_nx_1\}$ . If  $D = K_n^* - B$ , then we observe that  $\Delta^+(D) = n - 2$ . In addition, we see that there do not exist two different vertices u and v in D such that  $V(D) - \{u, v\} \subseteq N_D^+(u)$  and  $V(D) - \{u, v\} \subseteq N_D^+(v)$ . This can be seen as follows:

Suppose on the contrary that there exist two different vertices u and v in D such that  $V(D) - \{u, v\} \subseteq N_D^+(u)$  and  $V(D) - \{u, v\} \subseteq N_D^+(v)$ . If, without

loss of generality,  $u = x_1$ , then  $x_2 \notin N_D^+(x_1)$ . Therefore  $v = x_2$ . However, now  $x_3 \notin N_D^+(x_2)$ , a contradiction.

Applying Proposition C, we conclude that  $\gamma_{r2}(D) \geq 3$ . Since  $\gamma_{r2}(K_n^*) = 2$ , we deduce that  $b_{r2}(K_n^*) \leq n$ , and the proof is complete.

A tournament T = (V, E) is an orientation of a complete graph. A tournament is called *transitive* if  $p \to q$  and  $q \to r$  imply that  $p \to r$ .

**Theorem 15.** Let  $k \geq 2$  be an integer. If  $T_n$  is the transitive tournament of order  $n \geq k+1$ , then  $b_{rk}(T_n) = 1$ .

**Proof.** Let  $x_1x_2\cdots x_n$  be the unique directed Hamiltonian path of  $T_n$ . Then  $\deg_{T_n}^+(x_1) = n - 1$ , and therefore Propositions B and D imply that  $\gamma_{rk}(T_n) = k$ . Now let  $D = T_n - \{x_1 x_n\}$ , and let f be a  $\gamma_{rk}(D)$ -function. Assume first that  $f(x_n) = \emptyset$ . This implies that  $\bigcup_{u \in N_D^-(x_n)} f(u) = \{1, 2, \dots, k\}$ .

Since  $|f(x_1)| \ge 1$  and  $x_1 \notin N_D^-(x_n)$ , we obtain  $\omega(f) \ge k+1$ .

Next, assume that  $|f(x_n)| \ge 1$ . If  $|f(x_i)| \ge 1$  for each  $1 \le i \le n-1$ , then  $\omega(f) \ge n \ge k+1$ . So assume that  $f(x_i = \emptyset \text{ for an index } i \in \{1, 2, \dots, n-1\}$ . Then  $\bigcup_{u \in N_D^-(x_i)} f(u) = \{1, 2, \dots, k\}$ . Since  $x_n \notin N_D^-(x_i)$ , we obtain  $\omega(f) \ge k+1$ again.

Therefore  $\gamma_{rk}(D) \ge k+1$ . Since  $\gamma_{rk}(T_n) = k$ , we deduce that  $b_{rk}(T_n) = 1$ , and the proof is complete.

#### References

- [1] J. Amjadi, A. Bahremandpour, S.M. Sheikholeslami and L. Volkmann, The rainbow domination number of a digraph, Kragujevac J. Math. 37 (2013) 257–268.
- [2] B. Brešar, M.A. Henning and D.F. Rall, *Rainbow domination in graphs*, Taiwanese J. Math. 12 (2008) 213-225.
- [3] B. Brešar and T.K. Šumenjak, On the 2-rainbow domination in graphs, Discrete Appl. Math. 155 (2007) 2394–2400. doi:10.1016/j.dam.2007.07.018
- [4] G.J. Chang, J. Wu and X. Zhu, Rainbow domination on trees, Discrete Appl. Math. **158** (2010) 8–12. doi:10.1016/j.dam.2009.08.010
- [5] Ch. Tong, X. Lin, Y. Yang and M.Luo, 2-rainbow domination of generalized Petersen graphs P(n, 2), Discrete Appl. Math. **157** (2009) 1932–1937. doi:10.1016/j.dam.2009.01.020
- [6] N. Dehgardi, S.M. Sheikholeslami and L. Volkmann, The k-rainbow bondage number of a graph, Discrete Appl. Math. 174 (2014) 133–139. doi:10.1016/j.dam.2014.05.006

- J.F. Fink, M.S. Jacobson, L.F. Kinch and J. Roberts, *The bondage number of a graph*, Discrete Math. 86 (1990) 47–57. doi:10.1016/0012-365X(90)90348-L
- [8] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of Domination in Graphs (Marcel Dekker, Inc. New York, 1998).
- D. Meierling, S.M. Sheikholeslami and L. Volkmann, Nordhaus-Gaddum bounds on the k-rainbow domatic number of a graph, Appl. Math. Lett. 24 (2011) 1758–1761. doi:10.1016/j.aml.2011.04.046
- [10] S.M. Sheikholeslami and L. Volkmann, The k-rainbow domatic number of a graph, Discuss. Math. Graph Theory 32 (2012) 129–140. doi:10.7151/dmgt.1591
- [11] D.B. West, Introduction to Graph Theory (Prentice-Hall, Inc., 2000).
- Y. Wu and N. Jafari Rad, Bounds on the 2-rainbow domination number of graphs, Graphs Combin. 29 (2013) 1125–1133. doi:10.1007/s00373-012-1158-y
- [13] G. Xu, 2-rainbow domination in generalized Petersen graphs P(n, 3), Discrete Appl. Math. 157 (2009) 2570–2573. doi:10.1016/j.dam.2009.03.016

Received 10 February 2014 Revised 12 June 2014 Accepted 12 June 2014