

ON MINIMAL GEODETIC DOMINATION IN GRAPHS

HEARTY M. NUENAY¹

AND

FERDINAND P. JAMIL²

Department of Mathematics and Statistics
MSU-Iligan Institute of Technology
Iligan City, Philippines

e-mail: ferdinand.jamil@g.msuiit.edu.ph

Abstract

Let G be a connected graph. For two vertices u and v in G , a u - v geodesic is any shortest path joining u and v . The closed geodesic interval $I_G[u, v]$ consists of all vertices of G lying on any u - v geodesic. For $S \subseteq V(G)$, S is a geodesic set in G if $\bigcup_{u, v \in S} I_G[u, v] = V(G)$.

Vertices u and v of G are neighbors if u and v are adjacent. The closed neighborhood $N_G[v]$ of vertex v consists of v and all neighbors of v . For $S \subseteq V(G)$, S is a dominating set in G if $\bigcup_{u \in S} N_G[u] = V(G)$. A geodesic dominating set in G is any geodesic set in G which is at the same time a dominating set in G . A geodesic dominating set in G is a minimal geodesic dominating set if it does not have a proper subset which is itself a geodesic dominating set in G . The maximum cardinality of a minimal geodesic dominating set in G is the upper geodesic domination number of G . This paper initiates the study of minimal geodesic dominating sets and upper geodesic domination numbers of connected graphs.

Keywords: minimal geodesic dominating set, upper geodesic domination number.

2010 Mathematics Subject Classification: 05C12.

¹Research is fully supported by the Department of Science and Technology, Philippines, through the Accelerated Science and Technology Human Resource Development (ASTHRD) Scholarship Program.

²Corresponding author.

1. INTRODUCTION

Throughout this paper we consider only finite connected graphs with no loops or multiple edges. All basic graph theoretic terminologies and notations adapted here are taken from [11].

Let G and H be graphs with disjoint vertex sets. The *join* $G + H$ of G and H is the graph with vertex set $V(G + H) = V(G) \cup V(H)$ and edge set $E(G + H) = E(G) \cup E(H) \cup \{uv : u \in V(G), v \in V(H)\}$. The *composition* (or *lexicographic product*) $G[H]$ of G and H is the graph with vertex set $V(G[H]) = V(G) \times V(H)$ and $(u, v)(u', v') \in E(G[H])$ if and only if either $uu' \in E(G)$ or $u = u'$ and $vv' \in E(H)$.

Let G be a connected graph. For any two vertices u and v in G , a u - v *geodesic* refers to any shortest path in G joining u and v . The length of a u - v geodesic is called the *distance* between u and v , and is denoted by $d_G(u, v)$. The *eccentricity* $e_G(v)$ of a vertex v is defined by $e_G(v) = \max\{d_G(u, v) : u \in V(G)\}$ and the *diameter* of G is the number $\text{diam}(G) = \max\{d_G(u, v) : u, v \in V(G)\}$. The *closed geodesic interval* $I_G[u, v]$ is the set of all vertices lying on any u - v geodesic. For a subset S of the vertex set $V(G)$ of G , the *geodesic closure* of S is the set $I_G[S] = \bigcup_{u, v \in S} I_G[u, v]$. Various concepts inspired by geodesic closures are introduced in [7, 11]. A *geodesic set* in G is any set S of vertices in G satisfying $I_G[S] = V(G)$. The minimum cardinality $g(G)$ of a geodesic set is the *geodesic number* of G . Geodesic sets and geodesic numbers are studied in [1, 2, 3, 4, 5, 6]. A geodesic set S in G is a *minimal geodesic set* if S does not have a proper subset that is itself a geodesic set in G . The maximum cardinality of a minimal geodesic set in G is denoted by $g^+(G)$. Zhang *et al.* investigated a minimal geodesic set in a connected graph in [4].

We also define $I_G(u, v) = I_G[u, v] \setminus \{u, v\}$ and $I_G(S) = \bigcup_{u, v \in S} I_G(u, v)$. We call S a *2-path closure absorbing set* if for each $x \in V(G) \setminus S$, there exist $u, v \in S$ such that $d_G(u, v) = 2$ and $x \in I_G(u, v)$. The minimum cardinality of a 2-path closure absorbing set in G is denoted by $\rho_2(G)$. Since a 2-path closure absorbing set is always a geodesic set, $g(G) \leq \rho_2(G)$ for all connected graphs G . In [6], the geodesic numbers of some classes of graphs are described in terms of 2-path closure absorbing sets. A 2-path closure absorbing set S is a *minimal 2-path closure absorbing set* if S does not contain a proper subset that is itself 2-path closure absorbing. The maximum cardinality of a minimal 2-path closure absorbing set in G is denoted by $\rho_2^+(G)$.

The *open neighborhood* of a vertex v in G is the set $N_G(v) = \{u \in V(G) : uv \in E(G)\}$. The *degree*, $\text{deg}_G(v)$, of a vertex v refers to the value $|N_G(v)|$, and we define $\Delta(G) = \max\{\text{deg}_G(v) : v \in V(G)\}$. The *closed neighborhood* of v is the set $N_G[v] = N_G(v) \cup \{v\}$. A vertex v is an *extreme vertex* if the induced subgraph $\langle N_G[v] \rangle$ is a complete graph. The symbol $\text{Ext}(G)$ denotes the

set of all extreme vertices in G . For $S \subseteq V(G)$, we define $N_G(S) = \bigcup_{v \in S} N_G(v)$ and $N_G[S] = N_G(S) \cup S$. If $N_G[S] = V(G)$, then S is a *dominating set* in G . The minimum cardinality among dominating sets in G is called the *domination number* of G , and is denoted by $\gamma(G)$. A considerable number of studies have been dedicated in obtaining variations of the concept (see [12, 13, 14, 15]). The authors in [9] cited over 75 variations of domination and listed over 1,200 papers related to domination in graphs. An application to electrical power networks is being studied in [10].

A subset S of $V(G)$ is a *geodetic dominating set* in G if S is a geodetic set and at the same time a dominating set in G . The minimum cardinality of a geodetic dominating set is called the *geodetic domination number* of G , and is denoted by $\gamma_g(G)$. The study of geodetic domination was initiated by Escuardo, Gera, Hansberg, Jafari Rad and Volkman [8] in 2011. Some other interesting results can also be found in [16].

Customarily or as used in several literatures, the symbols like g -set, ρ_2 -set, ρ_2^+ -set, γ -set, and γ_g -set in a graph G would refer to a geodetic set of cardinality $g(G)$, a 2-path closure absorbing set with cardinality $\rho_2(G)$, a minimal 2-path closure absorbing set with cardinality $\rho_2^+(G)$, a dominating set with cardinality $\gamma(G)$, and a geodetic dominating set with cardinality $\gamma_g(G)$, respectively.

Since a 2-path closure absorbing set is also a geodetic dominating set, $g(G) \leq \gamma_g(G) \leq \rho_2(G)$ for all connected graphs G of order $n \geq 2$. In particular, if $\text{diam}(G) = 2$, then $\gamma_g(G) = \rho_2(G)$.

The following is found in [8].

Theorem 1.1 [8]. *Let G be a connected graph of order $n \geq 2$. Then*

- (i) $\gamma_g(G) = 2$ if and only if there exists a geodetic set $S = \{u, v\}$ such that $d_G(u, v) \leq 3$.
- (ii) $\gamma_g(G) = n$ if and only if G is the complete graph on n vertices.
- (iii) $\gamma_g(G) = n - 1$ if and only if there is a vertex v in G such that v is adjacent to every other vertex of G and $G - v$ is the union of at least two complete graphs.

2. MINIMAL GEODETIC DOMINATION

A geodetic dominating set S in a connected graph G of order $n \geq 2$ is a *minimal geodetic dominating set* in G if S does not have a proper subset which is itself a geodetic dominating set in G . The maximum cardinality of a minimal geodetic dominating set in G is the *upper geodetic domination number* of G , and is denoted by $\gamma_g^+(G)$. A minimal geodetic dominating set with cardinality $\gamma_g^+(G)$ is also called a γ_g^+ -set.

Example 2.1. (i) *If $m, n \geq 2$ and U and W are the partite sets of the complete bipartite graph $K_{m,n}$, then the minimal geodetic dominating sets in $K_{m,n}$ are U and W and all sets of the form $S = \{u, v, x, y\}$, where $u, v \in U$ and $x, y \in W$. More precisely,*

$$\gamma_g^+(K_{m,n}) = \begin{cases} 4, & \text{if } m = n = 3, \\ \max\{m, n\}, & \text{otherwise.} \end{cases}$$

(ii) *For $2 \leq n \leq 4$, $\gamma_g^+(P_n) = 2$, and for $n \geq 5$,*

$$\gamma_g^+(P_n) = \begin{cases} 2 \lfloor \frac{n}{4} \rfloor + 1, & \text{if } n \equiv 1 \pmod{4}, \\ 2 \lceil \frac{n}{4} \rceil, & \text{otherwise.} \end{cases}$$

Suppose that $n \equiv 1 \pmod{4}$, and $P_n = [u_1, u_2, \dots, u_n]$. Since the set $\{u_1, u_2, u_5, u_6, \dots, u_{4k-3}, u_{4k-2}, u_{4k+1}\}$ is a minimal geodetic dominating set in P_n , $\gamma_g^+(P_n) \geq 2 \lfloor \frac{n}{4} \rfloor + 1$. Let $S \subseteq V(P_n)$ be a minimal geodetic dominating set in P_n . For every $j = 1, 2, \dots, n - 3$, S contains at most two of the vertices u_j, u_{j+1}, u_{j+2} and u_{j+3} . Thus, $|S| \leq 2 \lfloor \frac{n}{4} \rfloor + 1$. Since S is arbitrary, $\gamma_g^+(P_n) \leq 2 \lfloor \frac{n}{4} \rfloor + 1$. Now, suppose that $n > 4$ but $n \neq 4a + 1$ for all positive integers a . Let k be the largest positive integer for which $4k + 1 < n$. Since the set of vertices $\{u_1, u_2, u_5, u_6, \dots, u_{4k+1}, u_n\}$ is a minimal geodetic dominating set in P_n , $\gamma_g^+(P_n) \geq 2 \lceil \frac{n}{4} \rceil$. Using similar arguments, if $S \subseteq V(P_n)$ is a minimal geodetic dominating set in P_n , then $|S| \leq 2 \lceil \frac{n}{4} \rceil$. This means that $\gamma_g^+(P_n) \leq 2 \lceil \frac{n}{4} \rceil$.

Theorem 2.2. *Let G be a connected graph of order $n \geq 2$. Then*

- (i) $\gamma_g^+(G) = 2$ *if and only if G is one of the following graphs: $P_2, C_4, \overline{K_2} + H$ where H is connected and either $H = K_{n-2}$ or $\rho_2^+(H) = 2$, G has a g -set $\{u, v\}$ with $u, v \in \text{Ext}(G)$ and $d_G(u, v) = 3$.*
- (ii) $\gamma_g^+(G) = n$ *if and only if $G = K_n$.*
- (iii) *For $n \geq 3$, $\gamma_g^+(G) = n - 1$ if and only if $G = K_1 + \bigcup_{j=1}^t K_{r_j}$, where $t \geq 2$.*

Proof. (i) Suppose that $\gamma_g^+(G) = 2$, and let $\{u, v\}$ be a γ_g^+ -set in G . Let $S = V(G) \setminus \{u, v\}$. Then $w \in I_G[u, v]$ for all $w \in S$ and $1 \leq d_G(u, v) \leq 3$ by Theorem 1.1. If $d_G(u, v) = 1$, then $S = \emptyset$ and $G = P_2$. Suppose that $d_G(u, v) = 2$. Then $G = \langle \{u, v\} \rangle + \langle S \rangle = \overline{K_2} + \langle S \rangle$. If $|S| = 1$, then $G = P_3 = \overline{K_2} + K_1$. If $|S| = 2$, then either $G = C_4$ or $G = \overline{K_2} + K_2$. Suppose that $|S| \geq 3$. Then $\langle S \rangle$ is connected. If $\langle S \rangle$ is the complete graph K_{n-2} , then $G = \overline{K_2} + K_{n-2}$. Suppose that $H = \langle S \rangle$ is not complete, and let T be a ρ_2^+ -set in H . Then T is a γ_g^+ -set in G . Thus $|T| = 2$. Hence, $\rho_2^+(H) = 2$. Finally, suppose that $d_G(u, v) = 3$. For each $x \in S$, either $ux \in E(G)$ or $xv \in E(G)$. Suppose that there exist $x, y \in N_G[u]$ with $d_G(x, y) = 2$. Consider $W = N_G(u) \cup \{v\}$. Let $z \in V(G) \setminus W$. If $z = u$, then $[x, z, y]$ is an x - y geodesic in G so that $z \in N_G[W]$ and $z \in I_G[W]$.

Suppose that $z \neq u$. Since $z \in I_G[u, v]$, there exist $a, b \in V(G)$ such that z lies on the u - v geodesic $[u, a, b, v]$. This means that $a \in N_G(u)$ and $z = b$. Thus $z \in N_G[W]$ and $z \in I_G[W]$. Accordingly, W is a geodesic dominating set in G . Let $T \subseteq W$ be a minimal geodesic dominating set in G . Since $v \notin N_G[N_G(u)]$, $v \in T$. Moreover, if $|T \cap N_G(u)| = 1$, then $u \notin I_G[T]$, a contradiction. Thus, $|T \cap N_G(u)| \geq 2$ so that $\gamma_g^+(G) \geq |T| \geq 3$, a contradiction. Therefore, $\langle N_G[u] \rangle$ is complete and $u \in Ext(G)$. Similarly, $v \in Ext(G)$.

Conversely, if G is P_2 or C_4 or $\overline{K_2} + K_{n-2}$, then $\gamma_g^+(G) = 2$. Suppose that $G = \overline{K_2} + H$, where H is connected and noncomplete with $\rho_2^+(H) = 2$. Then $diam(G) = 2$ and $T = V(\overline{K_2})$ is a minimal geodesic dominating set in G . Put $T = \{u, v\}$, and let Z be a minimal geodesic dominating set in G distinct from T . Then $|Z \cap T| \leq 1$. Suppose that $Z \cap T = \{u\}$. Since $ux \in E(G)$ for all $x \in V(H)$, $v \in I_G[Z \setminus \{u\}]$ and $V(H) \subseteq I_G[Z \setminus \{u\}]$ so that $Z \setminus \{u\}$ is a geodesic dominating set in G , a contradiction. Thus $Z \subseteq V(H)$ and, consequently, Z is a minimal 2-path closure absorbing set in H . Thus, $2 \leq |Z| \leq \rho_2^+(H) = 2$ so that $|Z| = 2$. Since Z is arbitrary, $\gamma_g^+(G) = |Z| = 2$. Now, let G have a g -set $\{u, v\}$ with $d_G(u, v) = 3$ and where the induced subgraphs $\langle N_G[u] \rangle$ and $\langle N_G[v] \rangle$ are complete. Then $Ext(G) = \{u, v\}$, which is the unique γ_g^+ -set in G . The conclusion follows.

(ii) If $G = K_n$, then $\gamma_g(G) = n$, by Theorem 1.1. Hence $\gamma_g^+(G) = n$. Suppose that $\gamma_g^+(G) = n$. Then each proper subset of $V(G)$ is not a geodesic dominating set in G . Let $v \in V(G)$, and set $S = V(G) \setminus \{v\}$. Then $v \notin N_G[S]$ or $v \notin I_G[S]$. If $v \notin N_G[S]$, then v is an isolated vertex, a contradiction. Thus $v \notin I_G[S]$ so that $v \in Ext(G)$. Since v is arbitrary, $V(G) = Ext(G)$ and $G = K_n$.

(iii) If $G = K_1 + \bigcup_{j=1}^t K_{r_j}$ for some $t \geq 2$, then $\gamma_g(G) = n-1$, by Theorem 1.1. By statement (ii), $\gamma_g^+(G) < n$. Thus $\gamma_g^+(G) = n-1$. Suppose that $\gamma_g^+(G) = n-1$. Let $S = V(G) \setminus \{v\}$, where $v \in V(G)$, be a γ_g^+ -set in G . We claim that $uv \in E(G)$ for all $u \in S$. Since v is not an endvertex, there exist $x, y \in S$ such that $[x, v, y]$ is an x - y geodesic in G . Suppose that, in the contrary, there exists $u \in S$ with $d_G(u, v) = 2$. Let $[u, w, v]$ be a u - v geodesic in G . If $x = w$ or $y = w$, then $S \setminus \{w\}$ is a geodesic dominating set in G , which is impossible. Suppose that $x \neq w$ and $y \neq w$. If $uy \notin E(G)$, then $S \setminus \{w\}$ is a geodesic dominating set in G . If $uy \in E(G)$ and $wy \notin E(G)$, then $S \setminus \{u\}$ is a geodesic dominating set in G . If $uy, wy \in E(G)$ and $ux \in E(G)$, then $S \setminus \{u\}$ is a geodesic dominating set in G . If $uy, wy \in E(G)$ and $ux \notin E(G)$, then $S \setminus \{w\}$ is a geodesic dominating set in G . Any of the above cases yields a contradiction. This proves the claim. Therefore, $G = K_1 + H$ for some graph H . Next, we show that $H = \bigcup_{j=1}^t K_{r_j}$, where $t \geq 2$. Suppose that H has a component K which is not a complete graph. Then K , consequently G , has a geodesic $[x, y, z]$ of length 2. Then $S \setminus \{y\}$ is a geodesic dominating set in G , a contradiction. Therefore, $H = \bigcup_{j=1}^t K_{r_j}$. Since

G is not a complete graph, $t \geq 2$. ■

Now follows a Nordhaus-Gaddum-type result. Let the symbol Ξ denote the infinite collection of all connected graphs G such that \overline{G} is also connected.

Theorem 2.3. *For all $G \in \Xi$ of order $n \geq 4$,*

$$4 \leq \gamma_g^+(G) + \gamma_g^+(\overline{G}) \leq 2n - 4.$$

In particular, $\gamma_g^+(G) + \gamma_g^+(\overline{G}) = 4$ if and only if $n = 4$.

Proof. Let $G \in \Xi$ be of order $n \geq 4$. Note that if G is either K_n or $K_1 + \bigcup_{j=1}^t K_{r_j}$ with $t \geq 2$, then \overline{G} is disconnected, a contradiction. In view of Theorem 2.2,

$$\gamma_g^+(G) + \gamma_g^+(\overline{G}) \leq (n - 2) + (n - 2) = 2n - 4.$$

The inequality at the left side is obvious.

In particular, if $n = 4$, then $\gamma_g^+(G) + \gamma_g^+(\overline{G}) = 4$. Conversely, suppose that $\gamma_g^+(G) + \gamma_g^+(\overline{G}) = 4$. Necessarily, $\gamma_g^+(G) = 2$ and $\gamma_g^+(\overline{G}) = 2$. By Theorem 2.2, G has a g -set $\{u, v\}$ with $u, v \in \text{Ext}(G)$ and $d_G(u, v) = 3$. Similarly, \overline{G} has a g -set $\{x, y\}$ with $x, y \in \text{Ext}(\overline{G})$ and $d_{\overline{G}}(x, y) = 3$. Assume that $x \in N_G[u]$. Suppose that $x = u$. Note that $N_{\overline{G}}(x) = N_G[v]$, and $\langle N_G[v] \rangle$ is not complete in \overline{G} . This means that $x \notin \text{Ext}(\overline{G})$, a contradiction. Suppose that $xu \in E(G)$. Since $xy \notin E(\overline{G})$, $xy \in E(G)$. If $vy \notin E(G)$, then $xv, vy \in E(\overline{G})$ so that $[x, v, y]$ is a geodesic in \overline{G} , a contradiction. Thus $[u, x, y, v]$ is a u - v geodesic in G . Suppose that $n \geq 5$, and let $z \in V(G)$ be distinct from u, x, y and v . Assume $xz \in E(\overline{G})$. Since $x \in \text{Ext}(\overline{G})$ and $xv \in E(\overline{G})$, $zv \in E(\overline{G})$ and, consequently, $zu \in E(G)$. Since $u \in \text{Ext}(G)$, $xz \in E(G)$, a contradiction. Therefore, $n = 4$. ■

Corollary 2.4. *If $G \in \Xi$ is of order $n \geq 4$, then $\gamma_g^+(G) + \gamma_g^+(\overline{G}) = 4$ if and only if $G = P_4$.*

Theorem 2.3 implies that if $G \in \Xi$ of order $n \geq 5$, then

$$5 \leq \gamma_g^+(G) + \gamma_g^+(\overline{G}) \leq 2n - 4.$$

Since $\gamma_g^+(C_5) = \gamma_g^+(\overline{C_5}) = 3$, this upper bound is sharp. Consider the graph G obtained from the cycle $C_4 = [v_1, v_2, v_3, v_4, v_1]$ by adding to C_4 two vertices x and y and the edges xv_1, xv_4, yv_2 and yv_3 . For this G , $\gamma_g^+(G) + \gamma_g^+(\overline{G}) = 5$, showing that the lower bound is sharp.

3. REALIZATION PROBLEMS

For nontrivial connected graphs G ,

$$2 \leq \gamma_g(G) \leq \gamma_g^+(G) \leq \rho_2^+(G).$$

In particular, $\gamma_g^+(K_{n,n}) = \rho_2^+(K_{n,n})$ for $n \geq 4$.

Theorem 3.1. *For every pair of positive integers a and b with $2 \leq a \leq b$, there exists a connected graph G such that $\gamma_g^+(G) = a$ and $\rho_2^+(G) = b$.*

Proof. If $a = b$, then we pick $G = K_{a,a}$. Suppose that $b = a + 1$. Obtain the graph G from $P_3 = [v_1, v_2, v_3]$ by adding $(a - 1)$ pendant edges v_3x_j , $j = 1, 2, \dots, a - 1$. Then $\gamma_g^+(G) = a$ and $\rho_2^+(G) = b$, which are determined by the sets $\{v_1, x_1, x_2, \dots, x_{a-1}\}$ and $\{v_1, v_2, x_1, x_2, \dots, x_{a-1}\}$, respectively.

Suppose that $b = a + k$, where $k \geq 2$. Write $V(K_k) = \{u_1, u_2, \dots, u_k\}$. Obtain G by joining $P_2 = [v_1, v_2]$ and $K_k + \overline{K_{a-1}}$ using new k edges v_2u_j , $j = 1, 2, \dots, k$. Note that $Ext(G) = \{v_1\} \cup V(\overline{K_{a-1}})$, and is a γ_g^+ -set in G . Thus, $\gamma_g^+(G) = a$. On the other hand, if $k \geq 2$, then $\rho_2^+(G) = a + k = b$ and is determined by the set $V(K_k) \cup Ext(G)$. ■

Corollary 3.2. *For every pair of positive integers a and b with $2 \leq a < b$, the smallest possible order of a connected graph G for which $\gamma_g^+(G) = a$ and $\rho_2^+(G) = b$ is $b + 1$.*

Theorem 3.3. *For all positive integers a, b, c with $2 \leq a \leq b < c$ and $c \geq b + 2$, there exists a connected graph G such that $\gamma_g(G) = a$, $\gamma_g^+(G) = b$ and $|V(G)| = c$.*

Proof. Suppose that $c = b + 2$. Write $a = 2 + k$ and $b = r + k$, $r \geq 2$ and $k = 0, 1, 2, \dots$. If $k = 0$, then we take $G = K_{2,r}$. In this case, $\gamma_g(G) = 2$ and $\gamma_g^+(G) = r$. Suppose that $k \geq 1$. Consider the graph G as in Figure 1.




Figure 1

If $r = 2$, G is obtained by adjoining to path $[y, y_1, x]$ $(k + 1)$ pendant edges xx_j , $j = 1, 2, \dots, k + 1$. Then $Ext(G) = \{y, x_1, x_2, \dots, x_{k+1}\}$ is the unique minimal geodetic dominating set in G . Here we have $\gamma_g(G) = \gamma_g^+(G) = 2 + k$. Now, suppose that $r \geq 3$. G is obtained from $K_{2,r-1}$ (with partite sets $U = \{x, y\}$ and $W = \{y_1, y_2, \dots, y_{r-1}\}$) by adding to $K_{2,r-1}$ $(k + 1)$ pendant edges xx_j , $j = 1, 2, \dots, k + 1$. The minimal geodetic dominating sets in G are $\{y, x_1, x_2, \dots, x_{k+1}\}$ and $W \cup \{x_1, x_2, \dots, x_{k+1}\}$. Thus $\gamma_g(G) = 2 + k$ and $\gamma_g^+(G) = r + k$.

Suppose that $c = b + 3$. Write $a = 2 + k$ and $b = r + k$, $k = 0, 1, \dots$ and $r \geq 2$. Suppose that $k = 0$. Consider the graph $G = G_1$ as in Figure 2.




Figure 2

If $r = 2$, then $\{y, z\}$ is the unique minimal geodetic dominating set in G so that $\gamma_g(G) = \gamma_g^+(G) = 2$. If $r \geq 3$, then $\gamma_g(G) = 2$ and $\gamma_g^+(G) = (r - 1) + 1 = r$, the latter being determined by $\{z, y_1, y_2, \dots, y_{r-1}\}$. Suppose that $k \geq 1$. Obtain G as the graph G_2 in Figure 2 by taking the union of $K_{1,k+1}$ (with partite sets $\{z\}$ and $\{x, z_1, z_2, \dots, z_k\}$) and $K_{2,r}$ (with partite sets $\{x, y\}$ and $\{y_1, y_2, \dots, y_r\}$). Note that $\{z_1, z_2, \dots, z_k\}$ is always contained in a geodetic dominating set in G . Thus $\gamma_g(G) = 2 + k$ and $\gamma_g^+(G) = r + k$.

Finally, suppose that $c = b + d$, where $d \geq 4$. Write $a = 2 + k$ and $b = r + k$, where $r \geq 2$ and $k = 0, 1, 2, \dots$, and put $l = c - b - 3$. For $k = 0$, we obtain G as the graph G_1^* in Figure 3 by joining $K_{l+1} + \overline{K_2}$ and $K_{r-1,2}$ using the common vertices x and y_1 . Note that $Ext(G) = \{z\}$, and $S = \{z, y\}$ is a minimal geodetic dominating in G and every minimal geodetic dominating set that contains y coincides S . Thus, aside from S , the other minimal geodetic dominating set in G is $\{z, y_1, y_2, \dots, y_{r-1}\}$. Consequently, $\gamma_g(G) = 2 = a$ and $\gamma_g^+(G) = r = b$. Now, suppose that $k \geq 1$. Consider G as the graph G_2^* in Figure 3 obtained from G_1^* in Figure 3 by adjoining pendant edges zz_j . $Ext(G) = \{z_1, z_2, \dots, z_k\}$ and if S is a γ_g^+ -set that contains y , then $|S| = k + 2$. In this case, $\gamma_g(G) = k + 2 = a$ and $\gamma_g^+(G) = r + k = b$. ■




Figure 3

The next corollary follows from Theorem 2.2 and the existence proof of Theorem 3.3.

Corollary 3.4. *For every pair of positive integers a and b with $2 \leq a < b$, the minimum order of a connected graph G for which $\gamma_g(G) = a$ and $\gamma_g^+(G) = b$ is $b + 2$.*

4. ρ_3 -SETS

Let G be a connected graph of order $n \geq 2$ and $S \subseteq V(G)$. S is said to be a ρ_3 -set in G if for every $w \in V(G) \setminus S$ there exist $u, v \in S$ such that $d_G(u, v) \leq 3$ and $w \in I_G[u, v]$. We denote by $\rho_3(G)$ the minimum cardinality of a ρ_3 -set in G . Since every 2-path closure absorbing set is a ρ_3 -set, $\rho_3(G) \leq \rho_2(G)$. In particular, if $\text{diam}(G) = 2$, then $\rho_2(G) = \rho_3(G)$. Since a ρ_3 -set is a geodetic dominating set, $\gamma_g(G) \leq \rho_3(G)$. If $\text{diam}(G) \leq 3$, then $\gamma_g(G) = \rho_3(G)$. However, in general, $\gamma_g(G)$ and $\rho_3(G)$ are not necessarily equal.

Graph G is said to be K_3 -free (resp. C_4 -free) if G does not contain K_3 (resp. C_4) as a subgraph.

Theorem 4.1. *Let G be a connected graph of order $n \geq 2$, and let $S \subseteq V(G)$.*

- (i) *If S is a ρ_3 -set in G , then for all $v \in S$, $\min\{d_G(u, v) : u \in S\} \leq 3$.*

- (ii) If G is K_3 -free and C_4 -free and S is a geodetic dominating set in G , then S is a ρ_3 -set in G .

Proof. The conclusions in statements (i) and (ii) are trivially satisfied for cases where $n = 2$, $n = 3$ and $n = 4$. Assume that $n \geq 5$, and let $S \subseteq V(G)$. To prove statement (i), suppose that S is a ρ_3 -set in G , and suppose that there is $v \in S$ such that $d_G(v) = \min\{d_G(u, v) : u \in S\} \geq 4$. Let $w \in S$ be such that $d_G(w, v) = d_G(v)$. Then there exists $u \in V(G) \setminus S$ lying on a w - v geodesic with $d_G(u, v) = 2$. Since S is dominating in G , there exists $z \in S$ such that $uz \in E(G)$. Observe that $d_G(z, v) \leq 3$, a contradiction. Therefore, $d_G(v) \leq 3$ for all $v \in S$.

Next, we prove statement (ii). Suppose that G is K_3 -free and C_4 -free, and let $v \in V(G) \setminus S$. If S is a dominating set in G , then there exists $x \in S$ such that $xv \in E(G)$. If S is a geodetic set, then v is not an endvertex of G . Pick $u \in N_G(v)$ with $u \neq x$. Since G is K_3 -free, $[x, v, u]$ is an x - u geodesic in G . If $u \in S$, then x and u are the desired vertices in S for v . Suppose that $u \notin S$. Pick $y \in S$ such that $uy \in E(G)$. Since G is K_3 -free and C_4 -free, $xy, vy \notin E(G)$. Consequently, $[x, v, u, y]$ is an x - y geodesic in G with $d_G(x, y) = 3$. ■

If G is a connected graph of order $n \geq 2$ which is K_3 -free and C_4 -free, then $\rho_3(G) = \gamma_g(G)$. In particular, if T is a tree of order $n \geq 2$, then $\rho_3(T) = \gamma_g(T)$.

Theorem 4.2. Let G be a connected K_3 -free graph of order $n \geq 2$. Then

$$\rho_3(G) \leq \gamma_g^+(G).$$

Proof. Suppose that $Ext(G) \neq \emptyset$. Put $Ext(G) = \{x_1, x_2, \dots, x_k\}$ for some positive integer k . For each $j = 1, 2, \dots, k$, define $S_j = \{x_1, x_2, \dots, x_j\}$. If $N_G[S_k] \neq V(G)$, choose $x_{k+1} \in V(G) \setminus N_G[S_k]$, and put $S_{k+1} = \{x_1, x_2, \dots, x_k, x_{k+1}\}$. If $N_G[S_{k+1}] \neq V(G)$, then choose $x_{k+2} \in V(G) \setminus N_G[S_{k+1}]$, and put $S_{k+2} = \{x_1, x_2, \dots, x_{k+1}, x_{k+2}\}$. Continuing in this way, there is a smallest positive integer m such that $N_G[S_m] = V(G)$. If $Ext(G) = \emptyset$, then construct $S_m = \{x_1, x_2, \dots, x_m\}$ by choosing any $x_1 \in V(G)$ and put $S_1 = \{x_1\}$, and for $j \geq 2$, $x_j \in V(G) \setminus N_G[S_{j-1}]$, where $S_{j-1} = \{x_1, x_2, \dots, x_{j-1}\}$. In any case, we claim that $S = S_m$ is a minimal geodetic dominating set and at the same time a ρ_3 -set in G . Clearly, S is a dominating set in G . Let $u \in V(G) \setminus S$. Then there exists $w \in S$ such that $uw \in E(G)$. Since $u \notin Ext(G)$ and G is K_3 -free, there exists $v \in V(G)$ such that $[v, u, w]$ is a v - w geodesic in G . Suppose that $v \notin S$. There exists $z \in S$ such that $zv \in E(G)$. Since G is K_3 -free, $uz \notin E(G)$. Also, by the construction of S , $zw \notin E(G)$. Thus, $[w, u, v, z]$ is a w - z geodesic in G . Here, $d_G(w, z) \leq 3$ and $u \in I_G[w, z] \subseteq I_G[S]$. Since u is arbitrary, S is a ρ_3 -set and a geodetic dominating set in G . Now let $S^* = S \setminus \{x_j\}$, $j = 1, 2, \dots, m$. We will show that S^* is not a dominating set in G . Suppose that $Ext(G) \neq \emptyset$. If $j \leq k$, then $x_j \in Ext(G)$ and S^* is not a geodetic set in G . Suppose that $j > k$. Since $x_j \notin Ext(G)$, there exist

$u, v \in V(G)$ such that $[u, x_j, v]$ is a u - v geodesic in G . Since $x_j \in S$, $u, v \notin S$. In fact, $x \notin S$ for all $x \in N_G[x_j]$. Thus $x_j \notin N_G[S^*]$, and S^* is not a dominating set in G . The case where $Ext(G) = \emptyset$ is handled similarly. Since j is arbitrary, S is a minimal geodetic dominating set in G . Therefore, $\rho_3(G) \leq |S| \leq \gamma_g^+(G)$. ■

It is easy to verify that $\rho_3(P_5) = 3 = \gamma_g^+(P_5)$. Hence the bound given in Theorem 4.2 is sharp.

5. JOIN AND COMPOSITION OF GRAPHS

For connected graphs G and H , if $S \subseteq V(G)$ is a 2-path closure absorbing set in G , then S is a geodetic dominating set in $G + H$.

Theorem 5.1. *For noncomplete connected graphs G , $\gamma_g^+(G + K_n) = \rho_2^+(G)$.*

Proof. First, we claim that if $S \subseteq V(G + K_n)$ is a geodetic dominating set in $G + K_n$, then $A = S \cap V(G)$ is a 2-path closure absorbing set in G . Let $S \subseteq V(G + K_n)$ be a geodetic dominating set in $G + K_n$. Let $x \in V(G) \setminus A$, and let $u, v \in S$ such that $x \in I_G[u, v]$. Necessarily, $u, v \in V(G)$. Since $diam(G + K_n)$ is 2, $[u, x, v]$ is a u - v geodesic in G . Thus $d_G(u, v) = 2$ and A is a 2-path closure absorbing set in G .

Now let $S \subseteq V(G + K_n)$ be a minimal geodetic dominating set in $G + K_n$. The above result implies that $A = S \cap V(G)$ is a 2-path closure absorbing set in G , and consequently, A is a geodetic dominating set in $G + K_n$. Since S is a minimal geodetic dominating set, $S = A$ so that S is a minimal 2-path closure absorbing set in G . Since S is arbitrary, $\gamma_g^+(G + K_n) \leq \rho_2^+(G)$.

Conversely, let $S \subseteq V(G)$ be a ρ_2^+ -set in G . Then S is a geodetic dominating set in $G + K_n$. That S is, in fact, a minimal geodetic dominating set in $G + K_n$ follows from the claim above. This yields $\rho_2^+(G) \leq \gamma_g^+(G + K_n)$. ■

Theorem 5.2. *For all noncomplete connected graphs G and H ,*

$$\gamma_g^+(G + H) = \max\{4, \rho_2^+(G), \rho_2^+(H)\}.$$

Proof. Let $S \subseteq V(G + H)$ be a minimal geodetic dominating set in $G + H$. If $S \subseteq V(G)$, then S is a minimal 2-path closure absorbing set in G since $diam(G + H) = 2$. This means that $|S| \leq \rho_2^+(G)$. Similarly, if $S \subseteq V(H)$, then $|S| \leq \rho_2^+(H)$. Suppose that $A = S \cap V(G) \neq \emptyset$ and $B = S \cap V(H) \neq \emptyset$. Then $|A| \geq 2$ and $|B| \geq 2$, and $V(H) \subseteq I_{G+H}[A]$ and $V(G) \subseteq I_{G+H}[B]$. The minimality of S implies that $|A| = |B| = 2$ and $|S| = 4$. Hence $\gamma_g^+(G + H) \leq \max\{4, \rho_2^+(G), \rho_2^+(H)\}$.

To prove the other inequality, note that if $S \subseteq V(G)$, then S is a minimal geodetic dominating set in $G + H$ if and only if S is a minimal 2-path closure

absorbing set in G . This means that $\max\{\rho_2^+(G), \rho_2^+(H)\} \leq \gamma_g^+(G + H)$. Since G and H are noncomplete, we can pick $u, v \in V(G)$ and $x, y \in V(H)$ such that $d_G(u, v) = 2$ and $d_H(x, y) = 2$. Then $\{u, v, x, y\}$ is a minimal geodetic dominating set in $G + H$. This means that $4 \leq \gamma_g^+(G + H)$. This completely establishes the desired inequality. ■

Next, we investigate the minimal geodetic domination in the composition of graphs $G + K_n$.

For $A \subseteq V(G)$, we define $A^g = A \cap I_G(A)$, and for $S \subseteq V(G[H])$, $S_G = \{u \in V(G) : (u, v) \in S \text{ for some } v \in V(H)\}$.

It is known (see [16]) that if $S \subseteq V(G[H])$ is a geodetic dominating set in $G[H]$, then S_G is a geodetic dominating set in G .

Theorem 5.3. [16] *Let G be a noncomplete connected graph and $n \geq 2$. Then $S \subseteq V(G[K_n])$ is a geodetic dominating set in $G[K_n]$ if and only if $S = [(A \setminus A^g) \times V(K_n)] \cup T$, where $A = S_G$ and $T_G = A^g$.*

Corollary 5.4. *For all noncomplete connected graphs G and $n \geq 2$,*

$$\gamma_g^+(G[K_n]) \geq \max\{n|A| - (n - 1)|A^g| : A \text{ is a minimal geodetic dominating set in } G\}.$$

Proof. Let

$$\alpha = \max\{n|A| - (n - 1)|A^g| : A \text{ is a minimal geodetic dominating set in } G\}.$$

Let $A \subseteq V(G)$ be a minimal geodetic dominating set in G , and let $S = [(A \setminus A^g) \times V(K_n)] \cup [A^g \times \{v\}]$, where $v \in V(K_n)$. By Theorem 5.3, S is a geodetic dominating set in $G[K_n]$. Suppose that there exists $S^* \subseteq S$ such that S^* is a geodetic dominating set in $G[K_n]$. By Theorem 5.3, $S^* = [(B \setminus B^g) \times V(K_n)] \cup T$, where B is a geodetic dominating set in G and $T_G = B^g$. Since $S^* \subseteq S$, $B \subseteq A$. Since A is a minimal geodetic dominating set in G , $A = B$. Therefore, $S = S^*$ and S is a minimal geodetic dominating set in $G[K_n]$. Thus, $\gamma_g^+(G[K_n]) \geq |S| = n|A| - (n - 1)|A^g|$. Since A is arbitrary, $\gamma_g^+(G[K_n]) \geq \alpha$. ■

Lemma 5.5. *Let G be a noncomplete connected graph and $n \geq 2$.*

- (i) *If $S \subseteq V(G)$ is a geodetic dominating set (respectively, minimal geodetic dominating set) in G , then $\{u\} \times S$ is a geodetic dominating set (resp. minimal geodetic dominating set) in $K_n[G]$ for all $u \in V(K_n)$.*
- (ii) *If $S \subseteq V(G)$ is a geodetic set (resp. minimal geodetic set but not dominating) in G , then $\{(w, z)\} \cup (\{u\} \times S)$ is a geodetic dominating set (respectively, minimal geodetic dominating set) in G for all $z \in V(G)$ and for all distinct $w, u \in V(K_n)$.*

Proof. Let S be a geodetic dominating set in G and $u \in V(K_n)$. Let $(x, y) \in V(K_n[G]) \setminus (\{u\} \times S)$. Suppose that $x \neq u$. Then $(x, y)(u, v) \in E(K_n[G])$ for all $v \in S$. Thus, $(x, y) \in N_{K_n[G]}[\{u\} \times S]$. Choose $v_1, v_2 \in S$ such that $d_G(v_1, v_2) \geq 2$. Then $(x, y) \in I_{K_n[G]}[(u, v_1), (u, v_2)] \subseteq I_{K_n[G]}[\{u\} \times S]$. Suppose that $x = u$. Then $y \notin S$. Since S is a geodetic dominating set in G , $y \in N_G[S] \cap I_G[S]$. Thus, $(x, y) \in N_{K_n[G]}[\{u\} \times S]$ and $(x, y) \in I_{K_n[G]}[\{u\} \times S]$. This proves that $\{u\} \times S$ is a geodetic dominating set in $K_n[G]$. Finally, let $\{u\} \times T \subseteq \{u\} \times S$ be a geodetic dominating set in $K_n[G]$. Then $T \subseteq S$ and T is a geodetic dominating set in G . If S is a minimal geodetic dominating set in G , then $T = S$, and this proves statement (i).

To prove statement (ii), let $C = \{(w, z)\} \cup (\{u\} \times S)$, where $S \subseteq V(G)$ is a geodetic set in G , $z \in V(G)$ and $u, w \in V(K_n)$ with $u \neq w$. Let $(a, b) \in V(K_p[G]) \setminus C$. Suppose that $a = u$. Then $b \notin S$ and $aw \in E(K_n)$ so that $(a, b)(w, z) \in E(K_p[G])$. Since S is a geodetic set in G , there exist $x, y \in S$ such that $b \in I_G[x, y]$. Then $(u, x), (u, y) \in C$ and $(a, b) \in I_{K_p[G]}[(u, x), (u, y)]$. Suppose that $a \neq u$. Then $au \in E(K_n)$ and $(a, b) \in N_{K_p[G]}[\{u\} \times S]$. Choose $x, y \in S$ such that $d_G(x, y) \geq 2$. Then $(u, x), (u, y) \in C$ and $(a, b) \in I_{K_p[G]}[(u, x), (u, y)]$. In any case, $(a, b) \in N_{K_p[G]}[C]$ and $(a, b) \in I_{K_p[G]}[C]$. Since (a, b) is arbitrary, C is a geodetic dominating set in $K_p[G]$. Suppose that S is a minimal geodetic set in G but not dominating. Let $(a, b) \in C$, and put $C^* = C \setminus \{(a, b)\}$. If $a = u$, then $b \in S$ and $S \setminus \{b\}$ is not a geodetic set in G . This case means that C^* is not a geodetic set in $K_p[G]$. On the other hand, if $a \neq u$, then $(a, b) = (w, z)$ and C^* is not a dominating set in $K_p[G]$. Therefore, C is a minimal geodetic dominating set in $K_p[G]$. ■

Theorem 5.6. *Let G be a noncomplete connected graph and $n \geq 2$, and let $C \subseteq V(K_n[G])$. Then C is a minimal geodetic dominating set in $K_n[G]$ if and only if one of the following is true:*

- (i) $C = \{u\} \times S$ for some minimal geodetic dominating set in G and $u \in V(K_n)$;
- (ii) $C = \{(w, z)\} \cup (\{u\} \times S)$ for some nondominating but minimal geodetic set S in G , for some $z \in V(G)$ and distinct $w, u \in V(K_n)$;
- (iii) $C = \{(u_1, v_1), (u_1, v_2), (u_2, w_1), (u_2, w_2)\}$ for some distinct $u_1, u_2 \in V(K_n)$ and some $v_1, w_1, v_2, w_2 \in V(G)$ with $d_G(v_1, v_2) \geq 2$ and $d_G(w_1, w_2) \geq 2$.

Proof. By Lemma 5.5, if property (i) or property (ii) holds, then C is a minimal geodetic dominating set in $K_n[G]$. It can also be readily verified that if property (ii) holds, then C is a minimal geodetic dominating set.

Let $C \subseteq V(K_n[G])$ be a minimal geodetic dominating set in $K_n[G]$. Then C contains distinct vertices (u, v) and (u, y) . For if it were false and $(u, v) \in C$, then for all $y \in V(G) \setminus \{v\}$, $(u, y) \notin I_{K_n[G]}[C]$, a contradiction. Moreover, since G is noncomplete, we may choose v and y such that $d_G(v, y) \geq 2$. Suppose that $C = \{u\} \times S$ for some $S \subseteq V(G)$. Let $z \in V(G) \setminus S$. Since $(u, z) \in N_{K_n[G]}[C]$,

$z \in N_G[S]$. Similarly, $z \in I_G[S]$. Accordingly, S is a geodetic dominating set in G . Let $T \subseteq S$ be a geodetic dominating set in G . Then $\{u\} \times T \subseteq C$ and is a geodetic dominating set in $K_n[G]$ by Lemma 5.5. By the definition of C , $T = S$ and S is a minimal geodetic dominating set in G . This establishes property (i).

Now suppose that $C \neq \{u\} \times S$ for any $S \subseteq V(G)$. Let $S = \{t \in V(G) : (u, t) \in C\}$. Note that $(a, b) \in N_{K_n[G]}[(u, v), (u, y)] \cap I_{K_n[G]}[(u, v), (u, y)]$ for all $a \neq u$ and all $b \in V(G)$. Since C is a minimal geodetic dominating set in $K_n[G]$, $\{u\} \times S$ is not a geodetic dominating set in $K_p[G]$. Thus, $N_G[S] \neq V(G)$ or $I_G[S] \neq V(G)$. Suppose that $I_G[S] = V(G)$. Then S is not a dominating set in G . Let $b \in V(G) \setminus N_G[S]$. There exists $(w, z) \in C$ such that $(u, b)(w, z) \in E(K_p[G])$. Necessarily, $w \neq u$. Since $\{(w, z)\} \cup (\{u\} \times S)$ is a geodetic dominating set in $K_p[G]$, $C = \{(w, z)\} \cup (\{u\} \times S)$. In view of Lemma 5.5, S is a minimal geodetic set in G , and property (ii) is established. Finally suppose that $I_G[S] \neq V(G)$, and let $b \in V(G) \setminus I_G[S]$. Then there exists $w \in V(K_n)$ distinct from u and some $z, r \in V(G)$ with $d_G(z, r) \geq 2$ such that $(u, b) \in I_{K_n[G]}[(w, z), (w, r)]$. Since $\{(u, v), (u, y), (w, z), (w, r)\} \subseteq C$ is a geodetic dominating set, $C = \{(u, v), (u, y), (w, z), (w, r)\}$. ■

Corollary 5.7. *Let G be a noncomplete connected graph with $g^+(G) < \gamma_g^+(G)$ and $n \geq 2$. Then*

$$\gamma_g^+(K_n[G]) = \max\{4, \gamma_g^+(G)\}.$$

6. γ_g^+ -SUBGRAPH

A graph H is a γ_g^+ -subgraph if there exists a connected graph G containing H as an induced subgraph such that $V(H)$ is a γ_g^+ -set in G .

The idea of the following result is taken from [4].

Theorem 6.1. *Let H be a connected graph. Then H is a γ_g^+ -subgraph if and only if either H is complete or H has no vertex v with $e_H(v) = 1$.*

Proof. Let there be a connected graph G containing H as an induced subgraph and such that $V(H)$ is a γ_g^+ -set in G . Suppose that H is noncomplete and suppose that $v \in V(H)$ with $e_H(v) = 1$. We claim that $S = V(H) \setminus \{v\}$ is a geodetic dominating set in G . Let $w \in V(G) \setminus S$. Suppose that $w = v$. Since H is noncomplete, there exist $a, b \in V(H)$ such that $d_G(a, b) = d_H(a, b) = 2$. Necessarily, $a \neq v$ and $b \neq v$ so that $a, b \in S$. Since $av, bv \in E(H) \subseteq E(G)$, $w \in I_G[a, b] \subseteq I_G[S]$ and $w \in N_G[a] \subseteq N_G[S]$. Suppose that $w \neq v$. Since $V(H)$ is a geodetic set in G , there exist $a, b \in V(H)$ such that $w \in I_G[a, b]$. If $a = v$ or $b = v$, then $d_H(a, b) = d_G(a, b) = 1$, a contradiction. Thus $a, b \in S$. Since $av, bv \in E(G)$, $d_G(a, b) = 2$ and $aw, bw \in E(G)$. Hence $w \in I_G[S]$ and $w \in N_G[S]$. Thus, S is a geodetic dominating set in G . This is a contradiction

since $V(H)$ is a minimal geodetic dominating set in G and S is a proper subset of $V(H)$.

By Theorem 2.2, if H is complete, then $V(H)$ is the γ_g^+ -set in $G = H$. Suppose that H is noncomplete having no vertex u with $e_H(u) = 1$. For each $u \in V(H)$, choose $v \in V(H)$ such that $d_H(u, v) = 2$. Corresponding to each pair u and v , add to H the vertex $x_{u,v}$ and the edges $ux_{u,v}$ and $vx_{u,v}$. Let G be the resulting graph of minimum order obtained in this way. Then $|V(G) \setminus V(H)| \leq |V(H)|$. We claim that $V(H)$ is a γ_g^+ -set in G . Let $x \in V(G) \setminus V(H)$. Then $x = x_{u,v}$ for some $u, v \in V(H)$ with $d_H(u, v) = d_G(u, v) = 2$. More precisely, $xu, xv \in E(G)$. Thus, $x \in I_G[u, v]$ and $x \in N_G[u]$. In other words, $V(H)$ is a geodetic dominating set in G .

Let $u \in V(H)$, and let $v \in V(H)$ such that $d_H(u, v) = 2$. Corresponding to u and v is a $x_{u,v} \in V(G) \setminus V(H)$. By its construction, $x_{u,v} \notin I_G[V(H) \setminus \{u\}]$. Since u is arbitrary, $V(H)$ is a minimal geodetic dominating set in G . Finally, let $S \subseteq V(G)$ be a minimal geodetic dominating set in G . For the triple $u, v, x_{u,v}$, if $u, v \in S$, then $x_{u,v} \notin S$, or, equivalently, if $x_{u,v} \in S$, then $u \notin S$ or $v \notin S$. Thus

$$|S| = |S \setminus V(H)| + |V(H) \cap S| \leq |V(H) \setminus S| + |V(H) \cap S| = |V(H)|.$$

Since S is arbitrary, $V(H)$ is a γ_g^+ -set in G . ■

Acknowledgment

The authors would like to thank the referees for their corrections and recommendations that contributed to the improvement of the paper.

REFERENCES

- [1] S. Canoy Jr. and G. Cagaanan, *On the geodesic and hull numbers of graphs*, Congr. Numer. **161** (2003) 97–104.
- [2] S. Canoy Jr. and G. Cagaanan, *On the geodetic covers and geodetic bases of the composition $G[K_m]$* , Ars Combin. **79** (2006) 33–45.
- [3] G. Chartrand, F. Harary and P. Zhang, *Geodetics sets in graphs*, Discuss. Math. Graph Theory **20** (2000) 129–138.
doi:10.7151/dmgt.1112
- [4] G. Chartrand, F. Harary and P. Zhang, *On the geodetic number of a graph*, Networks **39** (2002) 1–6.
doi:10.1002/net.10007
- [5] I. Aniversario, F. Jamil and S. Canoy Jr., *The closed geodetic numbers of graphs*, Util. Math. **74** (2007) 3–18.
- [6] S. Canoy Jr., G. Cagaanan and S. Gervacio, *Convexity, geodetic and hull numbers of the join of graphs*, Util. Math. **71** (2007) 143–159.

- [7] S. Canoy Jr. and I.J. Garces, *Convex sets under some graph operations*, Graphs Combin. **18** (2002) 787–793.
doi:0.1007/s003730200065
- [8] H. Escuardo, R. Gera, A. Hansberg, N. Jafari Rad and L. Volkmann, *Geodetic domination in graphs*, J. Combin. Math. Combin. Comput. **77** (2011) 89–101.
- [9] T. Haynes, S.T. Hedetniemi and P. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, Inc. New York, 1998).
- [10] T. Haynes, S.M. Hedetniemi, S.T. Hedetniemi and M. Henning, *Domination in graphs applied to electrical power networks*, SIAM J. Discrete Math. **15** (2000) 519–529.
doi:10.1137/S0895480100375831
- [11] F. Buckley and F. Harary, *Distance in Graphs* (Redwood City, CA: Addison-Wesley, 1990).
- [12] M. Lemańska, *Weakly convex and convex domination numbers*, Opuscula Math. **24** (2004) 181–188.
- [13] J. Bondy and G. Fan, *A sufficient condition for dominating cycles*, Discrete Math. **67** (1987) 205–208.
doi:10.1016/0012-365X(87)90029-X
- [14] E. Cockayne and S.T. Hedetniemi, *Towards a theory of domination in graphs*, Networks **7** (1977) 247–261.
doi:10.1002/net.3230070305
- [15] H. Walikar, B. Acharya and E. Samathkumar, *Recent Developments in the Theory of Domination in Graphs* (Allahabad, 1979).
- [16] T.L. Tachobo, F.P.Jamil and S. Canoy Jr., *Monophonic and geodetic domination in the join, corona and composition of graphs*, Ars Combin. **112** (2013) 13–32.

Received 28 December 2013

Revised 25 June 2014

Accepted 1 August 2014