

IMPROVED SUFFICIENT CONDITIONS FOR HAMILTONIAN PROPERTIES

JENS-P. BODE¹, ANIKA FRICKE²

AND

ARNFRIED KEMNITZ¹

¹*Computational Mathematics*
Technische Universität Braunschweig
38092 Braunschweig, Germany

²*Zentrum für erfolgreiches Lehren und Lernen*
Ostfalia Hochschule für angewandte Wissenschaften
38302 Wolfenbüttel, Germany

e-mail: jp.bode@tu-bs.de
Anika.Fricke@ostfalia.de
a.kemnitz@tu-bs.de

Abstract

In 1980 Bondy [2] proved that a $(k+s)$ -connected graph of order $n \geq 3$ is traceable ($s = -1$) or Hamiltonian ($s = 0$) or Hamiltonian-connected ($s = 1$) if the degree sum of every set of $k+1$ pairwise nonadjacent vertices is at least $((k+1)(n+s-1)+1)/2$. It is shown in [1] that one can allow exceptional $(k+1)$ -sets violating this condition and still implying the considered Hamiltonian property. In this note we generalize this result for $s = -1$ and $s = 0$ and graphs that fulfill a certain connectivity condition.

Keywords: Hamiltonian, traceable, Hamiltonian-connected.

2010 Mathematics Subject Classification: 05C45.

1. INTRODUCTION

Let $G = (V(G), E(G))$ be a finite simple graph. A path or cycle of a graph is called Hamiltonian if it contains each vertex of the graph exactly once. A graph is called traceable if it contains a Hamiltonian path, Hamiltonian if it contains a Hamiltonian cycle, and Hamiltonian-connected if it contains a Hamiltonian

path between each pair of vertices. The p -closure $\text{Cl}_p(G)$ is obtained from G by successively joining pairs of nonadjacent vertices with degree sum at least p as long as such pairs do exist.

The following result shows that some Hamiltonian properties are preserved by generating specific closures.

Theorem 1 [3]. *Let G be a graph of order n . The graph G is traceable, Hamiltonian, or Hamiltonian-connected if and only if $\text{Cl}_{n-1}(G)$ is traceable, $\text{Cl}_n(G)$ is Hamiltonian, or $\text{Cl}_{n+1}(G)$ is Hamiltonian-connected, respectively.*

The next theorem gives sufficient conditions for Hamiltonian properties depending on the independence number $\alpha(G)$ and the connectivity $\kappa(G)$.

Theorem 2 [4]. *If G is a graph of order $n \geq 3$ with independence number $\alpha(G)$ and connectivity $\kappa(G)$ with $\alpha(G) \leq \kappa(G) - s$, then G is traceable for $s = -1$, Hamiltonian for $s = 0$, and Hamiltonian-connected for $s = 1$.*

Bondy proved sufficient conditions for the Hamiltonicity of graphs with respect to independent sets of arbitrary size. We present a short proof of a generalization of this result which also contains the traceable and the Hamiltonian-connected case.

Let $\sigma_k(G)$ be the minimum sum of the degrees of any k independent vertices, that is,

$$\sigma_k(G) = \min \left\{ \sum_{v \in S} d(v) : S \subseteq V(G), S \text{ is independent, } |S| = k \right\}.$$

If G does not contain k independent vertices, then we set $\sigma_k(G) = \infty$.

Theorem 3 [2]. *Let G be a $(k + s)$ -connected graph of order $n \geq 3$. If*

$$\sigma_{k+1}(G) \geq \frac{1}{2}((k+1)(n+s-1) + 1),$$

then G is traceable for $s = -1$, Hamiltonian for $s = 0$, and Hamiltonian-connected for $s = 1$.

Proof. Assume that G contains a set of $k + 1$ pairwise independent vertices v_0, \dots, v_k with $d(v_i) + d(v_j) \leq n + s - 1$ for all $0 \leq i < j \leq k$. Then

$$k \cdot \sum_{0 \leq i \leq k} d(v_i) = \sum_{0 \leq i < j \leq k} d(v_i) + d(v_j) \leq \binom{k+1}{2} (n+s-1) = \frac{1}{2} k(k+1)(n+s-1),$$

which implies $\sigma_{k+1}(G) \leq \sum_{i=0}^k d(v_i) \leq \frac{1}{2} k(k+1)(n+s-1)$, a contradiction. It follows that there exists a pair of vertices having degree sum at least $n + s$ in each set of $k + 1$ independent vertices, that is, the closure $\text{Cl}_{n+s}(G)$ does not contain an

independent set of $k + 1$ vertices. Therefore, $\alpha(\text{Cl}_{n+s}(G)) \leq k \leq \kappa(\text{Cl}_{n+s}(G)) - s$ since $\text{Cl}_{n+s}(G)$ is $(k + s)$ -connected. Then, by Theorem 2, the graph $\text{Cl}_{n+s}(G)$ is traceable for $s = -1$, Hamiltonian for $s = 0$, and Hamiltonian-connected for $s = 1$. Application of Theorem 1 completes the proof. ■

Setting $k = 0$ and $k = 1$ in Theorem 3 results in the sufficient condition of Dirac [5] and Ore [7, 8], respectively.

In [1] it was asked whether it would be possible to allow $(k + 1)$ -sets of pairwise independent vertices in Theorem 3 violating the degree condition and still implying traceability, Hamiltonicity, or Hamiltonian-connectedness.

Let \bar{n}_{k+1}^s be the number of such sets, that is,

$$\bar{n}_{k+1}^s = \left| \left\{ \{v_0, \dots, v_k\} \text{ indep.} : \sum_{i=0}^k d(v_i) < \frac{1}{2}((k+1)(n+s-1)+1) \right\} \right|.$$

The following upper bounds for \bar{n}_{k+1}^s have been proved.

Theorem 4 [1]. *Let G be a $(k + s)$ -connected graph of order at least 3 and connectivity $\kappa(G) \geq 2$. If*

$$\bar{n}_{k+1}^s(G) < \binom{\kappa(G) - s + 1}{k + 1},$$

then G is traceable for $s = -1$, Hamiltonian for $s = 0$, and Hamiltonian-connected for $s = 1$.

If the minimum degree of the graph is sufficiently large, then this result can be improved.

Theorem 5 [1]. *Let G be a $(k + s)$ -connected graph, $k + s \geq 2$, of order $n \geq 3$ and minimum degree $\delta(G) \geq \frac{n+2(s+1)}{3}$. If*

$$\bar{n}_{k+1}^s(G) < \binom{\delta(G) - s + 1}{k + 1},$$

then G is traceable for $s = -1$, Hamiltonian for $s = 0$, and Hamiltonian-connected for $s = 1$.

The upper bounds for \bar{n}_{k+1}^s in Theorems 4 and 5 are tight for odd $k + s$ (see [1]).

In this note we generalize the result of Theorem 5 for $s = -1$ and $s = 0$ and graphs that fulfill a certain connectivity condition.

2. RESULTS

A cycle C is called a D_λ -cycle of a graph G if every component of $G - V(C)$ has less than λ vertices.

We use the following result on D_λ -cycles for the proof of Theorem 7.

Lemma 6 [6]. *Let G be a λ -connected graph of order $n \geq 3$ and minimum degree δ . If*

$$\delta \geq \frac{n + (\lambda - 1)\lambda}{\lambda + 1},$$

then G has a D_λ -cycle.

Theorem 7. *Let G be a λ -connected graph of order $n \geq 3$ and minimum degree $\delta = \delta(G) \geq \frac{n + (\lambda - 1)\lambda}{\lambda + 1}$. If*

$$\bar{n}_{k+1}^0(G) < \binom{\delta - \lambda + 3}{k + 1} + (\lambda - 2) \binom{\delta - \lambda + 2}{k},$$

then G is Hamiltonian.

Proof. If $\lambda = 1$, then G is Hamiltonian by Dirac's condition $\delta \geq \frac{n}{2}$ ($k = s = 0$ in Theorem 3). Therefore, let $\lambda \geq 2$.

Let $G' \cong \text{Cl}_n(G)$ be the n -closure of G . Assume that G' is not Hamiltonian. Due to Lemma 6, there is a D_λ -cycle in G' since $\delta' = \delta(G') \geq \delta \geq \frac{n + (\lambda - 1)\lambda}{\lambda + 1}$. Let C be a longest D_λ -cycle in G' . By the assumption, there exists a vertex $x \in V(G') \setminus V(C)$ which has $t \leq \lambda - 2$ neighbors in its component of $G' - V(C)$ and at least $\delta' - t$ neighbors on C . Let $v_1, \dots, v_{\delta' - t}$ be neighbors of x on C and v_i^+ be the successor of v_i on C with respect to a given orientation of the vertices of C . If there were adjacent vertices v_i^+ and v_j^+ or adjacent vertices x and v_i^+ , then this would imply the existence of a cycle being longer than C . Therefore, $\{x, v_1^+, \dots, v_{\delta' - t}^+\}$ is an independent set in G' . By the same reason none of the other at least t vertices of the component of $G' - V(C)$ containing x is adjacent to a vertex v_i^+ , $1 \leq i \leq \delta' - t$. Therefore, the number of $(k + 1)$ -sets of independent vertices of G' is at least $f(t) = \binom{\delta' - t + 1}{k + 1} + t \binom{\delta' - t}{k}$. In $\text{Cl}_n(G)$ the degree sum of every pair of nonadjacent vertices is at most $n - 1$ and hence, as in the proof of Theorem 3, the degree sum of every $(k + 1)$ -set of independent vertices is at most $\frac{1}{2}(k + 1)(n - 1)$. Hence $\bar{n}_{k+1}^0(G') \geq f(t)$. Since $f(t) - f(t + 1) = (t + 1) \binom{\delta' - t - 1}{k - 1} \geq 0$, the function $f(t)$ is decreasing and hence has its minimum for the largest possible t , that is, for $t = \lambda - 2$. Therefore,

$$\begin{aligned} \bar{n}_{k+1}^0(G) &\geq \bar{n}_{k+1}^0(G') \geq f(\lambda - 2) \geq \binom{\delta' - \lambda + 3}{k + 1} + (\lambda - 2) \binom{\delta' - \lambda + 2}{k} \\ &\geq \binom{\delta - \lambda + 3}{k + 1} + (\lambda - 2) \binom{\delta - \lambda + 2}{k}, \end{aligned}$$

which is a contradiction to the condition of the theorem.

Hence G' is Hamiltonian, which implies the Hamiltonicity of G according to Theorem 1. ■

Theorem 8. *Let G be a $(\lambda - 1)$ -connected graph of order $n = n(G) \geq 2$ and minimum degree $\delta = \delta(G) \geq \frac{n+(\lambda-2)\lambda}{\lambda+1}$. If*

$$\bar{n}_{k+1}^{-1}(G) < \binom{\delta - \lambda + 4}{k+1} + (\lambda - 2) \binom{\delta - \lambda + 3}{k},$$

then G is traceable.

Proof. Let $H \cong G + K_1$. We show that H fulfills all conditions of Theorem 7.

The graph H is λ -connected and has order $n(H) \geq 3$ and minimum degree $\delta(H) = \delta(G) + 1 \geq \frac{n(H)+(\lambda-1)\lambda}{\lambda+1}$. The $(k+1)$ -sets of independent vertices in G and H coincide since the vertex of K_1 is adjacent to all other vertices. Moreover, each $(k+1)$ -set of independent vertices in H having a degree sum less than $\frac{1}{2}((k+1)(n(H)-1)+1)$ has a degree sum less than $\frac{1}{2}((k+1)(n(G)-2)+1)$ since $d_H(u) = d_G(u) + 1$ for every vertex $u \in V(G)$ and $n(H) = n(G) + 1$. This implies

$$\begin{aligned} \bar{n}_{k+1}^0(H) &= \bar{n}_{k+1}^{-1}(G) < \binom{\delta(G)-\lambda+4}{k+1} + (\lambda - 2) \binom{\delta(G)-\lambda+3}{k} \\ &= \binom{\delta(H)-\lambda+3}{k+1} + (\lambda - 2) \binom{\delta(H)-\lambda+2}{k}. \end{aligned}$$

Therefore, H is Hamiltonian according to Theorem 7 implying the traceability of G . ■

Setting $\lambda = 2$ in Theorems 7 and 8 implies the result of Theorem 5 for the cases $s = 0$ and $s = -1$. Moreover, it turns out that the condition $k + s \geq 2$ in Theorem 5 can be omitted in these cases.

The preceding results can be summarized as follows.

Corollary 9. *Let G be a $(\lambda + s)$ -connected graph of order $n \geq 3 + s$ and minimum degree $\delta = \delta(G) \geq \frac{n+(\lambda+s-1)\lambda}{\lambda+1}$. If*

$$\bar{n}_{k+1}^s(G) < \binom{\delta - \lambda - s + 3}{k+1} + (\lambda - 2) \binom{\delta - \lambda - s + 2}{k},$$

then G is traceable for $s = -1$ and Hamiltonian for $s = 0$.

Up to now, we did not succeed in proving the statement of Corollary 9 for $s = 1$, that is, for Hamiltonian-connectedness.

REFERENCES

- [1] J.-P. Bode, A. Kemnitz, I. Schiermeyer and A. Schwarz, *Generalizing Bondy's theorems on sufficient conditions for Hamiltonian properties*, Congr. Numer. **203** (2010) 5–13.
- [2] J.A. Bondy, Longest paths and cycles in graphs of high degree, Research Report CORR 80-16 (Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, Canada, 1980).
- [3] J.A. Bondy and V. Chvátal, *A method in graph theory*, Discrete Math. **15** (1976) 111–135.
doi:10.1016/0012-365X(76)90078-9
- [4] V. Chvátal and P. Erdős, *A note on Hamiltonian circuits*, Discrete Math. **2** (1972) 111–113.
doi:10.1016/0012-365X(72)90079-9
- [5] G.A. Dirac, *Some theorems on abstract graphs*, Proc. London Math. Soc. **s3-2** (1952) 69–81.
doi:10.1112/plms/s3-2.1.69
- [6] P. Fraisse, D_λ -cycles and their applications for Hamiltonian graphs (LRI, Rapport de Recherche **276**, Centre d'Orsay, Université de Paris-Sud, 1986).
- [7] O. Ore, *Note on Hamiltonian circuits*, Amer. Math. Monthly **67** (1960) 55.
- [8] O. Ore, *Hamilton connected graphs*, J. Math. Pures Appl. **42** (1963) 21–27.

Received 20 January 2014

Revised 3 September 2014

Accepted 3 September 2014