Discussiones Mathematicae Graph Theory 35 (2015) 615–628 doi:10.7151/dmgt.1822

ON UNIQUE MINIMUM DOMINATING SETS IN SOME CARTESIAN PRODUCT GRAPHS

JASON T. HEDETNIEMI

Department of Mathematical Sciences Clemson University Clemson, South Carolina 29634, USA

e-mail: jhedetn@clemson.edu

Abstract

Unique minimum vertex dominating sets in the Cartesian product of a graph with a complete graph are considered. We first give properties of such sets when they exist. We then show that when the first factor of the product is a tree, consideration of the tree alone is sufficient to determine if the product has a unique minimum dominating set.

Keywords: vertex domination, graph products, trees.

2010 Mathematics Subject Classification: 05C69, 05C76, 05C05.

1. INTRODUCTION

In this paper, we show that if T is a nontrivial tree, then $T \Box K_n$ has a unique minimum dominating set if and only if T has a minimum dominating set D such that each vertex in D has at least n+1 external private neighbors with respect to D. The study of unique minimum vertex dominating sets began with Gunther, Hartnell, Markus and Rall in [12] where the authors established a method for recognizing unique γ -sets in trees, and provided a characterization of those trees which have a unique γ -set. Their work was later expanded upon by Fischermann in [3] where block graphs were considered, and by Fischermann and Volkmann in [8] where cactus graphs were considered. The maximum number of edges contained in graphs with unique γ -sets was studied in [5] and [11], and complexity results concerning unique γ -sets can be found in [6]. Uniqueness of other types of dominating sets has also been studied. For example, edge domination was studied in [17] and [7]. Distance k domination was analyzed in [7]. Total domination was first studied in [14] and later in [4]. Mixed domination was considered in [8], and paired domination was studied in [1]. Connections between unique minimum dominating sets and unique irredundant and independent dominating sets was studied in [10], while connections between maximum independent sets and unique upper dominating sets can be found in [9]. Finally, properties of unique domination were used in [16] and [15] to study properties of Roman dominating sets.

In the work to follow, we consider unique minimum dominating sets in graphs $G \square K_n$ where G is a connected, fnite, simple, nontrivial graph and K_n is the complete graph on n vertices. A characterization of the unique γ -sets in such graphs is considered in Section 3. Using this characterization, we then generalize a main result of [12] in Section 4, giving a method for recognizing a γ -set as unique when the first factor G is a tree. In Section 5, we consider the ways two such graphs, each having a unique minimum dominating set, can be combined while preserving a unique γ -set. Finally, in Section 6, we present the proof of our main result and characterize those trees whose Cartesian product with a complete graph has a unique γ -set.

2. NOTATION AND DEFINITIONS

Let G be a finite, simple graph with vertex set V(G) and edge set E(G). For any vertex u in G, the open neighborhood of u is the set N(u) defined by $N(u) = \{v : v \in V \}$ $uv \in E(G)$, and the closed neighborhood of u, denoted N[u], is the set $N(u) \cup \{u\}$. If S is a subset of V(G), then the open neighborhood of S is $\bigcup_{v \in S} N(v)$ and the closed neighborhood of S is $S \cup N(S)$; these are denoted by N(S) and N[S], respectively. Any subset D of V(G) with the property that N[D] = V(G) is called a *dominating set* of G. A dominating set of G of minimum cardinality is called a minimum dominating set or a γ -set of G, and its cardinality is denoted by $\gamma(G)$. If D is a dominating set of G and $x \in D$, then a private neighbor of x with respect to D (or just a private neighbor if the dominating set is clear from the context) is any vertex u that belongs to $N[x] - N[D - \{x\}]$. If $u \neq x$, then u is also called an *external* private neighbor of x with respect to D. We let epn(x, D)denote the set of external private neighbors of x with respect to D. A vertex in a dominating set need not have a private neighbor, but if the dominating set is minimal with respect to set inclusion, then each of its vertices has a private neighbor.

The Cartesian product of two graphs G_1 and G_2 is the graph $G_1 \square G_2$ whose vertex set is the Cartesian product of the sets $V(G_1)$ and $V(G_2)$ with two vertices. Two vertices (a_1, a_2) and (b_1, b_2) in $G_1 \square G_2$ adjacent if either $a_1 = b_1$ and $a_2b_2 \in E(G_2)$, or $a_2 = b_2$ and $a_1b_1 \in E(G_1)$. For i = 1, 2 we define the projections π_{G_i} : $G_1 \square G_2 \to G_i$ by $\pi_{G_i}((u_1, u_2)) = u_i$. Additionally, for $(u_1, u_2) \in V(G_1 \square G_2)$, we define the G_i -layer through (u_1, u_2) to be the induced subgraph

$$G_i^{(u_1,u_2)} = \left\langle \{(v_1,v_2) : \pi_{G_{3-i}}((v_1,v_2)) = \pi_{G_{3-i}}((u_1,u_2)) \} \right\rangle.$$

We note that if A is a dominating set of $G_1 \square G_2$, then $\pi_{G_i}(A)$ dominates G_i for i = 1 and i = 2. For other graph product terminology, we follow [13].

We consider graphs $G \square K_n$ where G is a connected, finite, simple graph. We assume that the vertex set of K_n is $\{1, 2, ..., n\}$ which we will denote by [n]. For $u \in V(G)$ and for $k \in [n]$, we denote the G-layer through (u, k) as G^k for notational convenience. We let \mathcal{U} denote the class of all finite simple graphs that have a unique minimum dominating set. If $G \in \mathcal{U}$, then we let UD(G) denote the unique γ -set for G.

Our main theorem, proven in Section 6, is as follows.

Theorem 1. Let n be a positive integer and let T be a nontrivial tree. The graph $T \Box K_n \in \mathcal{U}$ if and only if T has a minimum dominating set D such that for all $v \in D$, $|epn(v, D)| \ge n + 1$.

3. BASIC STRUCTURE

Suppose that $G \square K_n \in \mathcal{U}$. What can we say about $UD(G \square K_n)$? We begin with the following observation.

Lemma 2. If $G \square K_n \in U$, then there exists $S \subseteq V(G)$ such that $UD(G \square K_n) = S \times [n]$.

Proof. Denote $UD(G \Box K_n)$ by D. Without loss of generality, suppose that $(v, 1) \in D$ but $(v, 2) \notin D$. Let

$$D' = \{(x,1) : (x,2) \in D\} \cup \{(y,2) : (y,1) \in D\} \cup \{(w,j) : (w,j) \in D, 3 \le j \le n\}.$$

We claim that D' is also a γ -set for $G \square K_n$.

- If $x \in \pi_G(D)$, then by the definition of D', it follows that the K_n -layer through (x, 1) is contained in N[D'].
- If $x \notin \pi_G(D)$, then for $1 \leq j \leq n$, each (x, j) is dominated by some (v_j, j) in D. Thus, (x, 1) is dominated by $(v_2, 1)$ in D', (x, 2) is dominated by $(v_1, 2)$ in D', and (x, j) is dominated by (v_j, j) in D' for $3 \leq j \leq n$. Hence, every vertex in the K_n -layer through (x, 1) is contained in N[D'].

Thus, we see that D' is a γ -set of $G \square K_n$ distinct from D, proving our result. **Corollary 3.** If $G \square K_n \in \mathcal{U}$, then $\gamma(G \square K_n)$ is a multiple of n. Any subset A of $V(G \Box K_n)$ such that $A = S \times [n]$ for some subset S of V(G) is said to have the *stacked property*. Before proceeding to our next result, we recall the following lemma from [12].

Lemma 4 [12]. Let G be a graph with a unique γ -set D. Let [u, v] be any edge in G other than an edge connecting a vertex in D to one of its private neighbors. Let G^- be the graph obtained from G by deleting the edge [u, v]. Then G^- has D as the unique γ -set.

We now consider the following consequence of Lemma 2.

Proposition 5. If $G \square K_n \in \mathcal{U}$, then $G \in \mathcal{U}$. Moreover, $G \square K_m \in \mathcal{U}$ for $1 \le m \le n$.

Proof. Denote $UD(G \Box K_n)$ by D. By Lemma 2, there exists $S \subseteq V(G)$ such that $D = S \times [n]$. Thus, for any $(x, i) \in D$, the external private neighbors of (x, i) with respect to D all belong to G^i . Define H to be the graph

$$G \square K_n - \{(v, n)(v, j) : v \in V(G), 1 \le j \le n - 1\}.$$

We see that H is isomorphic to $(G \Box K_{n-1}) \cup G$. By Lemma 4, D is still the unique γ -set for H. The proposition follows by induction.

Suppose that $A \subseteq V(G \Box K_n)$ has the stacked property and that $\{v\} \times [n] \subseteq A$. If $(u, j) \in epn((v, j), A)$ for some j, then $(u, i) \in epn((v, i), A)$ for $1 \leq i \leq n$. Bearing this in mind, suppose that D is a γ -set of $G \Box K_n$ with the stacked property. Additionally, suppose that $(v, 1) \in D$ has $epn((v, 1), D) = \{(u_1, 1), (u_2, 1), \dots, (u_j, 1)\}$ for some $j \leq n$. This implies that $epn((v, i), D) = \{(u_1, i), (u_2, i), \dots, (u_j, i)\}$ for $2 \leq i \leq n$. The set D' defined by

$$D' = (D - \{(v, 1), (v, 2), \dots, (v, j)\}) \cup \{(u_1, 1), (u_2, 2), \dots, (u_j, j)\}$$

is a γ -set of $G \square K_n$ distinct from D. Thus, we have the following.

Lemma 6. If G is a connected, nontrivial graph such that $G \square K_n \in \mathcal{U}$, then for each element $v \in UD(G \square K_n)$,

$$|epn(v, UD(G \Box K_n))| \ge n+1.$$

The graph $K_{1,n+1} \square K_n$ demonstrates that this "bound" is sharp. The family of graphs $K_m \square K_n$, $m \ge n$, demonstrates that no condition on the number of external private neighbors for vertices in a minimum dominating set is, by itself, sufficient to force the product with K_n to have a unique γ -set. For use in the proof of Theorem 13 to follow, we note here the following. ON UNIQUE MINIMUM DOMINATING SETS IN SOME CARTESIAN PRODUCT ...619

Observation 7. If $v \in V(G)$ has at least n + 1 leaf neighbors, then $\{v\} \times [n]$ is contained in every γ -set of $G \square K_n$.

In [12], the authors prove the following lemma.

Lemma 8 [12]. Let D be a γ -set of a graph G. If for every $x \in D$, $\gamma(G - x) > \gamma(G)$, then D is the unique γ -set of G.

The following statement is a generalization of this result to our setting.

If $G \square K_n$ has a γ -set D satisfying the stacked property such that for every $v \in \pi_G(D)$, $\gamma(G \square K_n - (\{v\} \times [n])) > \gamma(G \square K_n)$, then D is the unique γ -set of $G \square K_n$.

This statement, however, does not hold for a general product $G \square K_n$. The graph G illustrated in Figure 1 provides a counterexample. Define H to be the graph $G \square K_2$. The set D defined by $D = \{1, 2, 3, 4, 5, 6\} \times \{1, 2\}$ is a γ -set satisfying the stacked property such that for every $v \in \pi_G(D), \gamma(H - \{(v, 1), (v, 2)\}) > \gamma(H)$. However, D is not a unique γ -set since the set $\{(1, 1), (1, 2), (2, 1), (2, 2), (3, 1), (3, 2), (4, 1), (5, 1), (6, 1), (10, 2), (14, 2), (18, 2)\}$ is also a γ -set of H.



In the next section, we will show that if G is a tree, then the conditions above do imply that $G \square K_n \in \mathcal{U}$. The following lemma will be used in the proof.

Lemma 9. If $G \square K_n$ has a γ -set D satisfying the stacked property such that for every $v \in \pi_G(D)$, $\gamma(G \square K_n - (\{v\} \times [n])) > \gamma(G \square K_n)$, then for all $y \in D$, $|epn(y, D)| \ge n + 1$.

Proof. Let $v \in \pi_G(D)$. Suppose for some $j \leq n$ that

$$epn((v, 1), D) = \{(u_1, 1), (u_2, 1), \dots, (u_j, 1)\}.$$

Since D satisfies the stacked property,

$$epn((v,i), D) = \{(u_1,i), (u_2,i), \dots, (u_j,i)\}$$

for $1 \leq i \leq n$. The set

$$(D - (\{v\} \times [n])) \cup \{(u_1, 1), (u_2, 2), \dots, (u_j, j), (u_j, j+1), \dots, (u_j, n)\}$$

is a dominating set of $G \square K_n - (\{v\} \times [n])$ of cardinality equal to |D|, a contradiction. Thus, our result follows.

Before we proceed to our first theorem, we need the following two lemmas, which are generalizations of Lemmas 3 and 4 from [12].

Lemma 10. Let $G \Box K_n \in \mathcal{U}$ and let $v \notin \pi_G(UD(G \Box K_n))$. For any subset B of $\{v\} \times [n], \gamma(G \Box K_n - B) = \gamma(G \Box K_n)$.

Proof. Suppose that $\gamma(G \Box K_n - B) < \gamma(G \Box K_n)$. This implies that $G \Box K_n - B$ is dominated by a set D' with $|D'| < |UD(G \Box K_n)|$. However, for any $(v, i) \in B$, $D' \cup \{(v, i)\}$ is a dominating set of $G \Box K_n$ distinct from $UD(G \Box K_n)$ of cardinality less than or equal to $|UD(G \Box K_n)|$, a contradiction. Thus, $\gamma(G \Box K_n - B) \ge \gamma(G \Box K_n)$. Since $UD(G \Box K_n)$ dominates $G \Box K_n - B$, we see that $\gamma(G \Box K_n - B) = \gamma(G \Box K_n)$.

Lemma 11. Let G be a connected, nontrivial graph, let $G \square K_n \in \mathcal{U}$, and let $v \in \pi_G(UD(G \square K_n))$. For any subset B of $\{v\} \times [n]$, $\gamma(G \square K_n - B) \ge \gamma(G \square K_n)$.

Proof. For the sake of contradiction, suppose that $\gamma(G \Box K_n - B) < \gamma(G \Box K_n)$ for some $B \subseteq \{v\} \times [n]$. If D' is a γ -set of $G \Box K_n - B$, then $|D'| < |UD(G \Box K_n)|$ and D' dominates all of the external private neighbors of the vertices in B with respect to $UD(G \Box K_n)$. However, for any $(v, i) \in B$, $D' \cup \{(v, i)\}$ is a γ -set of $G \Box K_n$ and $UD(G \Box K_n) \neq D' \cup \{(v, i)\}$, a contradiction.

4. Trees

In this section, we restrict our attention to graphs $T \Box K_n$ where T is a nontrivial tree. We prove a set of equivalences which can be used to determine whether a γ -set in $T \Box K_n$ is unique. This result, formulated as Theorem 13 below, is a generalization of the following theorem from [12], and as such, the notation and proof structure are similar.

Theorem 12 [12]. Let T be a tree of order at least 3. The following conditions are equivalent.

(1) T has a unique γ -set D.

620

- (2) T has a γ -set D for which every vertex $x \in D$ has at least two private neighbors other than itself.
- (3) T has a γ -set D for which every vertex $x \in D$ has the property that $\gamma(T-x) > \gamma(T)$.

Theorem 13. Let T be a nontrivial tree. The following conditions are equivalent.

- (1) $T \square K_n \in \mathcal{U}$.
- (2) $T \square K_n$ has a stacked γ -set D such that for all $v \in D$, $|epn(v, D)| \ge n + 1$.
- (3) $T \square K_n$ has a stacked γ -set A such that for every $v \in \pi_G(A)$, $\gamma(T \square K_n (\{v\} \times [n])) > \gamma(T \square K_n)$.

Proof. By Lemmas 2 and 6, we see that statement (1) implies statement (2). We first show that statement (2) implies statement (1). We proceed by induction on |V(T)|.

The base case is given by $T = K_{1,n+1}$ where the result holds. We note that for any other tree T on n+2 vertices, statement (2) does not hold for $T \square K_n$. Suppose then that the result has been shown whenever |V(T)| < r. Let T be a tree on r vertices for which there exists a subset $S \subseteq V(T)$ such that $S \times [n]$ is a γ -set for $T \square K_n$ and such that every element $v \in S \times [n]$ satisfies $|epn(v, S \times [n])| \ge n+1$. To simplify notation, we let $D = S \times [n]$ and $H = T \square K_n$. Suppose that H - D contains two vertices (u, 1), (v, 1) which are connected by the edge (u,1)(v,1). Let H(u) be the component of $(T-uv) \Box K_n$ containing (u, 1), and let H(v) be the component containing (v, 1). Let $D(u) = D \cap V(H(u))$ and $D(v) = D \cap V(H(v))$. We first claim that D(u) and D(v) are γ -sets for H(u)and H(v) respectively. To see this, note that D(u) and D(v) dominate H(u) and H(v). Additionally, if H(u), for example, had a γ -set A of cardinality smaller than |D(u)|, then $A \cup D(v)$ would be a dominating set of $T \square K_n$ smaller than D, a contradiction. Since all private neighbors with respect to D are preserved in the individual components, our induction hypothesis implies that D(u) and D(v) are the unique γ -sets for H(u) and H(v) respectively.

Assume now that D' is a γ -set of H distinct from D. If $D' \cap (\{u, v\} \times [n]) = \emptyset$ then $D' \cap V(H(u)) = D(u)$ and $D' \cap V(H(v)) = D(v)$, a contradiction. Thus, $D' \cap (\{u, v\} \times [n]) \neq \emptyset$.

If $D' \cap (\{u\} \times [n]) \neq \emptyset$, then $D' \cap V(H(u))$ dominates H(u) in which case $|D' \cap V(H(u))| > |D(u)|$. Similarly, if $D' \cap (\{v\} \times [n]) \neq \emptyset$, then $|D' \cap V(H(v))| > |D(v)|$.

If $D' \cap (\{u\} \times [n]) = \emptyset$ but $D' \cap (\{v\} \times [n]) \neq \emptyset$, then certainly $D' \cap V(H(u))$ dominates $H(u) - (\{u\} \times [n])$ in which case by Lemma 10, $|D' \cap V(H(u))| \ge |D(u)|$. Similarly, if $D' \cap (\{v\} \times [n]) = \emptyset$ but $D' \cap (\{u\} \times [n]) \neq \emptyset$, then $|D' \cap V(H(v))| \ge |D(v)|$. Thus, since $D' \cap (\{u, v\} \times [n]) \neq \emptyset$, we see that $|D'| = |D' \cap V(H(u))| + |D' \cap V(H(v))| > |D(u)| + |D(v)| = |D|$, a contradiction. Hence, in this case, D is the unique γ -set for H.

Our last case assumes there are no edges in H of the form (u, 1)(v, 1) with $(u, 1), (v, 1) \in V(H) - D$. In this case, let $(x, i) \in D$. If (y, i) is an external private neighbor of (x, i) with respect to D, then y is a leaf of T. Hence, $x \in V(T)$ has at least n + 1 leaf neighbors. As observed above, this implies that $\{x\} \times [n]$ is contained in every γ -set of H. Since $(x, i) \in D$ was arbitrary, we see that D is the unique γ -set of H. Hence, we have now shown that (1) and (2) are equivalent.

Assume now that statement (3) holds. By Lemma 9, statement (2) holds. Our work above then implies that statement (1) also holds. Thus, we next prove that statement (1) implies statement (3).

Let $T \square K_n \in \mathcal{U}$. Let $D = UD(T \square K_n)$ and let $H = T \square K_n$. By Lemma 2, there exists $S \subseteq V(T)$ such that $D = S \times [n]$. Suppose that $\{v\} \times [n] \subseteq D$. Partition $N((v, 1)) \cap V(G^1)$ as $epn((v, 1), D) \cup Q((v, 1))$. Let

$$epn((v, 1), D) = \{(p_1, 1), (p_2, 1), \dots, (p_m, 1)\}$$

and

$$Q((v,1)) = \{(q_1,1), (q_2,1), \dots, (q_k,1)\}\$$

We know that $m \ge n+1$ and that $k \ge 0$. Let $H(p_i)$, respectively $H(q_j)$, be the component of $H - (\{v\} \times [n])$ containing $(p_i, 1)$, respectively $(q_j, 1)$. For $1 \le i \le m$, let $D(p_i) = D \cap V(H(p_i))$ and define $D(q_j)$ similarly. Since T is a tree, we see that

$$\gamma(H) = |D| = n + \sum_{i=1}^{m} |D(p_i)| + \sum_{j=1}^{k} |D(q_j)|.$$

Since $H - (\{v\} \times [n])$ is the disjoint union

$$\left[\bigcup_{i=1}^m H(p_i)\right] \cup \left[\bigcup_{j=1}^k H(q_j)\right],$$

we can calculate $\gamma(H - (\{v\} \times [n]))$ by calculating $\gamma(H(p_i))$ and $\gamma(H(q_j))$ for each i and j and summing the results.

First, we consider $H(p_i)$. If $V(H(p_i)) = \{p_i\} \times [n]$, then $D(p_i) = \emptyset$. In this case, it is easy to see that $\gamma(H(p_i)) = 1 = |D(p_i)| + 1$.

If $V(H(p_i)) \neq \{p_i\} \times [n]$, then $D(p_i) \neq \emptyset$. Moreover, for each j such that $1 \leq j \leq n$, no neighbor of (p_i, j) in the graph $H(p_i)$ is in $D(p_i)$, since $(p_i, j) \in epn((v, j), D)$. Thus, $D(p_i)$ is not a γ -set for $H(p_i)$ since it does not dominate $(p_i, 1)$. Nevertheless, suppose that $\gamma(H(p_i)) = |D(p_i)|$, and let B be a γ -set of $H(p_i)$. It follows that $(D - D(p_i)) \cup B$ is a dominating set of H of cardinality equal to |D|, contradicting the uniqueness of D. Hence, $\gamma(H(p_i)) > |D(p_i)|$. Since $D(p_i) \cup \{(p_i, 1)\}$ dominates $H(p_i)$, we see, once again, that $\gamma(H(p_i)) = |D(p_i)| + 1$.

622

ON UNIQUE MINIMUM DOMINATING SETS IN SOME CARTESIAN PRODUCT ...623

Next, we consider $H(q_j)$. Since $(q_j, i) \notin epn((v, i), D)$ for $1 \leq i \leq n$, we see that $D(q_j)$ is a γ -set of $H(q_j)$. Moreover, for each $v \in D(q_j)$, $|epn(v, D(q_j))| \geq n+1$. Thus, $D(q_j)$ is the unique γ -set of $H(q_j)$, giving us that $\gamma(H(q_j)) = |D(q_j)|$. Thus, we can now compute $\gamma(H_{n-1}(\{v\}) \times [n])$:

Thus, we can now compute $\gamma(H - (\{v\} \times [n]))$:

$$\gamma(H - (\{v\} \times [n])) = \sum_{i=1}^{m} \gamma(H(p_i)) + \sum_{j=1}^{k} \gamma(H(q_j))$$

= $\sum_{i=1}^{m} (|D(p_i)| + 1) + \sum_{j=1}^{k} |D(q_j)|$
= $\gamma(H) + m - n$
 $\geq \gamma(H) + (n+1) - n$
= $\gamma(H) + 1 > \gamma(H).$

Thus, we see that statement (1) implies statement (3), and our proof is complete.

In Section 6 to follow, we will use this result to show that finding a γ -set in $T \square K_n$ is not required to determine whether $T \square K_n \in \mathcal{U}$. We will show that analysis of a γ -set of T will suffice.

5. Combining Graphs with Unique γ -Sets

Suppose that $G_1 \square K_n$ and $G_2 \square K_n$ have unique minimum dominating sets. In this section, we consider the ways in which these two graphs can be combined to produce a new graph having a unique minimum dominating set. We discuss four operations. Throughout this section, $G_1 \square K_n$ and $G_2 \square K_n$, denoted H_1 and H_2 respectively, are nontrivial graphs in \mathcal{U} . Let D_1 and D_2 denote the sets $UD(G_1 \square K_n)$ and $UD(G_2 \square K_n)$ respectively.

Operation 1. If $x \notin \pi_{G_1}(D_1)$ and $y \notin \pi_{G_2}(D_2)$, then $((G_1 \cup G_2) + xy) \Box K_n \in \mathcal{U}$ and $UD(((G_1 \cup G_2) + xy) \Box K_n) = D_1 \cup D_2$.

Proof. Let H denote the graph $((G_1 \cup G_2) + xy) \Box K_n$. First, we see that $D_1 \cup D_2$ dominates all of H. Let D be a γ -set for H. It follows that

$$|D| \le |D_1 \cup D_2| = |D_1| + |D_2|.$$

Without loss of generality, suppose that $|D \cap V(H_1)| \leq |D_1|$. Since the only vertices of H_1 that could be dominated from outside of H_1 are elements of $\{x\} \times [n]$, we see that either $D \cap V(H_1)$ dominates all of H_1 , or $D \cap V(H_1)$ fails to dominate a subset B of $\{x\} \times [n]$.

First, suppose that $D \cap V(H_1)$ dominates all of H_1 . Since H_1 has a unique γ -set, and since we are assuming $|D \cap V(H_1)| \leq |D_1|$, we have that $D \cap V(H_1) =$

 D_1 . However, if $D \cap V(H_1) = D_1$, then we also have $D \cap V(H_2) = D_2$ since $x \notin \pi_{G_1}(D_1)$. Thus, in this case, we have that $D = D_1 \cup D_2$.

Now suppose that $D \cap V(H_1)$ fails to dominate a subset B of $\{x\} \times [n]$. By Lemma 10, we have that $|D \cap V(H_1)| \ge |D_1|$. Since $|D| \le |D_1| + |D_2|$, we have that $|D \cap V(H_2)| \le |D_2|$. Note, however, that $D \cap V(H_2)$ intersects $\{y\} \times [n]$, in which case we have a set of cardinality at most $|D_2|$ that is distinct from D_2 and dominates H_2 . This contradicts the uniqueness of D_2 . Our result now follows.

Operation 2. Let $x \in \pi_{G_1}(D_1)$ and $y \in \pi_{G_2}(D_2)$. If u is a new vertex in neither G_1 nor G_2 , then $((G_1 \cup G_2) + \{ux, uy\}) \Box K_n \in \mathcal{U}$ and $UD(((G_1 \cup G_2) + \{ux, uy\}) \Box K_n) = D_1 \cup D_2$.

Proof. Let H denote the graph $((G_1 \cup G_2) + \{ux, uy\}) \Box K_n$. First, note that $D_1 \cup D_2$ dominates H. If D is a γ -set of H with $|D| < |D_1| + |D_2|$, then $D \cap (\{u\} \times [n]) \neq \emptyset$. Suppose that $\{(u, i_1), (u, i_2), \dots, (u, i_k)\} \subseteq D$. Then $\{(x, i_1), (x, i_2), \dots, (x, i_k)\}$ and $\{(y, i_1), (y, i_2), \dots, (y, i_k)\}$ need not be dominated from H_1 and H_2 respectively. However, by Lemma 11, we know that $|D \cap V(H_1)| \geq |D_1|$ and that $|D \cap V(H_2)| \geq |D_2|$. Thus, $|D| \geq |D_1| + |D_2| + k > |D_1 \cup D_2|$. Thus, no γ -set of H intersects $\{u\} \times [n]$. Hence, any γ -set of H intersects each of $V(H_1)$ and $V(H_2)$ in a γ -set, in which case $D = D_1 \cup D_2$.

Before we discuss the next operation, we need the following lemma.

Lemma 14. Let T be a tree, and let $T \Box K_n \in \mathcal{U}$. If $(v, i) \notin UD(T \Box K_n)$ is adjacent to at least two elements of $UD(T \Box K_n)$, then $(T - v)\Box K_n \in \mathcal{U}$ and $UD((T - v)\Box K_n) = UD(T\Box K_n)$.

Proof. Let H' denote the graph $(T-v)\Box K_n$ and let D denote the set $UD(T\Box K_n)$. By Lemma 10, we know that $\gamma(H') = \gamma(T\Box K_n)$. Thus, D is a γ -set for H'. We must show that D is the only γ -set for H'. Note that the removal of $(v, 1), (v, 2), \ldots, (v, n)$ from $T\Box K_n$ breaks $T\Box K_n$ into $k \geq 2$ components; call them H_1, H_2, \ldots, H_k .

We claim that for i = 1, 2, ..., k, $D_i = D \cap V(H_i)$ is the unique γ -set for H_i . Without loss of generality, consider D_1 . Clearly D_1 is a dominating set for H_1 . If D'_1 were a smaller dominating set of H_1 , then $D'_1 \cup D_2 \cup \cdots \cup D_k$ would be a smaller γ -set for $T \square K_n$. Thus, D_1 will be a γ -set of H_1 . By the same logic, D_1 is the unique γ -set for H_1 .

Thus, each H_i has D_i as its unique minimum dominating set, in which case H' has $D = D_1 \cup D_2 \cup \cdots \cup D_k$ as its unique γ -set.

Operation 3. Let G_2 be a tree. If $x \in \pi_{G_1}(D_1)$, $y \notin \pi_{G_2}(D_2)$, and y is a neighbor of at least two vertices in $\pi_{G_2}(D_2)$, then $((G_1 \cup G_2) + xy) \Box K_n \in \mathcal{U}$ and $UD(((G_1 \cup G_2) + xy) \Box K_n) = D_1 \cup D_2$.

On Unique Minimum Dominating Sets in Some Cartesian Product $\dots 625$

Proof. Let H denote the graph $((G_1 \cup G_2) + xy) \Box K_n$. Note that $D_1 \cup D_2$ dominates H. Let D be a γ -set of H. Suppose that $(\{y\} \times [n]) \cap D \neq \emptyset$. This implies that some subset of $\{x\} \times [n]$ will be dominated from outside of H_1 . By Lemma 11, we still have that $|D \cap V(H_1)| \geq |D_1|$. Additionally, $D \cap V(H_2)$ dominates H_2 , in which case $|D \cap V(H_2)| > |D_2|$ since D_2 is the unique γ -set for H_2 and $y \notin \pi_{G_2}(D_2)$. Thus, we have $|D| > |D_1 \cup D_2|$. This implies that no γ -set of H intersects $\{y\} \times [n]$. Hence, if D is a γ -set for H, then $D \cap V(H_1) = D_1$. Lemma 6 then implies that $D \cap V(H_2) = D_2$. Thus, $D_1 \cup D_2$ is the unique γ -set for H.

Operation 4. Let G_1 and G_2 be trees. If $x \in \pi_{G_1}(D_1)$ and $y \in \pi_{G_2}(D_2)$, then $((G_1 \cup G_2) + \{xy\}) \Box K_n \in \mathcal{U}$ and $UD(((G_1 \cup G_2) + \{xy\}) \Box K_n) = D_1 \cup D_2$.

Proof. Once again, let H denote the graph $((G_1 \cup G_2) + \{xy\}) \Box K_n$. Since $D_1 \cup D_2$ dominates H, we have that $\gamma(H) \leq |D_1 \cup D_2| = |D_1| + |D_2|$. Let A denote the set $\{x\} \times [n]$, let B denote the set $\{y\} \times [n]$, and suppose that D is a γ -set for H.

If $A \subseteq D$ and $B \subseteq D$, then $D \cap V(H_1)$ and $D \cap V(H_2)$ are γ -sets for H_1 and H_2 , respectively, in which case $D = D_1 \cup D_2$.

Suppose that $D \cap A = \emptyset$. This implies that $D \cap V(H_1)$ dominates $H_1 - A$. However, by Theorem 13, we know that $\gamma(H_1 - A) > \gamma(H_1) = |D_1|$. Additionally, in this case $D \cap V(H_2)$ is a γ -set of H_2 implying that $D \cap H_2 = D_2$. Thus, we have $|D| > |D_1| + |D_2| = |D_1 \cup D_2|$. The same contradiction arises if $D \cap B = \emptyset$.

This leaves us with one case to consider. Without loss of generality, suppose that $0 < |D \cap A| < |A|$ and that $D \cap B \neq \emptyset$. Then $D \cap V(H_1)$ dominates H_1 and $D \cap V(H_2)$ dominates H_2 . However, since D_1 and D_2 are the unique γ -sets for H_1 and H_2 respectively, and since $A \subseteq D_1$, we have that $|D \cap V(H_1)| > |D_1|$ and that $|D \cap V(H_2)| \ge |D_2|$. Thus, we have $|D| > |D_1 \cup D_2|$, a contradiction.

Thus, we have $D = D_1 \cup D_2$, which implies $D_1 \cup D_2$ is the unique γ -set for H.

6. MAIN RESULT

We are now able to prove our main theorem, which we restate for your convenience.

Theorem 1. Let n be a positive integer and let T be a nontrivial tree. The graph $T \Box K_n \in \mathcal{U}$ if and only if T has a minimum dominating set D such that for all $v \in D$, $|epn(v, D)| \ge n + 1$.

Proof. Suppose that $T \square K_n$ has a unique γ -set denoted UD. By Lemma 2, we know that UD satisfies the stacked property. Thus, there exists $S \subseteq V(T)$

such that $UD = S \times [n]$. Additionally, by Lemma 6, for every element $v \in UD$, $|epn(v, UD)| \ge n + 1$. This implies that for every element $w \in S$, $|epn(w, S)| \ge n + 1$. Finally, by the proof of Proposition 5, S is a γ -set for T.

Now suppose that T is a tree, and suppose that T has a γ -set D' for which every element in D' has at least n + 1 external private neighbors with respect to D'. Define H to be the graph $T \Box K_n$, and let $D = D' \times [n]$. Clearly D is a dominating set for H. Furthermore, since every element of D' has at least n + 1external private neighbors with respect to D' in T, every element of D has at least n + 1 external private neighbors with respect to D in H. Thus, if we can prove that D is a γ -set for H, Theorem 13 will imply that D is the unique γ -set for H. We do this by induction on the cardinality of T.

The base case is given by $T = K_{1,n+1}$ where the result holds. Thus, assume the result holds whenever |V(T)| < r. Let T be a tree on r vertices having a γ -set D' for which every element in D' has at least n+1 external private neighbors with respect to D'. By Theorem 12, D' is the unique γ -set for T. Let H be $T \square K_n$, and let D be defined as above. Consider a diametral path in T, call it $v_1v_2\cdots v_k$. Note that v_k is a leaf and cannot be an element of D'. This implies that $v_{k-1} \in D'$. In order for v_{k-1} to have at least n+1 external private neighbors, v_{k-1} must be adjacent to at least n-1 other leaves. Let A be the set $\{v_{k-1}\} \cup epn(v_{k-1}, D') \text{ and let } B = \{v \in N(v_{k-1}) : v \notin D', |N(v) \cap D'| \ge 2\}.$ We note that B equals either the empty set or $\{v_{k-2}\}$. By Theorem 12, $\{v_{k-1}\}$ and $D' - \{v_{k-1}\}$ are the unique minimum dominating sets for $T\langle A \rangle$ and $T - (A \cup B)$ respectively. By our induction hypothesis, $\{v_{k-1}\} \times [n]$ and $D - (\{v_{k-1}\} \times [n])$ are the unique minimum dominating sets for $T\langle A \rangle \Box K_n$ and $(T - (A \cup B)) \Box K_n$ respectively. Our original graph H can be reconstructed from $T\langle A\rangle \Box K_n$ and $(T - (A \cup B)) \Box K_n$ by performing at least one of the operations discussed in Section 5 above. Hence, D is the unique γ -set for H.

Theorem 1 implies that in order to determine whether $T \Box K_n \in \mathcal{U}$ it is sufficient to consider T alone. That is, one need only find a γ -set in T and count the number of external private neighbors for each vertex in the set. Since finding a minimum dominating set in a tree can be done in linear time (see [2]), we see that the problem of determining for which K_n , $T \Box K_n \in \mathcal{U}$ is polynomial.

Acknowledgements

Before concluding, the author would like to thank the anonymous referees for their suggestions, which helped improve this paper. He would also like to thank his two advisors, Dr. Kevin James of Clemson University and Dr. Doug Rall of Furman University, for their guidance. ON UNIQUE MINIMUM DOMINATING SETS IN SOME CARTESIAN PRODUCT ...627

References

- M. Chellali and T. Haynes, Trees with unique minimum paired-dominating sets, Ars Combin. 73 (2004) 3–12.
- [2] E. Cockayne, S. Goodman and S. Hedetniemi, A linear algorithm for the domination number of a tree, Inform. Process. Lett. 4 (1975) 41–44. doi:10.1016/0020-0190(75)90011-3
- M. Fischermann, Block graphs with unique minimum dominating sets, Discrete Math. 240 (2001) 247–251. doi:10.1016/S0012-365X(01)00196-0
- [4] M. Fischermann, Unique total domination graphs, Ars Combin. 73 (2004) 289–297.
- M. Fischermann, D. Rautenbach and L. Volkmann, Maximum graphs with a unique minimum dominating set, Discrete Math. 260 (2003) 197–203. doi:10.1016/S0012-365X(02)00670-2
- [6] M. Fischermann, D. Rautenbach and L. Volkmann, A note on the complexity of graph parameters and the uniqueness of their realizations, J. Combin. Math. Combin. Comput. 47 (2003) 183–188.
- M. Fischermann and L. Volkmann, Unique minimum domination in trees, Australas. J. Combin. 25 (2002) 117–124.
- [8] M. Fischermann and L. Volkmann, Cactus graphs with unique minimum dominating sets, Util. Math. 63 (2003) 229–238.
- [9] M. Fischermann and L. Volkmann, Unique independence, upper domination and upper irredundance in graphs, J. Combin. Math. Combin. Comput. 47 (2003) 237– 249.
- [10] M. Fischermann, L. Volkmann and I. Zverovich, Unique irredundance, domination, and independent domination in graphs, Discrete Math. 305 (2005) 190–200. doi:10.1016/j.disc.2005.08.005
- [11] M. Fraboni and N. Shank, Maximum graphs with unique minimum dominating set of size two, Australas. J. Combin. 46 (2010) 91–99.
- [12] G. Gunther, B. Hartnell, L. Markus and D. Rall, Graphs with unique minimum dominating sets, in: Proc. 25th S.E. Int. Conf. Combin., Graph Theory, and Computing, Congr. Numer. 101 (1994) 55–63.
- [13] R. Hammack, W. Imrich and S. Klavžar, Handbook of Product Graphs (CRC Press, 2011).
- [14] T. Haynes and M. Henning, Trees with unique minimum total dominating sets, Discuss. Math. Graph Theory 22 (2002) 233–246. doi:10.7151/dmgt.1172
- [15] M. Henning, Defending the Roman Empire from multiple attacks, Discrete Math.
 271 (2003) 101–115. doi:10.1016/S0012-365X(03)00040-2

- [16] M. Henning and S. Hedetniemi, Defending the Roman Empire—a new strategy, Discrete Math. 266 (2003) 239–251. doi:10.1016/S0012-365X(02)00811-7
- [17] J. Topp, Graphs with unique minimum edge dominating sets and graphs with unique maximum independent sets of vertices, Discrete Math. 121 (1993) 199-210. doi:10.1016/0012-365X(93)90553-6

Received 15 May 2014 Revised 23 December 2014 Accepted 23 December 2014