

HARDNESS RESULTS FOR TOTAL RAINBOW CONNECTION OF GRAPHS

LILY CHEN

School of Mathematics Science
Huaqiao University
Quanzhou 362021, China

e-mail: lily60612@126.com

BOFENG HUO

Department of Mathematics
Qinghai Normal University
Xining, 810008, China

e-mail: hbf@qhnu.edu.cn

AND

YINGBIN MA

College of Mathematics and Information Science
Henan Normal University
Xinxiang 453007, China

e-mail: mayingbincw@htu.cn

Abstract

A total-colored path is *total rainbow* if both its edges and internal vertices have distinct colors. The *total rainbow connection number* of a connected graph G , denoted by $trc(G)$, is the smallest number of colors that are needed in a total-coloring of G in order to make G *total rainbow connected*, that is, any two vertices of G are connected by a total rainbow path. In this paper, we study the computational complexity of total rainbow connection of graphs. We show that deciding whether a given total-coloring of a graph G makes it total rainbow connected is NP-Complete. We also prove that given a graph G , deciding whether $trc(G) = 3$ is NP-Complete.

Keywords: total rainbow connection, computational complexity.

2010 Mathematics Subject Classification: 05C15, 05C40, 68Q25, 68R10.

1. INTRODUCTION

All graphs considered in this paper are simple, finite and undirected. We follow the notation and terminology of Bondy and Murty [1] for those not described here.

An *edge-coloring* of a graph G is a mapping from the edges of G to some finite set of colors, where adjacent edges may have the same color. An edge-colored graph is *rainbow connected* if any two vertices are connected by a path whose edges have distinct colors. This concept of rainbow connection in graphs was introduced by Chartrand *et al.* in [4]. The *rainbow connection number* of a connected graph G , denoted by $rc(G)$, is the smallest number of colors that are needed in an edge-coloring of G in order to make G rainbow connected. Observe that $diam(G) \leq rc(G) \leq n - 1$, where $diam(G)$ denotes the diameter of G and n is the order of G . It is easy to verify that $rc(G) = 1$ if and only if G is a complete graph, and that $rc(G) = n - 1$ if and only if G is a tree. For an overview of the rainbow connection topic, we refer the reader to some new papers [7, 8, 13], and the survey and the monograph [14, 15].

A *vertex-coloring* of a graph G is a mapping from the vertices of G to some finite set of colors. If all pairs of adjacent vertices have distinct colors, we called the vertex-coloring *proper*. In [9], Krivelevich and Yuster proposed the concept of rainbow vertex-connection. A vertex-colored graph, not necessarily proper, is *rainbow vertex-connected* if any two vertices are connected by a path whose internal vertices have distinct colors. The *rainbow vertex-connection number* of a connected graph G , denoted by $rvc(G)$, is the smallest number of colors that are needed in a vertex-coloring of G in order to make G rainbow vertex-connected. An easy observation is that if G is of order n , then $rvc(G) \leq n - 2$ and $rvc(G) = 0$ if and only if G is a complete graph. Notice that $rvc(G) \geq diam(G) - 1$ with equality if the diameter is 1 or 2. There are some approaches to study the bounds of $rvc(G)$, we refer to [9, 12, 16].

Uchizawa *et al.* [18] and Liu *et al.* [16] introduced an analogous definition using total-colorings. A *total-coloring* of a graph G is a mapping from the vertices and edges of G to some finite set of colors. A total-colored graph is *total rainbow connected* if any two vertices are connected by a path whose edges and internal vertices have distinct colors. The *total rainbow connection number* of a connected graph G , denoted by $trc(G)$, is the smallest number of colors that are needed in a total-coloring of G in order to make G total rainbow connected. Observe that $trc(G) = 1$ if and only if G is a complete graph, and $trc(G) \geq 3$ if and only if G is not complete.

For the rainbow connection number and the rainbow vertex-connection number, some examples were given to show that there is no upper bound for one of the parameters in terms of the other, see [9]. In [16], Liu *et al.* compared $trc(G)$

with $rc(G)$ and $rvc(G)$. Notice that $trc(G) \geq \max\{rc(G), rvc(G)\}$. Liu *et al.* showed that for every sufficiently large s , there exists an example of a graph G with $trc(G) = rvc(G) = s$.

The computational complexity of rainbow connectivity and rainbow vertex connectivity has been studied. In [2], Caro *et al.* conjectured that computing $rc(G)$ is an NP-Hard problem, as well as that even deciding whether a graph has $rc(G) = 2$ is NP-Complete. In [3], Chakraborty *et al.* confirmed this conjecture and obtained the following theorems.

Theorem 1. *Given a graph G , deciding if $rc(G) = 2$ is NP-Complete. In particular, computing $rc(G)$ is NP-Hard.*

Given an edge-coloring of the graph, if the coloring is arbitrary, they showed that checking whether the coloring makes the graph rainbow connected is NP-Complete.

Theorem 2. *The following problem is NP-Complete: Given an edge-colored graph G , check whether the given coloring makes G rainbow connected.*

In [10], Li *et al.* considered bipartite graphs, and obtained the computational complexity results for bipartite graphs. Chen *et al.* [5] investigated the computational complexity of rainbow vertex-connection, and obtained the similar results.

Since computing $rc(G)$ and $rvc(G)$ is NP-Hard, a natural conjecture is that computing $trc(G)$ is also NP-Hard. In this paper, we consider the computational complexity of total rainbow connection of graphs and give some similar results. In Section 2, we show that deciding whether a given total-coloring of a graph G makes it total rainbow connected is NP-Complete. In Section 3, we prove that given a graph G , deciding whether $trc(G) = 3$ is NP-Complete.

2. TOTAL RAINBOW CONNECTION

Now, we give our first theorem.

Theorem 3. *The following problem is NP-Complete: Given a total-colored graph G , check whether the given coloring makes G total rainbow connected.*

We define the problem in Theorem 3 as TOTAL RAINBOW CONNECTION, and the problem in Theorem 2 as RAINBOW CONNECTION.

Problem 1. TOTAL RAINBOW CONNECTION.

Given: Total-colored graph G .

Decide: Whether G is total rainbow connected under the coloring?

Problem 2. RAINBOW CONNECTION.

Given: Edge-colored graph G .

Decide: Whether the coloring makes G rainbow connected?

Clearly, the problem of TOTAL RAINBOW CONNECTION is in NP. By Theorem 2, the problem of RAINBOW CONNECTION is NP-Complete. We reduce Problem 2 to Problem 1, which shows that Problem 1 is NP-Complete, concluding the proof of Theorem 3.

Lemma 4. *Problem 2 $\preceq$ Problem 1.*

Proof. Let G be a given graph with an edge-coloring c . We want to construct a graph G' with a total-coloring c' such that G' is total rainbow connected under the coloring c' if and only if G is rainbow connected under the coloring c .

Let $V = \{v_1, v_2, \dots, v_{n-1}, v_n\}$ be the vertex set of G , and $\alpha_1, \alpha_2, \dots, \alpha_n$ be n new colors that are not used in c . Let G' be a graph isomorphic to G . We define the coloring c' as follows: $c'(e) = c(e)$, for $e \in E(G)$ and $c'(v_i) = \alpha_i$, for $1 \leq i \leq n$.

Since $\alpha_1, \alpha_2, \dots, \alpha_n$ are n new colors, one can easily check that G is rainbow connected under the coloring c if and only if G' is total rainbow connected under the coloring c' . ■

3. TOTAL RAINBOW CONNECTION NUMBER 3

Notice that $trc(G) = 1$ if and only if G is a complete graph, and $trc(G) \geq 3$ if and only if G is not complete. By Theorem 1, the problem of deciding if $rc(G) = 2$ is NP-Complete. Correspondingly, we show that the problem of deciding if $trc(G) = 3$ is NP-Complete.

Theorem 5. *Given a graph G , deciding whether $trc(G) = 3$ is NP-Complete. Thus, computing $trc(G)$ is NP-Hard.*

We define the problem above as TOTAL RAINBOW CONNECTION NUMBER 3.

Problem 3. TOTAL RAINBOW CONNECTION NUMBER 3.

Given: Graph $G = (V, E)$.

Decide: Whether there is a total-coloring of G with three colors such that all pairs $\{u, v\} \in V^{(2)}$ ($V^{(2)}$ means the unordered pairs from V) are connected by a total rainbow path?

Clearly, Problem 3 is in NP. To show that it is NP-Complete, we need to define another two problems.

Problem 4. SUBSET TOTAL RAINBOW CONNECTION NUMBER 3.

Given: Graph $G = (V, E)$ and a set of pairs $P \subseteq V^{(2)}$.

Decide: Whether there is a total-coloring of G with three colors such that all pairs $\{u, v\} \in P$ are connected by a total rainbow path?

Problem 5. 3-COLORABILITY.

Given: Graph $G = (V, E)$.

Decide: Whether there is a vertex-coloring of G with three colors such that all pairs of adjacent vertices are assigned different colors?

In the following, we will reduce Problem 4 to Problem 3 and then reduce Problem 5 to Problem 4. Since 3-COLORABILITY is NP-Complete [6], we have that the problem of TOTAL RAINBOW CONNECTION NUMBER 3 is NP-Complete, which proves Theorem 5.

Lemma 6. *Problem 4 $\preceq$ Problem 3.*

Proof. Given a graph $G = (V, E)$ and a set of pairs $P \subseteq V^{(2)}$, we construct a graph $G' = (V', E')$ as follows.

For each vertex $v \in V$, we introduce a new vertex x_v ; for every pair $\{u, v\} \in V^{(2)} \setminus P$, we introduce a new vertex $x_{\{u, v\}}$. Set

$$V' = V \cup \{x_v : v \in V\} \cup \{x_{\{u, v\}} : \{u, v\} \in V^{(2)} \setminus P\}$$

and

$$E' = E \cup \{vx_v : v \in V\} \cup \{ux_{\{u, v\}}, vx_{\{u, v\}} : \{u, v\} \in V^{(2)} \setminus P\} \cup \{xy : x, y \in V' \setminus V\}.$$

In the following, we will prove that there exists a total-coloring of G' with three colors which makes G' total rainbow connected if and only if there is a total-coloring of G with three colors such that all pairs $\{u, v\} \in P$ are connected by a total rainbow path.

First suppose there is a total-coloring of G' with three colors which makes G' total rainbow connected. Observe that G is a subgraph of G' . For each pair $\{u, v\} \in P$, the paths of length not more than 2 that connect u and v have to be in G . Thus, under the coloring, all pairs in P are connected by a total rainbow path.

On the other hand, assume that $c : V \cup E \rightarrow \{1, 2, 3\}$ is a total-coloring of G such that all pairs $\{u, v\} \in P$ are connected by a total rainbow path. We extend the coloring c to a total-coloring $c' : V' \cup E' \rightarrow \{1, 2, 3\}$ in the following way: $c'(vx_v) = 1$ for all $v \in V$, $c'(ux_{\{u, v\}}) = 1$ and $c'(vx_{\{u, v\}}) = 2$ for all $\{u, v\} \in V^{(2)} \setminus P$, $c'(x_v) = c'(x_{\{u, v\}}) = 3$, $c'(xy) = 2$ for all $x, y \in V' \setminus V$. Now we show that G' is indeed total rainbow connected under this coloring. For any two vertices u and v , if $\{u, v\} \in P$, the total rainbow path connecting u and v in G is also the total rainbow path in G' . If $\{u, v\} \in V^{(2)} \setminus P$, then $ux_{\{u, v\}}v$ is a total rainbow path connecting u and v . If $u \in V$, $v \in V' \setminus V$, then ux_vv is a total

rainbow path connecting u and v . If $u, v \in V' \setminus V$, then uv is the total rainbow path. Hence, G' is total rainbow connected under the coloring c' .

This completes the proof of the lemma. ■

Lemma 7. *Problem 5 $\preceq$ Problem 4.*

Proof. Let $G = (V, E)$ be an instance of the 3-COLORABILITY problem. We construct an instance $G' = (V', E')$ of the SUBSET TOTAL RAINBOW CONNECTION NUMBER 3 problem.

Let $V = \{v_1, v_2, \dots, v_{n-1}, v_n\}$ be the vertex set of G . For each $v_i \in V$, we introduce three new vertices v_i^1, v_i^2, v_i^3 . For each edge $e = v_i v_j \in E$, we introduce a new vertex v_{ij} . We set

$$V' = \{v_i^1, v_i^2, v_i^3 : 1 \leq i \leq n\} \cup \{v_{ij} : v_i v_j \in E\}$$

and

$$E' = \{v_i^1 v_i^2, v_i^2 v_i^3 : 1 \leq i \leq n\} \cup \{v_i^2 v_{ij}, v_{ij} v_j^2 : v_i v_j \in E\}.$$

Now we define the set $P \subseteq V'^{(2)}$ as follows.

$$P = \{\{v_i^1, v_i^3\}, \{v_i^3, v_{ij}\}, \{v_j^1, v_j^3\}, \{v_j^3, v_{ij}\}, \{v_i^2, v_j^2\} : v_i v_j \in E\}.$$

If there is a vertex-coloring $c : V(G) \rightarrow \{1, 2, 3\}$ such that the adjacent vertices are assigned different colors, then we define a total-coloring c' of G' as follows. For $1 \leq i \leq n$, let $c'(v_i^1 v_i^2) = c(v_i)$, $c'(v_i^2) \in \{1, 2, 3\} \setminus \{c(v_i)\}$, $c'(v_i^2 v_i^3) \in \{1, 2, 3\} \setminus \{c(v_i), c'(v_i^2)\}$. For each $v_i v_j \in E$, let $c'(v_i^2 v_{ij}) = c(v_i)$, $c'(v_j^2 v_{ij}) = c(v_j)$, $c'(v_{ij}) \in \{1, 2, 3\} \setminus \{c(v_i), c(v_j)\}$. For all other vertices, we assign them the colors arbitrarily. It is easy to check that all $\{u, v\} \in P$ are connected by a total rainbow path.

On the other hand, suppose there is a total-coloring c' of G' with three colors such that all pairs $\{u, v\} \in P$ are connected by a total rainbow path, we define a vertex-coloring c of G as follows. For $1 \leq i \leq n$, $c(v_i) = c'(v_i^1 v_i^2)$. We show that this coloring is proper, that is, if $v_i v_j \in E$, then $c(v_i) \neq c(v_j)$. Indeed, for $v_i v_j \in E$, since $\{v_i^2, v_j^2\} \in P$, there is a total rainbow path connecting v_i^2 and v_j^2 , this path must be $v_i^2 v_{ij} v_j^2$, thus $c'(v_i^2 v_{ij}) \neq c'(v_j^2 v_{ij})$. Since $\{v_i^1, v_i^3\} \in P$, $\{v_j^1, v_j^3\} \in P$, and the total rainbow paths connecting v_i^1 and v_i^3 , v_j^1 and v_j^3 are $v_i^1 v_i^2 v_i^3$, $v_j^1 v_j^2 v_j^3$, we have $c'(v_i^1 v_i^2) = c'(v_i^2 v_{ij})$ since only three colors are used. Similarly, $c'(v_j^1 v_j^2) = c'(v_j^2 v_{ij})$. Then $c'(v_i^1 v_i^2) \neq c'(v_j^1 v_j^2)$. Therefore, $c(v_i) \neq c(v_j)$, thus c is a proper coloring of G , concluding the proof. ■

Acknowledgments

We thank the referees for their useful comments and suggestions which helped to improve the presentation of the manuscript.

This work was supported by NSFC (No. 11261047), the Scientific Research Funds of Huaqiao University (No. 14BS311, No. 14BS316), and Quanzhou Science and Technology Project (No. 2015Z115), and the Scientific Research Foundation for Ph.D. of Henan Normal University (No. qd14143).

REFERENCES

- [1] J.A. Bondy and U.S.R. Murty, *Graph Theory* (GTM 244, Springer, 2008).
- [2] Y. Caro, A. Lev, Y. Roditty, Zs. Tuza and R. Yuster, *On rainbow connection*, Electron J. Combin. **15** (2008) R57.
- [3] S. Chakraborty, E. Fischer, A. Matsliah and R. Yuster, *Hardness and algorithms for rainbow connectivity*, J. Comb. Optim. **21** (2011) 330–347.
doi:10.1007/s10878-009-9250-9
- [4] G. Chartrand, G.L. Johns, K.A. McKeon and P. Zhang, *Rainbow connection in graphs*, Math. Bohem. **133** (2008) 85–98.
- [5] L. Chen, X. Li and Y. Shi, *The complexity of determining the rainbow vertex-connection of graphs*, Theoret. Comput. Sci. **412** (2011) 4531–4535.
doi:10.1016/j.tcs.2011.04.032
- [6] M. Garey, D.S. Johnson and L.J. Stockmeyer, *Some simplified NP-complete graph problems*, Theoret. Comput. Sci. **1** (1976) 237–267.
doi:10.1016/0304-3975(76)90059-1
- [7] X. Huang, X. Li and Y. Shi, *Note on the hardness of rainbow connections for planar and line graphs*, Bull. Malays. Math. Sci. Soc. **88** (2015) 1235–1241.
doi:10.1007/s40840-014-0077-x
- [8] X. Huang, X. Li, Y. Shi, J. Yue and Y. Zhao, *Rainbow connections for outerplanar graphs with diameter 2 and 3*, Appl. Math. Comput. **242** (2014) 277–280.
doi:10.1016/j.amc.2014.05.066
- [9] M. Krivelevich and R. Yuster, *The rainbow connection of a graph is (at most) reciprocal to its minimum degree*, J. Graph Theory **63** (2010) 185–191.
- [10] S. Li, X. Li and Y. Shi, *Note on the complexity of deciding the rainbow (vertex-) connectedness for bipartite graphs*, Appl. Math. Comput. **258** (2015) 155–161.
doi:10.1016/j.amc.2015.02.015
- [11] X. Li, Y. Mao and Y. Shi, *The strong rainbow vertex-connection of graphs*, Util. Math. **93** (2014) 213–223.
- [12] X. Li and Y. Shi, *On the rainbow vertex-connection*, Discuss. Math. Graph Theory **33** (2013) 307–313.
doi:10.7151/dmgt.1664
- [13] X. Li and Y. Shi, *Rainbow connection in 3-connected graphs*, Graphs Combin. **29** (2013) 1471–1475.
doi:10.1007/s00373-012-1204-9

- [14] X. Li, Y. Shi and Y. Sun, *Rainbow connections of graphs: A survey*, Graphs Combin. **29** (2013) 1–38.
doi:10.1007/s00373-012-1243-2
- [15] X. Li and Y. Sun, *Rainbow Connections of Graphs* (New York, Springer Briefs in Math., Springer, 2012).
- [16] H. Liu, A. Mestre and T. Sousa, *Total rainbow k -connection in graphs*, Discrete Appl. Math. **174** (2014) 92–101.
doi:10.1016/j.dam.2014.04.012
- [17] I. Schiermeyer, *Rainbow connection in graphs with minimum degree three*, IWOCA 2009, Lecture Notes in Comput. Sci. **5874** (2009) 432–437.
- [18] K. Uchizawa, T. Aoki, T. Ito, A. Suzuki and X. Zhou, *On the rainbow connectivity of graphs: Complexity and FPT algorithms*, Algorithmica **67** (2013) 161–179.
doi:10.1007/s00453-012-9689-4

Received 10 March 2015

Revised 7 July 2015

Accepted 7 July 2015