

## THE EXISTENCE OF QUASI REGULAR AND BI-REGULAR SELF-COMPLEMENTARY 3-UNIFORM HYPERGRAPHS

LATA N. KAMBLE

*Department of Mathematics*  
*Abasaheb Garware College,*  
*Karve Road, Pune-411004*

**e-mail:** lata7429@gmail.com

CHARUSHEELA M. DESHPANDE

AND

BHAGYASHREE Y. BAM

*Department of Mathematics*  
*College of Engineering Pune*  
*Pune-411006*

**e-mail:** dcm.maths@coep.ac.in

### Abstract

A  $k$ -uniform hypergraph  $H = (V; E)$  is called self-complementary if there is a permutation  $\sigma : V \rightarrow V$ , called a complementing permutation, such that for every  $k$ -subset  $e$  of  $V$ ,  $e \in E$  if and only if  $\sigma(e) \notin E$ . In other words,  $H$  is isomorphic with  $H' = (V; V^{(k)} - E)$ . In this paper we define a bi-regular hypergraph and prove that there exists a bi-regular self-complementary 3-uniform hypergraph on  $n$  vertices if and only if  $n$  is congruent to 0 or 2 modulo 4. We also prove that there exists a quasi regular self-complementary 3-uniform hypergraph on  $n$  vertices if and only if  $n$  is congruent to 0 modulo 4.

**Keywords:** self-complementary hypergraph, uniform hypergraph, regular hypergraph, quasi regular hypergraph, bi-regular hypergraph.

**2010 Mathematics Subject Classification:** 05C65.

## 1. INTRODUCTION

Sachs [8] and Ringel [7] proved that a graph of order  $n$  is self-complementary if and only if  $n$  is congruent to 0 or 1 modulo 4. They also proved that a regular graph of order  $n$  is self-complementary if and only if  $n$  is congruent to 1 modulo 4.

Szymański and Wojda [9] proved that “A self-complementary 3-uniform hypergraph of order  $n$  exists if and only if  $n$  is congruent to 0 or 1 or 2 modulo 4.”

Potočník, and Šajana [6] raised the following question strengthening Hartman’s conjecture [2, 3] about the existence of large sets of (not necessarily isomorphic) designs.

**Question [6].** *Is it true that for every triple of integers  $t < k < n$  such that  $\binom{n-i}{k-i}$  is even for all  $i = 0, \dots, t$ , there exists a self-complementary  $t$ -subset-regular  $k$ -uniform hypergraph of order  $n$ ?*

The answer to the above question is affirmative for  $k = 2$  and  $t = 1$  (see [8]). The answer was proved affirmative also for the case  $k = 3$  and  $t = 1$  (see [6]). And in [4] it is shown that the answer to the question above is affirmative for the remaining case of 3-uniform hypergraphs, namely for the case  $k = 3, t = 2$ .

In this paper we digress a little from the case  $k = 3$  and  $t = 1$  to prove that a quasi-regular self-complementary 3-uniform hypergraph of order  $n$  exists if and only if  $n \geq 4$  and  $n$  is congruent to 0 modulo 4, and a bi-regular self-complementary 3-uniform hypergraph of order  $n$  exists if and only if  $n$  is congruent to 0 or 2 modulo 4.

## 2. PRELIMINARY DEFINITIONS AND RESULTS

**Definition** ( $k$ -uniform hypergraph). Let  $V$  be a finite set with  $n$  vertices. By  $V^{(k)}$  we denote the set of all  $k$ -subsets of  $V$ . A  $k$ -uniform hypergraph is a pair  $H = (V; E)$ , where  $E \subset V^{(k)}$ .  $V$  is called the vertex set, and  $E$  the edge set of  $H$ .

**Definition** (Degree of a vertex). The degree of a vertex  $v$  in a hypergraph  $H$  is the number of edges containing the vertex  $v$  and is denoted as  $d_H(v)$ .

**Definition** (Regular hypergraph). A hypergraph  $H$  is said to be regular if all vertices have the same degree.

**Definition** (Bi-regular hypergraph). A hypergraph  $H$  is said to be bi-regular if there exist two distinct positive integers  $d_1$  and  $d_2$  such that the degree of each vertex is either  $d_1$  or  $d_2$ .

**Definition** (Quasi regular hypergraph). A hypergraph  $H$  is said to be quasi regular if the degree of each vertex is either  $r$  or  $r - 1$  for some positive integer  $r$ .

It is clear that every quasi regular hypergraph is bi-regular.

**Definition** (Self-complementary  $k$ -uniform hypergraph). A  $k$ -uniform hypergraph  $H = (V; E)$  is called self-complementary if there exists a permutation  $\sigma : V \rightarrow V$ , called a complementing permutation, such that for every  $k$ -subset  $e$  of  $V$ ,  $e \in E$  if and only if  $\sigma(e) \notin E$ .

In other words,  $H$  is isomorphic to  $H' = (V; V^{(k)} - E)$ .

**Definition** (Tournament). A tournament is a directed graph  $(V, A)$  with the property that for all pairs of distinct vertices  $u, v \in V$ , either  $(u, v) \in A$  or  $(v, u) \in A$ .

Further, a tournament is said to be *self-converse* if there exists a bijection  $\varphi : V \rightarrow V$  such that for all distinct  $u, v \in V$ , we have  $(u, v) \in A$  if and only if  $(\varphi(u), \varphi(v)) \notin A$ .

Kocay [5] proved the following result on complementing permutations of self-complementary 3-uniform hypergraphs.

**Proposition 1** [5]. *A permutation  $\sigma$  is a complementing permutation of a self-complementary 3-uniform hypergraph if and only if*

- (i) *every cycle of  $\sigma$  has even length, or*
- (ii)  *$\sigma$  has 1 or 2 fixed points, and the length of all other cycles is a multiple of 4.*

Szymański and Wojda [9] proved the following result on the order of a self-complementary uniform hypergraph.

**Proposition 2** [9]. *Let  $k$  and  $n$  be positive integers,  $k \leq n$ . A  $k$ -uniform self-complementary hypergraph of order  $n$  exists if and only if  $\binom{n}{k}$  is even.*

**Remark 3.** For 3-uniform self-complementary hypergraph the Proposition 2 can be stated as “A 3-uniform self-complementary hypergraph of order  $n$  exists if and only if  $n \equiv 0$  or 1 or 2 (mod 4).”

The following remark is obvious and hence is stated without proof.

**Remark 4.** If  $H$  is a self-complementary 3-uniform hypergraph of order  $n$  with complementing permutation  $\sigma$ , then

- (i) for any vertex  $v$  in  $H$ ,  $d_H(v) + d_H(\sigma(v)) = \binom{n-1}{2}$ ,
- (ii) for any vertex  $v$  in  $H$ ,  $d_H(v) = d_H(\sigma^2(v)) = d_H(\sigma^4(v)) = \dots$  and  $d_H(\sigma(v)) = d_H(\sigma^3(v)) = d_H(\sigma^5(v)) = \dots$

Further, if  $x$  is a fixed point of  $\sigma$ , then  $d_H(x) = \frac{1}{2} \binom{n-1}{2}$ .

**Lemma 5.** *If  $H$  is a self-complementary 3-uniform hypergraph on  $n$  vertices, where  $n$  is congruent to 1 modulo 4 and  $n \geq 5$ , then  $H$  cannot be bi-regular.*

**Proof.** Let  $H$  be a self-complementary 3-uniform hypergraph on  $n$  vertices where  $n$  is congruent to 1 modulo 4, i.e.,  $n = 4m + 1$ ,  $m \in \mathbb{N}$ . Let  $\sigma : V(H) \rightarrow V(H)$  be its complementing permutation. By Proposition 1,  $\sigma$  necessarily has one fixed point, say  $x$ .

From Remark 4(ii)  $d_H(x) = m(4m - 1)$ . For  $H$  to be bi-regular either  $d_1 = m(4m - 1)$  or  $d_2 = m(4m - 1)$ . Without loss of generality let  $d_1 = m(4m - 1)$ . Since there are only two types of degrees  $d_1$  and  $d_2$ , for any other vertex  $v$ ,  $d_v(H)$  is  $d_1$  or  $d_2$ . By Remark 4(i) we have,  $d_1 + d_2 = \frac{4m(4m-1)}{2}$  which gives  $d_2 = 2m(4m - 1) - m(4m - 1) = m(4m - 1) = d_1$ . Hence  $H$  cannot be bi-regular. ■

### 3. EXISTENCE OF A QUASI REGULAR AND BI-REGULAR SELF-COMPLEMENTARY 3-UNIFORM HYPERGRAPH

The following theorem gives a necessary and sufficient condition on the order  $n$  of a quasi regular self-complementary 3-uniform hypergraph. This theorem actually gives a construction of a quasi regular self-complementary 3-uniform hypergraph of desirable order.

**Theorem 6.** *There exists a quasi regular self-complementary 3-uniform hypergraph of order  $n$  if and only if  $n \geq 4$  and  $n \equiv 0 \pmod{4}$ .*

**Proof.** Let  $H$  be a quasi regular self-complementary 3-uniform hypergraph on  $n$  vertices such that degree of each vertex is either  $r$  or  $r - 1$  for some positive integer  $r$ .




Figure 1. The types of triples making up the edge set of a quasi regular self-complementary 3-uniform hypergraph on  $n = 4m$  vertices.

Let  $\sigma : V(H) \rightarrow V(H)$  be a complementing permutation of  $H$ . By Proposition 1,  $\sigma$  has (i) every cycle of even length, or (ii) 1 or 2 fixed points and the

length of all the other cycles is a multiple of 4. By Remark 3, we know that a self-complementary 3-uniform hypergraph exists if and only if  $n \equiv 0 \pmod{4}$  or  $n \equiv 1 \pmod{4}$ , or  $n \equiv 2 \pmod{4}$ . Lemma 5 shows that  $n$  is not congruent to 1 modulo 4.

If  $n \equiv 2 \pmod{4}$ , i.e.,  $n = 4m + 2$ ,  $m \in \mathbb{N}$ , then either  $\sigma$  has 2 fixed points and the length of all other cycles is a multiple of 4 or  $\sigma$  has all cycles of even length.

If  $\sigma$  has 2 fixed points, then both must have the same degree and for some other vertex  $v$ ,  $d_H(v) \neq d_H(\sigma(v))$  otherwise  $H$  will be regular. Since there are only two possible degrees  $r$  and  $r - 1$ , from Remark 4 we get that  $r + r - 1 = \binom{n-1}{2} = \binom{4m+1}{2}$ , i.e.,  $2r - 1 = 2m(4m + 1)$ , a contradiction.

If  $\sigma$  has all cycles of even length, then again we get the same contradiction.

Hence, if there exists a quasi regular self-complementary 3-uniform hypergraph on  $n$  vertices, then  $n \equiv 0 \pmod{4}$ .

For the converse, we construct a quasi regular self-complementary 3-uniform hypergraph on  $n$  vertices where  $n \equiv 0 \pmod{4}$ .

Let  $m$  be a positive integer such that  $n = 4m$  and  $V = V_0 \cup V_1 \cup V_2 \cup V_3$ , where  $V_i = \{v_j^i : j \in \mathbb{Z}_m\}$ ,  $i \in \mathbb{Z}_4$ .

For every pairwise distinct triple  $i, i', i'' \in \mathbb{Z}_4$  we define the following subsets of  $V^{(3)}$ :

$$\begin{aligned} E_i &= V_i^{(3)}, \\ E_{(i,i')} &= \{\{v_{j_1}^i, v_{j_2}^i, v_{j'}^{i'}\} : j_1, j_2, j' \in \mathbb{Z}_m, j_1 \neq j_2\}, \\ E_{(i,i',i'')} &= \{\{v_j^i, v_{j'}^{i'}, v_{j''}^{i''}\} : j, j', j'' \in \mathbb{Z}_m\}. \end{aligned}$$

Let us denote

$$E = E_0 \cup E_1 \cup E_{(2,1)} \cup E_{(2,3)} \cup E_{(3,0)} \cup E_{(3,2)} \cup E_{(1,3)} \cup E_{(0,2)} \cup E_{(0,1,3)} \cup E_{(0,1,2)}.$$

Let  $H$  be the 3-uniform hypergraph with vertex set  $V$  and edge set  $E$ . Figure 1 explains the construction of the hypergraph  $H$ . We show that  $H$  is quasi regular. Take any vertex  $v_j^i$ .

*Case (i)* If  $i \in \{0, 1\}$ , then the vertex  $v_j^i$  lies in  $\binom{m-1}{2}$  triples of  $E_i$ ,  $(m-1)m$  triples of  $E_{(i,i')}$ ,  $\binom{m}{2}$  triples of  $E_{(i',i)}$  and  $2m^2$  triples of  $E_{(i,i',i'')}$ . Hence, for every vertex  $v_j^i$  in  $H$  with  $i \in \{0, 1\}$ , we have

$$d_H(v_j^i) = \binom{m-1}{2} + \binom{m}{2} + m(m-1) + 2m^2 = 4m^2 - 3m + 1.$$

*Case (ii)* If  $i \in \{2, 3\}$ , then the vertex  $v_j^i$  lies in  $2(m-1)m$  triples of  $E_{(i,i')}$ ,  $2\binom{m}{2}$  triples of  $E_{(i',i)}$  and  $m^2$  triples of  $E_{(i,i',i'')}$ . Hence for every vertex  $v_j^i$  in  $H$  with  $i \in \{2, 3\}$ , we obtain

$$d_H(v_j^i) = 2(m-1)m + 2\binom{m}{2} + m^2 = 4m^2 - 3m.$$

Thus  $H$  is quasi regular with degrees  $r = 4m^2 - 3m + 1$  and  $r - 1 = 4m^2 - 3m$ . To prove that  $H$  is self-complementary, we define a permutation  $\phi : V \rightarrow V$  by  $\phi(v_j^0) = v_j^3$ ,  $\phi(v_j^1) = v_j^2$ ,  $\phi(v_j^2) = v_j^1$  and  $\phi(v_j^3) = v_j^0$ , for all  $j \in \mathbb{Z}_m$ . Then  $\phi$  is a complementing permutation of  $H$  and  $H$  is self-complementary. ■

In the next theorem we give a necessary and sufficient condition on the order  $n$  of a bi-regular 3-uniform hypergraph to be self-complementary. In this theorem we shall use the following result by Alspach [1] on existence of a regular self-converse tournament.

**Theorem 7** (Alspach [1]). *There exists a regular self-converse tournament with  $n$  vertices for every odd integer  $n$ .*

**Theorem 8.** *There exists a bi-regular self-complementary 3-uniform hypergraph of order  $n$  if and only if either  $n \equiv 0 \pmod{4}$  or  $n \equiv 2 \pmod{4}$  and  $n \geq 4$ .*

**Proof.** Necessity follows from Lemma 5 and Remark 3. Conversely, let  $n \equiv 0 \pmod{4}$ . The self-complementary 3-uniform hypergraph constructed in Theorem 6 is quasi regular and hence biregular.

Let  $n \equiv 2 \pmod{4}$ . Then  $n = 4m + 2 = 2k$  where  $k = 2m + 1$  is odd. Let  $V = V_0 \cup V_1$ , where  $V_i = \{v_j^i : j \in \mathbb{Z}_k\}$ ,  $i \in \mathbb{Z}_2$ . By Theorem 7, there exists a regular self-converse tournament  $T = (\mathbb{Z}_k, A)$ .

For  $i \in \mathbb{Z}_2$ , we define the following subsets of  $V^{(3)}$ :

$$\begin{aligned} E_i &= V_i^{(3)}, \\ E_{(i,i+1)} &= \{\{v_{j_1}^i, v_{j_2}^i, v_j^{i+1}\} : j_1, j_2, j \in \mathbb{Z}_k, j_1, j_2, j \text{ pairwise distinct}\}, \\ E_A &= \{\{v_{k_1}^i, v_{k_2}^i, v_{k_1}^{i+1}\} : (k_1, k_2) \in A, i \in \mathbb{Z}_2\}. \end{aligned}$$

Let

$$E = E_0 \cup E_{(0,1)} \cup E_A.$$

Let  $H$  be the 3-uniform hypergraph with vertex set  $V$  and edge set  $E$ . Figure 2 explains the construction of the hypergraph  $H$ . We show that  $H$  is bi-regular. Let  $v_j^i$  be an arbitrary vertex of  $H$ .

*Case (i)* If  $i = 0$ , then the vertex  $v_j^0$  lies in  $\binom{k-1}{2}$  triples of  $E_0$ ,  $(k-1)(k-2)$  triples of  $E_{(0,1)}$  and  $\frac{3(k-1)}{2}$  triples of  $E_A$ . Hence

$$d_H(v_j^0) = \binom{k-1}{2} + (k-1)(k-2) + \frac{3(k-1)}{2} = \frac{3(k-1)^2}{2}.$$

Case (ii) If  $i = 1$ , then the vertex  $v_j^1$  lies in  $\binom{k-1}{2}$  triples of  $E_{(0,1)}$ ,  $\frac{3(k-1)}{2}$  triples of  $E_A$ . Therefore,

$$d_H(v_j^1) = \binom{k-1}{2} + \frac{3(k-1)}{2} = \frac{k^2-1}{2}.$$




Figure 2. The types of triples making up the edge set of a bi-regular self-complementary 3-uniform hypergraph on  $n = 4m + 2$  vertices.

This proves that  $H$  is bi-regular with degrees  $d_1 = \frac{3(k-1)^2}{2}$  and  $d_2 = \frac{k^2-1}{2}$ .

Let  $\varphi : \mathbb{Z}_k \rightarrow \mathbb{Z}_k$  be an arc-reversing mapping of the tournament  $T$ ; that is,  $\varphi$  is a bijection on  $\mathbb{Z}_k$  such that  $\varphi(a) \notin A$  for all  $a \in A$ .

To prove that  $H$  is self-complementary, we define a permutation  $\phi : V \rightarrow V$  by  $\phi(v_j^i) = v_{\varphi(j)}^{i+1}$  for  $i \in \mathbb{Z}_2$  and  $j \in \mathbb{Z}_k$ .  $\phi$  interchanges the sets  $E_1$  and  $E_0$ , and also the sets  $E_{(0,1)}$  and  $E_{(1,0)}$ . Furthermore, for all  $(k_1, k_2) \in A$  and  $i \in \mathbb{Z}_2$ , since  $\varphi$  is arc-reversing,  $\phi$  maps the triple  $\{v_{k_1}^i, v_{k_2}^i, v_{k_1}^{i+1}\} \in E_A$  to the triple  $\{v_{\varphi(k_1)}^{i+1}, v_{\varphi(k_2)}^{i+1}, v_{\varphi(k_1)}^i\} \notin E_A$ . It follows that  $\phi$  is a complementing permutation of  $H$  and therefore  $H$  is self-complementary. ■

### Acknowledgment

We are grateful to the referees for their helpful comments and Professor N.S. Bhavne for many helpful discussions.

### REFERENCES

- [1] B. Alspach, *On point-symmetric tournaments*, Canad. Math. Bull. **13** (1970) 317–323.  
doi:10.4153/CMB-1970-061-7
- [2] C.J. Colbourn and J.H. Dinitz, *The CRC Handbook of Combinatorial Designs* (CRC Press, Boca Raton, 1996).

- [3] A. Hartman, *Halving the complete design*, Ann. Discrete Math. **34** (1987) 207–224.  
doi:10.1016/s0304-0208(08)72888-3
- [4] M. Knor and P. Potočník, *A note on 2-subset-regular self-complementary 3-uniform hypergraphs*, Ars Combin. **11** (2013) 33–36.
- [5] W. Kocay, *Reconstructing graphs as subsumed graphs of hypergraphs, and some self-complementary triple systems*, Graphs Combin. **8** (1992) 259–276.  
doi:10.1007/BF02349963
- [6] P. Potočník and M. Šajana, *The existence of regular self-complementary 3-uniform hypergraphs*, Discrete Math. **309** (2009) 950–954.  
doi:10.1016/j.disc.2008.01.026
- [7] G. Ringel, *Selbstkomplementäre Graphen*, Arch. Math. **14** (1963) 354–358.  
doi:10.1007/BF01234967
- [8] H. Sachs, *Über selbstkomplementäre Graphen*, Publ. Math. Debrecen **9** (1962) 270–288.
- [9] A. Szymański and A.P. Wojda, *A note on  $k$ -uniform self-complementary hypergraphs of given order*, Discuss. Math. Graph Theory **29** (2009) 199–202.  
doi:10.7151/dmgt.1440

Received 24 November 2014

Revised 1 August 2015

Accepted 1 August 2015