

ON THE H -FORCE NUMBER OF HAMILTONIAN GRAPHS AND CYCLE EXTENDABILITY

ERHARD HEXEL

Department of Mathematics
Technische Universität Ilmenau
Postfach 0565, D-98684 Ilmenau, Germany
e-mail: erhard.hexel@tu-ilmenau.de

Abstract

The H -force number $h(G)$ of a hamiltonian graph G is the smallest cardinality of a set $A \subseteq V(G)$ such that each cycle containing all vertices of A is hamiltonian. In this paper a lower and an upper bound of $h(G)$ is given. Such graphs, for which $h(G)$ assumes the lower bound are characterized by a cycle extendability property. The H -force number of hamiltonian graphs which are exactly 2-connected can be calculated by a decomposition formula.

Keywords: cycle, hamiltonian graph, H -force number, cycle extendability.

2010 Mathematics Subject Classification: 05C45.

1. INTRODUCTION

Throughout this paper, only finite graphs without loops or multiple edges are considered. The number of vertices of a graph G , i.e., its *order* will be denoted by n . We use the standard graph terminology according to [3].

Let G be a hamiltonian graph with vertex set $V = V(G)$ and edge set $E = E(G)$. A nonempty vertex set $X \subseteq V(G)$ is called a *hamiltonian cycle enforcing set* (for short, *H -force set*) of G if every X -cycle of G (i.e., a cycle of G containing all vertices of X) is a hamiltonian one. Let $h(G)$ denote the smallest cardinality of an H -force set of G and call it the *H -force number* of G . The concepts of H -force set and H -force number were first given by Fabrici *et al.* (see [4]) and studied there for several special families of hamiltonian graphs. Timková (see [9]) determined the H -force number of generalized dodecahedral graphs. Note also, that the concepts of H -force set and H -force number were extended to hamiltonian digraphs and hypertournaments in [10] and [7], respectively.

The authors in [4] observed that the H -force number $h(G)$ of a hamiltonian graph G satisfies

- $h(G) = 1$ if and only if G is a cycle,
- $h(G) = n$ if and only if G is 1-hamiltonian (that is, if G is hamiltonian and $G - v$ is hamiltonian for every $v \in V$).

For a hamiltonian graph G , we define sets $S = S(G) = \{x \in V \mid G - x \text{ is hamiltonian}\}$ and $T = T(G) = \{x \in V \mid G - x \text{ is 2-connected}\}$. Then, we have $S \subseteq T$. Let $s(G) = |S(G)|$ and $t(G) = |T(G)|$.

Proposition 1. *Let G be a hamiltonian graph and P be a path of G containing no branch vertex of G , i.e., no vertex of degree at least 3 in G . Then, every smallest H -force set $F \subseteq V(G)$ contains at most one vertex of P .*

Let $\mathcal{H}$ be the family of hamiltonian graphs that do not contain adjacent vertices of degree 2. Also, let G' be the graph formed from a hamiltonian graph G by replacing each maximal path not containing a branch vertex by a single vertex. Then, G' is hamiltonian and has no adjacent vertices of degree 2, so $G' \in \mathcal{H}$. Because $h(G') = h(G)$, it is sufficient to restrict our study to the family $\mathcal{H}$.

The main results of this paper are Theorems 2, 7, 8 and 11. Theorem 2 shows that $s(G)$ and $t(G)$ form bounds for the H -force number $h(G)$. After this theorem, we discuss some consequences. Theorem 7 contains a decomposition formula for the H -force number of hamiltonian graphs which are exactly 2-connected. In Theorem 8 hamiltonian graphs G for which $S(G)$ is an H -force set are characterized by a cycle extendability property. Eventually, a sum formula for hamiltonian graphs G with $s(G) < h(G)$ is proved in Theorem 11.

2. RESULTS AND PROOFS

Theorem 2. *Let $G \in \mathcal{H}$. Then*

$$s(G) \leq h(G) \leq t(G).$$

The proof of this theorem requires the following exchange property.

Lemma 3. *Let $G \in \mathcal{H}$ and let $F \subseteq V$ be a smallest H -force set of G . Then, for every vertex $v \in F \setminus T$ there exists a vertex $u \in T$ such that $(F \setminus \{v\}) \cup \{u\}$ is an H -force set of G .*

Proof. Suppose there exists a vertex $v \in V \setminus T$. Then G is exactly 2-connected. Let C be any fixed hamiltonian cycle of G and w be a cut-vertex of $G - v$. Then, C consists of two v - w -paths P_1 and P_2 both of which have at least one inner vertex but no inner vertex in common. Since G is not a cycle, C has a chord.

But, there is no chord connecting an inner vertex of P_1 with an inner vertex of P_2 . Let $F \subseteq V$ be a smallest H -force set of G (i.e., $|F| = h(G)$) and suppose $v \in F$.

Case 1. The cut-vertex w of $G - v$ can be chosen so that each P_i , for $i = 1, 2$, has a chord of C , say $x_i y_i$. Then, the subpath (x_i, y_i) of P_i contains an inner vertex z_i such that $z_i \in F$. Otherwise, the x_i - y_i -path on C which passes v forms together with $x_i y_i$ a non-hamiltonian F -cycle. By the choice of F , $F \setminus \{v\}$ is not an H -force set of G , i.e., G contains a non-hamiltonian $(F \setminus \{v\})$ -cycle C' not passing v . Since z_1 and z_2 belong to different components of $G - \{v, w\}$ and since w is a cut-vertex of $G - v$, every z_1 - z_2 -path of $G - v$ is passing w which contradicts the fact that C' is a cycle.

Case 2. By any choice of the cut-vertex w of $G - v$ only one of P_1 and P_2 has a chord. Suppose for a fixed w that P_1 has no chord. Then P_1 has only one inner vertex u where $d_G(u) = 2$. Since every hamiltonian cycle of G passes the edge uv , $F' := (F \setminus \{v\}) \cup \{u\}$ is also an H -force set of G . Moreover, we have $u \in T$ because otherwise there exists a cut-vertex z of $G - u$ which is also a cut-vertex of $G - v$. Hence, C consists of two v - z -paths (with no common inner vertices) such that both of them have at least one chord, a contradiction. That proves the assertion. ■

Proof of Theorem 2. Let $F \subseteq V$ be any smallest H -force set of G . Suppose that S contains a vertex x such that $x \notin F$. A hamiltonian cycle C of $G - x$ is, obviously, a non-hamiltonian F -cycle of G . That is a contradiction and proves $S \subseteq F$ and, consequently, $s(G) \leq h(G)$.

Let $F \subseteq V$ be a smallest H -force set of G . If $F \subseteq T$ then $h(G) \leq t(G)$ trivially holds. Otherwise, there exists an $x \in F \setminus T$. By Lemma 3 there is a $y \in T$ such that $(F \setminus \{x\}) \cup \{y\}$ is an H -force set of G , too. The repeated use of the above exchange property finally yields a smallest H -force set $F' \subseteq T$ and proves the upper bound. ■

From the proof of Theorem 2, we have $S \subseteq F$ and we can choose F such that $F \subseteq T$.

Corollary 4. *Let $G \in \mathcal{H}$. Then,*

- (i) $s(G) = n$ if and only if $h(G) = n$.
- (ii) If $s(G) = n - 1$, then $h(G) = n - 1$.

Proof. Statement (i) is an immediate consequence of the lower bound in Theorem 2.

If $s(G) = n - 1$, then the lower bound of Theorem 2 implies $h(G) \geq n - 1$, and by (i) we have $h(G) \neq n$ which proves (ii). ■




Figure 1

The graph G of order 20 shown in Figure 1 is hamiltonian (the bold painted edges form a hamiltonian cycle) with $S = V \setminus \{x, y\}$ and with $V \setminus \{x\}$ as a smallest H -force set confirms that the converse of statement (ii) does not hold.

Theorem 2 has the following two consequences. A planar graph is called *outerplanar* if it can be embedded in the plane in such a way that every vertex is incident with the unbounded face.

Theorem 5. *Let $G \in \mathcal{H}$ be outerplanar. Then $h(G)$ corresponds to the number of vertices of degree 2 whose two neighbours are adjacent.*

Proof. Let $G \in \mathcal{H}$ be outerplanar and let $x \in V$. If $d_G(x) \geq 3$ then $x \notin T$ and also $x \notin S$. Assume otherwise $d_G(x) = 2$ and let $y, z \in V$ denote the neighbours of x . If $yz \notin E$ then $x \notin T$ and also $x \notin S$. If $yz \in E$ then $G - x$ is hamiltonian which yields $x \in S$ and, consequently, $x \in T$. Hence, $S = T$ and the statement can be deduced from Theorem 2. ■

In [4], the H -force number of an outerplanar hamiltonian graph G different from a cycle was proved to be equal to the number of leafs of the weak dual of G . The *weak dual* of an outerplanar graph G is a tree and is obtained from the dual of G by removing the vertex corresponding to the unbounded face.

Theorem 6. *For $G \in \mathcal{H}$, $h(G) = 2$ if and only if $t(G) = 2$.*

Proof. Suppose first $h(G) = 2$. Then by Lemma 3 there exists a smallest H -force set $F = \{x, y\}$ of G such that $F \subseteq T$. Assume that there exists a vertex

$v \in T \setminus F$ which means that $G - v$ is 2-connected. Then, $G - v$ and, consequently, G has two different x - y -paths with no common inner vertices. Hence, G has an F -cycle not passing v , a contradiction. That proves $F = T$ and $t(G) = 2$.

Suppose now $t(G) = 2$. Since G is not a cycle we have $h(G) \geq 2$. And, by Theorem 2 we have $h(G) \leq 2$ which completes the proof. ■

In [4], hamiltonian graphs with H -force number 2 have been characterized already by a condition on crossed chords of a hamiltonian cycle. In [4] they also noted that every hamiltonian graph with $h(G) = 2$ is planar.

Now, we give a decomposition formula with respect to the H -force number of a hamiltonian graph which is exactly 2-connected. To that end, let $G \in \mathcal{H}$ be a graph with vertices $u, v \in V$ such that $G - \{u, v\}$ is disconnected, i.e., $u, v \notin T$. Any given hamiltonian cycle C of G can be divided into two u - v -paths P_1 and P_2 which have no inner vertices in common. For $i = 1, 2$, let G_i denote the graph which results from $G[V(P_i)]$ (the subgraph of G induced by $V(P_i)$) by introducing an additional vertex w_i ($w_1 \neq w_2$) and edges uw_i, vw_i . Obviously, G_i is also a member of $\mathcal{H}$.

Theorem 7. *Let $G \in \mathcal{H}$ with $u, v \in V(G)$ such that $G - \{u, v\}$ is disconnected, and let G_1, G_2 be graphs derived from G as described above. Then,*

$$h(G) = h(G_1) + h(G_2) - 2.$$

Proof. On the one hand, from $u, v \notin T(G_i)$ and Lemma 3 it follows that G_i has a smallest H -force set $F_i \subseteq V(G_i)$ such that $u, v \notin F_i$. F_i contains w_i because $G_i - w_i$ is hamiltonian. Let $F := (F_1 \setminus \{w_1\}) \cup (F_2 \setminus \{w_2\})$ and let C_F denote an F -cycle of G . $F_i \setminus \{w_i\}$ is not empty for $i = 1, 2$ which implies that neither G_1 nor G_2 contains C_F as a cycle. Suppose that C_F is not a hamiltonian cycle of G . Then, without loss of generality, there exists a vertex $x \in V(G) \setminus V(G_2)$ which is not contained in F . Let $P_{F,1}$ denote the u - v -path of C_F which is completely contained in G_1 . Then, the cycle obtained by connecting $P_{F,1}$ with the u - v -path (u, w_1, v) is an F_1 -cycle of G_1 which is not hamiltonian, a contradiction. Consequently, F is an H -force set of G and

$$\begin{aligned} h(G) &\leq |F| = |F_1 \setminus \{w_1\}| + |F_2 \setminus \{w_2\}| = (|F_1| - 1) + (|F_2| - 1) \\ &= h(G_1) + h(G_2) - 2. \end{aligned}$$

On the other hand, Lemma 3 implies that G has an H -force set $F \subseteq V(G)$ where $|F| = h(G)$ and $u, v \notin F$. Clearly, $F_i := (F \cap V(G_i)) \cup \{w_i\}$ is a subset of $V(G_i)$. If C_i denotes an F_i -cycle of G_i , then C_i contains w_i and also the vertices u and v . Hence, $C_i - w_i$ is a u - v -path of G_i and also of G . By connecting the u - v -paths $C_1 - w_1$ and $C_2 - w_2$ we obtain an F -cycle $\tilde{C}$ in G . If C_i for $i = 1$ or 2 would not be hamiltonian in G_i , then $\tilde{C}$ could not be hamiltonian in G .

This contradicts the fact that F is an H -force set of G and implies that F_i is an H -force set of G_i . Hence,

$$h(G) = |F| = (|F_1| - 1) + (|F_2| - 1) \geq (h(G_1) - 1) + (h(G_2) - 1) = h(G_1) + h(G_2) - 2$$

which proves the statement of Theorem 7 ■

If, for example, G_t denotes the hamiltonian graph which consists of a “chain” of $t \geq 1$ cube graphs (see Figure 2) then by induction and using Theorem 7 we obtain for the H -force-number $h(G_t) = 2t + 2$.




Figure 2

Next, we will give a characterization of hamiltonian graphs G such that $S(G)$ is an H -force set of G and, consequently, $h(G) = s(G)$. To this end, let us consider the concept of cycle extendable graphs (which was first investigated by Hendry in [5]) and weaken it in a suitable sense.

A cycle C of a graph G is called *extendable* if G contains a $V(C)$ -cycle C' which has exactly one vertex more than C . A graph G is called *cycle extendable* if G contains a cycle and if every non-hamiltonian cycle is extendable. Cycle extendable graphs are obviously hamiltonian ones.

In [5], Hendry raised the problem whether every hamiltonian chordal graph is cycle extendable or not. Jiang proved in [6] that every planar hamiltonian chordal graph is also cycle extendable. Moreover, a hamiltonian graph which is an interval graph or a split graph has been proved to be cycle extendable, see [1] and also [2].

Now, we call a non-hamiltonian cycle C of a graph G *weakly extendable* if G contains a $V(C)$ -cycle of length $n - 1$. And, a graph G is called *weakly cycle extendable* if G is hamiltonian and if every non-hamiltonian cycle is weakly extendable. Trivially, every cycle extendable graph is weakly cycle extendable. Every outerplanar graph which belongs to $\mathcal{H}$ is also weakly cycle extendable.

Theorem 8. *Let $G \in \mathcal{H}$. Then, the following conditions are equivalent.*

- (i) $S(G)$ is an H -force set, i.e., $h(G) = s(G)$.
- (ii) G is weakly cycle extendable.

Proof. Suppose that $S = S(G)$ is an H -force set and that G contains a cycle C which is not weakly extendable. Then, $G - x$ is not hamiltonian for each $x \in V(G) \setminus V(C)$ which implies $x \notin S$. Hence, C is an S -cycle which contradicts our claim that S is an H -force set. Thus, G is weakly cycle extendable.

Now, let G be weakly cycle extendable and suppose that S is not an H -force set. If S is empty then $G - x$ is not hamiltonian for each $x \in V(G)$. Since G is not a cycle, there exists a cycle C in G of length at most $n - 2$, and C is not weakly extendable, a contradiction. So, suppose that S is not empty and let C be a non-hamiltonian S -cycle of G . Then, C is weakly extendable, i.e., G has a $V(C)$ -cycle C' of length $n - 1$. Suppose C' does not contain a vertex $x \in V(G)$. Then $G - x$ is hamiltonian and, consequently, $x \in S$. That together with

$$x \in V(G) \setminus V(C') \subseteq V(G) \setminus V(C) \subseteq V(G) \setminus S$$

yields a contradiction which proves that S is an H -force set. ■

Hence, every weakly cycle extendable graph $G \in \mathcal{H}$ has a uniquely determined smallest H -force set. In Figure 3, a not weakly cycle extendable graph with a unique smallest H -force set (the two black vertices) is presented.




Figure 3

Theorem 9. Let $G \in \mathcal{H}$.

- (i) If $s(G) \geq n - 1$, then G is weakly cycle extendable.
- (ii) If $s(G) \leq 1$, then G is not weakly cycle extendable.

Proof. (i) If $s(G) = n$ then G is 1-hamiltonian which implies that every non-hamiltonian cycle of G is weakly extendable. If $s(G) = n - 1$ then every S -cycle is hamiltonian. For every other non-hamiltonian cycle C of G , there is an $x \in S$ which is not contained in C . Since $G - x$ is hamiltonian, C is a cycle of $G - x$ and, consequently, weakly extendable in G .

(ii) If $s(G) = 0$ then G has no cycle of length $n - 1$, i.e., every non-hamiltonian cycle is not weakly extendable. If $s(G) = 1$ then, obviously, G has at least five vertices. Let be $S = \{x\}$ and let C be a hamiltonian cycle of $G - x$. Moreover, let y and z be two neighbors of x . Then, C passes y and z and consists of two y - z -paths P_1 and P_2 with no common inner vertex. At least one of these paths has more than one inner vertex. Otherwise, because of $n \geq 5$, each of P_1 and

P_2 would have exactly one inner vertex which implies $s(G) > 1$, a contradiction. Suppose, now, that P_1 has at least two inner vertices. Then, $V(P_2) \cup \{x\}$ is the vertex set of a cycle C' of length at most $n - 2$. C' cannot be weakly extendable in G because otherwise there would exist a $V(C')$ -cycle of length $n - 1$ in G which is different from C . That contradicts the claim $S(G) = \{x\}$. ■

For every integer $n \geq 9$ and all k with $2 \leq k \leq n - 2$ we were able to construct a weakly cycle extendable graph of order n with H -force number k .

Now, let $\mathcal{F} = \mathcal{F}(G)$ for a given graph $G \in \mathcal{H}$ denote the family of all H -force sets of G . As is easily seen, $\bar{\mathcal{F}} = \{X \subseteq V \mid X \notin \mathcal{F}\}$ is an independence system on V which means that $\bar{\mathcal{F}}$ satisfies the following two properties.

(M1) $\emptyset \in \bar{\mathcal{F}}$.

(M2) $X \in \bar{\mathcal{F}}, Y \subseteq X$ implies $Y \in \bar{\mathcal{F}}$.

In general, the independence system $(V, \bar{\mathcal{F}})$ is not also a matroid which means that the property

(M3) If $X, Y \in \bar{\mathcal{F}}$ and $|X| = |Y| + 1$, then there exists an $x \in X \setminus Y$ such that $Y \cup \{x\} \in \bar{\mathcal{F}}$.

is not satisfied for every graph $G \in \mathcal{H}$ (see, also [8]). Consider the hamiltonian graph G with vertex set $V = \{1, 2, \dots, 7\}$ which consists of the cycle $(1, 2, \dots, 7)$ and the chords 14 and 36. For G we have $\{1, 2, 3, 4\} \in \bar{\mathcal{F}}$ and $\{1, 2, 3, 6, 7\} \in \bar{\mathcal{F}}$ but, property (M3) is not satisfied for these two sets.

Theorem 10. *If G is a weakly cycle extendable graph, then $(V, \bar{\mathcal{F}})$ is a matroid.*

Proof. Let $X, Y \in \bar{\mathcal{F}}$ be two sets where $|X| = |Y| + 1$. As G is weakly cycle extendable, G contains a Y -cycle C of length $n - 1$. Let $v \in V$ be the only vertex which does not belong to C . Hence, $X \setminus \{v\}$ is a subset of $V(C)$. If there is a vertex $x \in X \setminus \{v\}$ with $x \notin Y$, then we have $Y \cup \{x\} \in \bar{\mathcal{F}}$ and, consequently, $Y \setminus \{x\} \in \bar{\mathcal{F}}$. Otherwise, we have $Y = X \setminus \{v\}$. That yields $Y \cup \{v\} = X \in \bar{\mathcal{F}}$ and proves the property (M3). ■

The maximal independent sets of the matroid $(V, \bar{\mathcal{F}})$, which are the members of $\bar{\mathcal{F}}$ of maximal cardinality, are just the vertex sets of the cycles of length $n - 1$ of G .

If $\mathcal{C} = \mathcal{C}(G)$ denotes the set of all cycles in G which are not weakly extendable, then let $(\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m)$ denote a partition of $\mathcal{C}$, i.e., $\mathcal{C}$ is the union of $m \geq 1$ nonempty and disjoint subsets $\mathcal{C}_i$ of $\mathcal{C}(G)$. We call a partition $(\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m)$ *vertex-unsaturated* (for short, *unsaturated*) if $V(\mathcal{C}_i)$ where

$$V(\mathcal{C}_i) := \bigcup_{C \in \mathcal{C}_i} V(C)$$

is different from $V(G)$ for $i = 1, 2, \dots, m$. Now, let $p(G)$ denote the smallest integer m for which there exists an unsaturated partition $(\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m)$ of $\mathcal{C}(G)$.

Theorem 11. *Let $G \in \mathcal{H}$ be a graph that is not weakly cycle extendable. Then,*

$$h(G) = s(G) + p(G).$$

Proof. First, let $(\mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_m)$ be an unsaturated partition of $\mathcal{C}(G)$ such that $m = p(G)$. For $i = 1, 2, \dots, m$ let $v_i \in V(G) \setminus V(\mathcal{C}_i)$ be any fixed vertex. We prove that $X := S(G) \cup \{v_1, \dots, v_m\}$ is an H -force set which implies $h(G) \leq s(G) + p(G)$. For this purpose, let C be any non-hamiltonian cycle of G .

If there exists a $V(C)$ -cycle C' of length $n - 1$ in G , then $S(G)$ contains a vertex v such that $\{v\} = V(G) \setminus V(C')$. Hence, $v \notin V(C)$ and, consequently, $X \not\subseteq V(C)$. If there is no $V(C)$ -cycle of length $n - 1$ in G , then G contains a $V(C)$ -cycle $C'' \in \mathcal{C}(G)$. In this case there exists a partition set $\mathcal{C}_i$, $1 \leq i \leq m$, such that $C'' \in \mathcal{C}_i$. Then

$$v_i \in V(G) \setminus V(\mathcal{C}_i) \subseteq V(G) \setminus V(C'') \subseteq V(G) \setminus V(C)$$

implies $X \not\subseteq V(C)$. Thus, every X -cycle is hamiltonian and X is an H -force set.

Assume now that there exists an H -force set X of G with less than $s(G) + p(G)$ vertices. Since, by Theorem 8, $S(G)$ is not an H -force set, there exists a nonempty subset $Y \subseteq V(G) \setminus S(G)$ such that $X = S(G) \cup Y$. Because of the assumption we have $|Y| < p(G)$. Note that every cycle $C \in \mathcal{C}(G)$ is an $S(G)$ -cycle because otherwise there would exist an $x \in S(G) \setminus V(C)$ such that $V(G) \setminus \{x\}$ is the vertex set of a cycle C' of length $n - 1$ in G with $V(C) \subseteq V(C')$, a contradiction with respect to $C \in \mathcal{C}(G)$. Since, moreover, every X -cycle is hamiltonian, we have that for every $C \in \mathcal{C}(G)$ there exists a vertex $y \in Y$ such that $y \notin V(C)$.

For every $y \in Y$, let us define $\mathcal{D}_y = \{C \in \mathcal{C}(G) \mid y \notin V(C)\}$. Then, we have

$$\mathcal{C}(G) = \bigcup_{y \in Y} \mathcal{D}_y$$

and, because of $\mathcal{C}(G) \neq \emptyset$, there exists a vertex $y_1 \in Y$ such that $\mathcal{D}_{y_1} \neq \emptyset$. Now, we are able to construct an unsaturated partition of $\mathcal{C}(G)$. To this end, let $\mathcal{C}_1 := \mathcal{D}_{y_1}$ and $Y_1 := Y \setminus \{y_1\}$. We may assume that the partition sets $\mathcal{C}_1, \dots, \mathcal{C}_k$ with $k \geq 1$ are already constructed. If Y_k contains a vertex y_{k+1} such that the set

$$\mathcal{D}_{y_{k+1}} \setminus \bigcup_{i=1}^k \mathcal{C}_i$$

is not empty, then let

$$\mathcal{C}_{k+1} := \mathcal{D}_{y_{k+1}} \setminus \bigcup_{i=1}^k \mathcal{C}_i.$$

This procedure terminates after at most $|Y| - 1$ steps and yields an unsaturated partition $(\mathcal{C}_1, \dots, \mathcal{C}_m)$ with $m < p(G)$ which contradicts the definition of $p(G)$. ■

As an immediate consequence of Theorem 11 we have

Corollary 12. *Let $G \in \mathcal{H}$ be a not weakly cycle extendable graph. Then, the following conditions are equivalent.*

- (1) $h(G) = s(G) + 1$,
- (2) $(\mathcal{C}(G))$ is unsaturated.

REFERENCES

- [1] A. Abueida and R. Sritharan, *Cycle extendability and Hamiltonian cycles in chordal graph classes*, SIAM J. Discrete Math. **20** (2006) 669–681.
doi:10.1137/S0895480104441267
- [2] G. Chen, R.J. Faudree, R.J. Gould and M.S. Jacobson, *Cycle extendability of Hamiltonian interval graphs*, SIAM J. Discrete Math. **20** (2006) 682–689.
doi:10.1137/S0895480104441450
- [3] R. Diestel, *Graph Theory* (Springer, Graduate Texts in Mathematics **173**, 2005).
- [4] I. Fabrici, E. Hexel and S. Jendrol', *On vertices enforcing a Hamiltonian cycle*, Discuss. Math. Graph Theory **33** (2013) 71–89.
doi:10.7151/dmgt.1653
- [5] G.R.T. Hendry, *Extending cycles in graphs*, Discrete Math. **85** (1990) 59–72.
doi:10.1016/0012-365X(90)90163-C
- [6] T. Jiang, *Planar Hamiltonian chordal graphs are cycle extendable*, Discrete Math. **257** (2002) 441–444.
doi:10.1016/S0012-365X(02)00505-8
- [7] R. Li, X. Zhang, S. Li, Q. Guo and Y. Guo, *The H -force set of a hypertournament*, Discrete Appl. Math. **169** (2014) 168–175.
doi:10.1016/j.dam.2013.12.020
- [8] J.G. Oxley, *Matroid Theory* (Oxford University Press, Oxford, 1992).
- [9] M. Timková, *Enforced Hamiltonian cycles in generalized dodecahedra*, Electron. J. Graph Theory Appl. **1** (2013) 77–88.
doi:10.5614/ejgta.2013.1.2.1
- [10] X. Zhang, R. Li and S. Li, *H -force sets of locally semicomplete digraphs*, Discrete Appl. Math. **160** (2012) 2491–2496.
doi:10.1016/j.dam.2012.06.014

Received 27 July 2015
 Revised 23 February 2016
 Accepted 23 February 2016