Discussiones Mathematicae Graph Theory 37 (2017) 649–663 doi:10.7151/dmgt.1949

# FORBIDDEN PAIRS AND (k, m)-PANCYCLICITY

CHARLES BRIAN CRANE

Department of Mathematics Marygrove College 8425 W McNichols Rd, Detroit, MI 48221, USA

e-mail: ccrane@marygrove.edu

#### Abstract

A graph G on n vertices is said to be (k, m)-pancyclic if every set of k vertices in G is contained in a cycle of length r for each  $r \in \{m, m+1, \ldots, n\}$ . This property, which generalizes the notion of a vertex pancyclic graph, was defined by Faudree, Gould, Jacobson, and Lesniak in 2004. The notion of (k, m)-pancyclicity provides one way to measure the prevalence of cycles in a graph. We consider pairs of subgraphs that, when forbidden, guarantee hamiltonicity for 2-connected graphs on  $n \ge 10$  vertices. There are exactly ten such pairs. For each integer  $k \ge 1$  and each of eight such subgraph pairs  $\{R, S\}$ , we determine the smallest value m such that any 2-connected  $\{R, S\}$ -free graph on  $n \ge 10$  vertices is guaranteed to be (k, m)-pancyclic. Examples are provided that show the given values are best possible. Each such example we provide represents an infinite family of graphs.

Keywords: hamiltonian, pancyclic, forbidden subgraph, cycle, claw-free. 2010 Mathematics Subject Classification: 05C38, 05C45.

### 1. INTRODUCTION

Let G = (V, E) denote a simple graph of order  $n \ge 3$ . We say G is *pancyclic* if G contains a cycle of each possible length, from 3 up to n. The notion of vertex pancyclicity was defined by Bondy in [2]. The graph G is *vertex pancyclic* if every vertex of G is contained in a cycle of each possible length. We consider the property (k, m)-pancyclicity, defined in 2004 by Faudree *et al.* [8], which is a generalization of vertex pancyclicity.

**Definition** (Faudree, Gould, Jacobson and Lesniak [8]). Given integers k and m with  $0 \le k \le m \le n$ , a graph G of order n is said to be (k, m)-pancyclic if for any k-set  $S \subseteq V$  and any integer r with  $m \le r \le n$ , there exists a cycle of length r in G that contains S.

Whenever m > n or k > n, we define (k, m)-pancyclicity to be the same as hamiltonicity. Note that (k, n)-pancyclicity is hamiltonicity, (0, 3)-pancyclicity represents pancyclicity, and (1, 3)-pancyclicity is vertex pancyclicity. Note also that whenever a graph is (k, m)-pancyclic for some  $k \ge 1$ , then it must also be (k - 1, m)-pancyclic and (k, m + 1)-pancyclic.

Relationships between hamiltonian-type properties and bounds on the quantity  $\sigma_2(G) = \min\{d(x) + d(y) : xy \notin E(G)\}$  have been studied extensively. Ore proved in [9] that if  $\sigma_2(G) \ge n$ , then G is hamiltonian. In 1971, Bondy showed in [3] that the condition  $\sigma_2(G) \ge n+1$  guarantees G is pancyclic. Then in 2004, Faudree *et al.* showed that this bound ensures much more than pancyclicity. Their result uses the notion of (k, m)-pancyclicity, and provides insight into the prevalence of cycles in such a graph.

**Theorem A** (Faudree, Gould, Jacobson and Lesniak [8]). Let G be a graph of order  $n \ge 3$ . If  $\sigma_2(G) \ge n+1$ , then G is (k, 2k)-pancyclic for each integer  $k \ge 2$ .

Another technique that has been employed to ensure hamiltonian-type properties is the forbidding of a subgraph or subgraphs. Given a graph H, we say Gis H-free if G does not contain H as an induced subgraph. In this context, H is called a *forbidden subgraph*. If  $\mathcal{F}$  is a family of graphs, we say G is  $\mathcal{F}$ -free if G is F-free for each  $F \in \mathcal{F}$ .

In 2015, it was shown in [5] that if only claw-free graphs are considered, we may lower the  $\sigma_2(G)$  bound to n in Theorem A and simultaneously guarantee (k, k+3)-pancyclicity as opposed to (k, 2k)-pancyclicity.

**Theorem B** [5]. Let G be a claw-free graph of order  $n \ge 3$ . If  $\sigma_2(G) \ge n$ , then G is (k, k+3)-pancyclic for each integer  $k \ge 1$ .

### 1.1. Pairs of forbidden subgraphs

A number of hamiltonian-type results have been obtained involving forbidden families of subgraphs. The claw is often a member of these forbidden families, as well as the graphs  $Z_1, Z_2, Z_3, B, N$ , and W, which are pictured in Figure 1.

A characterization of all pairs of subgraphs that, when forbidden, imply hamiltonicity in 2-connected graphs of order  $n \ge 10$  was achieved in [7] by Faudree and Gould. Their result extended an earlier characterization by Bedrossian [1], which used graphs of small order to eliminate the pair  $\{K_{1,3}, Z_3\}$ .

**Theorem C** (Faudree and Gould [7]). Let R and S be connected graphs  $(R, S \neq P_3)$  and let G be a 2-connected graph of order  $n \geq 10$ . Then G is  $\{R, S\}$ -free implies G is hamiltonian if, and only if, without loss of generality  $R = K_{1,3}$  and S is one of the graphs  $C_3, P_4, P_5, P_6, Z_1, Z_2, Z_3, B, N$ , or W.

They went on to characterize the forbidden pairs that guarantee pancyclicity in 2-connected graphs of order  $n \ge 10$ .

650



Figure 1. Some graphs that are commonly forbidden.

**Theorem D** (Faudree and Gould [7]). Let R and S be connected graphs  $(R, S \neq P_3)$  and let G  $(G \neq C_n)$  be a 2-connected graph of order  $n \geq 10$ . Then G is  $\{R, S\}$ -free implies G is pancyclic if, and only if, without loss of generality  $R = K_{1,3}$  and S is one of the graphs  $P_4, P_5, P_6, Z_1$ , or  $Z_2$ .

In this paper, we investigate pairs of forbidden subgraphs that guarantee (k, m)-pancyclicity for integers  $k \leq m$  in 2-connected graphs. Since a (k, m)-pancyclic graph must be hamiltonian, we need only consider the forbidden pairs from Theorem C. For each integer  $k \geq 1$  and each pair  $\{K_{1,3}, S\}$  where  $S \in \{C_3, P_4, Z_1, Z_2, Z_3, B, N, W\}$ , we determine the smallest integer m such that any 2-connected  $\{K_{1,3}, S\}$ -free graph is guaranteed to be (k, m)-pancyclic. A summary of results is provided at the end of the paper (see Theorem 7). We also give examples that show the provided values are best possible. Each such example represents an infinite family of graphs.

## 1.2. Notation

For terms and notation not defined here, we refer the reader to [4]. For a vertex  $v \in V$ , we denote by d(v) the degree of v, and by N(v) the neighborhood of v. Given a subgraph H of G and a vertex  $v \in V(G)$ , we let  $N_H(v) = N(v) \cap V(H)$ , and  $d_H(v) = |N_H(v)|$ . For  $S \subseteq V$ , let  $N(S) = \{v \in V - S : vh \in E(G) \text{ for some } v \in V(G)\}$   $h \in S$ . The independence number of G will be denoted by  $\alpha(G)$ . For an integer  $i \geq 1$  we let  $P_i$  denote a path on i vertices, and for an integer  $j \geq 3$  we let  $C_j$  denote a j-cycle. Given distinct vertices  $u, v \in V(G)$ , a (u, v)-path is any path from u to v in G. Given a vertex u and a subgraph H such that  $u \notin V(H)$ , a (u, H)-path is any path in G from u to a vertex  $v \in V(H)$ .

Given a path P, we denote by (P) the set of all internal vertices of P, that is V(P) minus the end vertices of P. Given a cycle C and a vertex  $v \in V(C)$ , we impose an orientation on C and let  $v^-(v^+)$  denote the vertex that appears directly before (after) v on C. We let xCy denote the path from x to y along C in the direction of the imposed orientation, while  $xC^-y$  will denote the path from x to y in the opposite direction along C. For a set of vertices  $S \subseteq V$ , we let  $\langle S \rangle_G$  denote the subgraph induced by S in G. If the context is clear, we denote this induced subgraph by  $\langle S \rangle$ .

# 2. Forbidden Pairs That Guarantee Cycle Extendability

A non-hamiltonian cycle C is *extendable* if there exists a cycle C' in G such that  $V(C') = V(C) \cup \{v\}$  for some  $v \in V(G) - V(C)$ . We say G is *cycle extendable* if every non-hamiltonian cycle in G is extendable. We will first examine the pairs of forbidden subgraphs that guarantee cycle extendability in 2-connected graphs. These pairs were completely characterized in [7].

**Theorem E** (Faudree and Gould [7]). Let R and S be connected graphs  $(R, S \neq P_3)$  and let G be a 2-connected graph of order  $n \geq 10$ . Then G is  $\{R, S\}$ -free implies G is cycle extendable if, and only if, without loss of generality  $R = K_{1,3}$  and S is one of the graphs  $C_3, P_4, Z_1$ , or  $Z_2$ .

If  $S = C_3$  then we must have  $G = C_n$ , and so G is (k, n)-pancyclic for all k in this case. If  $S = Z_1$ , it is easy to show that either  $G = C_n$  or G is a complete graph minus at most a matching (see [7]). Thus we make the following observation.

**Theorem 1.** Let  $G \neq C_n$  be a 2-connected  $\{K_{1,3}, Z_1\}$ -free graph of order  $n \geq 5$ . Then each of the following hold:

- (i) G is (1,3)-pancyclic;
- (ii) G is (k, 4)-pancyclic for  $k \in \{2, 3\}$ ;
- (iii) G is (k, k)-pancyclic for each  $k \ge 4$ .

These results are best possible.

**Proof.** Since G is isomorphic to  $K_n$  minus at most a matching, it is easy to see that for all  $k \ge 4$ , each k-set lies on a k-cycle. Since G is cycle extendable, part

(iii) follows. Parts (i) and (ii) follow similarly. The graph  $H = K_n - uv$  shows that part (ii) is best possible, since any 2-set or 3-set of vertices containing  $\{u, v\}$  in H is not contained in a triangle.

# 2.1. The pair $\{K_{1,3}, P_4\}$

Concerning the pair  $\{K_{1,3}, P_4\}$ , we show the following.

**Theorem 2.** Let G be a 2-connected  $\{K_{1,3}, P_4\}$ -free graph with  $n \ge 10$  vertices. Then the following hold:

(i) G is (1, 4)-pancyclic;

(ii) G is (k, k+2)-pancyclic for each integer  $k \ge 2$ .

These results are best possible.

We will use two lemmas from [6] in the proof.

**Lemma 3** (Egawa, Fujisawa, Fujita and Ota [6]). Suppose G is a connected noncomplete  $P_4$ -free graph, and let S be a minimum cutset of G. Then for every two vertices u and v with  $u \in S$  and  $v \in V - S$ , we have  $uv \in E$ .

The following lemma follows immediately from Lemma 3.

**Lemma 4** (Egawa, Fujisawa, Fujita and Ota [6]). Let  $r \ge 2$ , and suppose G is a connected P<sub>4</sub>-free graph. Then G is  $K_{1,r}$ -free if and only if  $\alpha(G) \le r - 1$ .

We now prove Theorem 2.

**Proof of Theorem 2.** Let  $a \in V$ , and pick a shortest cycle C that contains a. Such a cycle must exist since G is hamiltonian. Now G is  $P_4$ -free, so the length of C must be 3 or 4. Since G is cycle extendable, we have shown that G is (1, 4)-pancyclic.

To prove part (ii) we first show that, given two vertices  $u, v \in V$ , there exists a 4-cycle in G that contains u and v. Since G is 2-connected, we may pick two internally vertex-disjoint (u, v)-paths P and Q so that (a) |(P)| is as small as possible, and (b) subject to condition (a), |(Q)| is as small as possible. Then  $|(P)| \leq |(Q)| \leq 2$  because G is P<sub>4</sub>-free. For the same reason, if |(Q)| = 2, then |(P)| = 0 and we have a 4-cycle containing u and v. Otherwise if  $|(P)| \leq |(Q)| \leq 1$ , then there exists a 3-cycle or 4-cycle that contains u and v, and as G is cycle extendable by Theorem E, there must be a 4-cycle containing u and v.

Now let  $S \subset V$  and  $|S| = k \geq 2$ . We have shown there exists a 4-cycle that contains at least two vertices from S. We wish to obtain a cycle of length  $t \geq 4$  that contains exactly two vertices which are not in S. Now G is cycle extendable, so if the 4-cycle contains fewer than two vertices that are not in S,

we may perform cycle extensions until we obtain a cycle  $C = v_1 v_2 \cdots v_t v_1$  with  $4 \le t \le k+2$  that contains exactly two vertices which are not in S.

If t = k + 2 we are done, since G is cycle extendable. So assume t < k + 2, and let  $w \in S - V(C)$ . We show we may obtain a cycle  $\hat{C}$  of length t + 1 that contains w and  $S \cap V(C)$ . Note that such a cycle can contain at most two vertices which are not in S. Note also that, since G is  $K_{1,3}$ -free, we have  $\alpha(G) \leq 2$  by Lemma 4.

First, suppose  $d_C(w) \geq 2$ . In the following, indices that are not in the set  $\{1, 2, \ldots, t\}$  are taken modulo t. Let  $v_i, v_j \in N_C(w)$  be such that i < j and  $N(w) \cap \{v_{i+1}, v_{i+2}, \ldots, v_{j-1}\} = \emptyset$ . Now  $wv_{j+1} \notin E$ , or else we have the desired cycle  $\hat{C}$ . Hence  $v_{i+1}v_{j+1} \in E$ , or else  $\{w, v_{i+1}, v_{j+1}\}$  is an independent set, contradicting the fact that  $\alpha(G) \leq 2$ . But now  $v_i C^- v_{j+1} v_{i+1} C v_j w v_i$  is the desired cycle  $\hat{C}$ . Thus we assume  $d_C(w) \leq 1$ .

Suppose  $d_C(w) = 1$ . Without loss of generality, assume  $wv_1 \in E$ . Now  $wv_j \notin E$  for all  $2 \leq j \leq t$ , so  $\{v_2, v_3, \ldots, v_t\}$  induces a clique since  $\alpha(G) \leq 2$ . Since G is 2-connected, by Lemma 3 there exists a vertex  $z \neq v_1$  such that  $zw \in E$  and  $zv_j \in E$  for all  $j \in \{2, 3, \ldots, t\}$ . There also exists a vertex  $v_m \in V(C) - \{v_1\}$  that is not in S. If  $m \neq t$ , then  $v_1wzv_{m+1}Cv_tv_{m-1}C^-v_1$  is the desired cycle  $\hat{C}$ . If m = t, then  $v_1wzv_{t-1}C^-v_1$  is the cycle  $\hat{C}$ .

Lastly, suppose  $d_C(w) = 0$ . Since  $\alpha(G) \leq 2$ , V(C) induces a clique. As G is 2-connected, by Lemma 3 there exist distinct vertices  $z_1$  and  $z_2$  such that  $wz_1, wz_2 \in E$  and  $z_1x, z_2x \in E$  for all  $x \in V(C)$ . Let  $V(C) - S = \{a, b\}$ . As  $\langle V(C) \rangle$  is a clique, we may modify the indices of C if necessary so that  $a = v_1$  and  $b = v_2$ . Now  $v_3Cv_tz_1wz_2v_3$  is the desired cycle  $\hat{C}$ .

We may repeat this argument as necessary to obtain a cycle containing S whose length is k, k+1, or k+2. Since G is cycle extendable, this completes the proof of part (ii).

To see that part (ii) is best possible, let  $n \ge k+2$  and construct the graph  $H_1$  in Figure 2 by removing k-1 edges incident to some vertex v in a copy of  $K_n$ . Now  $H_1$  is 2-connected and  $\{K_{1,3}, P_4\}$ -free, but the k-set  $S = V(H_1) - N(v)$  is not contained in a cycle of length k + 1.

To see that part (i) is best possible, construct a graph G as follows. Let  $H = K_{n-3}$ , and add vertices x, y, and a such that  $N_G(x) = N_G(y) = V(H) \cup \{a\}$  and  $N_G(a) = \{x, y\}$ . Then G is 2-connected and  $\{K_{1,3}, P_4\}$ -free, but the vertex a is not contained in a triangle.

### 2.2. The pair $\{K_{1,3}, Z_2\}$

We now consider  $\{K_{1,3}, Z_2\}$ , the only remaining pair from Theorem E that guarantees cycle extendability in 2-connected graphs. We prove the following.



Figure 2. The k-set  $V(H_1) - N(v)$  is not contained in a (k+1)-cycle.

**Theorem 5.** Let  $G \neq C_n$  be a 2-connected  $\{K_{1,3}, Z_2\}$ -free graph with  $n \geq 10$  vertices. Then the following hold:

- (i) G is (1, 4)-pancyclic;
- (ii) G is (k, 3k)-pancyclic for each integer  $k \ge 2$ .

These results are best possible.

**Proof.** We first prove part (i). Let  $w \in V$ . If w is contained in a triangle, then w is contained in a 4-cycle since G is cycle extendable. Thus we assume that w is not contained in a triangle. Since G is 2-connected and  $G \neq C_n$ , there exists a vertex of degree at least 3. This implies G contains a triangle, since G is claw-free.

As G is 2-connected, we let s be the smallest integer such that there exists a triangle H and a pair of (w, H)-paths (P, Q) which are vertex-disjoint (except for w) with P having length s. Now pick a triangle T so that there exists a pair of (w, T)-paths  $(P_T, Q_T)$  that are vertex-disjoint (except for w),  $P_T$  has length s, and  $Q_T$  has minimum possible length.

Let T = axya, and without loss of generality let  $P_T = xx_1x_2\cdots x_s$  and  $Q_T = yy_1y_2\cdots y_t$ , where  $x_s = w = y_t$ . Note that  $P_T$  and  $Q_T$  are each induced paths and  $s \leq t$ .

Suppose  $s \ge 2$ . Now  $yx_1 \notin E$ , for otherwise w is connected to the triangle  $xyx_1x$  by  $Q_T$  and a path  $x_1x_2\cdots x_s$  that is shorter than  $P_T$ , violating the minimality of s. Similarly,  $ax_1 \notin E$ . Also  $ax_2 \notin E$  by the minimality of s. Now we must have  $yx_2 \in E$  or else  $\{y, a, x, x_1, x_2\}$  induces a  $Z_2$ . But then  $x_2 \neq w$  by the minimality of  $t \ge 2$ . Therefore  $s \ge 3$ , and avoiding a claw centered at  $x_2$ , we must have  $yx_3 \in E$ . This is a contradiction, since the triangle  $yx_2x_3y$  now violates the minimality of s.

Hence we must have s = 1. We may assume  $t \ge 3$ , for otherwise w is contained in a  $C_3$  or  $C_4$ . First note that  $xy_1 \notin E$ , or else the minimality of t is violated via the triangle  $xyy_1x$ . Also  $ay_i \notin E$  for all  $2 \le i \le t$ , by the minimality of t. Furthermore  $ay_1 \notin E$ , for otherwise  $\{a, y, y_1, y_2, y_3\}$  induces a  $Z_2$ . But now  $xy_2 \in E$  since otherwise  $\{x, a, y, y_1, y_2\}$  induces a  $Z_2$ . This implies t = 3, for otherwise  $\{x, w, y_2, y\}$  induces a claw centered at x. We now have a contradiction, since w is contained in a triangle.

Therefore w must be contained in a 4-cycle. Since G is cycle extendable, we have shown that G is (1, 4)-pancyclic. To see that part (i) is best possible, construct a graph G as follows. Remove an edge uv from a copy of  $K_{n-1}$ , and add a new vertex w such that  $N_G(w) = \{u, v\}$ . Now G fulfills the assumptions of Theorem 5, but G is not (1, 3)-pancyclic since w is not contained in a triangle.

Our proof of part (ii) will be by induction on k. We begin with a claim that will serve as the base case.

#### **Claim.** Any set of two vertices is contained in a 6-cycle in G.

**Proof.** Let  $w, z \in V$ . We consider two cases.

Case 1. Suppose z is contained in a triangle. If there is a triangle in G that contains both w and z, then the claim holds as G is cycle extendable. Thus suppose there is no triangle in G that contains both w and z.

As G is 2-connected, we let s be the smallest integer such that there exists a triangle H containing z and a pair of (w, H)-paths (P, Q) which are vertexdisjoint (except for w) with P having length s. Now pick a triangle T containing z so that there exists a pair of (w, T)-paths  $(P_T, Q_T)$  that are vertex-disjoint (except for w),  $P_T$  has length s, and  $Q_T$  has minimum possible length.

We may assume

(1) 
$$|(P_T)| + |(Q_T)| \ge 3$$
,

since otherwise we have the desired 6-cycle (using the fact that G is cycle extendable). Let T = axya, and let  $P_T = xx_1x_2\cdots x_s$  and  $Q_T = yy_1y_2\cdots y_t$ , where  $x_s = w = y_t$ . Note that  $P_T$  and  $Q_T$  are each induced paths and  $s \leq t$ .

For each *i* with  $2 \le i \le s$ , we have  $ax_i \notin E$ , for otherwise the minimality of *s* is violated. Similarly  $ay_i \notin E$  for all *i* with  $2 \le i \le t$ , by the minimality of *t*. By inequality (1), we know  $t \ge 3$ . Now  $ay_1 \notin E$  or else  $\{a, y, y_1, y_2, y_3\}$  induces a  $Z_2$ .

Suppose  $yx_i \in E$  for some  $i \in \{2, 3, ..., s\}$ . Note that i < s since  $yw \notin E$ . Then  $yx_{i+1} \in E$ , or else  $\{a, x, y, x_i, x_{i+1}\}$  induces a  $Z_2$ . But by induction, this implies that  $yx_{i+2}, yx_{i+3}, ..., yx_s \in E$ , which is a contradiction since  $yx_s \notin E$ . Therefore  $yx_i \notin E$  for each  $i \in \{2, 3, ..., s\}$ . We now consider four subcases.

Case 1.1. Let  $|(P_T)| = 0$  and  $z \in \{x, y\}$ . By our choice of T and the minimality of t, we have  $xy_1 \notin E$ . Inequality (1) implies  $t \ge 4$ . Now  $xy_2 \notin E$  or else  $\{x, y, y_2, w\}$  induces a claw centered at x. But then  $\{a, x, y, y_1, y_2\}$  induces a  $Z_2$ .

656

Case 1.2. Suppose  $|(P_T)| = 0$  and  $z \notin \{x, y\}$ . We have z = a, and again  $t \ge 4$ . Now  $aw \notin E$  or else z and w are contained in the triangle axwa. Note that  $xy_{t-1} \in E$ , for otherwise  $\{a, y, x, w, y_{t-1}\}$  induces a  $Z_2$ . Also  $xy_{t-2} \notin E$  since  $\{x, w, y_{t-2}, y\}$  cannot induce a claw. Avoiding a  $Z_2$  induced by the set  $\{x, w, y_{t-1}, y_{t-2}, y_{t-3}\}$ , we must have  $xy_{t-3} \in E$ . But now  $\{x, a, y_{t-3}, w\}$  induces a claw centered at x.

Case 1.3. Let  $|(P_T)| \ge 1$  and  $z \in \{x, y\}$ . First suppose z = x. Note that  $ax_1 \notin E$ , or else the minimality of s is violated by the triangle  $axx_1a$ . Similarly  $yx_1 \notin E$ . This is a contradiction, since  $\{y, a, x, x_1, x_2\}$  now induces a  $Z_2$ .

Therefore we must have z = y. Now  $yx_1 \notin E$ , or else the minimality of s is violated. Thus  $ax_1 \in E$  since otherwise  $\{y, a, x, x_1, x_2\}$  induces a  $Z_2$ . Then  $w = x_2$ , since  $\{a, x, x_1, x_2, x_3\}$  cannot induce a  $Z_2$ . Now  $xy_{t-1} \notin E$ , or else the paths  $P = xy_{t-1}w$  and  $Q = ax_1w$  violate the minimality of  $t \geq 3$ . Then  $x_1y_{t-1} \in E$  or else  $\{a, x, x_1, w, y_{t-1}\}$  induces a  $Z_2$ . But now  $\{y_{t-1}, w, x_1, x, y\}$  induces a  $Z_2$ .

Case 1.4. Suppose  $|(P_T)| \ge 1$  and  $z \notin \{x, y\}$ . We have z = a. Now  $ax_1 \notin E$ , since otherwise the minimality of s is violated via the triangle  $axx_1a$ . Hence  $yx_1 \in E$ , or else  $\{y, a, x, x_1, x_2\}$  induces a  $Z_2$ . This implies  $w = x_2$ , since otherwise  $\{y, x, x_1, x_2, x_3\}$  induces a  $Z_2$ . Now  $xy_{t-1} \notin E$ , or else the paths  $P = xy_{t-1}w$  and  $Q = yx_1w$  violate the minimality of  $t \ge 3$ . Thus  $x_1y_{t-1} \in E$ , since  $\{y, x, x_1, x_2, x_3\}$  cannot induce a  $Z_2$ . This is a contradiction, since now  $\{y_{t-1}, w, x_1, x_2, a_3\}$  induces a  $Z_2$ . This is a contradiction of  $y_{t-1}$ .

Case 2. Suppose neither z nor w is contained in a triangle. By part (i), we know that z is contained in a 4-cycle. Also, since G is 2-connected and claw-free, we must have d(w) = d(z) = 2. Now let s be the smallest integer such that there exists a 4-cycle H containing z and a pair of (w, H)-paths (P, Q) which are vertex-disjoint (except for w) with P having length s. Now pick a 4-cycle C containing z so that there exists a pair of (w, C)-paths  $(P_C, Q_C)$  that are vertex-disjoint (except for w),  $P_C$  has length s, and  $Q_C$  has minimum possible length.

We may assume

(2) 
$$|(P_C)| + |(Q_C)| \ge 2$$

since otherwise we clearly have the desired 6-cycle. Since z is not contained in a triangle, C must be an induced 4-cycle. Let  $P_C = xx_1x_2\cdots x_s$  and  $Q_C = yy_1y_2\cdots y_t$ , where  $x_s = w = y_t$ . Note that  $P_C$  and  $Q_C$  are induced paths. By inequality (2), we have  $t \ge 2$ . Since d(z) = 2, we know  $z \notin \{x, y\}$ . Let  $V(C) - \{x, y, z\} = \{a\}$ . For each *i* with  $2 \le i \le s$ , we have  $ax_i \notin E$ , for otherwise the minimality of *s* is violated. Similarly  $ay_i \notin E$  for all  $2 \le i \le t$ , by the minimality of *t*. We now consider three subcases corresponding to the possible positions of *x* and *y* relative to *z* on *C*. Case 2.1. Suppose C = zaxyz. Now  $xy_1 \in E$  or else  $\{y, z, x, y_1\}$  induces a claw. Note that  $xy_i \notin E$  for all  $2 \leq i \leq t$ , since otherwise  $\{x, a, y, y_i\}$  induces a claw. But now  $w = y_2$ , for otherwise  $\{x, y, y_1, y_2, y_3\}$  induces a  $Z_2$ . By inequality (2) and the minimality of s, we must have  $w = x_2$ . Since w is not contained in a triangle, we know  $x_1y_1 \notin E$ . Also  $ay_1 \notin E$ , or  $\{y_1, y, a, w\}$  induces a claw centered at  $y_1$ . Then  $ax_1 \in E$ , for otherwise  $\{x, a, y_1, x_1\}$  induces a claw centered at x. But now  $ax_1w_1y_2a$  is the desired 6-cycle.

Case 2.2. Suppose C = zayxz. We have  $yx_1 \in E$ , for otherwise  $\{x, z, y, x_1\}$ induces a claw. Since  $yw \notin E$ , we know  $s \ge 2$ . If  $yx_i \in E$  for some  $i \ge 2$ , then  $\{y, a, x, x_i\}$  induces a claw. Hence  $yx_i \notin E$  for all  $2 \le i \le s$ . But now  $w = x_2$ , for otherwise  $\{y, x, x_1, x_2, x_3\}$  induces a  $Z_2$ .

Since w is not contained in a triangle, we have  $x_1y_{t-1} \notin E$ . Also  $xy_{t-1} \notin E$ , or  $\{x, z, x_1, y_{t-1}\}$  induces a claw centered at x. But then we must have  $w = y_2$ , for otherwise  $\{y, x, x_1, w, y_{t-1}\}$  induces a  $Z_2$ . Then  $ay_1 \in E$ , since otherwise  $\{y, a, x, y_1\}$  induces a claw centered at y. But now  $ay_1wx_1xz_1$  is the desired 6-cycle.

Case 2.3. Suppose C = zxayz. We know  $s \ge 2$  since  $\{x, z, a, w\}$  cannot induce a claw. Now  $ax_1 \in E$ , or else  $\{x, z, a, x_1\}$  induces a claw. But then  $w = x_2$ , for otherwise  $\{a, x, x_1, x_2, x_3\}$  induces a  $Z_2$ . This implies  $t \ge 3$ , or else  $yy_1wx_1xzy$  is the desired cycle of length six.

Now  $x_1y_{t-1} \notin E$  since w is not contained in a triangle. Also  $xy_{t-1} \notin E$ , since otherwise the pair of paths  $Q = xy_{t-1}w$  and  $P = ax_1w$  violate the minimality of t. But now  $\{a, x, x_1, w, y_{t-1}\}$  induces a  $Z_2$ . This contradiction completes Case 2, and the proof of the claim.

We now proceed with the inductive step of the proof for part (ii) of Theorem 5. Given an integer  $k \ge 2$  with  $3k \le n$ , suppose that any set of k vertices is contained in a cycle of length 3k in G. Let  $S \subset V$  be a set of k + 1 vertices with  $3(k+1) \le n$ , and let C be a 3k-cycle that contains at least k vertices of S. We wish to prove there exists a (3k+3)-cycle containing S. Now if  $S \subset V(C)$ , then since G is cycle extendable, there exists a cycle of length 3k + 3 that contains S. Therefore we assume  $S - V(C) = \{w\}$ .

Let  $\mathcal{F}$  be the set of pairs of paths in G defined by:  $\{P, Q\} \in \mathcal{F}$  if and only if Pand Q are (w, C)-paths that are vertex-disjoint (except for w) and |(P)| + |(Q)|is minimal among all such pairs of (w, C)-paths. Since G is 2-connected,  $\mathcal{F}$  is nonempty. Now pick a pair of paths  $\{P, Q\} \in \mathcal{F}$  so that P and Q have endpoints that are as close as possible on C. Let x and y denote the endpoints of P and Q, respectively, on C. Let  $P = xx_1x_2\cdots x_s$  and  $Q = yy_1y_2\cdots y_t$ , where  $x_s = w = y_t$ . By their definition, P and Q are induced paths. We now consider two cases.

Case 1. Assume x and y occur consecutively on C. Let  $y = x^+$ . If  $|(P)| + |(Q)| \le 2$ , then since G is cycle extendable, we may clearly extend C to obtain a

658

cycle of length 3(k+1) that contains S. Therefore suppose  $|(P)| + |(Q)| \ge 3$ .

Without loss of generality we may assume that  $|(P)| \ge |(Q)|$ , and hence  $|(P)| \ge 2$ . For each *i* with  $2 \le i \le s$ , we have  $x^-x_i, y^+x_i \notin E$  by the minimality of |(P)| + |(Q)|. Now  $x^-x_1 \notin E$ , or else  $\{x^-, x, x_1, x_2, x_3\}$  induces a  $\mathbb{Z}_2$ .

Suppose  $|(P)| \geq 3$ . Further, suppose  $yx_2 \in E$ . Then  $xy^+ \in E$ , since otherwise  $\{y, x, y^+, x_2\}$  induces a claw. But now  $yx_3 \in E$ , or else  $\{x, y^+, y, x_2, x_3\}$  induces a  $Z_2$ . Also  $yx_4 \in E$ , for otherwise  $\{x, y^+, y, x_3, x_4\}$  induces a  $Z_2$ . This yields a contradiction since now  $\{y, x, x_2, x_4\}$  induces a claw centered at y. Thus  $yx_2 \notin E$ .

Now suppose  $yx_1 \in E$ . Then  $yx_3 \in E$  or else the set  $\{y, x, x_1, x_2, x_3\}$  induces a  $Z_2$ . But now  $xy^+ \in E$ , since otherwise  $\{y, x, y^+, x_3\}$  induces a claw centered at y. This yields a contradiction, since now  $\{x, y^+, y, x_3, x_2\}$  induces a  $Z_2$ . Hence  $yx_1 \notin E$ . Avoiding a claw centered at x, we then have  $yx^- \in E$ . But this is a contradiction, since  $\{x^-, y, x, x_1, x_2\}$  now induces a  $Z_2$ .

Therefore |(P)| = 2. Also  $|(Q)| \ge 1$  since  $|(P)| + |(Q)| \ge 3$ . Note that  $w = x_3$ . First suppose  $yx_1 \in E$ . Then  $yx_2 \in E$ , for otherwise  $\{y, x, x_1, x_2, w\}$  induces a  $Z_2$ . In order to avoid a claw induced by  $\{y, x, y^+, x_2\}$ , we must have  $xy^+ \in E$ . This is a contradiction, since now  $\{x, y^+, y, x_2, w\}$  induces a  $Z_2$ . Therefore  $yx_1 \notin E$ must hold. Then  $x^-y \in E$ , since otherwise  $\{x, x^-, y, x_1\}$  induces a claw. We must also have  $yx_2 \in E$ , for otherwise  $\{x^-, y, x, x_1, x_2\}$  induces a  $Z_2$ . But now  $\{x^-, x, y, x_2, w\}$  induces a  $Z_2$ . This contradiction completes Case 1.

Case 2. Assume x and y do not occur consecutively on C. Suppose the length of xCy is less than or equal to the length of yCx.

For each *i* with  $2 \leq i \leq s$ , we have  $x^+x_i, x^-x_i \notin E$  or else the minimality of |(P)| + |(Q)| is violated. We must have  $x^+x_1 \notin E$ , for otherwise the pair  $\{x^+x_1Pw, Q\} \in \mathcal{F}$  has endpoints that are closer together on *C* than *x* and *y*.

Suppose  $|(P)| \ge 2$ . Now  $x^-x_1 \notin E$  or else  $\{x, x^-, x_1, x_2, x_3\}$  induces a  $Z_2$ . Since G is claw-free, we must have  $x^-x^+ \in E$ . But now  $\{x^-, x^+, x, x_1, x_2\}$  induces a  $Z_2$ . So we must have  $|(P)| \le 1$ . By symmetry we also have  $|(Q)| \le 1$ .

Suppose |(P)| = 1. If  $x^-x_1 \notin E$ , then  $x^-x^+ \in E$  in order to avoid a claw, and then  $\{x^-, x^+, x, x_1, w\}$  induces a  $Z_2$ . Hence  $x^-x_1 \in E$ . If |(Q)| = 1, then by symmetry  $y^+y_1 \in E$ . Therefore in any case, it is clear we may extend C to obtain a cycle C' with vertex set  $V(C) \cup (P) \cup (Q)$  that satisfies  $|V(C') - V(C)| \leq 2$  and  $|N(w) \cap V(C')| \geq 2$ .

Relabel the vertices so that  $C' = u_1 u_2 \cdots u_m u_1$ , where  $u_1, u_i \in N(w)$  for some i > 1 with  $N(w) \cap \{u_2, u_3, \ldots, u_{i-1}\} = \emptyset$  and i as small as possible. If  $u_1$  and  $u_i$  occur consecutively on C', then  $u_1 w u_2 C' u_m u_1$  is a cycle of length at most 3(k+1) that contains S. Thus we assume that  $u_1$  and  $u_i$  do not occur consecutively on C'.

Since G is claw-free, we must have  $u_m u_2, u_{i-1}u_{i+1} \in E$ . If i = 3, then  $u_1 w u_3 C' u_m u_2 u_1$  is a cycle of length at most 3(k+1) that contains S. If i = 4,

then  $u_1wu_4u_3u_5C'u_mu_2u_1$  is a cycle of length at most 3(k+1) that contains S. Hence we assume  $i \ge 5$ .

Now  $u_i u_2 \notin E$ , since otherwise  $u_1 w u_i u_2 C' u_{i-1} u_{i+1} C' u_m u_1$  is the desired cycle. Also  $u_i u_m \notin E$  by symmetry. Therefore  $u_1 u_i \in E$ , since  $\{u_m, u_2, u_1, w, u_i\}$ cannot induce a  $Z_2$ . Note that  $u_i u_3 \notin E$ , or else  $u_1 w u_i u_3 C' u_{i-1} u_{i+1} C' u_m u_2 u_1$ is the desired cycle. But now  $u_1 u_3 \in E$ , or else  $\{u_i, w, u_1, u_2, u_3\}$  induces a  $Z_2$ . We may assume i > 5, for otherwise  $u_1 w u_5 u_4 u_6 C' u_m u_2 u_3 u_1$  is the desired cycle. Note that  $u_i u_4 \notin E$ , or else  $\{u_i, w, u_1, u_3, u_4\}$  induces a  $Z_2$ . An easy inductive argument now verifies that  $u_1 u_j \in E$  for all  $2 \leq j \leq i - 2$ . But then  $u_1 w u_i u_{i-1} u_{i+1} C' u_m u_2 C' u_{i-2} u_1$  is the desired cycle.

We have shown that there must exist a cycle of length at most 3(k+1) which contains S. Since G is cycle extendable, this completes Case 2.

We have thus shown by induction that every set S of  $k \ge 2$  vertices is contained in a 3k-cycle whenever  $3k \le n$ . Since G is cycle extendable, this completes the proof of part (ii).



Figure 3. The set  $\{v_1, v_2, \ldots, v_k\}$  is not contained in a (3k - 1)-cycle.

To see that part (ii) is best possible, construct the graph  $H_2$  in Figure 3 as follows. Given  $n \ge 3k$ , take a copy of  $K_{n-k}$  with vertex set  $\{u_1, u_2, \ldots, u_{n-k}\}$ . Let  $H_2$  be the graph obtained by adding k distinct vertices  $v_1, v_2, \ldots, v_k$  such that  $N(v_i) = \{u_{2i-1}, u_{2i}\}$  for all i with  $1 \le i \le k$ . Then  $H_2$  is 2-connected and  $\{K_{1,3}, Z_2\}$ -free, but the set  $S = \{v_1, v_2, \ldots, v_k\}$  is not contained in a cycle of length 3k - 1. Forbidden Pairs and (k, m)-Pancyclicity

# 3. Four Pairs for Which Hamiltonicity Is Best Possible

Given an integer  $k \ge 1$ , the forbidden pairs in the following theorem do not guarantee (k, m)-pancyclicity for any integer m < n under the given conditions.

**Theorem 6.** Let G be a 2-connected  $K_{1,3}$ -free graph on n vertices.

- (i) If  $n \ge 5$  and G is M-free for some  $M \in \{B, N, W\}$ , then G is (k, n)-pancyclic for all  $k \ge 1$ .
- (ii) If  $n \ge 10$  and G is Z<sub>3</sub>-free, then G is (k, n)-pancyclic for all  $k \ge 1$ .

These results are best possible.

**Proof.** For part (i), we construct a graph  $H_3$  (see Figure 4) with  $|V(H_3)| = n \ge k \ge 1$  as follows. Let  $r \ge 1$  be an integer. Pick integers  $l_1, l_2, \ldots, l_r \ge 1$  such that  $\sum_{i=1}^r l_i \le k$ , and let  $A_i = K_{l_i}$  for each i with  $1 \le i \le r$ . Add 2r distinct vertices  $x_1, y_1, x_2, y_2, \ldots, x_r, y_r$  such that for each  $i \in \{1, 2, \ldots, r\}$ , we have  $x_i w, y_i w \in E$  for all  $w \in V(A_i)$ . For each i with  $1 \le i \le r-1$ , add a path  $Q_i$  with  $|V(Q_i)| \ge 2$  such that  $y_i Q_i x_{i+1}$  is an induced path. Lastly, add a path  $Q_r$  with  $|V(Q_r)| \ge 2$  such that  $y_r Q_r x_1$  is an induced path. Now  $H_3$  is 2-connected and  $\{K_{1,3}, M\}$ -free for each  $M \in \{B, N, W\}$ , but any k-set  $S \subseteq V(H_3)$  with  $\bigcup_{i=1}^r V(A_i) \subseteq S$  is not contained in a cycle of length n-1.



Figure 4. The set  $\bigcup_{i=1}^{r} V(A_i)$  is not contained in an (n-1)-cycle.

Regarding part (ii), for some integer t with  $0 \le t \le k-1$ , construct the graph  $H_4$  in Figure 5 as follows. Let  $G_1 = K_{t+2r}$  for some integer r with  $4r \ge k-t$ , and let  $V(G_1) = \{u_1, u_2, \ldots, u_{t+2r}\}$ . Add a matching  $M = \{x_1y_2, x_3y_4, \ldots, x_{2r-1}y_{2r}\}$  such that  $x_iu_{t+i} \in E$  for all odd i with  $1 \le i \le 2r-1$ , and  $y_iu_{t+i} \in E$  for all even i with  $2 \le i \le 2r$ . Let  $S \subseteq V(H_4)$  be any k-set satisfying  $\{u_1, u_2, \ldots, u_{t+1}\} \subseteq S$ .

The graph  $H_4$  is 2-connected and  $\{K_{1,3}, Z_3\}$ -free, but there is no cycle of length n-1=t+4r-1 that contains S.



Figure 5. The set  $\{u_1, u_2, \ldots, u_{t+1}\}$  is not contained in an (n-1)-cycle.

The following theorem summarizes results from this paper.

**Theorem 7.** Let G be a 2-connected  $K_{1,3}$ -free graph of order  $n \ge 10$ , and let  $k \ge 2$ .

- (i) If  $G \neq C_n$  is  $Z_1$ -free, then G is (k, k)-pancyclic for all  $k \geq 4$ .
- (ii) If G is  $P_4$ -free, then G is (k, k+2)-pancyclic.
- (iii) If  $G \neq C_n$  is Z<sub>2</sub>-free, then G is (k, 3k)-pancyclic.
- (iv) If G is  $C_3$ -free, then G is (k, n)-pancyclic.
- (v) If G is  $Z_3$ -free, then G is (k, n)-pancyclic.
- (vi) If G is B-free, then G is (k, n)-pancyclic.
- (vii) If G is N-free, then G is (k, n)-pancyclic.
- (viii) If G is W-free, then G is (k, n)-pancyclic.

These results are best possible under the given conditions.

#### References

 P. Bedrossian, Forbidden subgraph and minimum degree conditions for Hamiltonicity (Ph.D. Thesis, Memphis State University, 1991).

- [2] J.A. Bondy, *Pancyclic graphs*, in: Proceedings of the Second Louisiana Conference on Combinatorics, Graph Theory, and Computing, Ray C. Mullin (Ed(s)), (Louisiana State University, Baton Rouge, LA, 1971) 167–172. doi:10.1016/0095-8956(71)90016-5
- [3] J.A. Bondy, *Pancyclic graphs I*, J. Combin. Theory Ser. B **11** (1971) 80–84. doi:10.1016/0095-8956(71)90016-5
- [4] G. Chartrand, L. Lesniak and P. Zhang, Graphs and Digraphs, 5<sup>th</sup> Edition (Chapman and Hall/CRC, Boca Raton, FL, 2011).
- [5] C.B. Crane, Generalized pancyclic properties in claw-free graphs, Graphs Combin. 31 (2015) 2149–2158. doi:10.1007/s00373-014-1510-5
- [6] Y. Egawa, J. Fujisawa, S. Fujita and K. Ota, On 2-factors in r-connected {K<sub>1,k</sub>, P<sub>4</sub>}free graphs, Tokyo J. Math. **31** (2008) 415–420. doi:10.3836/tjm/1233844061
- [7] R.J. Faudree and R.J. Gould, Characterizing forbidden pairs for Hamiltonian properties, Discrete Math. 173 (1997) 45–60. doi:10.1016/S0012-365X(96)00147-1
- [8] R.J. Faudree, R.J. Gould, M.S. Jacobson and L. Lesniak, Generalizing pancyclic and k-ordered graphs, Graphs Combin. 20 (2004) 291–310. doi:10.1007/s00373-004-0576-x
- [9] O. Ore, Note on hamilton circuits, Amer. Math. Monthly 67 (1960) 55. doi:10.2307/2308928

Received 22 July 2015 Revised 3 June 2016 Accepted 4 June 2016