

## CONSTANT SUM PARTITION OF SETS OF INTEGERS AND DISTANCE MAGIC GRAPHS

SYLWIA CICHACZ AND AGNIESZKA GÖRLICH

AGH University of Science and Technology  
Faculty of Applied Mathematics

**e-mail:** cichacz@agh.edu.pl  
forys@agh.edu.pl

### Abstract

Let  $A = \{1, 2, \dots, tm + tn\}$ . We shall say that  $A$  has the  $(m, n, t)$ -balanced constant-sum-partition property ( $(m, n, t)$ -BCSP-property) if there exists a partition of  $A$  into  $2t$  pairwise disjoint subsets  $A^1, A^2, \dots, A^t, B^1, B^2, \dots, B^t$  such that  $|A^i| = m$  and  $|B^i| = n$ , and  $\sum_{a \in A^i} a = \sum_{b \in B^j} b$  for  $1 \leq i \leq t$  and  $1 \leq j \leq t$ . In this paper we give sufficient and necessary conditions for a set  $A$  to have the  $(m, n, t)$ -BCSP-property in the case when  $m$  and  $n$  are both even. We use this result to show some families of distance magic graphs.

**Keywords:** constant sum partition, distance magic labeling, product of graphs.

**2010 Mathematics Subject Classification:** 05C78.

### 1. INTRODUCTION

Let  $A = \{1, 2, \dots, tm + tn\}$ . We shall say that  $A$  has the  $(m, n, t)$ -balanced constant-sum-partition property ( $(m, n, t)$ -BCSP-property) if there exists a partition of  $A$  into pairwise disjoint subsets  $A^1, A^2, \dots, A^t, B^1, B^2, \dots, B^t$  such that  $|A^i| = m$  and  $|B^i| = n$ , and  $\sum_{a \in A^i} a = \sum_{b \in B^j} b$  for  $1 \leq i, j \leq t$ . A positive integer  $\mu = \sum_{a \in A^i} a = \sum_{b \in B^j} b$  is called a *balanced constant*.

All graphs considered in this paper are simple finite graphs. Given a graph  $G$ , we denote its order by  $|G|$ , its size by  $||G||$ , its vertex set by  $V(G)$  and the edge set by  $E(G)$ . The *neighborhood*  $N(x)$  of a vertex  $x$  is the set of vertices adjacent to  $x$ , and the *degree*  $d(x)$  of  $x$  is  $|N(x)|$ , the size of the neighborhood of  $x$ .

*Distance magic labeling* (also called *sigma labeling*) of a graph  $G = (V, E)$  of order  $n$  is a bijection  $l: V \rightarrow \{1, 2, \dots, n\}$  with the property that there is a positive integer  $k$  (called *magic constant*) such that  $w(x) = \sum_{y \in N_G(x)} l(y) = k$  for every  $x \in V$ . If a graph  $G$  admits a distance magic labeling, then we say that  $G$  is a *distance magic graph* (see [29]). It was proved recently that the magic constant is unique ([27]).

The concept of distance magic labeling has been motivated by the construction of magic rectangles. Magic rectangles are a natural generalization of magic squares which have long intrigued mathematicians and the general public [17]. A *magic  $(m, n)$ -rectangle*  $S$  is an  $m \times n$  array in which the first  $mn$  positive integers are placed so that the sum over each row of  $S$  is constant and the sum over each column of  $S$  is another (different if  $m \neq n$ ) constant. Harmuth proved the following theorem.

**Theorem 1** [19, 20]. *For  $m, n > 1$  there is a magic  $(m, n)$ -rectangle  $S$  if and only if  $m \equiv n \pmod{2}$  and  $(m, n) \neq (2, 2)$ .*

As in the case of magic squares, we can construct a distance magic complete  $m$  partite graph with each part size equal to  $n$  by labeling the vertices of each part by the columns of the magic rectangle. Moreover, observe that constant sum partition of  $\{1, 2, \dots, n\}$  leads to complete multipartite distance magic labeled graphs. For instance, the partition  $\{1, 4\}, \{2, 3\}$  of the set  $\{1, 2, 3, 4\}$  with constant sum 5 leads to distance magic labeling of the complete bipartite graph  $K_{2,2}$ , see [6]. Beena proved the following.

**Theorem 2** [6]. *Let  $m$  and  $n$  be two positive integers such that  $m \leq n$ . The complete bipartite graph  $K_{m,n}$  is a distance magic graph if and only if*

- $m + n \equiv 0$  or  $3 \pmod{4}$ , and
- either  $n \leq \lfloor (1 + \sqrt{2})m - \frac{1}{2} \rfloor$  or  $2(2n + 1)^2 - (2m + 2n - 1)^2 = 1$ .

Moreover, Kotlar recently gave necessary and sufficient conditions for complete 4-partite graph to be distance magic (see [22]). He also posted the following open problem.

**Problem 1.1** [22]. Let  $n, k$  and  $p_1, p_2, \dots, p_k$  be positive integers such that  $p_1 + p_2 + \dots + p_k = n$  and  $\binom{n+1}{2}/k$  is an integer. When is it possible to find a partition of the set  $\{1, 2, \dots, n\}$  into  $k$  subsets of sizes  $p_1, p_2, \dots, p_k$ , respectively, such that the sum of the elements in each subset is  $\binom{n+1}{2}/k$ ?

A similar problem was also considered in [2, 7, 9, 12, 14, 23, 24]. Namely, a non-increasing sequence  $\langle m_1, \dots, m_k \rangle$  of positive integers is said to be  *$n$ -realizable* if the set  $\{1, 2, \dots, n\}$  can be partitioned into  $k$  mutually disjoint subsets  $X_1, X_2, \dots, X_k$  such that  $\sum_{x \in X_i} x = m_i$  for each  $1 \leq i \leq k$ . The study of  $n$ -realizable sequences was motivated by the ascending subgraph decomposition

problem posed by Alavi, Boals, Chartrand, Erdős and Oellerman [1], which asks for a decomposition of a given graph  $G$  of size  $\binom{n+1}{2}$  by subgraphs  $H_1, H_2, \dots, H_n$ , where  $H_i$  has size  $i$  and is a subgraph of  $H_{i+1}$  for each  $i = 1, 2, \dots, n-1$ . These authors conjectured that a forest of stars of size  $\binom{n+1}{2}$  with each component having at least  $n$  edges admits an ascending subgraph decomposition by stars. This is equivalent to the fact that every non-increasing sequence  $\langle m_1, \dots, m_k \rangle$  with  $\sum_{i=1}^k m_i = \binom{n+1}{2}$  and  $m_k \geq n$  is  $n$ -realizable, a result which was proved by Ma, Zhou and Zhou [25]. Although the general ascending subgraph decomposition conjecture is unsolved so far, some partial results have been obtained [10, 11, 13].

We recall two out of four standard graph products (see [21]). Both, the *lexicographic product*  $G \circ H$  and the *direct product*  $G \times H$  are graphs with the vertex set  $V(G) \times V(H)$ . Two vertices  $(g, h)$  and  $(g', h')$  are adjacent in:

- $G \circ H$  if and only if either  $g$  is adjacent to  $g'$  in  $G$  or  $g = g'$  and  $h$  is adjacent to  $h'$  in  $H$ ;
- $G \times H$  if  $g$  is adjacent to  $g'$  in  $G$  and  $h$  is adjacent to  $h'$  in  $H$ .

The graph  $G \circ H$  is also called the *composition* and denoted by  $G[H]$  (see [18]). The product  $G \times H$ , also known as *Kronecker product*, *tensor product*, *categorical product* and *graph conjunction*, is the most natural graph product.

Some graphs which are distance magic among (some) products can be found in [3, 4, 6, 8, 16, 26, 28].

The following problem was posted in [5].

**Problem 1.2** [5]. If  $G$  is non-regular graph, determine if there is a distance magic labeling of  $G \circ C_4$ .

Anholcer and Cichacz proved the following.

**Theorem 3** [3]. *Let  $m$  and  $n$  be integers such that  $1 \leq m < n$ . Then  $K_{m,n} \circ C_4$  is distance magic if and only if the following conditions hold.*

- (1) *The numbers*

$$a = \frac{(m+n)(4m+4n+1)(2m-1)}{4mn-m-n}$$

*and*

$$b = \frac{(m+n)(4m+4n+1)(2n-1)}{4mn-m-n}$$

*are integers.*

- (2) *There exist integers  $p, q, t \geq 1$  such that*

$$p + q = (b - a),$$

$$4n = pt,$$

$$4m = qt.$$

Moreover, they showed that a product  $C_3^{(t)} \circ C_4$  is not distance magic, where  $C_3^{(t)}$ , called a *Dutch Windmill Graph*, is the graph obtained by taking  $t > 1$  copies of the cycle graph  $C_3$  with a vertex in common [15]. We prove that also the product  $C_3^{(t)} \times C_4$  is not distance magic.

Thus we state a problem similar to Problem 1.2 for direct product.

**Problem 1.3.** If  $G$  is a non-regular graph, determine if there is a distance magic labeling of  $G \times C_4$ .

The paper is organized as follows. In the next section we focus on sets having an  $(m, n, t)$ -BCSP-property. We give the necessary and sufficient conditions for a set  $A = \{1, 2, \dots, tm + tn\}$  to have the  $(m, n, t)$ -BCSP-property in the case when  $m$  and  $n$  are both even. In the third section we generalize the Beena's result ([6]) by showing necessary and sufficient conditions for  $t$  copies of  $K_{m,n}$  ( $tK_{m,n}$ ) to be distance magic, if  $m$  and  $n$  are both even. We use this result to give necessary and sufficient conditions for the direct product  $K_{m,n} \times C_4$  to be distance magic.

## 2. CONSTANT SUM PARTITION

**Theorem 4.** *Let  $m$  and  $n$  be two positive integers such that  $m \leq n$ . If the set  $A = \{1, 2, \dots, tm + tn\}$  has the  $(m, n, t)$ -BCSP-property, then the conditions hold:*

- $m + n \equiv 0 \pmod{4}$  or  $tm + tn \equiv 3 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ .

**Proof.** Suppose that  $A^1, A^2, \dots, A^t, B^1, B^2, \dots, B^t$  is an  $(m, n, t)$ -constant sum partition of the set  $A$ . Let  $A^i = \{a_0^i, a_1^i, \dots, a_{m-1}^i\}$  and  $B^i = \{b_0^i, b_1^i, \dots, b_{n-1}^i\}$  for  $i = 1, 2, \dots, t$ . Since for the balanced constant  $\mu$  we have  $\mu = \sum_{i=0}^{m-1} a_i^j = \sum_{l=0}^{n-1} b_l^j$ , for  $j = 1, 2, \dots, t$ , it is easy to observe that

$$\mu = \frac{1}{2t} \sum_{i=1}^{tn+tm} i = \frac{(tm + tn)(tm + tn + 1)}{4t},$$

which implies that  $m + n \equiv 0 \pmod{4}$  or  $tm + tn \equiv 3 \pmod{4}$ . Notice that  $\sum_{i=0}^{m-1} \sum_{j=1}^t a_i^j \leq \sum_{i=1}^{tm} (i + tn) = \frac{tm(tm+2tn+1)}{2}$ , thus  $\mu \leq \frac{m(tm+2tn+1)}{2}$ . This implies  $(m + n)(tm + tn + 1) \leq 2m(tm + 2tn + 1)$  and therefore

$$\left[ tm + \left( tn + \frac{1}{2} \right) \right]^2 \geq t^2 n^2 + tn + \left( tn + \frac{1}{2} \right)^2 = \frac{(2tn + 1)^2}{2} - \frac{1}{4}.$$

That is

$$1 \geq 2(2tn + 1)^2 - (2tm + 2tn + 1)^2.$$

Therefore,  $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ . ■

**Theorem 5.** *Let  $m$  and  $n$  be two positive integers such that  $m \leq n$ . If the conditions hold:*

- $m + n \equiv 0 \pmod{4}$  or  $tm + tn \equiv 3 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$ ,

*then the set  $A = \{1, 2, \dots, tm + tn\}$  has the  $(m, n, t)$ -BCSP-property.*

**Proof.** Using the same arguments as in the proof of Theorem 4, the condition  $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  relates to the solution when the  $tm$  elements in  $A^1 \cup A^2 \cup \dots \cup A^t$  have to be the  $tm$  largest integers  $1 + tn, 2 + tn, \dots, tn + tm$  (because then  $\sum_{i=0}^{m-1} \sum_{j=1}^t a_i^j = \sum_{i=1}^{tm} (i + tn) = \frac{tm(tm+2tn+1)}{2}$ ), whereas the  $tn$  elements in  $B^1 \cup B^2 \cup \dots \cup B^t$  have to be the  $tn$  smallest integers  $1, 2, \dots, tn$  and  $\mu = \frac{m(tm+2tn+1)}{2} = \frac{n(tn+1)}{2}$ . Notice that if  $m$  or  $n$  is odd, then  $t$  is odd since the constant  $\mu$  is an integer.

If  $m$  is odd, then there exists a magic  $(t, m)$ -rectangle by Theorem 1. Let  $a_{i,j}$  be an  $(i, j)$ -entry of the  $(t, m)$ -rectangle,  $0 \leq i \leq t-1$  and  $0 \leq j \leq m-1$ . Notice that  $\sum_{j=0}^{m-1} a_{i,j} = \frac{m(1+tm)}{2}$ . Let  $a_j^i = a_{i,j} + tn$ , for  $j = 0, 1, \dots, m-1$  and  $i = 0, 1, \dots, t-1$ .

If  $n$  is odd, then there exists a magic  $(t, n)$ -rectangle by Theorem 1. Let  $b_{i,j}$  be an  $(i, j)$ -entry of the  $(t, n)$ -rectangle,  $0 \leq i \leq t-1$  and  $0 \leq j \leq n-1$ . Notice that  $\sum_{i=0}^{n-1} b_{i,j} = \frac{n(1+tn)}{2}$ . Let  $b_j^i = b_{i,j}$ , for  $j = 0, 1, \dots, n-1$  and  $i = 0, 1, \dots, t-1$ .

If  $m$  is even, then  $a_{2j}^i = tn + i\frac{m}{2} + j + 1$ ,  $a_{2j+1}^i = tn + tm - i\frac{m}{2} - j$ , for  $j = 0, 1, \dots, m/2 - 1$  and  $i = 0, 1, \dots, t-1$ .

If  $n$  is even, then  $b_{2j}^i = i\frac{n}{2} + j + 1$ ,  $b_{2j+1}^i = tn - i\frac{n}{2} - j$ , for  $j = 0, 1, \dots, n/2 - 1$  and  $i = 0, 1, \dots, t-1$ . ■

**Theorem 6.** *Let  $m$  and  $n$  be two positive even integers such that  $m \leq n$ . The set  $A = \{1, 2, \dots, tm + tn\}$  has the  $(m, n, t)$ -BCSP-property if and only if the conditions hold:*

- $m + n \equiv 0 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ .

**Proof.** The necessity is obvious by Theorem 4. Suppose now that  $m$  and  $n$  are positive even integers satisfying above assumptions. We can also assume that  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$  (which in these case is equivalent to  $m > -n + \frac{\sqrt{2(2tn+1)^2-1}-1}{2t}$  since  $\sqrt{2(2tn+1)^2-1} > \sqrt{2(2tn+1)^2-1}$ ), by Theorem 5.

Let us partition the set  $A$  into  $t$  disjoint sets  $V_i = \{i + 2tj, 2t - i + 1 + 2tj, j \in \{0, 1, \dots, \frac{m+n-2}{2}\}\}$  for  $i \in \{1, \dots, t\}$  with cardinality  $m + n$ . For every  $a \in V_i$  let  $\bar{a}$  denote the element in  $V_i$  such that  $a + \bar{a} = tm + tn + 1$ . Observe that for every element  $a \in V_i$  there exists  $\bar{a} \in V_i$ . The sum of integers in each set  $V_i$  is  $K = \frac{(1+tm+tn)(m+n)}{2}$ . Obviously, a balanced constant is  $\mu = \frac{K}{2}$ .

Let  $W_i$  be the sequence of  $m$  greatest integers in  $V_i$  for every  $i \in \{1, \dots, t\}$ , so  $W_i = (tn + i, tm + tn - (m - 2)t - i + 1, \dots, tm + tn - 2t + i, tm + tn - i + 1)$ . Denote the  $j$ -th element in a sequence  $W_i$  by  $w_i^j$ . Then for each  $i$  we obtain that

$$\sum_{j=1}^m w_i^j = \frac{m(1 + tm + 2tn)}{2} =: S.$$

Since  $m > -n + \frac{\sqrt{2(2tn+1)^2-1}-1}{2t}$ , observe that  $S - \mu > 0$ . Hence, there exist nonnegative integers  $k$  and  $d$  such that  $S - \mu = km + d$ , where  $0 \leq d < m$ . Therefore,  $S - \mu = \frac{tm^2}{4} + \frac{tmn}{2} + \frac{m-n}{4} - \frac{tn^2}{4} = km + d \leq \frac{tmn}{2}$ , since  $m \leq n$ . Hence, we obtain that  $k \leq \frac{tn}{2}$ . Furthermore,  $tn - k > 0$ .

If  $d = 0$  we create sets  $A_1, \dots, A_t$  putting  $A_i = \{w_i^1 - k, \dots, w_i^m - k\}$ . Note that  $A_i \cap A_j = \emptyset$  for every  $i \neq j$ . Moreover,  $\sum_{a \in A_i} a = S - mk = \mu$  for  $i \in \{1, \dots, t\}$ .

Let  $B'_i = \{\overline{w_i^1 - k}, \dots, \overline{w_i^m - k}\}$ . Observe that the set

$$B = A \setminus \left( \bigcup_{i=1}^t A_i \cup \bigcup_{i=1}^t B'_i \right)$$

has cardinality  $t(n - m)$ . Indeed, we can part it into  $\frac{t(n-m)}{2}$  pairs with type  $\{a, \bar{a}\}$  (see Example 7). Then we part the set  $B$  into  $t$  disjoint subsets  $B_1'', \dots, B_t''$  with cardinality  $n - m$  so that the elements of every set  $B_i''$  create exactly  $\frac{n-m}{2}$  pairs with type  $\{a, \bar{a}\}$ . Let  $B_i = B'_i \cup B_i''$  for  $i \in \{1, \dots, t\}$ . Then each set  $B_i$  contains  $n$  elements and  $B_i \cap B_j = \emptyset$  for  $i \neq j$ . Furthermore,

$$\sum_{b \in B_i} b = \frac{(n - m)(tm + tn + 1)}{2} + m(tm + tn + 1) - \mu = \mu.$$

**Example 7.** Let  $m = n = t = 2$ . Then  $A = \{1, 2, \dots, 8\}$ ,  $S = 13$ ,  $\mu = 9$ . Since  $V_1 = \{1, 4, 5, 8\}$  and  $V_2 = \{2, 3, 6, 7\}$ , we have  $W_1 = \{5, 8\}$  and  $W_2 = \{6, 7\}$ . Observe that  $d = 0$  and then  $A_1 = \{3, 6\}$ ,  $A_2 = \{4, 5\}$ ,  $B'_1 = \{6, 3\}$  and  $B'_2 = \{5, 4\}$ . Therefore  $B = \{1, 2, 7, 8\}$  and elements of it create two pairs with type  $\{a, \bar{a}\}$ , namely  $\{1, 8\}$  and  $\{2, 7\}$ .

If  $d > 0$  we create sets  $A_i$  as follows. We subtract 1 from each of the first  $d$  labels:  $A_i = \{w_i^1 - k - 1, \dots, w_i^d - k - 1, w_i^{d+1} - k, \dots, w_i^m - k\}$  for

$i \in \{1, \dots, t\}$ . Then  $\sum_{a \in A_i} a = S - mk - d = \mu$  for  $i \in \{1, \dots, t\}$ . Then  $B'_i = \{\overline{w_i^1 - k - 1}, \dots, \overline{w_i^d - k - 1}, \overline{w_i^{d+1} - k}, \dots, \overline{w_i^m - k}\}$  and elements of a set  $B = A \setminus (\bigcup_{i=1}^t A_i \cup \bigcup_{i=1}^t B'_i)$  create  $\frac{t(n-m)}{2}$  pairs with type  $\{a, \bar{a}\}$ . As above, we part the set  $B$  into  $t$  disjoint subsets  $B''_1, \dots, B''_t$  with cardinality  $n - m$  so that the elements of every set  $B''_i$  create exactly  $\frac{n-m}{2}$  pairs with type  $\{a, \bar{a}\}$  and define pairwise disjoint sets  $B_i = B'_i \cup B''_i$  for  $i \in \{1, \dots, t\}$ . Each set  $B_i$  contains  $n$  elements and  $\sum_{b \in B_i} b = \mu$ .

Hence  $A$  has the  $(m, n, t)$ -BCSP-property. ■

Notice that although the numbers  $m = 3$ ,  $n = 6$ ,  $t = 3$  satisfy the necessary conditions of Theorem 4, they do not satisfy the sufficient conditions either of Theorem 5 or 6. Let  $A_1 = \{10, 26, 27\}$ ,  $A_2 = \{14, 24, 25\}$ ,  $A_3 = \{18, 22, 23\}$ ,  $B_1 = \{1, 4, 7, 13, 17, 21\}$ ,  $B_2 = \{2, 5, 8, 12, 16, 20\}$ ,  $B_3 = \{3, 6, 9, 11, 15, 19\}$ . Thus, the set  $A = \{1, 2, \dots, 27\}$  has the  $(3, 6, 3)$ -BCSP-property. Therefore, we conclude this section by stating the following.

**Conjecture 2.1.** *Let  $m$  and  $n$  be two positive integers such that  $m \leq n$ . The set  $A = \{1, 2, \dots, tm + tn\}$  has the  $(m, n, t)$ -BCSP-property if and only if the conditions hold:*

- $m + n \equiv 0 \pmod{4}$  or  $tm + tn \equiv 3 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ .

Recall that the conjecture is true for  $t = 1$  by Theorem 2. Moreover, one can verify that the conjecture is also true for  $t = 2$  (see e.g. [22], Theorem 2).

### 3. DISTANCE MAGIC GRAPHS

We obtain the following corollaries by Theorem 6.

**Corollary 1.** *Let  $m$  and  $n$  be two positive even integers such that  $m \leq n$ . The graph  $tK_{m,n}$  is distance magic if and only if the conditions hold:*

- $m + n \equiv 0 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ .

Let  $K_{m[a], n[b]} \cong \underbrace{K_{m, \dots, m}}_a \underbrace{K_{n, \dots, n}}_b$ .

**Corollary 2.** *Let  $m$  and  $n$  be two positive even integers such that  $m \leq n$ . The graph  $K_{m[t], n[t]}$  is distance magic if and only if the conditions hold:*

- $m + n \equiv 0 \pmod{4}$ , and
- $1 = 2(2tn + 1)^2 - (2tm + 2tn + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{2t}$ .

**Corollary 3.** Let  $m$  and  $n$  be two positive integers such that  $m \leq n$ . The graph  $K_{m,n} \times C_4$  is a distance magic graph if and only if the following conditions hold:

- $m + n \equiv 0 \pmod{2}$ , and
- $1 = 2(8n + 1)^2 - (8m + 8n + 1)^2$  or  $m \geq (\sqrt{2} - 1)n + \frac{\sqrt{2}-1}{8}$ .

**Proof.** Since  $K_{m,n} \times C_4 \cong 2K_{2m,2n}$  we are done by Theorem 2.  $\blacksquare$

We now show that there does not exist a distance magic labeling for  $C_3^{(t)} \times C_4$ .

**Theorem 8.** *The graph  $C_3^{(t)} \times C_4$  is not a distance magic graph.*

**Proof.** Let  $C_3^{(t)}$  have the central vertex  $x$  and let vertices  $y_i, z_i$  belong to  $i$ th copy of a cycle  $C_3$ . Let  $C_4 = v^0 v^1 v^2 v^3 v^0$ . Suppose that  $l$  is a distance magic labeling of the graph  $H = C_3^{(t)} \times C_4$  and  $k = w(x)$ , for all vertices  $x \in V(C_3^{(t)} \times C_4)$ . Let

- $l(x, v^0) + l(x, v^2) = s_1$ ,
- $l(x, v^1) + l(x, v^3) = s_2$ ,
- $l(y_i, v^0) + l(y_i, v^2) = a_i^1$ ,
- $l(y_i, v^1) + l(y_i, v^3) = a_i^2$ ,
- $l(z_i, v^0) + l(z_i, v^2) = b_i^1$ ,
- $l(z_i, v^1) + l(z_i, v^3) = b_i^2$ ,

for  $0 \leq i \leq t - 1$ .

Since  $k = a_i^1 + s_2 = b_i^1 + s_2$  and  $k = a_i^2 + s_1 = b_i^2 + s_1$ , we observe that  $l(y_i, v^0) + l(y_i, v^2) = l(z_i, v^0) + l(z_i, v^2) = a_1$  and  $l(y_i, v^1) + l(y_i, v^3) = l(z_i, v^1) + l(z_i, v^3) = a_2$  for  $0 \leq i \leq t - 1$ . Furthermore, since  $k = w(x, v^0) = 2ta_2 = w(x, v^1) = 2ta_1$ , we have  $a_1 = a_2 = a$  and hence  $s_1 = s_2 = s$ .

Notice that  $2s + 4ta = \sum_{x \in V(H)} l(x) = \sum_{i=1}^{8t+4} i = (4t + 2)(8t + 5)$ . Since  $k = 2ta = a + s$ , we obtain that  $(4t - 1)a = (2t + 1)(8t + 5)$ . Recall that  $a$  needs to be an integer, hence  $(4t - 1)$  needs to divide  $(22t + 5)$ . Therefore we obtain that  $t \in \{1, 2\}$ . Suppose that  $t = 2$  then  $|V(H)| = 20$ ,  $a = 15$ ,  $s = 45$ , then  $l(x, v^i) = 15$  for some  $i = 0, 1, 2, 3$  and thus  $l(x, v^{i+2}) = 30 > 20$ , a contradiction.  $\blacksquare$

Notice that if we want to find the values of  $m$  and  $n$  such that  $K_{m,n} \times C_4$  is a distance magic graph we need to solve the Diophantine equation

$$(1) \quad \alpha = 2(4n + 1)^2 - (4m + 4n + 1)^2$$

for some integer  $\alpha \leq 1$ . For instance if  $\alpha = 1$ , then the equation (1) is a Pell's equation, thus for example  $K_{102,246} \times C_4$  is a distance magic graph.



## REFERENCES

- [1] Y. Alavi, A.J. Boals, G. Chartrand, P. Erdős and O.R. Oellerman, *The ascending subgraph decomposition problem*, Congr. Numer. **58** (1987) 7–14.
- [2] K. Ando, S. Gervacio and M. Kano, *Disjoint subsets of integers having a constant sum*, Discrete Math. **82** (1990) 7–11.  
doi:10.1016/0012-365X(90)90040-O
- [3] M. Anholcer and S. Cichacz, *Note on distance magic products  $G \circ C_4$* , Graphs Combin. **31** (2015) 1117–1124.  
doi:10.1007/s00373-014-1453-x
- [4] M. Anholcer, S. Cichacz, I. Peterin and A. Tepeh, *Distance magic labeling and two products of graphs*, Graphs Combin. **31** (2015) 1125–1136.  
doi:10.1007/s00373-014-1455-8
- [5] S. Arumugam, D. Froncek and N. Kamatchi, *Distance magic graphs — A survey*, J. Indones. Math. Soc., Special Edition (2011) 11–26.
- [6] S. Beena, *On  $\Sigma$  and  $\Sigma'$  labelled graphs*, Discrete Math. **309** (2009) 1783–1787.  
doi:10.1016/j.disc.2008.02.038
- [7] F.L. Chen, H.L. Fu, Y. Wang and J. Zhou, *Partition of a set of integers into subsets with prescribed sums*, Taiwanese J. Math. **9** (2005) 629–638.
- [8] S. Cichacz, D. Froncek, E. Krop and C. Raridan, *Distance magic Cartesian products of graphs*, Discuss. Math. Graph Theory **36** (2016) 299–308.  
doi:10.7151/dmgt.1852
- [9] H. Enomoto and M. Kano, *Disjoint odd integer subsets having a constant even sum*, Discrete Math. **137** (1995) 189–193.  
doi:10.1016/0012-365X(93)E0128-Q
- [10] R.J. Faudree, A. Gyárfás and R.H. Schelp, *Graphs which have an ascending subgraph decomposition*, Congr. Numer. **59** (1987) 49–54.
- [11] H.L. Fu and W.H. Hu, *A note on ascending subgraph decompositions of complete multipartite graphs*, Discrete Math. **226** (2001) 397–402.  
doi:10.1016/S0012-365X(00)00171-0
- [12] H.L. Fu and W.H. Hu, *A special partition of the set  $I_n$* , Bull. Inst. Combin. Appl. **6** (1992) 57–61.
- [13] H.L. Fu and W.H. Hu, *Ascending subgraph decompositions of regular graphs*, Discrete Math. **253** (2002) 11–18.  
doi:10.1016/S0012-365X(01)00445-9
- [14] H.L. Fu and W.H. Hu, *Disjoint odd integer subsets having a constant odd sum*, Discrete Math. **128** (1994) 143–150.  
doi:10.1016/0012-365X(94)90108-2
- [15] J.A. Gallian, *A dynamic survey of graph labeling*, Electron. J. Combin. (2016) #DS6.

- [16] P. Gregor and P. Kovář, *Distance magic labelings of hypercubes*, Electron. Notes Discrete Math. **40** (2013) 145–149.  
doi:10.1016/j.endm.2013.05.027
- [17] T.R. Hagedorn, *Magic rectangles revisited*, Discrete Math. **207** (1999) 65–72.  
doi:10.1016/S0012-365X(99)00041-2
- [18] F. Harary, *Graph Theory* (Addison-Wesley, Reading, MA, 1994).
- [19] T. Harmuth, *Über magische Quadrate und ähnliche Zahlenfiguren*, Arch. Math. Phys. **66** (1881) 286–313.
- [20] T. Harmuth, *Über magische Rechtecke mit ungeraden Seitenzahlen*, Arch. Math. Phys. **66** (1881) 413–447.
- [21] R. Hammack, W. Imrich and S. Klavžar, *Handbook of Product Graphs*, Second Edition (CRC Press, Boca Raton, FL, 2011).
- [22] D. Kotlar, *Distance magic labeling in complete 4-partite graphs*, Graphs Combin. **32** (2016) 1027–1038.  
doi:10.1007/s00373-015-1627-1
- [23] A. Lladó and J. Moragas, *On the sumset partition problem*, Electron. Notes Discrete Math. **34** (2009) 15–19.  
doi:10.1016/j.endm.2009.07.003
- [24] A. Lladó and J. Moragas, *On the modular sumset partition problem*, European J. Combin. **33** (2012) 427–434.  
doi:10.1016/j.ejc.2011.09.001
- [25] K. Ma, H. Zhou and J. Zhou, *On the ascending star subgraph decomposition of star forests*, Combinatorica **14** (1994) 307–320.  
doi:10.1007/BF01212979
- [26] M. Miller, C. Rodger and R. Simanjuntak, *Distance magic labelings of graphs*, Australas. J. Combin. **28** (2003) 305–315.
- [27] A. O’Neal and P.J. Slater, *Uniqueness of vertex magic constants*, SIAM J. Discrete Math. **27** (2013) 708–716.  
doi:10.1137/110834421
- [28] S.B. Rao, T. Singh and V. Prameswaran, *Some sigma labelled graphs I*, in: *Graphs, Combinatorics, Algorithms and Applications*, S. Arumugam, B.D. Acharya and S.B. Rao, (Eds.), (Narosa Publishing House, New Delhi, 2004) 125–133.
- [29] V. Vilfred,  *$\Sigma$ -Labelled Graphs and Circulant Graphs* (Ph.D. Thesis, University of Kerala, Trivandrum, India, 1994).

Received 25 February 2016

Revised 7 October 2016

Accepted 8 October 2016