

A CHARACTERIZATION FOR 2-SELF-CENTERED GRAPHS

MOHAMMAD HADI SHEKARRIZ

MADJID MIRZAVAZIRI

Department of Pure Mathematics
Ferdowsi University of Mashhad
P.O. Box 1159, Mashhad 91775, Iran

e-mail: mh.shekariz@mail.um.ac.ir
mirzavaziri@um.ac.ir

AND

KAMYAR MIRZAVAZIRI

National Organization for Development of Exceptional Talents
Mashhad, Iran

e-mail: mirzavaziri@gmail.com

Abstract

A graph is called 2-self-centered if its diameter and radius both equal to 2. In this paper, we begin characterizing these graphs by characterizing edge-maximal 2-self-centered graphs via their complements. Then we split characterizing edge-minimal 2-self-centered graphs into two cases. First, we characterize edge-minimal 2-self-centered graphs without triangles by introducing *specialized bi-independent covering* (SBIC) and a structure named *generalized complete bipartite graph* (GCBG). Then, we complete characterization by characterizing edge-minimal 2-self-centered graphs with some triangles. Hence, the main characterization is done since a graph is 2-self-centered if and only if it is a spanning subgraph of some edge-maximal 2-self-centered graphs and, at the same time, it is a spanning supergraph of some edge-minimal 2-self-centered graphs.

Keywords: self-centered graphs, specialized bi-independent covering (SBIC), generalized complete bipartite graphs (GCB).

2010 Mathematics Subject Classification: 05C12, 05C69.

1. INTRODUCTION

Let $G = (V, E)$ be a connected finite simple graph. For $u, v \in V$ the *distance* of u and v , denoted by $d_G(u, v)$ or $d(u, v)$, is the length of a shortest path between u and v . The *eccentricity* of a vertex v , $\text{ecc}(v)$, is $\max\{d(u, v) : u \in V\}$. The maximum and minimum eccentricity of vertices of G are called *diameter* and *radius* of G and are denoted by $\text{diam}(G)$ and $\text{rad}(G)$, respectively. *Center* of a graph G is the subgraph induced by vertices with eccentricity $\text{rad}(G)$. A graph is called *self-centered* if it is equal to its center, or equivalently, its diameter equals its radius.

A graph G is called *k-self-centered* if $\text{diam}(G) = \text{rad}(G) = k$. The terminology *k-equi-eccentric graph* is also used by some authors. For studies on these graphs see [1–4, 7] and [6].

Clearly, a graph G is 1-self-centered if and only if G is a complete graph. In this paper, we try to characterize 2-self-centered graphs. An edge-maximal 2-self-centered graph can be easily characterized via a condition on its complement. We do this in Section 2, using a lemma which gives a necessary and sufficient condition for a graph to be 2-self-centered. For edge-minimal 2-self-centered graphs we need to divide the discussion into two cases: triangle-free or not. We do these in Section 3. Then, the characterization is done in sight of the following theorem.

Theorem 1. *A finite graph G is 2-self-centered if and only if there is an edge-minimal 2-self-centered graph G' and an edge-maximal 2-self-centered graph G'' such that G' is a spanning subgraph of G while G is itself a spanning subgraph of G'' .*

Proof. The proof is clear. Note that $G' \subseteq G$ implies $\text{rad}(G) \geq \text{rad}(G') = 2$ and $G \subseteq G''$ implies $\text{diam}(G) \leq \text{diam}(G'') = 2$. ■

Throughout this paper G is a connected finite simple graph and its complement is denoted by \overline{G} . If G is a graph and e is an edge in G , then $G \setminus e$ is the graph obtained from G by omitting e . Moreover, the graph obtained by adding an edge $e \notin E(G)$ to G is denoted by $G + e$. Whenever two vertices u and v are adjacent, we might write $u \sim v$. For concepts and notations of graph theory, the reader is referred to [5].

2. EDGE-MAXIMAL 2-SELF-CENTERED GRAPHS

In this section, we present a characterization for edge-maximal 2-self-centered graphs. The following lemma is not only essential to do so, but it is also going to be used all over this paper.

Lemma 2. *Let $G = (V, E)$ be a graph with n vertices. Then G is 2-self-centered if and only if the following two conditions are true:*

- (i) $2 \leq \deg(v) \leq n - 2$ for all $v \in V$;
- (ii) for each $u, v \in V$ with $uv \notin E$ there is a $w \in V$ such that $uw, vw \in E$.

Proof. The proof is obvious. Note that if G has a vertex v with $\deg(v) = n - 1$ then $\text{rad}(G) = 1$ and if there is a vertex u with $\deg(u) = 1$ then its neighbour should be adjacent to any vertex of G , since otherwise $\text{ecc}(u) > 2$. ■

Remark 3. If we show that for a graph G item (ii) of Lemma 2 holds and no vertex is adjacent to all vertices, then we can deduce that no vertex has degree 1 and therefore G is 2-self-centered.

A 2-self-centered graph G is said to be *edge-maximal* if there are no non-adjacent $u, v \in V(G)$ such that $G + uv$ is 2-self-centered. The following theorem is a characterization for edge-maximal 2-self-centered graphs.

Theorem 4. *Let G be a 2-self-centered graph. Then G is edge-maximal if and only if \overline{G} is disconnected and each connected component of \overline{G} is a star with at least two vertices.*

Proof. Let H_1, \dots, H_r be the connected components of \overline{G} , where r is a positive integer. At first, note that each H_i should be a tree with at least two vertices. To see this, if H_i has only one vertex v , then the degree of v in \overline{G} is zero and thus its degree should be $n - 1$ in G which contradicts to (i) of Lemma 2. Furthermore, if the connected component H_i is not a tree, then there is an edge e with end vertices u_0 and v_0 in H_i which is not a cut edge. Let $H = G + e$. Since G is edge-maximal, H cannot be 2-self-centered. Using Lemma 2, we can deduce that the degree of u_0 or v_0 in G must be $n - 2$. This means that the degree of u_0 or v_0 in \overline{G} is 1 and consequently e is a cut edge, a contradiction.

Now, we show that each connected component H_i is a star. Let u be a vertex with maximum degree k in H_i . If $k = 1$ then H_i is $K_{1,1}$. Let $k \geq 2$. If H_i is not $K_{1,k}$ then one of the neighbours of u , say v , has a neighbour $w \neq u$. Let e' be the edge between u and v in \overline{G} and $H' = G + e'$. Since G is edge-maximal, H' cannot be 2-self-centered. Using Lemma 2, we can again deduce that the degree of u or v in G should be $n - 2$. This means that the degree of u or v in \overline{G} is 1; which is a contradiction.

Conversely, suppose that \overline{G} is a disconnected graph whose connected components are all stars, each of which has at least two vertices. Then, $2 \leq \deg(v) \leq n - 2$ for all $v \in V(G)$ and whenever u and v are two non-adjacent vertices of G , there must be a $w \in V(G)$ such that u and v are both adjacent to w . Therefore, by Lemma 2 G is a 2-self-centered graph. Moreover, since every connected component of \overline{G} is a star with at least two vertices, adding an edge between two

non-adjacent vertices in G makes the complement to have a singleton as a connected component, which means that the resulted graph is not 2-self-centered. ■

3. EDGE-MINIMAL 2-SELF-CENTERED GRAPHS

A 2-self-centered graph G is said to be *edge-minimal* if for each $e \in E(G)$, $G \setminus e$ is not a 2-self-centered graph. In this section, we determine all edge-minimal 2-self-centered graphs. To do so, let at first suppose that \overline{G} is disconnected.

Proposition 5. *Let G be a graph. Then G is an edge-minimal 2-self-centered graph such that \overline{G} is disconnected if and only if it is the complete bipartite graph $K_{k,\ell}$ for some $k, \ell \geq 2$.*

Proof. Let H_1, \dots, H_r be the connected components of \overline{G} , where $r \geq 2$. At first we prove that each H_i is a clique in \overline{G} , or in another word, each H_i is an independent set in G . Let e be an edge in G between two vertices u and v of H_i . If $H = G \setminus e$, then edge minimality of G implies that H cannot be 2-self-centered.

Let u' and v' be two non-adjacent vertices of H . Then u' and v' are belonged to a connected component H_j of \overline{G} . Let w' be any vertex of $H_{j'}$, where $j' \neq j$. Thus $u'w', w'v' \in E(H)$. This shows that H satisfies part (ii) of Lemma 2.

Since H is not 2-self-centered, Lemma 2 implies that the degree of u or v in H is 1. Let the degree of u in H be 1. Thus u has a neighbour w in H . This implies that all other vertices of G are in H_i . We know that v is also in H_i . Thus H_i contains all vertices except w and w is itself a component. Hence, the degree of w in G is $n - 1$ which contradicts to Lemma 2.

Now we show that $r = 2$. Let $r \geq 3$. Choose x, y and z in three different components. Let $e = xy$ and $H = G \setminus e$. Due to the existence of z , H is clearly 2-self-centered which contradicts the edge-minimality of G .

Conversely, the complete bipartite graph $K_{k,\ell}$ for $k, \ell \geq 2$ is an edge-minimal 2-self-centered graph such that its complement is disconnected. ■

For those 2-self-centered graphs that have connected complements, Proposition 5 is not useful. So, we may develop the characterization in some separate propositions for them, or, we can prove a more general statement which covers this case as a special case. In this paper, we do the later one, for which some preliminaries are needed.

Definition 6. Let G be a 2-self-centered graph. A vertex x in G is called *critical* for u and v if $uv \notin E$ and x is the only common neighbour of u and v .

Lemma 7. *Let G be an edge minimal 2-self-centered graph with no critical vertex for any pair of vertices. Then G is triangle-free. Furthermore, every triangle-free 2-self-centered graph is edge-minimal.*

Proof. Suppose in contrary that there are $u, v, w \in V(G)$ such that $uv, vw, wu \in E(G)$. If $\deg(u) = \deg(v) = \deg(w) = 2$, then G is itself a triangle which contradicts to $\text{rad}(G) = 2$. If $\deg(u) = \deg(v) = 2$, then $\text{diam}(G) = 2$ implies that all other vertices of G are neighbours of w . Thus $\deg(w) = n - 1$ which contradicts the fact that $\text{rad}(G) = 2$. Hence, at most one of the vertices u, v and w has degree 2. Suppose that $\deg(u)$ and $\deg(v)$ are both greater than 2.

Let $e = uv$ and $H = G \setminus e$. Then edge-minimality of G implies that H is not a 2-self-centered graph. Since $\deg_G(u)$ and $\deg_G(v)$ are at least 3, this happens only if there are two vertices x and y such that $d_H(x, y) > 2$. Since $d_G(x, y) \leq 2$ it can be deduced that $\{x, y\} \cap \{u, v\} \neq \emptyset$. The cases $x = u$ and $y = v$ cannot happen at the same time because we have the path $x \sim w \sim y$ in H . If $x = u$ and y is a vertex other than v , then there is a path xt, ty in G for some vertex t , since v is not critical for u and y . For the case $y = v$ and any other vertex x the argument is similar. Therefore, H is a 2-self-centered graph, a contradiction.

Moreover, let G be triangle-free. If G is not edge-minimal, then there is an edge e with ends u and v such that $G \setminus e$ is still a 2-self-centered graph. Thus there is a path of length 2 between u and v in $G \setminus e$. This gives a triangle in G . ■

Nevertheless, there are examples of edge-minimal 2-self-centered graphs possessing some critical vertices *with* or *without* triangles.

Example 8. Let G be the graph with vertex set $V = \{0, 1, 2, 3, 4, 5, 6, 7\}$ and edge set $E = \{01, 23, 12, 14, 15, 23, 36, 37, 46, 57, 67\}$. Then G is an edge-minimal 2-self-centered graph possessing the critical vertex 6 for the vertices 4 and 7, *with* a triangle on 3, 6, 7, see Figure 1.





Figure 1. The graph G of Example 8.

Example 9. Let H be a graph constructed in the following way: consider the graph $K_{3,3}$ with two vertices y and z in different parts connected by the edge e . Omit e and add a vertex x with two edges xy and xz to obtain H . Then H is an edge minimal 2-self-centered graph possessing the critical vertex x for the vertices y and z , *without* any triangle.

Figure 2. The graph H of Example 9.

Definition 10. A graph G is called to have a *specialized bi-independent covering* via $(\mathbb{A}_r, \mathbb{B}_s)$ if

- (i) G is triangle-free,
- (ii) there are two families $\mathbb{A}_r = \{A_1, \dots, A_r\}$ and $\mathbb{B}_s = \{B_1, \dots, B_s\}$ of not necessarily distinct independent subsets of G such that we have $V(G) = \bigcup_{i=1}^r A_i = \bigcup_{j=1}^s B_j$,
- (iii) for all $u, v \in V(G)$ if $d(u, v) \geq 3$, then there is an $1 \leq i \leq r$ such that $u, v \in A_i$ or there is $1 \leq j \leq s$ such that $u, v \in B_j$,
- (iv) for all $u \in V(G)$ and $i \in \{1, \dots, r\}$ if $d(u, A_i) \geq 2$, then there is a $j \in \{1, \dots, s\}$ such that $A_i \cap B_j = \emptyset$ and $u \in B_j$, and
- (v) for all $u \in V(G)$ and $j \in \{1, \dots, s\}$ if $d(u, B_j) \geq 2$, then there is an $i \in \{1, \dots, r\}$ such that $A_i \cap B_j = \emptyset$ and $u \in A_i$.

To make it easy, we shorten the name “specialized bi-independent covering” to *SBIC*. It is straightforward to check that every triangle-free graph G has two families of independent sets $\mathbb{A}_r, \mathbb{B}_s$ such that G has a SBIC via $(\mathbb{A}_r, \mathbb{B}_s)$. To see this, fix two independent coverings of G , and by adding enough independent sets to them, we can always satisfy items (iii) to (v) of Definition 10.

We need the following definition to complete our characterization of triangle-free 2-self-centered graphs.

Definition 11. A graph G is called an *X-generalized complete bipartite*, denoted by $\text{GCB}_X(k, \ell, \mathbb{A}_r, \mathbb{B}_s)$, if X has an SBIC via $(\mathbb{A}_r, \mathbb{B}_s)$ and G is constructed in the following way:

- (1) $V(G) = K \cup L \cup Y \cup Z \cup V(X)$ where $|K| = k$, $|L| = \ell$, $Y = \{y_1, \dots, y_r\}$ and $Z = \{z_1, \dots, z_s\}$.
- (2) $a \sim t$ for all $a \in K$ and $t \in L \cup Y$.
- (3) $b \sim t$ for all $b \in L$ and $t \in K \cup Z$.
- (4) $y_i \sim t$ for all $t \in A_i$ and $1 \leq i \leq r$.
- (5) $z_j \sim t$ for all $t \in B_j$ and $1 \leq j \leq s$.
- (6) $y_i \sim z_j$ if and only if $A_i \cap B_j = \emptyset$.

Moreover, there are some special cases that must be treated separately:

- (7) If $k = 0$ then every member of Y has a neighbour in Z and for all $i, j \in \{1, \dots, r\}$ we have $A_i \cap A_j \neq \emptyset$ or there is a $p \in \{1, \dots, s\}$ such that $A_i \cap B_p = A_j \cap B_p = \emptyset$.
- (8) If $\ell = 0$ then every member of Z has a neighbour in Y and for all $i, j \in \{1, \dots, r\}$ we have $A_i \cap A_j \neq \emptyset$ or there is a $p \in \{1, \dots, s\}$ such that $A_i \cap B_p = A_j \cap B_p = \emptyset$.
- (9) If $r = 0$ then $k \neq 0$ and if $s = 0$ then $\ell \neq 0$.
- (10) $r = s = 0$ if and only if $X = \emptyset$ and $k, \ell \geq 2$.
- (11) If $|X| = 1$ then at least one of k or ℓ is non-zero.

Proposition 12. *Any generalized complete bipartite graph is a triangle-free 2-self-centered graph.*

Proof. Let $G = \text{GCB}_X(k, \ell, \mathbb{A}_r, \mathbb{B}_s)$ and $t = |X|$. Then $n := |V(G)| = k + \ell + r + s + t$. We show that G has no vertex of degree $n - 1$ and then we show that item (ii) of Lemma 2 holds for G . Then by Remark 3 we deduce that G is 2-self-centered. By proving that G has no triangle and using Lemma 7, we actually show that G is also edge-minimal.

For $a \in K$, $\deg(a) = \ell + r = n - s - k - t$. Thus if $r = s = 0$ then by item (10) of Definition 11 we have $k \geq 2$ and hence $\deg(a) \leq n - 2$. If r or s is non-zero, then by item (10) we have $t \neq 0$ and therefore we have $\deg(a) \leq n - 2$ (by items (2) and (9) of Definition 11, and because no element of K is adjacent to a vertex of X).

For $b \in L$, by a similar proof to the case $a \in K$ we can deduce that $\deg(b) \leq n - 2$.

For $y_i \in Y$, if $\ell \neq 0$ then $\deg(y_i) \leq n - 2$ because no element of L is adjacent to y_i . If $\ell = 0$ then either y_i is not adjacent to all vertices of X or if y_i is adjacent to all vertices of x then it is not adjacent to z_j for some $j \in \{1, \dots, s\}$ (which its existence is supported by item (8) of Definition 11), each of which cases yields to $\deg(y_i) \leq n - 2$.

For $z_j \in Z$ we have the same argument to $y_i \in Y$.

Finally, for each $x \in X$, item (11) of Definition 11 guarantees that $\deg(x) \leq n - 2$ whenever X has only one vertex. So, assume that $t \geq 2$. Therefore, there are two possibilities: either there is $\hat{x} \in X$ such that x is not adjacent to \hat{x} , or, x is adjacent to all other vertices of X . If the former case is true then $\deg(x) \leq n - 2$. For the later case, since x is not in any independent set with other vertices of X , we have there is some $i \in \{1, \dots, r\}$ and $j \in \{1, \dots, s\}$ such that $\{x\} \cap A_i = \{x\} \cap B_j = \emptyset$. Thus, by items (4) and (5) of Definition 11, we have x is not adjacent to y_i and z_j and hence $\deg(x) \leq n - 2$.

To show that (ii) of Lemma 2 is also satisfied, we should choose two vertices u and v in G and show that whenever they are not adjacent, they have at least

one common neighbour. There are 15 different ways for choosing u and v from $G = K \cup L \cup Y \cup Z \cup X$.

If $(u, v) \in (K \times L) \cup (K \times Y) \cup (L \times Z)$ then u and v are adjacent to each other.

If $(u, v) \in (K \times K) \cup (L \times L)$ then there is a path of length 2 between u and v via one of the sets $L \cup Y$ or $K \cup Z$.

If $(u, v) \in (Y \times Y)$ then if $k \neq 0$ there is a path of length 2 between u and v via any member of K . If $k = 0$ then, by item (7) of Definition 11, we have either there is a $z_p \in Z$ which is a common neighbour of u and v , or, u and v are both adjacent to a vertex $x \in X$. The case $(u, v) \in (Z \times Z)$ is also similar.

If $(u, v) \in (L \times Y)$ then if $k \neq 0$ there is a path of length 2 between u and v via any member of K . If $k = 0$ then, by item (7) of Definition 11, we have every member of Y has a neighbour in Z , so Z is non-empty and v has a neighbour in Z , namely \hat{z} . Since u is also adjacent to \hat{z} by item (3) of Definition 11, there is a path of length 2 between u and v . The case $(u, v) \in (K \times Z)$ is also similar.

If $(u, v) \in Y \times Z$ then $u = y_i$ and $v = z_j$ for some i and j . If $A_i \cap B_j \neq \emptyset$ then we can choose a c in $A_i \cap B_j$ such that there is a path of length 2 between u and v via c . If $A_i \cap B_j = \emptyset$ then u is adjacent to v , by item (6) of Definition 11.

If $(u, v) \in X \times X$ then either $d_X(u, v) \leq 2$ or by item (iii) of Definition 10 there is an $i \in \{1, \dots, r\}$ or a $j \in \{1, \dots, s\}$ such that both u and v are adjacent to y_i or z_j .

If $(u, v) \in (K \times X) \cup (L \times X)$ then there is an $i \in \{1, \dots, r\}$ or a $j \in \{1, \dots, s\}$ such that v is adjacent to y_i and z_j . Then, since u is adjacent to y_i or z_j , we have $d_G(u, v) = 2$.

If $(u, v) \in (Y \times X)$ then there is an $i \in \{1, \dots, r\}$ such that $u = y_i$. Then, either $d(v, A_i) \leq 1$ which means that $d(u, v) \leq 2$, or, if $d(v, A_i) \geq 2$ then by item (iv) of Definition 10 there is a $j \in \{1, \dots, s\}$ such that $A_i \cap B_j = \emptyset$ and $v \in B_j$. Hence by items (5) and (6) of Definition 11 we have z_j is adjacent to both u and v . The case $(u, v) \in (Z \times X)$ is also similar.

So, for each of 15 ways of choosing u and v from vertices of G we have $d(u, v) \leq 2$.

We finally show that G is triangle-free. On contrary, suppose that u, v, w are vertices of a triangle in G . The case that none of u, v and w is a vertex of X cannot happen because $K \cup Z$ and $L \cup Y$ are independent sets. Since X is triangle-free, u, v and w are not all together vertices of X . Meanwhile, if only two vertices of $\{u, v, w\}$ are in X , then the third is not adjacent to the other two because they cannot be in the same independent set in X . So, at most one of $\{u, v, w\}$ is a vertex of X . Let for instance w be a vertex of X . Then u and v are not members of Y or Z at the same time, because otherwise they are not adjacent together. The case that one of u and v is in Y and the other in Z is also impossible because it is contrary to item (6) of Definition 11. ■

Theorem 13. *A graph G is a triangle-free 2-self-centered graph if and only if there are positive integers k, ℓ, r, s and a graph X which has a SBIC via $(\mathbb{A}_r, \mathbb{B}_s)$ such that $G = \text{GCB}_X(k, \ell, \mathbb{A}_r, \mathbb{B}_s)$.*

Proof. Let Y' be a maximal independent subset of G , let Z' be a maximal independent subset of $G \setminus Y'$ and let $X = G \setminus (Y' \cup Z')$. Suppose that K (respectively L) is the set of all vertices in Z' (respectively Y') which are not adjacent to any member of X and put $Y = Y' \setminus L, Z = Z' \setminus K$.

Let $a \in K$ and $y' \in Y'$. We claim that $ay' \in E$. Suppose on the contrary that $ay' \notin E$. Since $\text{diam}(G) = 2$ there is a u in G such that $au, uy' \in E$. The vertex u cannot be in Y' or Z' since Y' and Z' are independent sets. Hence $u \in X$. This contradicts to the definition of K .

A similar argument shows that each member of L is adjacent to each member of Z' .

Let $k = |K|, \ell = |L|, r = |Y|, s = |Z|, Y = \{y_1, \dots, y_r\}$ and $Z = \{z_1, \dots, z_s\}$. Now put $A_i = N_X(y_i)$ and $B_j = N_X(z_j)$. We show that A_i 's and B_j 's are independent subsets of X and X has a SBIC via $(\mathbb{A}_r, \mathbb{B}_s)$.

Let x be an arbitrary member of X . Since Y' and Z' are maximal independent, there should be neighbours for x in Y' and Z' . We know that these neighbours are in Y and Z . Let y_i and z_j be adjacent to x . Thus $x \in A_i$ and $x \in B_j$. This shows that $X = \bigcup_{i=1}^r A_i = \bigcup_{j=1}^s B_j$.

Each A_i and each B_j is independent, since G is triangle-free. Moreover, if y_i and z_j are not adjacent to each other, then since $\text{diam}(G) = 2$, there should be an $x \in X$ with $y_i x, x z_j \in E(X)$. Thus $x \in A_i \cap B_j$. If y_i is adjacent to z_j then there must not be such an x , so we have $y_i \sim z_j$ if and only if $A_i \cap B_j = \emptyset$.

Furthermore, X is triangle-free since X is a subgraph of the triangle-free graph G .

Items (iii), (iv) and (v) of Definition 10 must hold because G is a triangle-free 2-self-centered graph. Therefore, X has an SBIC via $(\mathbb{A}_r, \mathbb{B}_s)$.

Items (1) to (6) of Definition 11 have already hold. Moreover, items (7) to (11) of Definition 11 must also hold because G is a triangle-free 2-self-centered graph. Hence $G = \text{GCB}_X(k, \ell, \mathbb{A}_r, \mathbb{B}_s)$.

Since the converse is evident by Proposition 12, we are done with the proof. ■

The reader should note that every complete bipartite graph $K_{k,\ell}$ with $k, \ell \geq 2$ is a generalized complete bipartite graph $\text{GCB}_\emptyset(k, \ell, \emptyset, \emptyset)$.

Now, we can consider edge-minimal 2-self-centered graphs with some triangles. We need the following procedure to proceed.

Procedure 14. *Let G be a graph, u, v, w form a triangle in G and suppose that v is a critical vertex for u and v_1, \dots, v_p and/or u is a critical vertex for v and u_1, \dots, u_q . Remove the edge uv and add edges uv_1, \dots, uv_p and vu_1, \dots, vu_q .*

The following theorem characterizes edge-minimal 2-self-centered graphs with triangles, which completes the characterization of all 2-self-centered graphs.

Theorem 15. *Let G be a graph. Then G is an edge-minimal 2-self-centered graph with some triangle if and only if the following two conditions are true:*

- (i) *for each edge of every triangle in G , at least one end-vertex is a critical vertex (for the other end-vertex of that edge and some other vertices of G), and*
- (ii) *iteration of Procedure 14 on G (at most to the number of triangles of G) transforms G to a triangle-free 2-self-centered graph.*

Proof. Assume that u, v, w form a triangle in G . Since G is edge-minimal, if we omit the edge uv , then the resulting graph is not 2-self-centered. This shows that u or v is a critical vertex. Let u be a critical vertex. Thus there are vertices u_1, \dots, u_q such that u is the common neighbour of v and each of the u_i 's. Moreover, if v is also a critical vertex for u and some other vertices, then we suppose that v_1, \dots, v_p are the vertices such that v is a common neighbour of u and each of the v_j 's.

If we omit uv and add edges $u_1v, \dots, u_qv, uv_1, \dots, uv_p$, then the resulting graph G' is clearly 2-self-centered and the number of triangles of G' is less than the number of triangles of G . To see this, note that edges of a triangle on u, v and w are omitted and no new triangle is added. In contrary, suppose that we have a new triangle. Then it should be of the form u_i, v, t (or v_j, u, s) which contradicts to the fact that u (or v) is a critical vertex for u_i and v (for v_j and u).

If G has still some triangle then we can proceed this process. Therefore, we finally transform G into a triangle-free 2-self-centered graph.

Conversely, if the two conditions are true for a graph G with some triangles, then G is an edge-minimal 2-self-centered graph because condition (ii) guarantees that G is 2-self-centered while condition (i) obligates G to be edge-minimal. ■

REFERENCES

- [1] J. Akiyama, K. Ando and D. Avis, *Miscellaneous properties of equi-eccentric graphs*, in: Convexity and Graph Theory (Jerusalem, 1981), North-Holland Math. Stud., Amsterdam **87** (1984) 13–23.
doi:10.1016/s0304-0208(08)72802-0
- [2] K. Balakrishnan, B. Brešar, M. Changat, S. Klavžar, I. Peterin and A.R. Subhamathi, *Almost self-centered median and chordal graphs*, Taiwanese J. Math. **16** (2012) 1911–1922.
- [3] F. Buckley, *Self-centered graphs*, in: Graph Theory and Its Applications: East and West (Jinan, 1986), Ann. New York Acad. Sci. **576** (1989) 71–78.
doi:10.1111/j.1749-6632.1989.tb16384.x

- [4] F. Buckley, Z. Miller and P.J. Slater, *On graphs containing a given graph as center*, J. Graph Theory **5** (1981) 427–434.
doi:10.1002/jgt.3190050413
- [5] J.L. Gross, J. Yellen and P. Zhang, Handbook of Graph Theory, Second Edition (CRC Press., 2014).
- [6] S. Klavžar, K.P. Narayankar and H.B. Walikar, *Almost self-centered graphs*, Acta Math. Sin. (Engl. Ser.) **27** (2011) 2343–2350.
- [7] S. Negami and G.H. Xu, *Locally geodesic cycles in 2-self-centered graphs*, Discrete Math. **58** (1986) 263–268.
doi:10.1007/s10114-011-9628-3

Received 30 November 2015

Revised 24 August 2016

Accepted 1 September 2016