Discussiones Mathematicae Graph Theory 39 (2019) 589–603 doi:10.7151/dmgt.2040

2-DISTANCE COLORINGS OF INTEGER DISTANCE GRAPHS

BRAHIM BENMEDJDOUB, ISMA BOUCHEMAKH

Faculty of Mathematics, Laboratory L'IFORCE University of Sciences and Technology Houari Boumediene (USTHB) B.P. 32 El-Alia, Bab-Ezzouar, 16111 Algiers, Algeria

> e-mail: brahimro@hotmail.com isma_bouchemakh2001@yahoo.fr

> > AND

ÉRIC SOPENA

Univ. Bordeaux, Bordeaux INP, CNRS, LaBRI, UMR5800 F-33400 Talence, France e-mail: Eric.Sopena@labri.fr

Abstract

A 2-distance k-coloring of a graph G is a mapping from V(G) to the set of colors $\{1, \ldots, k\}$ such that every two vertices at distance at most 2 receive distinct colors. The 2-distance chromatic number $\chi_2(G)$ of G is then the smallest k for which G admits a 2-distance k-coloring. For any finite set of positive integers $D = \{d_1, \ldots, d_\ell\}$, the integer distance graph G = G(D)is the infinite graph defined by $V(G) = \mathbb{Z}$ and $uv \in E(G)$ if and only if $|v - u| \in D$. We study the 2-distance chromatic number of integer distance graphs for several types of sets D. In each case, we provide exact values or upper bounds on this parameter and characterize those graphs G(D) with $\chi_2(G(D)) = \Delta(G(D)) + 1$.

Keywords: 2-distance coloring, integer distance graph.

2010 Mathematics Subject Classification: 05C15, 05C12.

1. INTRODUCTION

All the graphs we consider in this paper are simple and loopless undirected graphs. We denote by V(G) and E(G) the set of vertices and the set of edges of a graph G, respectively. For any two vertices u and v of G, we denote by $d_G(u, v)$ (or simply d(u, v) whenever the graph G is clear from the context) the *distance* between u and v, that is the length of a shortest path joining u and v. We denote by $\Delta(G)$ the maximum degree of G.

A (proper) k-coloring of a graph G is a mapping from V(G) to the set of colors $\{1, \ldots, k\}$ such that every two adjacent vertices receive distinct colors. The smallest k for which G admits a k-coloring is the chromatic number of G, denoted $\chi(G)$. A 2-distance k-coloring of a graph G is a mapping from V(G) to the set of colors $\{1, \ldots, k\}$ such that every two vertices at distance at most 2 receive distinct colors. 2-distance colorings are sometimes called L(1,1)-labelings (see [5] for a survey on L(h,k)-labelings) or square colorings in the literature. The smallest k for which G admits a 2-distance k-coloring is the 2-distance chromatic number of G, denoted $\chi_2(G)$.

The square G^2 of a graph G is the graph defined by $V(G^2) = V(G)$ and $uv \in E(G^2)$ if and only if $d_G(u, v) \leq 2$. Clearly, a 2-distance coloring of a graph G is nothing but a proper coloring of G^2 and, therefore, $\chi_2(G) = \chi(G^2)$ for every graph G.

The study of 2-distance colorings was initiated by Kramer and Kramer [8] (see also their survey on general distance colorings in [9]). The case of planar graphs has attracted a lot of attention in the literature (see e.g. [1–4, 6, 10, 14]), due to the conjecture of Wegner that suggests an upper bound on the 2-distance chromatic number of planar graphs depending on their maximum degree (see [15] for more details).

In this paper, we study 2-distance colorings of distance graphs. Although several coloring problems have been considered for distance graphs (see [11] for a survey), it seems that 2-distance colorings have not been considered yet. We present in Section 2 a few basic results on the chromatic number and the 2-distance chromatic number of distance graphs. We then consider specific sets D, namely $D = \{1, a\}, a \geq 3$ (in Section 3), $D = \{1, a, a + 1\}, a \geq 3$ (in Section 4), and $D = \{1, \ldots, m, a\}, 2 \leq m < a$ (in Section 5). We finally propose some open problems in Section 6.

2. Preliminaries

Let $D = \{d_1, \ldots, d_\ell\}$ be a finite set of positive integers. The *integer distance* graph (simply called *distance graph* in the following) G = G(D) is the infinite graph defined by $V(G) = \mathbb{Z}$ and $uv \in E(G)$ if and only if $|v - u| \in D$. The following proposition follows immediately.

Proposition 1. For every positive integers d_1, \ldots, d_ℓ with $gcd(\{d_1, \ldots, d_\ell\}) = p > 1$, the distance graph G(D) has p connected components, each of them being isomorphic to the distance graph G(D') with $D' = \{d_1/p, \ldots, d_\ell/p\}$.

In this situation, we then have $\chi_2(G(D)) = \chi_2(G(D'))$ so that we can always assume gcd(D) = 1 in the following.

It is easy to observe that the square of the distance graph G(D) is also a distance graph, namely the distance graph $G(D^2)$ where

$$D^{2} = D \cup \{d + d' : d, d' \in D\} \cup \{d - d' : d, d' \in D, d > d'\}.$$

For instance, for $D = \{1, 2, 5\}$, we get $D^2 = \{1, 2, 3, 4, 5, 6, 7, 10\}$. Note that if D has cardinality ℓ , then D^2 has cardinality at most $\ell(\ell + 1)$.

As observed in the previous section, $\chi_2(G) = \chi(G^2)$ for every graph G. Therefore, since $(G(D))^2 = G(D^2)$, determining the 2-distance chromatic number of the distance graph G(D) reduces to determining the chromatic number of the distance graph $G(D^2)$. The problem of determining the chromatic number of distance graphs has been extensively studied in the literature. When $|D| \leq 2$, this question is easily solved, thanks to the following general upper bounds.

Proposition 2. For every finite set of positive integers $D = \{d_1, \ldots, d_\ell\}$ and every positive integer p such that $d_i \not\equiv 0 \pmod{p}$ for every $i, 1 \leq i \leq \ell$, $\chi(G(D)) \leq p$.

Proof. Let $\lambda: V(G(D)) \longrightarrow \{1, \ldots, p\}$ be the mapping defined by

$$\lambda(x) = 1 + (x \mod p),$$

for every integer $x \in \mathbb{Z}$. Since $d_i \not\equiv 0 \pmod{p}$ for every $i, 1 \leq i \leq \ell$, the mapping λ is clearly a proper coloring of G(D).

Theorem 3 (Walther [13]). For every finite set of positive integers D,

$$\chi(G(D)) \le |D| + 1.$$

Proof. A (|D| + 1)-coloring of G(D) can be easily produced using the First-Fit greedy algorithm, starting from vertex 0, from left to right and then from right to left, since every vertex has exactly |D| neighbors on its left and |D| neighbors on its right.

Therefore, when $|D| \leq 2$, $\chi(G(D)) = 2$ if |D| = 1 or all elements in D are odd (since G(D) is then bipartite), and $\chi(G(D)) = 3$ otherwise (since G(D) then contains cycles of odd length). The case |D| = 3 has been settled by Zhu [16]. Whenever $|D| \geq 4$, only partial results have been obtained, namely for sets D having specific properties.

Another useful result is the following.

Theorem 4 (Voigt [12], cited in [7]). For every finite set of positive integers $D = \{d_1, \ldots, d_\ell\},\$

$$\chi(G(D)) \le \min_{n \in \mathbb{N}} n(|D_n| + 1),$$

where $D_n = \{ d_i : n | d_i, 1 \le i \le \ell \}.$

A coloring λ of a distance graph G(D) is *p*-periodic, for some integer $p \ge 1$, if $\lambda(x+p) = \lambda(x)$ for every $x \in \mathbb{Z}$. Walther also proved the following.

Theorem 5 (Walther [13]). For every finite set of positive integers D, if $\chi(G(D)) \leq k$, then G(D) admits a p-periodic k-coloring for some p.

The sequence $\lambda(x) \cdots \lambda(x+p-1)$ of such a *p*-periodic coloring λ is called the *pattern* of λ . In particular, the coloring defined in the proof of Proposition 2 was *p*-periodic with pattern $12 \cdots p$. In the following, we will describe such patterns using standard notation of Combinatorics on words. For instance, the pattern 121212345 will be denoted $(12)^3345$.

Finally, note that in any 2-distance coloring of a graph G, all vertices in the closed neighborhood of any vertex must be assigned distinct colors. Therefore, we have the following.

Observation 6. For every graph G, $\chi_2(G) \ge \Delta(G) + 1$.

In particular, this bound is attained by the distance graph G(D) with $D = \{1, \ldots, k\}, k \ge 2$.

Proposition 7. For every $k \geq 2$,

$$\chi_2(G(\{1,\ldots,k\})) = 2k + 1 = \Delta(G(\{1,\ldots,k\})) + 1.$$

Proof. This directly follows from Theorem 3, since $|\{1, \ldots, k\}^2| = 2k$.

3. The Case $D = \{1, a\}, a \ge 3$

We study in this section the 2-distance chromatic number of distance graphs G(D) with $D = \{1, a\}, a \geq 3$ (note that the case a = 2 is already solved by Proposition 7).

When $D = \{1, a\}, a \ge 3$, we have $\Delta(G(D)) = 4$ and

$$D^2 = \{1, 2, a - 1, a, a + 1, 2a\}.$$

The following theorem gives the 2-distance chromatic number of any such graph.

Theorem 8. For every integer $a \geq 3$,

$$\chi_2(G(\{1,a\})) = \begin{cases} 5 & \text{if } a \equiv 2 \pmod{5}, \text{ or } a \equiv 3 \pmod{5}, \\ 6 & \text{otherwise.} \end{cases}$$

Proof. Since $\{1, a\}^2 = \{1, 2, a - 1, a, a + 1, 2a\}$, we get $d \not\equiv 0 \pmod{5}$ for every $d \in \{1, a\}^2$ if and only if $a \equiv 2 \pmod{5}$ or $a \equiv 3 \pmod{5}$ and thus, by Proposition 2 and Observation 6, $\chi_2(G(\{1, a\})) = 5$.

Note that for every $x \in \mathbb{Z}$, the set of vertices

$$C(x) = \{x - a, x - 1, x, x + 1, x + a\}$$

induces a 5-clique in $G(\{1,a\}^2)$ (see Figure 1). We now claim that every 2distance 5-coloring λ of $G(\{1,a\})$ is necessarily 5-periodic, that is $\lambda(x+5) = \lambda(x)$ for every $x \in \mathbb{Z}$. To show that, it suffices to prove that any five consecutive vertices $x, \ldots, x+4$ must be assigned distinct colors. Assume to the contrary that this is not the case and, without loss of generality, let x = 0. Since vertices 0, 1 and 2 necessarily get distinct colors, we only have to consider two cases.



Figure 1. Subgraph of the distance graph $G(\{1, a\}), a \ge 3$.

Case 1. $\lambda(0) = \lambda(3) = 1$, $\lambda(1) = 2$, $\lambda(2) = 3$. Since C(1) induces a 5-clique in $G(\{1, a\}^2)$ (depicted in bold in Figure 1), we have

$$\{\lambda(1-a), \lambda(1+a)\} = \{4, 5\},\$$

which implies

$$\{\lambda(2-a),\lambda(2+a)\} = \{4,5\}$$

(More precisely, $\lambda(2-a) = 9 - \lambda(1-a)$ and $\lambda(2+a) = 9 - \lambda(1+a)$). This implies $\lambda(3-a) = \lambda(3+a) = 2$, a contradiction since d(3-a,3+a) = 2.

Case 2. $\lambda(0) = \lambda(4) = 1$, $\lambda(1) = 2$, $\lambda(2) = 3$, $\lambda(3) = 4$. As in the previous case we have

$$\{\lambda(1-a), \lambda(1+a)\} = \{4, 5\},\$$

which implies

 $\{\lambda(2-a), \lambda(2+a)\} = \{1, 5\}.$

We then get $\lambda(3-a) = \lambda(3+a) = 2$, again a contradiction.

Therefore, $\chi_2(G(\{1,a\})) = 5$ if and only if 5 does not divide any element of $\{1,a\}^2 = \{1,2,a-1,a,a+1,2a\}$. This is clearly the case if and only if $a \equiv 2 \pmod{5}$ or $a \equiv 3 \pmod{5}$.

We finally prove that there exists a 2-distance 6-coloring of $G(\{1, a\})$ for any value of a. We consider three cases, according to the value of a (mod 3).

Case 1. $a = 3k, k \ge 1$. Let λ be the (2a - 1)-periodic mapping defined by the pattern

$$(123)^k (456)^{k-1} 45.$$

If
$$\lambda(x) = \lambda(y) = c$$
, $1 \le c \le 5$, then

 $|x-y| \in \{3q, \ 0 \le q \le k-1\} \ \cup \ \{(2a-1)p+3q, \ p \ge 1, \ 1-k \le q \le k-1\}.$

If $\lambda(x) = \lambda(y) = 6$ (which occurs if and only if $k \ge 2$), then

 $|x-y| \in \{3q, \ 0 \le q \le k-2\} \cup \{(2a-1)p+3q, \ p \ge 1, \ 2-k \le q \le k-2\}.$

Therefore, in both cases, $|x - y| \notin \{1, 2, a - 1, a, a + 1, 2a\}$, and thus λ is a 2-distance 6-coloring of $G(\{1, a\})$.

Case 2. a = 3k + 1, $k \ge 1$. In that case, the result follows from Theorem 4 (taking n = 3), since the only element divisible by 3 in $\{1, 2, a - 1, a, a + 1, 2a\}$ is a - 1.

Case 3. $a = 3k + 2, k \ge 1$. Again, the result follows from Theorem 4 (taking n = 3), since the only element divisible by 3 in $\{1, 2, a - 1, a, a + 1, 2a\}$ is a + 1.

This concludes the proof.

4. The CASE $D = \{1, a, a+1\}, a \ge 3$

We study in this section the 2-distance chromatic number of distance graphs G(D) with $D = \{1, a, a + 1\}, a \geq 3$ (note that the case a = 2 is already solved by Proposition 7).

When $D = \{1, a, a + 1\}, a \ge 3$, we have $\Delta(G(D)) = 6$ and

$$D^{2} = \{1, 2, a - 1, a, a + 1, a + 2, 2a, 2a + 1, 2a + 2\}.$$

First we prove the following.

Theorem 9. For every integer $a, a \geq 3$,

$$\chi_2(G(\{1, a, a+1\})) = 7 = \Delta(G(\{1, a, a+1\})) + 1$$

if and only if $a \equiv 2 \pmod{7}$ or $a \equiv 4 \pmod{7}$.

Proof. Since $\{1, a, a+1\}^2 = \{1, 2, a-1, a, a+1, a+2, 2a, 2a+1, 2a+2\}$, we get $d \neq 0 \pmod{7}$ for every $d \in \{1, a, a+1\}^2$ if and only if $a \equiv 2 \pmod{7}$ or $a \equiv 4 \pmod{7}$ and thus, by Proposition 2 and Observation 6, $\chi_2(G(\{1, a, a+1\})) = 7$.

Note that for every $x \in \mathbb{Z}$, the set of vertices

$$C(x) = \{x - a - 1, x - a, x - 1, x, x + 1, x + a, x + a + 1\}$$

induces a 7-clique in $G(\{1, a, a + 1\}^2)$. We now claim that every 2-distance 7coloring λ of $G(\{1, a, a + 1\})$ is necessarily 7-periodic, that is $\lambda(x + 7) = \lambda(x)$ for every $x \in \mathbb{Z}$. To show that, it suffices to prove that any 7 consecutive vertices $x, \ldots, x + 6$ must be assigned distinct colors. Assume to the contrary that this is not the case and, without loss of generality, let x = 0. Since vertices 0, 1 and 2 necessarily get distinct colors, we only have to consider four cases (see Figure 2).



Figure 2. Subgraph of the distance graph $G(\{1, a, a + 1\}), a \ge 3$.

Case 1. Vertices 0, 1, 2, 3 are colored with the colors 1, 2, 3 and 1, respectively. We consider two subcases.

Subcase (a) $\lambda(4) = 2$. Since C(1) induces a 7-clique in $G(\{1, a, a + 1\}^2)$ (depicted in bold in Figure 2), we have

$$\{\lambda(-a), \lambda(1-a), \lambda(1+a), \lambda(2+a)\} = \{4, 5, 6, 7\}.$$

Since C(3) is also a 7-clique, we also have

 $\{\lambda(2-a),\lambda(3-a),\lambda(3+a),\lambda(4+a)\}=\{4,5,6,7\}.$

This implies $\lambda(-a) = \lambda(4-a)$ or $\lambda(1+a) = \lambda(5+a)$. Each of these cases thus corresponds to Case 2 below.

Subcase (b) $\lambda(4) \neq 2$. Note that we necessarily have $\lambda(4) \neq 3$ and $\lambda(4) \neq 1$, since vertex 4 is at distance 2 and 1 from vertices 2 and 3, respectively. We can thus assume $\lambda(4) = 4$, without loss of generality. Since d(5, 4) = 1 and d(5, 3) = 2, we have $\lambda(5) \notin \{1, 4\}$. Moreover, if $\lambda(5) = 2$, we get $\lambda(2) = \lambda(5)$, which corresponds to Case 2 below. We can thus suppose either $\lambda(5) = 3$ or $\lambda(5) > 4$, say $\lambda(5) = 5$ without loss of generality. We consider these two cases separately.

(i) $\lambda(5) = 3$. In that case, we necessarily have

$$\begin{aligned} &\{\lambda(-a),\lambda(1+a)\} \subseteq \{4,5,6,7\}, \ \{\lambda(1-a),\lambda(2+a)\} \subseteq \{4,5,6,7\}, \\ &\{\lambda(2-a),\lambda(3+a)\} \subseteq \{5,6,7\}, \ \{\lambda(3-a),\lambda(4+a)\} \subseteq \{2,5,6,7\}, \\ &\{\lambda(4-a),\lambda(5+a)\} \subseteq \{5,6,7\}. \end{aligned}$$

By setting $\{x, y, z\} = \{5, 6, 7\}$, we get

$$\begin{split} &\{\lambda(-a),\lambda(1+a)\} = \{x,y\}, \ \{\lambda(1-a),\lambda(2+a)\} = \{4,z\}, \\ &\{\lambda(2-a),\lambda(3+a)\} = \{x,y\}, \ \{\lambda(3-a),\lambda(4+a)\} = \{2,z\}, \\ &\{\lambda(4-a),\lambda(5+a)\} = \{x,y\}. \end{split}$$

Since $\lambda(-a), \lambda(2-a), \lambda(4-a) \in \{x, y\}$ and $\lambda(-a) \neq \lambda(2-a), \lambda(2-a) \neq \lambda(4-a)$, it follows that $\lambda(-a) = \lambda(4-a)$. That case corresponds to Case 2 below.

(ii) $\lambda(5) = 5$. In that case, we necessarily have

$$\begin{aligned} &\{\lambda(-a),\lambda(1+a)\} \subseteq \{4,5,6,7\}, \ \{\lambda(1-a),\lambda(2+a)\} \subseteq \{4,5,6,7\}, \\ &\{\lambda(2-a),\lambda(3+a)\} \subseteq \{5,6,7\}, \ \{\lambda(3-a),\lambda(4+a)\} \subseteq \{2,6,7\}, \\ &\{\lambda(4-a),\lambda(5+a)\} \subseteq \{3,6,7\}. \end{aligned}$$

By setting $\{x, y\} = \{6, 7\}$, we get

$$\begin{split} &\{\lambda(-a),\lambda(1+a)\} = \{5,x\}, \ \{\lambda(1-a),\lambda(2+a)\} = \{4,y\}, \\ &\{\lambda(2-a),\lambda(3+a)\} = \{5,x\}, \ \{\lambda(3-a),\lambda(4+a)\} = \{2,y\}, \\ &\{\lambda(4-a),\lambda(5+a)\} = \{3,x\}. \end{split}$$

We then necessarily have either $\lambda(1+a) = \lambda(5+a)$ or $\lambda(-a) = \lambda(4-a)$ and, in both cases, we are in the situation of Case 2 below.

Case 2. Vertices 0, 1, 2, 3, 4 are colored with the colors 1, 2, 3, 4 and 1, respectively. Again considering the 7-cliques C(1), C(2) and C(3) in $G(\{1, a, a + 1\}^2)$, we get

$$\{\lambda(1-a), \lambda(2+a)\} \subseteq \{5, 6, 7\},\$$

and

$$\{\lambda(2-a), \lambda(3+a)\} \subseteq \{5, 6, 7\},\$$

a contradiction, since vertices 1 - a, 2 - a, a + 2 and a + 3 induce a 4-clique in $G(\{1, a, a + 1\}^2)$.

Case 3. Vertices 0, 1, 2, 3, 4, 5 are colored with the colors 1, 2, 3, 4, 5 and 1, respectively. Considering the 7-cliques C(1), C(2) and C(3) in $G(\{1, a, a + 1\}^2)$, we get

$$\begin{aligned} &\{\lambda(-a), \lambda(1-a), \lambda(1+a), \lambda(2+a)\} = \{4, 5, 6, 7\}, \\ &\{\lambda(2-a), \lambda(3+a)\} \subseteq \{1, \lambda(-a), \lambda(1+a)\} \setminus \{4, 5\}, \\ &\{\lambda(3-a), \lambda(4+a)\} \subseteq \{2, \lambda(1-a), \lambda(2+a)\} \setminus \{4, 5\}, \end{aligned}$$

and thus

$$\{\lambda(2-a),\lambda(3+a)\}\subseteq \{1,6,7\} \ \text{ and } \ \{\lambda(3-a),\lambda(4+a)\}\subseteq \{2,6,7\}.$$

Assuming that none of Cases 1 or 2 occurs, we have to consider two subcases.

Subcase (a) $\lambda(6) = 2$. Considering the 7-clique C(4) in $G(\{1, a, a + 1\}^2)$, we get

 $\{\lambda(4-a), \lambda(5+a)\} \subseteq \{3, \lambda(2-a), \lambda(3+a)\} \setminus \{1, 2\} = \{3, 6, 7\}.$ If $\{\lambda(4-a), \lambda(5+a)\} = \{3, 6\}$, then

$$\{\lambda(3-a), \lambda(4+a)\} = \{2,7\}$$
$$\{\lambda(2-a), \lambda(3+a)\} = \{1,6\}$$
$$\{\lambda(1-a), \lambda(2+a)\} = \{5,7\}$$

and

$$\{\lambda(-a), \lambda(1+a)\} = \{4, 6\}.$$

If $\lambda(-a) = 6$, then $\lambda(2-a) = 1$ and thus $\lambda(4-a) = \lambda(-a) = 6$ which corresponds to Case 2. If $\lambda(1+a) = 6$, then $\lambda(3+a) = 1$ and thus $\lambda(5+a) = \lambda(1+a) = 6$ which again corresponds to Case 2.

The case $\{\lambda(4-a), \lambda(5+a)\} = \{3, 7\}$ is similar and leads to the same conclusion.

Finally, if $\{\lambda(4-a), \lambda(5+a)\} = \{6,7\}$, then $\lambda(3-a) = \lambda(4+a) = 2$, a contradiction since d(3-a, 4+a) = 2.

Subcase (b) $\lambda(6) = 6$. Considering the 7-clique C(4) in $G(\{1, a, a + 1\}^2)$, we get

$$\{\lambda(4-a), \lambda(5+a)\} \subseteq \{3, \lambda(2-a), \lambda(3+a)\} \setminus \{1, 6\} = \{3, 7\}.$$

This implies

$$\{\lambda(3-a), \lambda(4+a)\} = \{2, 6\},\$$

$$\{\lambda(2-a), \lambda(3+a)\} = \{1, 7\},\$$

$$\{\lambda(1-a), \lambda(2+a)\} = \{5, 6\}$$

and

$$\{\lambda(-a), \lambda(1+a)\} = \{4, 7\}$$

If $\lambda(-a) = 7$, then $\lambda(2-a) = 1$ and thus $\lambda(4-a) = \lambda(-a) = 7$ which corresponds to Case 2. If $\lambda(1+a) = 7$, then $\lambda(3+a) = 1$ and thus $\lambda(5+a) = \lambda(1+a) = 7$ which again corresponds to Case 2.

Case 4. Vertices 0, 1, 2, 3, 4, 5, 6 are colored with the colors 1, 2, 3, 4, 5, 6 and 1, respectively. Again considering the 7-cliques C(1), C(2) and C(3) in $G(\{1, a, a+1\}^2)$, we get

$$\begin{aligned} &\{\lambda(-a),\lambda(1-a),\lambda(1+a),\lambda(2+a)\} = \{4,5,6,7\},\\ &\{\lambda(2-a),\lambda(3+a)\} \subseteq \{1,\lambda(-a),\lambda(1+a)\} \setminus \{4,5\} \end{aligned}$$

and thus

$$\{\lambda(3-a), \lambda(4+a)\} \subseteq \{2, \lambda(1-a), \lambda(2+a)\} \setminus \{4, 5, 6\} = \{2, 7\}.$$

This implies

$$\{\lambda(2-a), \lambda(3+a)\} = \{1, 6\},\$$

$$\{\lambda(1-a), \lambda(2+a)\} = \{5, 7\}$$

and

$$\{\lambda(-a), \lambda(1+a)\} = \{4, 6\}.$$

Therefore,

$$\{\lambda(4-a),\lambda(5+a)\} \subseteq \{3,\lambda(2-a),\lambda(3+a)\} \setminus \{1,6\} = \{3\},$$

a contradiction since d(4 - a, 5 + a) = 2.

Therefore, every 2-distance 7-coloring λ of $G(\{1, a, a + 1\})$ is necessarily 7-periodic, and thus $\chi_2(G(\{1, a, a + 1\})) = 7$ if and only if 7 does not divide any element of $\{1, 2, a - 1, a, a + 1, a + 2, 2a, 2a + 1, 2a + 2\}$. This is clearly the case if and only if $a \equiv 2 \pmod{7}$ or $a \equiv 4 \pmod{7}$.

The following result provides an upper bound on $\chi_2(G(\{1, a, a+1\}))$ for any value of a.

Theorem 10. For every integer $a, a \ge 3$,

$$\chi_2(G(\{1, a, a+1\})) \le 9 = \Delta(G(\{1, a, a+1\})) + 3.$$

Proof. First recall that $\{1, a, a+1\}^2 = \{1, 2, a-1, a, a+1, a+2, 2a, 2a+1, 2a+2\}$. We consider three cases, according to the value of a (mod 3).

Case 1. $a = 3k, k \ge 1$. Since the only elements divisible by 3 in $\{1, a, a+1\}^2$ are a and 2a, the result follows by Theorem 4 (taking n = 3).

Case 2. $a = 3k + 1, k \ge 1$. Let λ be the (3a + 2)-periodic mapping defined by the pattern

$$(123)^k (456)^k 7123(789)^{k-1} 4568.$$

If $\lambda(x) = \lambda(y) = c$, $1 \le c \le 6$, then

|x|

$$\begin{aligned} |x-y| &\in \{3q, \ 0 \le q \le k-1\} \\ &\cup \{3q+2a-1, \ 1-k \le q \le 0\} \\ &\cup \{(3a+2)p+2a-1, \ p>0\} \\ &\cup \{(3a+2)p-2a+1, \ p>0\} \\ &\cup \{(3a+2)p+3q, \ p>0, \ 1-k \le q < 0\} \\ &\cup \{(3a+2)p+3q+2a-1, \ p>0, \ 1-k \le q < 0\} \\ &\cup \{(3a+2)p+3q, \ p>0, \ 0 < q \le k-1\} \\ &\cup \{(3a+2)p+3q-2a+1, \ p>0, \ 0 < q \le k-1\}. \end{aligned}$$

If $\lambda(x) = \lambda(y) = 7$ (which occurs if and only if $k \ge 2$), then

$$\begin{aligned} |x-y| &\in \{3q, \ 0 \le q \le k-2\} \\ &\cup \{3q+4, \ 0 \le q \le k-2\} \\ &\cup \{(3a+2)p+3q-4, \ p>0, \ 2-k \le q \le 0\} \\ &\cup \{(3a+2)p+3q+4, \ p>0, \ 0 \le q \le k-2\} \\ &\cup \{(3a+2)p+3q, \ p>0, \ 2-k \le q \le k-2\}. \end{aligned}$$

If $\lambda(x) = \lambda(y) = 8$ (which occurs if and only if $k \ge 2$), then

$$\begin{array}{l} - y | \in \{3q, \ 0 \le q \le k-2\} \\ \cup \{3q+a-2, \ 2-k \le q \le 0\} \\ \cup \{(3a+2)p+a-2, \ p>0\} \\ \cup \{(3a+2)p-a+2, \ p>0\} \\ \cup \{(3a+2)p+3q, \ p>0, \ 2-k \le q < 0\} \\ \cup \{(3a+2)p+3q+a-2, \ p>0, \ 2-k \le q < 0\} \\ \cup \{(3a+2)p+3q, \ p>0, \ 0 < q \le k-2\} \\ \cup \{(3a+2)p+3q-a+2, \ p>0, \ 0 < q \le k-2\}. \end{array}$$

If $\lambda(x) = \lambda(y) = 9$ (which occurs if and only if $k \ge 2$), then

$$|x-y| \in \{3q, \ 0 \le q \le k-2\} \ \cup \ \{(3a+2)p+3q, \ p \ge 1, \ 2-k \le q \le k-2\}.$$

Therefore, in all these cases, $|x-y| \notin \{1, 2, a-1, a, a+1, a+2, 2a, 2a+1, 2a+2\}$, and thus λ is a 2-distance 9-coloring of $G(\{1, a, a+1\})$.

Case 3. $a = 3k + 2, k \ge 1$. Since the only elements divisible by 3 in $\{1, a, a+1\}^2$ are a+1 and 2a+2, the result follows by Theorem 4 (taking n = 3). This concludes the proof.

From Theorems 9 and 10, we thus get the following.

Corollary 11. For every integer $a, a \ge 3, a \not\equiv 2, 4 \pmod{7}$,

$$8 \le \chi_2(G(\{1, a, a+1\})) \le 9.$$

5. The CASE $D = \{1, \dots, m, a\}, 2 \le m < a$

We study in this section the 2-distance chromatic number of distance graphs G(D) with $D = \{1, \ldots, m, a\}, 2 \leq m < a$ (note that the case a = m + 1 is already solved by Proposition 7).

When $D = \{1, ..., m, a\}$, we have $\Delta(G(D)) = 2m + 2$ and

$$D^2 = \{1, 2, \dots, 2m\} \cup \{a - m, a - m + 1, \dots, a + m\} \cup \{2a\}.$$

First we prove the following.

Theorem 12. For all integers m and $a, 2 \le m < a$,

$$\chi_2(G(\{1,\ldots,m,a\})) = 2m + 3 = \Delta(G(\{1,\ldots,m,a\})) + 1$$

if and only if $a \equiv m+1 \pmod{2m+3}$ or $a \equiv m+2 \pmod{2m+3}$.

Proof. Since $\{1, \ldots, m, a\}^2 = \{1, \ldots, 2m\} \cup \{a - m, a - m + 1, \ldots, a + m\} \cup \{2a\}$, we have $d \not\equiv 0 \pmod{2m+3}$ for every $d \in \{1, \ldots, m, a\}^2$ if and only if $a \equiv m+1 \pmod{2m+3}$ or $a \equiv m+2 \pmod{2m+3}$, and thus, by Proposition 2 and Observation 6, $\chi_2(G(\{1, \ldots, m, a\})) = 2m+3$.

We now claim that every 2-distance (2m + 3)-coloring λ of $G(\{1, \ldots, m, a\})$ is necessarily (2m + 3)-periodic, that is $\lambda(x + 2m + 3) = \lambda(x)$ for every $x \in \mathbb{Z}$. To show that, it suffices to prove that any 2m+3 consecutive vertices $x, \ldots, x+2m+2$ must be assigned distinct colors. Assume to the contrary that this is not the case and, without loss of generality, let x = 0. Since vertices $0, 1, \ldots, 2m$ necessarily get distinct colors, we only have to consider two cases.

Case 1. Vertices $0, 1, \ldots, 2m + 1$ are colored with the colors $1, 2, \ldots, 2m + 1$ and 1, respectively. Note that vertices m - a and m + a are both adjacent to all vertices $0, 1, \ldots, 2m$. Hence,

$$\{\lambda(m-a), \lambda(m+a)\} = \{2m+2, 2m+3\},\$$

which implies

$$\{\lambda(m+1-a), \lambda(m+1+a)\} = \{2m+2, 2m+3\}$$

(more precisely, $\lambda(m+1-a) = 4m+5-\lambda(m-a)$ and $\lambda(m+1+a) = 4m+5-\lambda(m+a)$). This implies $\lambda(m+2-a) = \lambda(m+2+a) = 2$, a contradiction since d(m+2-a,m+2+a) = 2.

Case 2. Vertices $0, 1, \ldots, 2m + 2$ are colored with the colors $1, 2, \ldots, 2m + 2$ and 1, respectively. As in the previous case we have

$$\{\lambda(m-a), \lambda(m+a)\} = \{2m+2, 2m+3\},\$$

which implies

$$\{\lambda(m+1-a), \lambda(m+1+a)\} = \{1, 2m+3\}.$$

We thus get $\lambda(m+2-a) = \lambda(m+2+a) = 2$, again a contradiction.

Therefore, every 2-distance (2m + 3)-coloring λ of $G(\{1, \ldots, m, a\})$ is necessarily (2m + 3)-periodic, and thus $\chi_2(G(\{1, \ldots, m, a\})) = 2m + 3$ if and only if 2m + 3 does not divide any element of $\{1, 2, \ldots, 2m\} \cup \{a - m, a - m + 1, \ldots, a + m\} \cup \{2a\}$. This is clearly the case if and only if $a \equiv m + 1 \pmod{2m + 3}$ or $a \equiv m + 2 \pmod{2m + 3}$.

For other values of a, we propose the following general upper bound.

Theorem 13. For all integers m and a, $2 \le m < a$,

$$\chi_2(G(\{1,\ldots,m,a\})) \le 4m+2 = 2\Delta(G(\{1,\ldots,m,a\})) - 2.$$

Proof. First note that $\{1, \ldots, m, a\}^2 = \{1, \ldots, 2m\} \cup \{a - m, \ldots, a + m\} \cup \{2a\}$. Therefore, if 2m + 1 does not divide a, then the set $\{1, \ldots, m, a\}^2$ contains only one element e divisible by 2m + 1 (with $e \in \{a - m, \ldots, a + m\}$). In that case, the result follows by Theorem 4 (taking n = 2m + 1).

Suppose now that a = k(2m+1), with $k \ge 1$. Let λ be the (2a-m)-periodic mapping defined by the pattern

$$[12\cdots(2m+1)]^{k}[(2m+2)(2m+3)\cdots(4m+2)]^{k-1}(2m+2)(2m+3)\cdots(3m+2).$$

If $\lambda(x) = \lambda(y) = c$, $1 \le c \le 3m + 2$, then

$$\begin{aligned} |x-y| &\in \{q(2m+1), \ 0 \leq q \leq k-1\} \\ &\cup \{p(2a-m)+q(2m+1), \ p \geq 1, \ 1-k \leq q \leq k-1\}. \end{aligned}$$

If $\lambda(x) = \lambda(y) = c$, $3m + 3 \le c \le 4m + 2$ (which occurs if and only if $k \ge 2$), then

$$\begin{aligned} |x-y| &\in \{q(2m+1), \ 0 \leq q \leq k-2\} \\ &\cup \{p(2a-m)+q(2m+1), \ p \geq 1, \ 2-k \leq q \leq k-2\}. \end{aligned}$$

Therefore, in both cases, $|x - y| \notin \{1, \dots, m, a\}^2$, and thus λ is a 2-distance (4m + 2)-coloring of $G(\{1, \dots, m, a\})$. This concludes the proof.

From Theorems 12 and 13, we thus get the following.

Corollary 14. For all integers m and $a, 2 \le m < a, a \not\equiv m+1, m+2 \pmod{2m+3}$,

$$2m + 4 \le \chi_2(G(\{1, \dots, m, a\})) \le 4m + 2.$$

6. Discussion

In this paper, we studied 2-distance colorings of several types of distance graphs. In each case, we characterized those distance graphs that admit an optimal 2distance coloring, that is distance graphs G(D) with $\chi_2(G(D)) = \Delta(G(D)) + 1$. We also provided general upper bounds for the 2-distance chromatic number of the considered graphs. Note here that all our results can be extended to a larger class of integer distance graphs, thanks to Proposition 1, by multiplying all the elements of the set D by the same constant k > 1.

We leave as open problems the question of completely determining the 2distance chromatic number of distance graphs G(D) when $D = \{1, a, a + 1\}$, $a \ge 3$, or $D = \{1, \ldots, m, a\}, 2 \le m < a$.

Considering other types of sets D would certainly be also an interesting direction for future research.

Acknowledgement

Most of this work has been done while the first author was visiting LaBRI, thanks to a grant from University of Sciences and Technology Houari Boumediene (USTHB). The second author was partially supported by the Cluster of excellence CPU, from the Investments for the future Programme IdEx Bordeaux (ANR-10-IDEX-03-02).

References

- G. Agnarsson and M.M. Halldórsson, Coloring powers of planar graphs, SIAM J. Discrete Math. 16 (2003) 651–662. doi:10.1137/S0895480100367950
- M. Bonamy, B. Lévêque and A. Pinlou, 2-distance coloring of sparse graphs, J. Graph Theory 77 (2014) 190–218. doi:10.1002/jgt.21782
- [3] M. Bonamy, B. Lévêque and A. Pinlou, Graphs with maximum degree Δ ≥ 17 and maximum average degree less than 3 are list 2-distance (Δ + 2)-colorable, Discrete Math. **317** (2014) 19–32. doi:10.1016/j.disc.2013.10.022
- [4] O.V. Borodin and A.O. Ivanova, 2-distance (Δ + 2)-coloring of planar graphs with girth six and Δ ≥ 18, Discrete Math. **309** (2009) 6496–6502. doi:10.1016/j.disc.2009.06.029

- T. Calamoneri, The L(h,k)-labelling problem: An updated survey and annotated bibliography, Comput. J. 54 (2011) 1344-1371. doi:10.1093/comjnl/bxr037
- [6] Z. Dvořák, D. Král, P. Nejedlý and R. Škrekovski, Coloring squares of planar graphs with girth six, European J. Combin. 29 (2008) 838–849. doi:10.1016/j.ejc.2007.11.005
- [7] A. Kemnitz and H. Kolberg, *Coloring of integer distance graphs*, Discrete Math. 191 (1998) 113–123. doi:10.1016/S0012-365X(98)00099-5
- [8] F. Kramer and H. Kramer, Un probleme de coloration des sommets d'un graphe, C. R. Acad. Sci. Paris A 268 (1969) 46–48.
- F. Kramer and H. Kramer, A survey on the distance-colouring of graphs, Discrete Math. 308 (2008) 422–426. doi:10.1016/j.disc.2006.11.059
- [10] K.-W. Lih and W.-F. Wang, *Coloring the square of an outerplanar graph*, Taiwanese J. Math. **10** (2006) 1015–1023. doi:10.11650/twjm/1500403890
- D.D.-F. Liu, From rainbow to the lonely runner: A survey on coloring parameters of distance graphs, Taiwanese J. Math. 12 (2008) 851–871. doi:10.11650/twjm/1500404981
- [12] M. Voigt, Über die chromatische Zahl einer speziellen Klasse unendlicher Graphen, Ph.D. Thesis (Techn. Univ. Ilmenau, 1992).
- [13] H. Walther, Uber eine spezielle Klasse unendlicher Graphen, in: Graphentheorie, K. Wagner and R. Bodendiek (Ed(s)), (Bibl. Inst., Mannheim, 1990) 2, 268–295.
- [14] W.-F. Wang and K.-W. Lih, Labeling planar graphs with conditions on girth and distance two, SIAM J. Discrete Math. 17 (2003) 264-275. doi:10.1137/S0895480101390448
- [15] G. Wegner, Graphs with given diameter and a coloring problem (Technical Report, University of Dortmund, 1977). doi:10.17877/DE290R-16496
- [16] X. Zhu, Circular chromatic number of distance graphs with distance sets of cardinality 3, J. Graph Theory 41 (2002) 195–207. doi:10.1002/jgt.10062

Received 5 December 2016 Revised 12 November 2017 Accepted 13 November 2017