

TOTAL COLORINGS OF EMBEDDED GRAPHS WITH NO 3-CYCLES ADJACENT TO 4-CYCLES

BING WANG

Department of Mathematics
Zaozhuang University, Shandong, 277160, China

e-mail: hellobingzi@163.com.

JIAN-LIANG WU

School of Mathematics
Shandong University, Jinan, 250100, China

AND

LIN SUN

School of Mathematics
Shandong University, Jinan, 250100, China
Department of Mathematics
Changji University, Changji, 831100, China

Abstract

A *total- k -coloring* of a graph G is a coloring of $V \cup E$ using k colors such that no two adjacent or incident elements receive the same color. The *total chromatic number* $\chi''(G)$ of G is the smallest integer k such that G has a total- k -coloring. Let G be a graph embedded in a surface of Euler characteristic $\varepsilon \geq 0$. If G contains no 3-cycles adjacent to 4-cycles, that is, no 3-cycle has a common edge with a 4-cycle, then $\chi''(G) \leq \max\{8, \Delta + 1\}$.

Keywords: total coloring, embedded graph, cycle.

2010 Mathematics Subject Classification: 05C15.

1. INTRODUCTION

All graphs considered in this paper are simple, finite and undirected, and we follow [2] for the terminologies and notations not defined here. Let G be a graph.

We use $V(G)$, $E(G)$, $\Delta(G)$ and $\delta(G)$ (or simply V , E , Δ and δ) to denote the vertex set, the edge set, the maximum degree and the minimum degree of G , respectively. A *total- k -coloring* of a graph G is a coloring of $V \cup E$ using k colors such that no two adjacent or incident elements receive the same color. The *total chromatic number* $\chi''(G)$ of G is the smallest integer k such that G has a total- k -coloring. Clearly, $\chi''(G) \geq \Delta + 1$. Behzad [1] and Vizing [18] posed independently the following famous conjecture, which is known as the Total Coloring Conjecture (TCC).

Conjecture. *For any graph G , $\chi''(G) \leq \Delta + 2$.*

This conjecture was confirmed for all graphs with $\Delta \leq 3$ independently by Vijayaditya and Rosenfeld in 1971, and in [13, 14], Kostochka proved that if $4 \leq \Delta \leq 5$, then $\chi''(G) \leq \Delta + 2$. Later, Kostochka [15] renewed the proof for $\Delta = 5$. We summary these result to the following lemma.

Lemma 1. *Let G be a graph with $\Delta(G) \leq 5$. Then $\chi''(G) \leq 7$.*

But for planar graphs, the famous conjecture was first proved by Borodin [4] for $\Delta \geq 11$ and then for $\Delta \geq 9$ [3], which was extended to $\Delta \geq 8$ by Jensen and Toft [9] and to $\Delta \geq 7$ by Sanders and Zhao [17]. So the only open case is $\Delta = 6$.

Interestingly, planar graphs with high maximum degree allow a stronger assertion, that is, every planar graph with high maximum degree Δ has a total- $(\Delta + 1)$ -coloring. This result was first established in [4] for $\Delta \geq 16$, which was extended to $\Delta \geq 14$ [3], $\Delta \geq 12$ [5], $\Delta \geq 11$ [6], $\Delta \geq 10$ [25] and finally $\Delta \geq 9$ [10]. However, for $\Delta \in \{4, 5, 6, 7, 8\}$, it is not known if the assertion still holds true. Such a study has attracted a considerable amount of attention. Recently, Shen *et al.* [11] proved that if G is a planar graph with $\Delta = 8$ and G contains no chordal 5-cycles or no chordal 6-cycles, then $\chi''(G) = \Delta + 1$. Wang and Wu [19] proved that if G is a planar graph with $\Delta \geq 7$ and every vertex is incident with at most one triangle, then $\chi''(G) = \Delta + 1$. Wang and Wu [20] proved that if G is a planar graph with $\Delta \geq 7$ with no 4-cycles, then $\chi''(G) = \Delta + 1$ (later, it is extended to $\Delta \geq 6$ by Shen and Wang [12]). Chang *et al.* [7] proved that if G is a planar graph with $\Delta \geq 7$ and every vertex v has an integer $k_v \in \{3, 4, 5, 6\}$, such that v is not in any k_v -cycle, then $\chi''(G) = \Delta + 1$.

Let G be a graph embedded in a surface of Euler characteristic ε , where *surfaces* in this paper are compact, connected 2-dimensional manifolds without boundary. All embeddings considered in this paper are *2-cell embeddings*. Wu and Wang [24] proved that if $\varepsilon < 0$ and $\Delta(G) \geq \sqrt{25 - 24\varepsilon} + 10$, then $\chi'_{list}(G) = \Delta(G)$ and $\chi''_{list}(G) = \Delta(G) + 1$, which extends a result of Borodin, Kostochka and Woodall in [5]. They also proved that $\chi''(G) = \Delta(G) + 1$ if $\varepsilon \geq 0$, $\Delta(G) \geq 9$ and no two triangles have a common edge, or if $\varepsilon \geq 0$, $\Delta(G) \geq 8$ and no two triangles have a common vertex. Wang *et al.* [22] proved that if $\varepsilon \geq 0$ and $\Delta(G) \geq 7$,

then $\chi''(G) \leq \Delta + 2$. Wang *et al.* [23] proved that if $\varepsilon \geq 0$ and $\Delta \geq 9$, then $\chi''(G) = \Delta + 1$. In this paper, we shall prove the following result.

Theorem 2. *Let G be a graph embedded in a surface of Euler characteristic $\varepsilon \geq 0$. If G contains no 3-cycles adjacent to 4-cycles, then $\chi''(G) \leq \max\{8, \Delta(G)+1\}$.*

The theorem shows that if a graph G can be embedded in a surface of Euler characteristic $\varepsilon \geq 0$, and contains no 3-cycles adjacent to 4-cycles, and $\Delta \geq 7$, then $\chi''(G) = \Delta + 1$.

2. PROOF OF THEOREM 2

We will introduce some more notations and definitions here for convenience. Let $G = (V, E, F)$ be an embedded graph, where F is the face set of G . For a vertex $v \in V$, let $N(v)$ denote the set of vertices adjacent to v , and let $d(v) = |N(v)|$ denote the degree of v , and for a face f , the degree of a face f , denoted by $d(f)$, is the number of edges incident with it, where each cut-edge is counted twice. A k -vertex, a k^+ -vertex or a k^- -vertex is a vertex of degree k , at least k or at most k , respectively. Similarly, A k -face, a k^+ -face is a face of degree k or at least k , respectively. Let $n_t(v)$ be the number of t -vertices adjacent to a vertex v , and $f_k(v)$ the number of k -faces incident with v . Especially, let $f_3(v) = t$. Let $v_1, v_2, \dots, v_d$ be neighbors of v in an anticlockwise order. Let f_i be face incident with v , v_i and v_{i+1} , for all i such that $i \in \{1, 2, \dots, d\}$. Note that all the subscripts in the paper are taken modulo d . For convenience, $(d_1, d_2, \dots, d_n)$ denotes a cycle (or a face) whose boundary vertices are of degree $d_1, d_2, \dots, d_n$ in the anticlockwise order. Specially, (i, j^+, k^+) -face is a 3-face uvw such that $d(u) = i \leq j \leq d(v) \leq k \leq d(w)$.

Proof of Theorem 2. Let $m = \max\{7, \Delta\}$ and $G = (V, E, F)$ be a minimal counterexample to Theorem 2 with $|V| + |E|$ as small as possible. Then every proper subgraph of G has a total- $(m+1)$ -coloring, but G itself does not. First we show some known properties of G .

- (a) Every 3-cycle is not adjacent to a 4^- -face. It follows that $f_3(v) \leq \left\lfloor \frac{d(v)}{2} \right\rfloor$ for any $v \in V(G)$.
- (b) For any edge $uv \in E(G)$, if $\min\{d(u), d(v)\} \leq \left\lfloor \frac{m}{2} \right\rfloor$, then $d(u) + d(v) \geq m + 2$. So all neighbors of any 2-vertex are 7^+ -vertices and all neighbors of any 3-vertex are 6^+ -vertices (see [20]).
- (c) The subgraph G_2 of G induced by all edges incident with 2-vertices is a forest. So for any component of G_2 , we root it at a 7^+ -vertex. Then every 2-vertex has exactly one *parent* and exactly one *child* (see [3, 6]).

- (d) Each 3-face of G is not incident with two 4^- -vertices (see [16]).
 (e) If v is a vertex of G with $n_2(v) \geq 1$, then $n_{4^+}(v) \geq 1$ (see [7]).

Lemma 3 [21]. *Suppose v is a d -vertex of G with $d \geq 5$. Let $v_1, \dots, v_d$ be the neighbors of v and $f_1, \dots, f_d$ be the faces incident with v in clockwise order, where f_i is incident with v_i and v_{i+1} , $i = 1, 2, \dots, d$. Note that eventually v_1 and v_{d+1} is the same vertex. Then there does not exist an integer i ($2 \leq i \leq d$) such that $d(v_1) = d(v_i) = 2$, $d(v_k) = 3$ ($2 \leq k \leq i - 1$) and $d(f_t) = 4$ ($1 \leq t \leq i - 1$).*

Lemma 4. *G contains no subgraph isomorphic to one of the configurations in Figure 1, where the vertices marked by $\bullet$ have no other neighbors in G .*

Proof. The proof that G contains no subgraph isomorphic to one of the configurations in Figure 1(1)–(4) can be found in [8]. It remains to prove that G has no configurations depicted in Figure 1(5)–(13).

By the minimality of G , every proper subgraph of G has a total- $(m + 1)$ -coloring φ with the color set $C = \{1, 2, \dots, m + 1\}$. Erase the colors on all 3^- -vertices. Let $C(v) = \{\varphi(uv) : u \in N(v)\} \cup \{\varphi(v)\}$.

Suppose that G contains a configuration depicted in Figure 1(5). Then $G' = G - vv_6$ has a total-8-coloring φ . If $\varphi(v_6x_5) \in C(v)$ or $\varphi(v_6x_6) \in C(v)$, then the forbidden colors for vv_6 is at most 7, so vv_6 can be properly colored. By recoloring the erased vertices, we obtain a total-8-coloring of G , a contradiction. So we can assume that $\varphi(v_6x_5) \notin C(v)$ and $\varphi(v_6x_6) \notin C(v)$. Without loss of generality, assume that $\varphi(v) = 6$, $\varphi(v_6x_5) = 7$, $\varphi(v_6x_6) = 8$, and $\varphi(vv_j) = j$ for $j \in \{1, 2, \dots, 5\}$. Then we recolor v with 7 or 8, and color vv_6 with 6. By recoloring the erased vertices, we obtain a total-8-coloring of G , a contradiction.

Suppose that G contains a configuration depicted in Figure 1(6)–(13). Then $G' = G - vv_7$ has a total-8-coloring φ . If $\varphi(v_7x_7) \in C(v)$, then the forbidden colors for vv_7 is at most 7, so vv_7 can be properly colored. By recoloring the erased vertices, we obtain a total-8-coloring of G , a contradiction. So we can assume that $\varphi(vv_7) \notin C(v)$. Without loss of generality, assume that $\varphi(v) = 8$, $\varphi(v_7x_7) = 7$, and $\varphi(vv_j) = j$ for $j \in \{1, 2, \dots, 6\}$. Thus, for each 3^- -vertex v_k ($1 \leq k \leq 7$), there is an edge incident with v_k colored 7, otherwise we can recolor vv_k with 7, and color vv_7 with k to obtain a total-8-coloring of G , a contradiction.

For each 4-vertex v_i ($1 \leq i \leq 6$), suppose its adjacent vertices are v, x_{i-1}, x_i, x_j . If $\varphi(v_i) \neq 7$ ($1 \leq i \leq 6$), then recolor v with 7, and color vv_7 with 8. By recoloring the erased vertices, we obtain a total-8-coloring of G , a contradiction. Otherwise, there is at least one 4-vertex colored with 7. Suppose v is adjacent to only one 4-vertex v_i colored with 7. If $|C(v_i)| < 8$, then we recolor v_i with a color in $C \setminus C(v_i)$, recolor v with 7, and color vv_7 with 8. Otherwise, $|C(v_i)| = 8$. If $i \notin \{\varphi(x_{i-1}), \varphi(x_i), \varphi(x_j)\}$, then we recolor v_i with i , recolor vv_i with 7, and color




Figure 1. Reducible configurations.

vv_7 with i . Otherwise, $i \in \{\varphi(x_{i-1}), \varphi(x_i), \varphi(x_j)\}$. Without loss of generality, $\varphi(x_i) = i$, then $8 \notin \{C(v_i) \setminus \varphi(v)\}$. Therefore, we recolor v_i with 8, recolor v with 7, and color vv_7 with 8. Finally, we recolor the erased vertices, we obtain a total-8-coloring of G , a contradiction. Otherwise, v is adjacent to two or three 4-vertices colored with 7, then we take the same operations as above, respectively. Thus we can also obtain a total-8-coloring of G , a contradiction. ■

Let $G = (V, E, F)$ be a graph which is embedded in a surface of nonnegative Euler characteristic. By Euler's formula $|V| - |E| + |F| = \varepsilon$, we have

$$\sum_{v \in V} (2d(v) - 6) + \sum_{f \in F} (d(f) - 6) = -6(|V| - |E| + |F|) = -6\varepsilon \leq 0.$$

Now we define the initial charge function $ch(x)$ of $x \in V \cup F$ to be $ch(v) = 2d(v) - 6$ if $v \in V$ and $ch(f) = d(f) - 6$ if $f \in F$. It follows that $\sum_{x \in V \cup F} ch(x) \leq 0$. Now we design appropriate discharging rules and redistribute weights accordingly. Note that any discharging procedure preserves the total charge of G . If we can define suitable discharging rules to charge the initial charge function ch to the final charge function ch' on $V \cup F$ such that $\sum_{x \in V \cup F} ch'(x) > 0$, then we get an obvious contradiction.

Our discharging rules are defined as follows.

R1. Every 2-vertex receives $\frac{3}{2}$ from its child and $\frac{1}{2}$ from its parent.

R2. Let f be a 3-face. If f is incident with a 3^- -vertex, then it gets $\frac{3}{2}$ from each of its incident 6^+ -vertices. If f is incident with a 4-vertex, then it gets $\frac{1}{2}$ from the 4-vertex and gets $\frac{5}{4}$ from each of its incident 5^+ -vertices. If f is not incident with any 4^- -vertex, then it gets 1 from each of its incident 5^+ -vertices.

R3. Let f be a 4-face. If f is incident with two 3^- -vertices, then it gets 1 from each of its two incident 6^+ -vertices. If f is incident with only one 3^- -vertex and one 4-vertex, then it gets $\frac{1}{2}$ from the incident 4-vertex and gets $\frac{3}{4}$ from each of its two incident 6^+ -vertices. If f is incident with only one 3^- -vertex and no 4-vertex, then it gets $\frac{2}{3}$ from each of its incident 5^+ -vertices. If f is not incident with any 3^- -vertex, then it gets $\frac{1}{2}$ from each of its incident vertices.

R4. Every 5-face gets $\frac{1}{3}$ from each of its incident 4^+ -vertices.

First, we begin to check $ch'(x) \geq 0$ for all $x \in V \cup F$. By our discharging rules, it is easy to check that $ch'(f) \geq 0$ for all $f \in F$ and $ch'(v) \geq 0$ for all 2-vertices $v \in V$. If $d(v) = 3$, then $ch'(v) = ch(v) = 0$. So it suffices to check that $ch'(v) \geq 0$ for all 4^+ -vertices G .

Let v be a 4^+ -vertex of G . If $d(v) = 4$, then v sends at most $\frac{1}{2}$ to each of its incident faces by R2 and R3, and it follows that $ch'(v) \geq ch(v) - \frac{1}{2} \times 4 = 0$. Suppose $d(v) = 5$. Then v sends at most $\frac{5}{4}$ to each of its incident 3-faces by

R2, at most $\frac{2}{3}$ to each of its incident 4^+ -faces by R3, at most $\frac{1}{3}$ to each of its incident 5-faces by R4. By (a), $f_3(v) \leq 2$. If $f_3(v) = 2$, then v is incident with at least three 5^+ -faces, that is, $f_{5^+}(v) \geq 3$, and it follows that $ch'(v) \geq ch(v) - \frac{5}{4} \times 2 - \frac{1}{3} \times 3 = \frac{1}{2} > 0$. If $f_3(v) = 1$, then $f_{5^+}(v) \geq 2$ and $f_{4^+}(v) \leq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{5}{4} - \frac{1}{3} \times 2 - \frac{2}{3} \times 2 = \frac{3}{4} > 0$. If $f_3(v) = 0$, then $ch'(v) \geq ch(v) - \frac{2}{3} \times 5 = \frac{2}{3} > 0$. Suppose $d(v) = 6$. Then $f_3(v) \leq 3$ and v sends at most $\frac{3}{2}$ to each of its incident 3-faces by R2, at most 1 to each of its incident 4^+ -faces by R3, at most $\frac{1}{3}$ to each of its incident 5-faces by R4. Thus, if $1 \leq f_3(v) \leq 3$, then by the similar argument as above, we have $ch'(v) \geq ch(v) - \max\{\frac{3}{2} \times 3 + \frac{1}{3} \times 3, \frac{3}{2} \times 2 + \frac{1}{3} \times 3 + 1, \frac{3}{2} + \frac{1}{3} \times 2 + 1 \times 3\} = \frac{1}{2} > 0$. Otherwise, v is adjacent to at least one 4^+ -vertex or incident with at least one 5^+ -face by Lemma 4, configuration (5). If v is adjacent to at least one 4^+ -vertex, then $ch'(v) \geq ch(v) - 2 \times \frac{3}{4} - 4 = \frac{1}{2} > 0$ by R3. If v is incident with at least one 5^+ -face, then $ch'(v) \geq ch(v) - \frac{1}{3} - 5 = \frac{2}{3} > 0$ by R4.

Suppose $d(v) = d \geq 7$. Let $N(v) = \{v_1, v_2, \dots, v_d\}$ and $f_1, f_2, \dots, f_d$ be faces incident with v in the clockwise order, where f_i is incident with v_i and v_{i+1} , for $i \in \{1, 2, \dots, d\}$, where all the subscripts here are taken modulo d . If $n_2(v) \geq 1$, then v sends at most $\frac{n_2(v)+2}{2}$ to all its adjacent 2-vertices by R1, at most $\frac{3}{2}$ to each of its incident 3-faces by R2, at most 1 to each of its incident 4^+ -faces by R3, at most $\frac{1}{3}$ to each of its incident 5-faces by R4.

Lemma 5. Suppose that $d(v_i) = d(v_k) = 2$ and $d(v_j) \geq 3$ for all $j = i + 1, \dots, k - 1$. If $f_i, f_{i+1}, \dots, f_{k-1}$ are 4^+ -faces, then v sends at most $\frac{3}{2} + (k - 3)$ (in total) to $f_i, f_{i+1}, \dots, f_{k-1}$.


Proof. By Lemma 3, $\max\{d(v_{i+1}), \dots, d(v_{k-1})\} \geq 4$ or $\max\{d(f_1), \dots, d(f_{k-1})\} \geq 5$. If $\max\{d(v_{i+1}), \dots, d(v_{k-1})\} \geq 4$, then v sends at most $2 \times \frac{3}{4} + (k - 1 - 2)$ to $f_i, \dots, f_{k-1}$ by R2. If $\max\{d(f_1), \dots, d(f_{k-1})\} \geq 5$, then v sends at most $\frac{1}{3} + (k - 1 - 1)$ to $f_i, \dots, f_{k-1}$ by R3 and R4. Since $2 \times \frac{3}{4} > 1 + \frac{1}{3}$, v sends at most $\frac{3}{2} + (k - 3)$ to $f_i, f_{i+1}, \dots, f_{k-1}$. ■

Case 1. $\Delta(G) = 7$. Let v be a 7-vertex. Then $ch(v) = 2 \times 7 - 6 = 8$. We consider the following cases.

Subcase 1.1. $n_2(v) = 6$. By (c), any 2-vertex is not incident with a 4-face. Moreover, $t = f_3(v) = 0$ and $f_{6^+}(v) \geq 5$ by Lemma 4, configurations (1) and (4). So $ch'(v) \geq ch(v) - \frac{6+2}{2} - 2 = 2 > 0$.

Subcase 1.2. $n_2(v) = 5$. Then $t \leq 1$ by Lemma 4, configurations (1) and (4). If $t = 0$, then $f_{6^+}(v) \geq 3$ and $f_{4^+}(v) \leq 4$ by Lemma 4, configuration (4), and it follows that $ch'(v) \geq ch(v) - \frac{5+2}{2} - 4 \times 1 = \frac{1}{2} > 0$. Otherwise, $f_{6^+}(v) \geq 4$ and $f_{4^+}(v) \leq 2$. Thus $ch'(v) \geq ch(v) - \frac{5+2}{2} - \frac{3}{2} - 2 \times 1 = 1 > 0$.

Subcase 1.3. $n_2(v) = 4$. There are four possible configurations as shown in Figure 2.


Figure 2. $n_2(v) = 4$.

For Figure 2(a), $t \leq 1$ and $f_{6+}(v) \geq 3$ by Lemma 4, configurations (1) and (4). If $t = 1$, then $f_{5+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{4+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 1 = \frac{11}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{4+2}{2} - \frac{3}{2} - 2 = \frac{3}{2} > 0$ by Lemma 5.

For Figure 2(b) and (c), $t \leq 1$ and $f_{6+}(v) \geq 2$ by Lemma 4, configurations (1) and (4). If $t = 1$, then $f_{5+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{4+2}{2} - \frac{3}{2} - 3 \times \frac{1}{3} - \frac{3}{2} = \frac{4}{3} > 0$ by Lemma 5. Otherwise, $ch'(v) \geq ch(v) - \frac{4+2}{2} - \frac{3}{2} - \frac{3}{2} - 1 = 1 > 0$ by Lemma 5.

For Figure 2(d), $t = 0$ and $f_{6+}(v) \geq 1$ by Lemma 4, configurations (1) and (4). Then $ch'(v) \geq ch(v) - \frac{4+2}{2} - 3 \times \frac{3}{2} = \frac{1}{2} > 0$ by Lemma 5.

Subcase 1.4. $n_2(v) = 3$. There are four possible configurations as shown in Figure 3.

Figure 3. $n_2(v) = 3$.

For Figure 3(a), $t \leq 2$ and $f_{6+}(v) \geq 2$ by Lemma 4, configurations (1) and (4). If $t = 2$, then $f_{5+}(v) \geq 3$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - 2 \times \frac{3}{2} - 3 \times \frac{1}{3} = \frac{3}{2} > 0$. If $t = 1$, then $f_{5+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 2 = \frac{4}{3} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 3 = 1 > 0$ by Lemma 5.

For Figure 3(b), $t \leq 1$ and $f_{6+}(v) \geq 1$ by Lemma 4, configurations (1) and (4). If $t = 1$, then $f_{5+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 1 - \frac{3}{2} = \frac{5}{6} > 0$ by Lemma 5. Otherwise, $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - \frac{3}{2} - 2 = \frac{1}{2} > 0$ by Lemma 5.

For Figure 3(c), $t \leq 2$ and $f_{6+}(v) \geq 1$ by Lemma 4, configurations (1) and (4). If $t = 2$, then $f_{5+}(v) \geq 4$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - 2 \times$

$\frac{3}{2} - 4 \times \frac{1}{3} = \frac{7}{6} > 0$. If $t = 1$, then $f_{5^+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - \frac{3}{2} - 1 = \frac{5}{6} > 0$ by Lemma 5. Otherwise, $ch'(v) \geq ch(v) - \frac{3+2}{2} - 2 \times (\frac{3}{2} + 1) = \frac{1}{2} > 0$ by Lemma 5.

For Figure 3(d), $t \leq 1$ by Lemma 4, configuration (1). If $t = 1$, then $f_{5^+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 2 \times \frac{3}{2} = \frac{1}{3} > 0$ by Lemma 5. Otherwise, $t = 0$. If v is adjacent to at least four 4^+ -vertices, then $ch'(v) \geq ch(v) - \frac{3+2}{2} - \frac{3}{2} - \frac{3}{2} - 3 \times \frac{3}{4} = \frac{1}{4} > 0$ by Lemma 5. Otherwise, v is adjacent to at least one 5^+ -vertex or incident with at least one 5^+ -face by Lemma 4, configuration (6), and Lemma 3. If v is adjacent to at least one 5^+ -vertex, then $ch'(v) \geq ch(v) - \frac{3+2}{2} - \max\{2 \times \frac{2}{3} + 1 + 2 \times \frac{3}{2}, \frac{3}{2} + 1 + 2 \times \frac{2}{3} + \frac{3}{2}\} = \frac{1}{6} > 0$ by R3 and Lemma 5. If v is incident with at least one 5^+ -face, then $ch'(v) \geq ch(v) - \frac{3+2}{2} - \max\{\frac{1}{3} + 2 + 2 \times \frac{3}{2}, \frac{3}{2} + 1 + \frac{1}{3} + \frac{3}{2} + 1\} = \frac{1}{6} > 0$ by R4 and Lemma 5.

Subcase 1.5. $n_2(v) = 2$. There are three possible configurations as shown in Figure 4.




Figure 4. $n_2(v) = 2$.

For Figure 4(a), we have that the face f_6 is a 5^+ -face, and it follows that f_6 receives at most $\frac{1}{3}$ from v by R4. So $ch(v) - \frac{2+2}{2} - \frac{1}{3} = 8 - 2 - \frac{1}{3} = \frac{17}{3}$. Moreover, $t \leq 2$ by Lemma 4, configuration (1), and (a). If $1 \leq t \leq 2$, then $f_{5^+}(v) \geq (t+1)$ (except f_6), and it follows that $ch'(v) \geq \frac{17}{3} - t \times \frac{3}{2} - (t+1) \times \frac{1}{3} - (7-1-t-t-1) = \frac{2+t}{6} > 0$. Otherwise, $ch'(v) \geq \frac{17}{3} - (\frac{3}{2} + 4) = \frac{1}{6} > 0$.

For Figure 4(b), $t \leq 2$ by Lemma 4, configuration (1), and (a). If $t = 2$, then $f_{5^+}(v) \geq 3$, it follows that $ch'(v) \geq ch(v) - \frac{2+2}{2} - 2 \times \frac{3}{2} - 3 \times \frac{1}{3} - \frac{3}{2} = \frac{1}{2} > 0$. If $t = 1$, then $f_{5^+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - \frac{1}{3} \times 2 - 2 \times 1 - \frac{3}{2} = \frac{1}{3} > 0$. Suppose $t = 0$. If v is adjacent to at least three 4^+ -vertices, then $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - 3 \times \frac{3}{4} - 2 = \frac{1}{4} > 0$ by Lemma 5. Otherwise, v is adjacent to at least one 5^+ -vertex or incident with at least one 5^+ -face by Lemma 4, configurations (7)–(8). By the same argument as above, we can obtain $ch'(v) \geq ch(v) - \frac{2+2}{2} - \max\{\frac{3}{2} + 3 + 2 \times \frac{2}{3}, \frac{3}{2} + 4 + \frac{1}{3}\} = \frac{1}{6} > 0$.

For Figure 4(c), $t \leq 2$ by Lemma 4, configuration (1), and (a). If $t = 2$, then $f_{5^+}(v) \geq 4$, it follows that $ch'(v) \geq ch(v) - \frac{2+2}{2} - 2 \times \frac{3}{2} - 4 \times \frac{1}{3} - 1 = \frac{2}{3} > 0$. If $t = 1$, then $f_{5^+}(v) \geq 2$, and it follows that $ch'(v) \geq ch(v) - \frac{2+2}{2} -$

$\max \left\{ \frac{3}{2} + \frac{1}{3} \times 2 + \frac{3}{2} + 2, \frac{3}{2} + \frac{1}{3} \times 2 + 1 + \frac{3}{2} + 1 \right\} = \frac{1}{3} > 0$. Suppose $t = 0$. If v is adjacent to at least three 4^+ -vertices, then $ch'(v) \geq ch(v) - \frac{2+2}{2} - \frac{3}{2} - 3 \times \frac{3}{4} - 2 = \frac{1}{4} > 0$ by Lemma 5. Otherwise, v is adjacent to at least one 5^+ -vertex or incident with at least one 5^+ -face by Lemma 4, configurations (9)–(10). By the same argument as above, we can obtain $ch'(v) \geq ch(v) - \frac{2+2}{2} - \max \left\{ \frac{3}{2} + 3 + 2 \times \frac{2}{3}, \frac{3}{2} + 4 + \frac{1}{3} \right\} = \frac{1}{6} > 0$.

Subcase 1.6. $n_2(v) = 1$. Note that $n_{4^+}(v) \geq 1$ by (e) and $t \leq 3$ by (a). Suppose $t = 0$. If v is adjacent to at least two 4^+ -vertices, then $ch'(v) \geq ch(v) - \frac{1+2}{2} - 3 \times \frac{3}{4} - 4 = \frac{1}{4} > 0$. Otherwise, v is adjacent to at least one 5^+ -vertex or incident with at least one 5^+ -face by Lemma 4, configurations (11)–(13), then $ch'(v) \geq ch(v) - \frac{1+2}{2} - 2 \times \frac{2}{3} - 5 = \frac{1}{6} > 0$ by R3 or $ch'(v) \geq ch(v) - \frac{1+2}{2} - \frac{1}{3} - 6 = \frac{1}{6} > 0$ by R4. Suppose $1 \leq t \leq 3$. If v is incident with a $(2, 7, 7)$ -face, then the other face incident with the 2-vertex is a 6^+ -face. Moreover, the other 3-faces incident with v are $(4, 5^+, 5^+)$ -faces by Lemma 4, configuration (3), and v is incident with at least t 5^+ -faces. Then we can obtain that $ch'(v) \geq ch(v) - \frac{1+2}{2} - \frac{3}{2} - (t-1) \times \frac{5}{4} - \frac{1}{3} \times t - (7-1-t-t) = \frac{3+5t}{12} > 0$, where v sends at most $\frac{5}{4}$ to each of its incident $(4, 5^+, 5^+)$ -faces by R2. Otherwise, 2-vertex is not incident with any 3-face. Then v is incident with at least $(t+1)$ 5^+ -faces, and it follows that $ch'(v) \geq ch(v) - \frac{1+2}{2} - t \times \frac{3}{2} - \frac{1}{3} \times (t+1) - (7-t-t-1) = \frac{1+t}{6} > 0$.

Subcase 1.7. $n_2(v) = 0$. Note that $t \leq 3$ by (a). If $t = 0$, then $ch'(v) \geq ch(v) - 7 \times 1 = 1 > 0$. Otherwise, by the same argument as above, we can obtain that $ch'(v) \geq ch(v) - t \times \frac{3}{2} - \frac{1}{3} \times (t+1) - (7-t-t-1) = \frac{10+t}{6} > 0$.

Case 2. $\Delta(G) \geq 8$. In [23], Theorem 1 was established for $\Delta \geq 9$. So we assume that $\Delta = 8$. Then $ch(v) = 2 \times 8 - 6 = 10$. By the same argument as above, we consider the following cases.

Subcase 2.1. $n_2(v) = 7$. Then $t = 0$ and $f_{6^+}(v) \geq 6$ by Lemma 4, configurations (1) and (4). So $ch'(v) \geq ch(v) - \frac{7+2}{2} - 2 = \frac{7}{2} > 0$.

Subcase 2.2. $n_2(v) = 6$. Then $t \leq 1$ and $f_{6^+}(v) \geq 4$ by Lemma 4, configurations (1) and (4). If $t = 0$, then $ch'(v) \geq ch(v) - \frac{6+2}{2} - 4 = 2 > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{6+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} = \frac{23}{6} > 0$.

Subcase 2.3. $n_2(v) = 5$. Then $t \leq 1$ and $f_{6^+}(v) \geq 2$ by Lemma 4, configurations (1) and (4). If $t = 0$, then $ch'(v) \geq ch(v) - \frac{5+2}{2} - 6 = \frac{1}{2} > 0$. Otherwise, $f_{5^+}(v) \geq 2$. Thus $ch'(v) \geq ch(v) - \frac{5+2}{2} - \frac{3}{2} - 2 \times \frac{1}{3} - 3 = \frac{4}{3} > 0$.

Subcase 2.4. $n_2(v) = 4$. Then $t \leq 2$ by Lemma 4, configuration (1). If $1 \geq t \geq 2$, then $f_{6^+}(v) \geq 1$ and $f_{5^+}(v) \geq t+1$, and it follows that $ch'(v) \geq ch(v) - \frac{4+2}{2} - t \times \frac{3}{2} - (t+1) \times \frac{1}{3} - (8-1-t-t-1) = \frac{4+t}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{4+2}{2} - \max \left\{ \frac{3}{2} + 3, \frac{3}{2} \times 2 + 2, \frac{3}{2} \times 3 + 1, \frac{3}{2} \times 4 \right\} = 1 > 0$.

Subcase 2.5. $n_2(v) = 3$. Then $t \leq 2$ by Lemma 4 configuration (1). If $1 \geq t \geq 2$, then $f_{5^+}(v) \geq t+1$, and it follows that $ch'(v) \geq ch(v) - \frac{3+2}{2} - t \times \frac{3}{2} -$

$(t+1) \times \frac{1}{3} - (8-t-t-1) = \frac{1+t}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{3+t}{2} - \max\{\frac{3}{2} + 4, \frac{3}{2} \times 2 + 3, \frac{3}{2} \times 3 + 2\} = 1 > 0$.

Subcase 2.6. $n_2(v) = 2$. Then $t \leq 2$ by Lemma 4 configuration (1). If $1 \geq t \geq 2$, then $f_{5+}(v) \geq t+1$, and it follows that $ch'(v) \geq ch(v) - \frac{2+t}{2} - t \times \frac{3}{2} - (t+1) \times \frac{1}{3} - (8-t-t-1) = \frac{4+t}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{2+t}{2} - \max\{\frac{3}{2} + 5, \frac{3}{2} \times 2 + 4\} = 1 > 0$.

Subcase 2.7. $n_2(v) = 1$. Then $t \leq 4$ by (a). If $1 \geq t \geq 4$, then $f_{5+}(v) \geq t$, and it follows that $ch'(v) \geq ch(v) - \frac{1+t}{2} - t \times \frac{3}{2} - t \times \frac{1}{3} - (8-t-t) = \frac{3+t}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - \frac{1+t}{2} - 8 = \frac{1}{2} > 0$.

Subcase 2.8. $n_2(v) = 0$. Then $t \leq 4$ by (a). If $1 \geq t \geq 4$, then $f_{5+}(v) \geq t$, and it follows that $ch'(v) \geq ch(v) - t \times \frac{3}{2} - t \times \frac{1}{3} - (8-t-t) = \frac{12+t}{6} > 0$. Otherwise, $ch'(v) \geq ch(v) - 8 = 2 > 0$.

Finally, according to the above argument, we have checked $ch'(x) \geq 0$ for all $x \in V \cup F$ and $ch'(x) > 0$ for any 5^+ -vertex $x \in V$. By Lemma 1, we have $\Delta(G) \geq 6$. So $\sum_{x \in V \cup F} ch'(x) > 0$. Hence we complete the proof of Theorem 2.

REFERENCES

- [1] M. Behzad, *Graphs and Their Chromatic Numbers* (Ph.D. Thesis, Michigan State University, 1965).
- [2] J.A. Bondy and U.S.R. Murty, *Graph Theory with Applications* (Macmillan Press Ltd., London, 1976).
- [3] O.V. Borodin, *On the total coloring of planar graphs*, J. Reine Angew. Math. **394** (1989) 180–185.
- [4] O.V. Borodin, *Coupled colourings of graphs on a plane*, Metody Diskret. Anal. **45** (1987) 21–27, in Russian.
- [5] O.V. Borodin, A.V. Kostochka and D.R. Woodall, *List edge and list total colourings of multigraphs*, J. Combin. Theory Ser. B **71** (1997) 184–204.
doi:10.1006/jctb.1997.1780
- [6] O.V. Borodin A.V. Kostochka and D.R. Woodall, *Total colorings of planar graphs with large maximum degree*, J. Graph Theory **26** (1997) 53–59.
doi:10.1002/(SICI)1097-0118(199709)26:1<53::AID-JGT6>3.0.CO;2-G
- [7] G.J. Chang, J. Hou and N. Roussel, *Local condition for planar graphs of maximum degree 7 to be 8-totally colorable*, Discrete Appl. Math. **159** (2011) 760–768.
doi:10.1016/j.dam.2011.01.001
- [8] D. Du, L. Shen and Y. Wang, *Planar graphs with maximum degree 8 and without adjacent triangles are 9-totally-colorable*, Discrete Appl. Math. **157** (2009) 2778–2784.
doi:10.1016/j.dam.2009.02.011
- [9] T.R. Jensen and B. Toft, *Graph Coloring Problems* (Wiley Interscience, 1995).

- [10] L. Kowalik, J.-S. Sereni and R. Škrekovski, *Total-colorings of plane graphs with maximum degree nine*, SIAM J. Discrete Math. **22** (2008) 1462–1479.
doi:10.1137/070688389
- [11] L. Shen and Y.Q. Wang, *Total colorings of planar graphs with maximum degree at least 8*, Sci. China Ser A: Math. **52** (2009) 1733–1742.
doi:10.1007/s11425-008-0155-3
- [12] L. Shen and Y. Wang, *On the 7 total colorability of planar graphs with maximum degree 6 and without 4-cycles*, Graphs Combin. **25** (2009) 401–407.
doi:10.1007/s00373-009-0843-y
- [13] A.V. Kostochka, *The total coloring of a multigraph with maximal degree 4*, Discrete Math. **17** (1977) 161–163.
doi:10.1016/0012-365X(77)90146-7
- [14] A.V. Kostochka, *An analogue of Shannon's estimate for complete colorings*, Metody Diskret. Anal. **30** (1977) 13–22, in Russian.
- [15] A.V. Kostochka, *The total chromatic number of any multigraph with maximum degree five is at most seven*, Discrete Math. **162** (1996) 199–214.
doi:10.1016/0012-365X(95)00286-6
- [16] B. Liu, J.F. Hou, J.L. Wu and G.Z. Liu, *Total colorings and list total colorings of planar graphs without intersecting 4-cycles*, Discrete Math. **309** (2009) 6035–6043.
doi:10.1016/j.disc.2009.05.006
- [17] D.P. Sanders and Y. Zhao, *On total 9-coloring planar graphs of maximum degree seven*, J. Graph Theory **31** (1999) 67–73.
doi:10.1002/(SICI)1097-0118(199905)31:1<67::AID-JGT6>3.0.CO;2-C
- [18] V.G. Vizing, *Some unsolved problems in graph theory*, Uspekhi Mat. Nauk **23** (1968) 117–134, in Russian.
- [19] B. Wang and J.-L. Wu, *Total coloring of planar graphs with maximum degree seven*, Inform. Process. Lett. **111** (2011) 1019–1021.
doi:10.1016/j.ipl.2011.07.012
- [20] P. Wang and J.-L. Wu, *A note on total colorings of planar graphs without 4-cycles*, Discuss. Math. Graph Theory **24** (2004) 125–135.
doi:10.7151/dmgt.1219
- [21] H.J. Wang, L.D. Wu, W.L. Wu, P.M. Pardalos and J.L. Wu, *Minimum total coloring of planar graph*, J. Global Optim. **60** (2014) 777–791.
doi:10.1007/s10898-013-0138-y
- [22] H.J. Wang, B. Liu, J.L. Wu and G.Z. Liu, *Total coloring of embedded graphs with maximum degree at least seven*, Theoret. Comput. Sci. **518** (2014) 1–9.
doi:10.1016/j.tcs.2013.04.030
- [23] H.J. Wang, B. Liu, J.L. Wu and B. Wang, *Total coloring of graphs embedded in surfaces of nonnegative Euler characteristic*, Sci. China Math. **57** (2014) 211–220.
doi:10.1007/s11425-013-4576-2

- [24] J.L. Wu and P. Wang, *List-edge and list-total colorings of graphs embedded on hyperbolic surfaces*, Discrete Math. **308** (2008) 6210–6215.
doi:10.1016/j.disc.2007.11.044
- [25] W.F. Wang, *Total chromatic number of planar graphs with maximum degree ten*, J. Graph Theory **54** (2007) 91–102.
doi:10.1002/jgt.20195

Received 30 September 2016

Revised 30 March 2017

Accepted 30 March 2017