

ON SELKOW'S BOUND ON THE INDEPENDENCE NUMBER OF GRAPHS

JOCHEN HARANT

AND

SAMUEL MOHR

Ilmenau University of Technology
Department of Mathematics
Ilmenau, Germany

e-mail: jochen.harant@tu-ilmenau.de
samuel.mohr@tu-ilmenau.de

Abstract

For a graph G with vertex set $V(G)$ and independence number $\alpha(G)$, Selkow [A *Probabilistic lower bound on the independence number of graphs*, *Discrete Math.* 132 (1994) 363–365] established the famous lower bound $\sum_{v \in V(G)} \frac{1}{d(v)+1} \left(1 + \max \left\{ \frac{d(v)}{d(v)+1} - \sum_{u \in N(v)} \frac{1}{d(u)+1}, 0 \right\} \right)$ on $\alpha(G)$, where $N(v)$ and $d(v) = |N(v)|$ denote the neighborhood and the degree of a vertex $v \in V(G)$, respectively. However, Selkow's original proof of this result is incorrect. We give a new probabilistic proof of Selkow's bound here.

Keywords: graph, independence number.

2010 Mathematics Subject Classification: 05C69.

We consider a *finite, simple, and undirected* graph G with *vertex set* $V(G)$. Let $N_G(v)$ and $d_G(v) = |N_G(v)|$ denote the *neighborhood* and the *degree* of $v \in V(G)$, respectively. A set of vertices $I \subseteq V(G)$ is *independent* if no two vertices in I are adjacent. The *independence number* $\alpha(G)$ of G is the maximum cardinality of an independent set of G .

The independence number is one of the most fundamental and well-studied graph parameters. In view of its computational hardness, various bounds on the independence number have been proposed. The classical lower bound $CW(G) = \sum_{v \in V(G)} \frac{1}{d_G(v)+1}$ on $\alpha(G)$ is due to Caro [2] and Wei [4]. It is natural to ask whether improvements of $\alpha(G) \geq CW(G)$ are possible if more information about G is known than just its degrees. The following result takes not only the degree of every vertex but also the degree distribution in its neighborhood into account.

Theorem 1 (Selkow, [3]).

$$\alpha(G) \geq CW(G) + \sum_{v \in V(G)} \frac{1}{d_G(v) + 1} \max \left\{ \frac{d_G(v)}{d_G(v) + 1} - \sum_{u \in N_G(v)} \frac{1}{d_G(u) + 1}, 0 \right\}.$$

Unfortunately, Selkow’s original proof of Theorem 1 is not correct. To our best knowledge, this has not been discovered earlier, and we are not aware of an alternative, correct proof. In order to extract the problematic part of Selkow’s argument, let us repeat his proof.

For an event A and a random variable X let $P(A)$ and $E(X)$ denote the probability of A and the expectation of X , respectively. First, a uniformly chosen ordering $<$ of $V(G)$ is considered. Obviously, the set

$$I_1 = \{v \in V(G) \mid u \in N_G(v) \Rightarrow v < u\}$$

is independent and it is easy to show that $E(|I_1|) = CW(G)$ (e. g. see [1]).

Next, let the graph H (depending on the ordering $<$) be obtained from G by removing $I_1 \cup \bigcup_{x \in I_1} N_G(x)$ and consider the set

$$I_2 = \{v \in V(H) \mid u \in N_H(v) \Rightarrow v < u\}.$$

It follows that

$$\alpha(G) \geq E(|I_1| + |I_2|) = E(|I_1|) + E(|I_2|) = CW(G) + \sum_{v \in V(G)} P(v \in I_2),$$

since $I_1 \cap I_2 = \emptyset$ and $I_1 \cup I_2$ is an independent set of G . To finish the proof of Theorem 1 in [3], the inequality

$$P(v \in I_2) \geq \frac{1}{d_G(v) + 1} \max \left\{ \frac{d_G(v)}{d_G(v) + 1} - \sum_{u \in N_G(v)} \frac{1}{d_G(u) + 1}, 0 \right\}$$

for all $v \in V(G)$ ([3], page 364, lines 14–17) is used. This turns out to be false in the following general sense.

For every $\varepsilon > 0$, there is a graph G and a vertex $v \in V(G)$, such that

$$0 < P(v \in I_2) < \varepsilon \cdot \frac{1}{d_G(v) + 1} \left(\frac{d_G(v)}{d_G(v) + 1} - \sum_{u \in N_G(v)} \frac{1}{d_G(u) + 1} \right).$$

To see this, let n be a large positive integer. Consider an arbitrary graph F on $n - 3$ vertices and let the graph G on n vertices be obtained by adding three new vertices v, w, x and the edges vw, wx , and xy for all $y \in V(F)$. For

an arbitrary ordering $<$ of $V(G)$, $v \in I_2$ if and only if $x \in I_1$, $x < w < v$, and $x < y$ for all $y \in V(F)$. It is easy to see that there are exactly $\binom{n-1}{2}(n-3)!$ such orderings with the property $v \in I_2$, thus, $P(v \in I_2) = \frac{\binom{n-1}{2}(n-3)!}{n!} = \frac{1}{2n}$, however, $\frac{1}{d_G(v)+1} \left(\frac{d_G(v)}{d_G(v)+1} - \sum_{u \in N_G(v)} \frac{1}{d_G(u)+1} \right) = \frac{1}{2} \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{1}{12}$.

Eventually, we present a new probabilistic proof of Theorem 1.

Proof of Theorem 1. As in Selkow's proof, consider a uniformly chosen ordering $<$ of $V(G)$, the set $I_1 = \{v \in V(G) \mid u \in N_G(v) \Rightarrow v < u\}$, and the graph H induced by $V(G) \setminus (I_1 \cup \bigcup_{x \in I_1} N_G(x))$.

With $f(v) = 0$ if $v \notin V(H)$ and $f(v) = \frac{1}{d_G(v)+1}$ if $v \in V(H)$, it follows

$$\begin{aligned} \alpha(G) &\geq E(|I_1| + \alpha(H)) = E(|I_1|) + E(\alpha(H)) \geq E(|I_1|) + E(CW(H)) \\ &= CW(G) + E\left(\sum_{v \in V(H)} \frac{1}{d_H(v)+1}\right) \geq CW(G) + E\left(\sum_{v \in V(H)} \frac{1}{d_G(v)+1}\right) \\ &= CW(G) + E\left(\sum_{v \in V(G)} f(v)\right) = CW(G) + \sum_{v \in V(G)} \frac{1}{d_G(v)+1} P(v \in V(H)). \end{aligned}$$

Using $P(v \in V(H)) \geq 0$ and $P(v \notin V(H)) = P\left(v \in I_1 \vee \left(\bigvee_{u \in N_G(v)} u \in I_1\right)\right) \leq P(v \in I_1) + \sum_{u \in N_G(v)} P(u \in I_1) = \frac{1}{d_G(v)+1} + \sum_{u \in N_G(v)} \frac{1}{d_G(u)+1}$, Theorem 1 is proved. ■

REFERENCES

- [1] N. Alon and J.H. Spencer, *The Probabilistic Method* (Wiley, New York, 1992).
- [2] Y. Caro, *New Results on the Independence Number* (Technical Report, Tel-Aviv University, 1979).
- [3] S.M. Selkow, *A Probabilistic lower bound on the independence number of graphs*, *Discrete Math.* **132** (1994) 363–365. doi:10.1016/0012-365X(93)00102-B
- [4] V.K. Wei, *A Lower Bound on the Stability Number of a Simple Graph* (Technical Memorandum, TM 81 - 11217 - 9, Bell laboratories, 1981).

Received 20 November 2017
Accepted 18 December 2017