Discussiones Mathematicae
Graph Theory 39 (2019) 815–828
doi:10.7151/dmgt.2101

ON DECOMPOSING THE COMPLETE SYMMETRIC DIGRAPH INTO ORIENTATIONS OF $K_4 - e$

Ryan C. Bunge[1] Brian D. Darrow, Jr.[2] Toni M. Dubczuk[1]

Saad I. El-Zanati[1] Hanson H. Hao[3] Gregory L. Keller[4]

Genevieve A. Newkirk[1] and Dan P. Roberts[5]

[1]*Illinois State University, Normal, IL 61790-4520, USA*
[2]*Southern Connecticut State University, New Haven, CT 06515, USA*
[3]*Illinois Math and Science Academy, Aurora, IL 60506, USA*
[4]*Bloomington High School, Bloomington, IL 61701, USA*
[5]*Illinois Wesleyan University, Bloomington, IL 61701, USA*

e-mail: saad@ilstu.edu

Abstract

Let D be any of the 10 digraphs obtained by orienting the edges of $K_4 - e$. We establish necessary and sufficient conditions for the existence of a (K_n^*, D)-design for 8 of these digraphs. Partial results as well as some nonexistence results are established for the remaining 2 digraphs.

Keywords: digraph decompositions, orientations of $K_4 - e$.

2010 Mathematics Subject Classification: 05C51, 05B30.

1. Introduction

Let $\mathbb{Z}_m$ denote the group of integers modulo m. For integers a and b with $a \leq b$, let $[a, b] = \{a, a+1, \ldots, b\}$. For a graph (or digraph) H, let $V(H)$ and $E(H)$ denote the vertex set of H and the edge (or arc) set of H, respectively. The *order* and the *size* of a graph (or digraph) H are $|V(H)|$ and $|E(H)|$, respectively.

We denote the complete multipartite graph with parts of sizes a_i for $1 \leq i \leq m$ by $K_{a_1,a_2,\ldots,a_m}$. If $a_i = a$ for all $i \in \{1, 2, \ldots, m\}$, then we use the notation $K_{m\times a}$. Additionally, $K_{m\times a,b}$ denotes the complete multipartite graph with m parts of size a and one part of size b.

Let H be a graph and let $\mathcal{G}$ be a set of subgraphs of H. We will refer to a graph $G \in \mathcal{G}$ as a *G-block*. A *$\mathcal{G}$-decomposition* of H is a set $\Delta = \{G_1, G_2, \ldots, G_r\}$

of pairwise edge-disjoint subgraphs of H such that for every $i \in [1, r]$, $G_i \cong G$ for some $G \in \mathcal{G}$ and such that $E(H) = \bigcup_{i=1}^{r} E(G_i)$. Of particular importance is when $\mathcal{G} = \{G\}$, in which case we write "G-decomposition of H" instead of "$\{G\}$-decomposition of H." A G-decomposition of H is also known as an (H, G)-*design*. The set of all n for which K_n admits a G-decomposition is called the *spectrum of* G. The spectrum has been determined for many classes of graphs, including for all graphs on at most 4 vertices [4] and all graphs on 5 vertices (see [3] and [10]). We direct the reader to [2] and [5] for recent surveys on graph decompositions.

By *blowing up* the vertices of a graph G by some positive integer t, we mean replacing every vertex of G with t independent vertices and replacing every edge in G by a $K_{t,t}$. For example, assume we have a $(K_{x\times 2}, K_3)$-design. After blowing up the vertices of $K_{x\times 2}$ by 5, our corresponding $(K_{x\times 2}, K_3)$-design becomes a $(K_{x\times 10}, K_{3\times 5})$-design.

Similar concepts to the ones defined above for undirected graphs can be defined for digraphs. First, we introduce additional notation. For an undirected graph G, let G^* denote the digraph obtained from G by replacing each edge $\{u, v\} \in E(G)$ with the arcs (u, v) and (v, u). Thus K_n^*, the *complete symmetric digraph of order* n, is the digraph on n vertices with the arcs (u, v) and (v, u) between every pair of distinct vertices u and v.

Let D and H be digraphs such that D is a subgraph of H. The *reverse orientation of* D, denoted $\mathrm{Rev}(D)$, is the digraph with vertex set $V(D)$ and arc set $\{(v, u) : (u, v) \in E(D)\}$. A D-*decomposition* of H is a set $\Delta = \{D_1, D_2, \ldots, D_r\}$ of pairwise arc-disjoint subgraphs of H each of which is isomorphic to D and such that $E(H) = \bigcup_{i=1}^{r} E(D_i)$. As with the undirected case, a D-decomposition of H is also known as an (H, D)-*design*, and the set of all n for which K_n^* admits a D-decomposition is called the *spectrum of* D. Furthermore, we say D is *self-complementary in* H if D is isomorphic to the digraph with arc set $E(H) \setminus E(D)$. That is, D is self-complementary in H if H has size $2 \cdot |E(D)|$ and there exists an (H, D)-design.

The spectra for several digraphs of small order have been determined. This includes the spectra for all digraphs on at most 3 vertices [11] and all bipartite digraphs on 4 vertices with up to 5 arcs [7].

In this paper, we extend the known results on small digraphs by determining the spectrum for 8 of the 10 digraphs obtained by orienting the edges of $K_4 - e$, the graph obtained from removing a single edge from K_4. Some nonexistence results are proven for the remaining 2 such digraphs. We use the naming convention found in *An Atlas of Graphs* [13] by Read and Wilson. The digraphs under investigation are shown in Figures 1 and 2 with a key that denotes a labeled copy for each of the 10 digraphs of interest. For example, D75$[w, x, y, z]$ refers to the digraph with vertex set $\{w, x, y, z\}$ and arc set $\big\{(w, x), (w, y), (w, z), (x, y), (z, y)\big\}$.



Figure 1. The four orientations of $K_4 - e$ that are self-complementary in $(K_4 - e)^*$.



Figure 2. The six orientations of $K_4 - e$ that are not self-complementary in $(K_4 - e)^*$, shown paired with their reverse orientations.

Note that 6 of the digraphs of interest in this paper occur in pairs with respect to their reverse orientations (see Figure 2). Namely, D69 $\cong$ Rev(D90), D74 $\cong$ Rev(D89), and D85 $\cong$ Rev(D105). The remaining 4 digraphs of interest (see Figure 1) are isomorphic to their reverse orientations, e.g., D75 $\cong$ Rev(D75), which is shown in the proceeding section (see Lemma 4) to imply that these 4 digraphs are self-complementary in $(K_4 - e)^*$.

2. Some Basic Results

The necessary conditions for a digraph D to decompose K_n^* include

(A) $|V(D)| \leq n$,

(B) $|E(D)|$ divides $n(n-1)$, and
(C) both $\gcd\{\text{outdegree}(v) : v \in V(D)\}$ and $\gcd\{\text{indegree}(v) : v \in V(D)\}$ divide $n-1$.

Applying these necessary conditions to the 10 digraphs under consideration, we obtain the following necessary condition: For $D \in \{\text{D69}, \text{D74}, \text{D75}, \text{D85}, \text{D87}, \text{D89}, \text{D90}, \text{D101}, \text{D102}, \text{D105}\}$, a (K_n^*, D)-design exists only if $n \equiv 0$ or $1 \pmod 5$.

The following observation was stated in [6].

Observation 1. *Let D and H be digraphs. A D-decomposition of H exists if and only if a* $\mathrm{Rev}(D)$*-decomposition of* $\mathrm{Rev}(H)$ *exists.*

The fact that $K_n^* \cong \mathrm{Rev}(K_n^*)$ leads to our next observation, also stated in [6].

Observation 2. *Let D be a digraph. A (K_n^*, D)-design exists if and only if a $(K_n^*, \mathrm{Rev}(D))$-design exists.*

2.1. Results for self-complementary digraphs

We note that the existence of $(K_4 - e)$-decompositions of complete multigraphs (i.e., the spectrum of index λ) is known [12]. However, we present here the following theorem reduced to what is useful for characterizing the spectra of our 4 self-complementary digraphs.

Theorem 3 [4]**.** *There exists a $(K_4 - e)$-decomposition of K_n if and only if $n \equiv 0$ or $1 \pmod 5$ and $n \geq 6$.*

Since there does not exist a $(K_4 - e)$-decomposition of K_5, we must address decompositions of K_5^* (see Section 3). To make use of the known spectrum of $K_4 - e$, we present the following.

Lemma 4. *Let D be an orientation of a graph G. Then D is isomorphic to* $\mathrm{Rev}(D)$ *if and only if D is self-complementary in G^*.*

Proof. Let D' be the digraph with vertex set $V(G^*)$ and arc set $E(G^*) \setminus E(D)$. Note that $E(D') = \big\{(v,u) : (u,v) \in E(D)\big\}$, which implies that D' is both the reverse orientation of D and the complement of D in G^*. The result then follows. ■

Since there exists a (K, G)-design if and only if a (K^*, G^*)-design exists, we arrive at the following corollary of the above lemma.

Corollary 5. *Let D be an orientation of a simple graph G such that D is self-complementary in G^*. If there exists a (K, G)-design, then there exists a (K^*, D)-design.*

In light of Corollary 5, we can combine Theorem 3 and Example 11 (see Section 3) to characterize the spectra of the digraphs that are self-complementary in $(K_4 - e)^*$, namely D75, D87, D101, and D102 (as seen in Figure 1).

Theorem 6. *Let $D \in \{\mathrm{D75}, \mathrm{D87}, \mathrm{D101}, \mathrm{D102}\}$. There exists a (K_n^*, D)-design if and only if $n \equiv 0$ or $1 \pmod 5$ and $n \geq 5$.*

2.2. Results for non-self-complementary digraphs

Our general constructions also use some basic results concerning decompositions of both complete graphs and complete multipartite graphs into complete graphs of orders 3 and 5. These are sometimes stated in the language of group divisible designs and/or pairwise balanced designs. Note that these background results concern graphs, as opposed to digraphs. Theorems 7, 8, and 9 can be found in the *Handbook of Combinatorial Designs* [8] (see [1] and [9]).

Theorem 7. *If n is odd, then a $\{K_3, K_5\}$-decomposition of K_n exists.*

Theorem 8. *The necessary and sufficient conditions for the existence of a K_3-decomposition of $K_{u\times m}$ are* (i) $u \geq 3$, (ii) $(u-1)m \equiv 0 \pmod 2$, *and* (iii) $u(u-1)m^2 \equiv 0 \pmod 6$.

Theorem 9. *If $u \geq 3$ and $u \equiv 0 \pmod 3$, then there exists a K_3-decomposition of $K_{u\times 2,4}$.*

Our general constructions further rely on the following direct result of blowing up the vertices in the graphs of a decomposition. This well-known building block is a special case of Wilson's Fundamental Construction.

Lemma 10. *Let m, r, s, t, u_1, $u_2, \ldots, u_m$ all be positive integers. If there exists a $\{K_r, K_s\}$-decomposition of $K_{u_1,u_2,\ldots,u_m}$, then there exists a $\{K_{r\times t}, K_{s\times t}\}$-decomposition of $K_{tu_1,tu_2,\ldots,tu_m}$. In particular, if there exists a $(K_{u_1,u_2,\ldots,u_m}, K_r)$-design, then there exists a $(K_{tu_1,tu_2,\ldots,tu_m}, K_{r\times t})$-design.*

3. Examples of Small Designs

We now turn our attention to the designs of small order which will be used for the general constructions.

Given a digraph represented by the notation $D[a, b, c, d]$ and some $i \in \mathbb{Z}_n$, we define $D[a, b, c, d] + i = D[a + i, b + i, c + i, d + i]$ where all addition is performed in $\mathbb{Z}_n$. By convention, define $\infty + 1 = \infty$.

Example 11. There exists a (K_5^*, D)-design for $D \in \{\mathrm{D74}, \mathrm{D75}, \mathrm{D85}, \mathrm{D87}, \mathrm{D89}, \mathrm{D101}, \mathrm{D102}, \mathrm{D105}\}$.

Let $V(K_5^*) = \mathbb{Z}_4 \cup \{\infty\}$.
A $(K_5^*, \mathrm{D74})$-design is given by $\{\mathrm{D74}[0,\infty,2,1] + i : i \in \mathbb{Z}_4\}$.
A $(K_5^*, \mathrm{D75})$-design is given by $\{\mathrm{D75}[0,\infty,3,1] + i : i \in \mathbb{Z}_4\}$.
A $(K_5^*, \mathrm{D85})$-design is given by $\{\mathrm{D85}[0,\infty,2,1] + i : i \in \mathbb{Z}_4\}$.
A $(K_5^*, \mathrm{D87})$-design is given by

$$\{\mathrm{D87}[0,\infty,1,2], \mathrm{D87}[0,3,2,\infty], \mathrm{D87}[3,\infty,2,1], \mathrm{D87}[3,0,1,\infty]\}.$$

A $(K_5^*, \mathrm{D101})$-design is given by $\{\mathrm{D101}[0,\infty,3,1] + i : i \in \mathbb{Z}_4\}$.
A $(K_5^*, \mathrm{D102})$-design is given by

$$\{\mathrm{D102}[0,\infty,1,2], \mathrm{D102}[1,3,0,\infty], \mathrm{D102}[2,\infty,3,0], \mathrm{D102}[3,1,2,\infty]\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 12. There exists a (K_6^*, D)-design for $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$.

Let $V(K_6^*) = \mathbb{Z}_6$.
A $(K_6^*, \mathrm{D69})$-design is given by $\{\mathrm{D69}[0,2,1,4] + i : i \in \mathbb{Z}_6\}$.
A $(K_6^*, \mathrm{D74})$-design is given by $\{\mathrm{D74}[0,5,1,3] + i : i \in \mathbb{Z}_6\}$.
A $(K_6^*, \mathrm{D85})$-design is given by $\{\mathrm{D85}[0,3,5,4] + i : i \in \mathbb{Z}_6\}$.
Applying Observation 2, we obtain the remaining designs.

Example 13. There exists a (K_{10}^*, D)-design for $D \in \{\mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D105}\}$.

Let $V(K_{10}^*) = \mathbb{Z}_9 \cup \{\infty\}$.
A $(K_{10}^*, \mathrm{D74})$-design is given by

$$\{\mathrm{D74}[0,\infty,4,6] + i : i \in \mathbb{Z}_9\} \cup \{\mathrm{D74}[0,8,1,4] + i : i \in \mathbb{Z}_9\}.$$

A $(K_{10}^*, \mathrm{D85})$-design is given by

$$\{\mathrm{D85}[0,\infty,4,7] + i : i \in \mathbb{Z}_9\} \cup \{\mathrm{D85}[0,6,1,2] + i : i \in \mathbb{Z}_9\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 14. There exists a (K_{11}^*, D)-design for $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$.

Let $V(K_{11}^*) = \mathbb{Z}_{11}$.
A $(K_{11}^*, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[0,2,7,5] + i : i \in \mathbb{Z}_{11}\} \cup \{\mathrm{D69}[0,1,4,3] + i : i \in \mathbb{Z}_{11}\}.$$

A $(K_{11}^*, \mathrm{D74})$-design is given by

$$\{\mathrm{D74}[0,7,1,3] + i : i \in \mathbb{Z}_{11}\} \cup \{\mathrm{D74}[0,4,10,8] + i : i \in \mathbb{Z}_{11}\}.$$

A $(K_{11}^*, \mathrm{D85})$-design is given by

$$\{\mathrm{D85}[0,6,10,7] + i : i \in \mathbb{Z}_{11}\} \cup \{\mathrm{D85}[0,9,8,2] + i : i \in \mathbb{Z}_{11}\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 15. There exists a (K_{20}^*, D)-design for $D \in \{\mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D105}\}$.

Let $V(K_{20}^*) = \mathbb{Z}_{19} \cup \{\infty\}$.
A $(K_{20}^*, \mathrm{D74})$-design is given by

$$\{\mathrm{D74}[0, \infty, 13, 2] + i : i \in \mathbb{Z}_{19}\} \cup \{\mathrm{D74}[0, 12, 1, 10] + i : i \in \mathbb{Z}_{19}\}$$
$$\cup \{\mathrm{D74}[0, 14, 16, 12] + i : i \in \mathbb{Z}_{19}\} \cup \{\mathrm{D74}[0, 15, 18, 13] + i : i \in \mathbb{Z}_{19}\}.$$

A $(K_{20}^*, \mathrm{D85})$-design is given by

$$\{\mathrm{D85}[0, \infty, 2, 16] + i : i \in \mathbb{Z}_{19}\} \cup \{\mathrm{D85}[0, 12, 1, 6] + i : i \in \mathbb{Z}_{19}\}$$
$$\cup \{\mathrm{D85}[0, 13, 17, 7] + i : i \in \mathbb{Z}_{19}\} \cup \{\mathrm{D85}[0, 3, 18, 10] + i : i \in \mathbb{Z}_{19}\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 16. There exists a (K_{21}^*, D)-design for $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$.

Let $V(K_{21}^*) = \mathbb{Z}_{21}$.
A $(K_{21}^*, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[0, 4, 1, 12] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D69}[0, 7, 2, 16] + i : i \in \mathbb{Z}_{21}\}$$
$$\cup \{\mathrm{D69}[0, 6, 19, 13] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D69}[0, 17, 20, 9] + i : i \in \mathbb{Z}_{21}\}.$$

A $(K_{21}^*, \mathrm{D74})$-design is given by

$$\{\mathrm{D74}[0, 13, 4, 14] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D74}[0, 2, 5, 20] + i : i \in \mathbb{Z}_{21}\}$$
$$\cup \{\mathrm{D74}[0, 18, 16, 1] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D74}[0, 8, 17, 7] + i : i \in \mathbb{Z}_{21}\}.$$

A $(K_{21}^*, \mathrm{D85})$-design is given by

$$\{\mathrm{D85}[0, 18, 1, 15] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D85}[0, 17, 2, 7] + i : i \in \mathbb{Z}_{21}\}$$
$$\cup \{\mathrm{D85}[0, 16, 19, 11] + i : i \in \mathbb{Z}_{21}\} \cup \{\mathrm{D85}[0, 12, 20, 9] + i : i \in \mathbb{Z}_{21}\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 17. There exists a (K_{25}^*, D)-design for $D \in \{\mathrm{D69}, \mathrm{D90}\}$.

Let $V(K_{25}^*) = \mathbb{Z}_5 \times \mathbb{Z}_5$. A $(K_{25}^*, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[(1, i), (0, 1 + i), (1, 2 + i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2, i), (0, 1 + i), (2, 2 + i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3, i), (0, 1 + i), (3, 2 + i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(0, 1 + i), (1, i), (4, i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(0, 2 + i), (1, i), (4, 3 + i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(1, 3 + i), (1, i), (0, 3 + i), (0, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2, 1 + i), (1, 1 + i), (2, 4 + i), (1, i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3, i), (1, 1 + i), (3, 3 + i), (1, i)] : i \in \mathbb{Z}_5\}$$

$$\cup \{\mathrm{D69}[(0,4+i),(2,1+i),(2,2+i),(1,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(1,1+i),(2,i),(4,2+i),(1,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(0,2+i),(2,2+i),(0,1+i),(2,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(1,i),(2,1+i),(3,1+i),(2,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3,3+i),(2,1+i),(4,i),(2,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3,4+i),(4,i),(1,2+i),(2,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(0,3+i),(4,i),(3,1+i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(0,4+i),(4,2+i),(4,1+i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(1,1+i),(4,i),(1,i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(1,4+i),(4,1+i),(4,3+i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2,4+i),(4,3+i),(1,2+i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2,1+i),(4,1+i),(2,i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2,3+i),(4,1+i),(4,4+i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(2,2+i),(4,4+i),(4,i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3,4+i),(4,4+i),(0,i),(3,i)] : i \in \mathbb{Z}_5\}$$
$$\cup \{\mathrm{D69}[(3,3+i),(4,1+i),(0,i),(4,i)] : i \in \mathbb{Z}_5\}.$$

Applying Observation 2, we obtain a $(K^*_{25}, \mathrm{D90})$-design.

Example 18. There exists a (K^*_{30}, D)-design for $D \in \{\mathrm{D69}, \mathrm{D90}\}$.

Let $V(K^*_{30}) = \mathbb{Z}_5 \times \mathbb{Z}_6$.
A $(K^*_{30}, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[(1,i),(0,1+i),(1,2+i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,4+i),(0,1+i),(2,i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,2+i),(0,1+i),(2,4+i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,i),(0,1+i),(3,2+i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,4+i),(0,1+i),(4,i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,1+i),(1,i),(4,2+i),(0,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,i),(1,1+i),(0,3+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,4+i),(2,i),(4,1+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,1+i),(2,i),(2,1+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,4+i),(2,i),(2,3+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,3+i),(2,i),(3,i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,5+i),(2,i),(3,4+i),(1,i)] : i \in \mathbb{Z}_6\}$$

$$\cup \{\mathrm{D69}[(2,4+i),(2,i),(3,1+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,2+i),(2,i),(4,i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(4,3+i),(2,i),(4,4+i),(1,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,i),(2,1+i),(0,2+i),(2,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,5+i),(3,i),(2,5+i),(2,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,i),(3,1+i),(0,4+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,2+i),(4,1+i),(4,i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(1,4+i),(4,i),(4,3+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,4+i),(4,1+i),(0,1+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,1+i),(4,3+i),(1,3+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,2+i),(4,3+i),(4,1+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,3+i),(4,3+i),(4,5+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,1+i),(4,2+i),(1,1+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,3+i),(4,5+i),(1,2+i),(3,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(0,5+i),(4,2+i),(0,4+i),(4,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(2,2+i),(4,1+i),(1,2+i),(4,i)] : i \in \mathbb{Z}_6\}$$
$$\cup \{\mathrm{D69}[(3,3+i),(4,1+i),(3,1+i),(4,i)] : i \in \mathbb{Z}_6\}.$$

Applying Observation 2, we obtain a $(K^*_{30}, \mathrm{D90})$-design.

Example 19. There exists a $(K^*_{3\times 5}, D)$-design for $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$.

Let $V(K^*_{3\times 5}) = \mathbb{Z}_{15}$ with vertex partition $\{V_i : i \in \mathbb{Z}_3\}$, where $V_i = \{j \in \mathbb{Z}_{15} : j \equiv i \pmod 3\}$.
A $(K^*_{3\times 5}, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[0,8,10,11] + i : i \in \mathbb{Z}_{15}\} \cup \{\mathrm{D69}[0,4,5,7] + i : i \in \mathbb{Z}_{15}\}.$$

A $(K^*_{3\times 5}, \mathrm{D74})$-design is given by

$$\{\mathrm{D74}[0,2,10,11] + i : i \in \mathbb{Z}_{15}\} \cup \{\mathrm{D74}[0,13,5,4] + i : i \in \mathbb{Z}_{15}\}.$$

A $(K^*_{3\times 5}, \mathrm{D85})$-design is given by

$$\{\mathrm{D85}[0,7,5,1] + i : i \in \mathbb{Z}_{15}\} \cup \{\mathrm{D85}[0,8,10,14] + i : i \in \mathbb{Z}_{15}\}.$$

Applying Observation 2, we obtain the remaining designs.

Example 20. There exists a $(K^*_{5\times 5}, D)$-design for $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$.

First, let $V(K^*_{5\times 5}) = \mathbb{Z}_{25}$ with vertex partition $\{V_i : i \in \mathbb{Z}_5\}$, where $V_i = \{j \in \mathbb{Z}_{25} : j \equiv i \pmod 5\}$.
A $(K^*_{5\times 5}, \mathrm{D69})$-design is given by

$$\{\mathrm{D69}[1,14,0,24] + i : i \in \mathbb{Z}_{25}\} \cup \{\mathrm{D69}[2,8,0,18] + i : i \in \mathbb{Z}_{25}\}$$
$$\cup \{\mathrm{D69}[3,12,0,22] + i : i \in \mathbb{Z}_{25}\} \cup \{\mathrm{D69}[4,11,0,21] + i : i \in \mathbb{Z}_{25}\}.$$

A $(K^*_{5\times 5}, \mathrm{D90})$-design follows from Observation 2.
Next, let $D \in \{\mathrm{D74}, \mathrm{D89}, \mathrm{D105}, \mathrm{D85}\}$. A $(K_{5\times 5}, K_5)$-design can be obtained by removing one parallel class from an affine plane of order 5. Thus, there exists a $(K^*_{5\times 5}, K^*_5)$-design. Since a (K^*_5, D)-design exists by Example 11, the desired $(K^*_{5\times 5}, D)$-design exists.

4. Main Results

We first show some nonexistence results for $(\mathrm{D69}, K^*_n)$- and $(\mathrm{D90}, K^*_n)$-designs. Interestingly for $n \equiv 0 \pmod 5$, these designs do not exist for $n \in \{5, 10, 15, 20\}$ (see Theorem 21) but do exist for $n \in \{25, 30\}$ (see Examples 17 and 18). By Wilson's Theorem [14], there exists an integer n_0 such that for all $n \geq n_0$ that satisfy the necessary conditions there exists both a $(\mathrm{D69}, K^*_n)$-design and a $(\mathrm{D90}, K^*_n)$-design. We conjecture that $n_0 = 25$ for this pair of digraphs.

Theorem 21. *There does not exist a* D69- *or* D90-*decomposition of* K^*_{5k} *for* $1 \leq k \leq 4$.

Proof. We prove by contradiction that a D69-decomposition of K^*_{5k} cannot exist. Note that by Observation 2, a D90-decomposition must also not exist.

Let Δ be a D69-decomposition of K^*_{5k}. Given a vertex $v \in V(K^*_{5k})$, let $n_w(v)$ denote the number of D69-blocks in Δ where vertex w in $\mathrm{D69}[w,x,y,z]$ is identified with vertex v. Define $n_x(v)$, $n_y(v)$, and $n_z(v)$ similarly. Thus, the following must hold:

$$1n_w(v) + 2n_x(v) + 0n_y(v) + 2n_z(v) = 5k - 1,$$
$$2n_w(v) + 0n_x(v) + 3n_y(v) + 0n_z(v) = 5k - 1.$$

Substituting $\bar{n}(v) = n_x(v) + n_z(v)$, the above equations can be parameterized as

$$n_w(v) = 5k - 1 - 2\bar{n}(v), \tag{1}$$
$$n_y(v) = -\tfrac{1}{3}(5k - 1 - 4\bar{n}(v)). \tag{2}$$

Since $\bar{n}(v)$, $n_w(v)$, and $n_y(v)$ must all be nonnegative integers, we have that

$$\left.\begin{array}{l} 0 \leq 5k - 1 - 2\bar{n}(v) \\ 0 \leq -\frac{1}{3}(5k - 1 - 4\bar{n}(v)) \end{array}\right\} \implies \tfrac{1}{4}(5k-1) \leq \bar{n}(v) \leq \tfrac{1}{2}(5k-1).$$

Furthermore, equation (2) implies that $5k - 1 - 4\bar{n}(v)$ must be a multiple of 3; hence, $k + \bar{n}(v) + 1 \equiv 0 \pmod 3$.

Next, consider the case when $k = 1$. The above conditions require that for every $v \in V(K_5^*)$, we have $1 \leq \bar{n}(v) \leq 2$ and $\bar{n}(v) \equiv 1 \pmod 3$. Thus, $\bar{n}(v)$ can only equal 1, and by equation (1), $n_w(v) = 2$ for every $v \in V(K_5^*)$. However, this would imply $|\Delta| = 10$, which is a contradiction (because $|\Delta| = 4$ when $k = 1$). Similarly if k is 2, 3, or 4, then $\bar{n}(v)$ can only equal 3, 5, or 7, respectively, which further yields only one value for $n_w(v)$: 3, 4, or 5, respectively, for every $v \in V(K_{5k}^*)$. However, this would imply $|\Delta|$ is a multiple of $5k$, which is a contradiction because

$$|\Delta| = \frac{|E(K_{5k}^*)|}{|E(\mathrm{D69})|} = \frac{5k(5k-1)}{5} = k(5k-1),$$

which is not divisible by $5k$. ■

Next we turn our attention to developing the general constructions needed to piece together the small designs presented in Section 3 and show sufficiency of the necessary conditions for the remaining four non-self-complementary digraphs.

Theorem 22. *Let $D \in \{\mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D105}\}$. If $n \equiv 0 \pmod 5$ with $n \geq 5$, then a (K_n^*, D)-design exists.*

Proof. Let $D \in \{\mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D105}\}$ and let $n \equiv 0$ or $5 \pmod{10}$.

Case 1. $n \equiv 0 \pmod{10}$. Let $n = 10x = 5(2x)$ for some positive integer x. When x is 1 or 2 the result follows from Examples 13 and 15, respectively, so we now consider when $x \geq 3$. Let $H_1, H_2, \ldots, H_x$ be disjoint sets of 2 vertices each.

Subcase 1a. $x \equiv 0$ or $1 \pmod 3$. Let $K_{x\times 2}$ have vertex partition $\{H_i : 1 \leq i \leq x\}$. By Theorem 8, a $(K_{x\times 2}, K_3)$-design exists. Therefore, by Lemma 10 a $(K_{x\times 10}, K_{3\times 5})$-design exists. Let H_i' be the set obtained from H_i after blowing up each vertex in $K_{x\times 2}$ by 5. Now consider K_n^* to have vertex set $\bigcup_{i=1}^{x} H_i'$ where each H_i' induces a K_{10}^*. Thus, K_n^* decomposes into copies of K_{10}^* and $K_{3\times 5}^*$. Since both a (K_{10}^*, D)-design and a $(K_{3\times 5}^*, D)$-design exist by Examples 13 and 19, respectively, we have our desired (K_n^*, D)-design.

Subcase 1b. $x \equiv 2 \pmod 3$. Let $H_0 = H_{x-1} \cup H_x$ and let $K_{(x-2)\times 2,4}$ have vertex partition $\{H_i : 0 \leq i \leq x-2\}$. By Theorem 9, a $(K_{(x-2)\times 2,4}, K_3)$-design exists. Therefore, by Lemma 10 a $(K_{(x-2)\times 10,20}, K_{3\times 5})$-design exists. Let H_i' be the set obtained from H_i after blowing up each vertex in $K_{(x-2)\times 2,4}$ by 5. Now consider K_n^* to have vertex set $\bigcup_{i=0}^{x-2} H_i'$ where H_0' induces a K_{20}^* and, for $1 \leq i \leq x-2$, each H_i' induces a K_{10}^*. Thus, K_n^* decomposes into copies of K_{10}^*, K_{20}^*, and $K_{3\times 5}^*$. Since a (K_{10}^*, D)-design, a (K_{20}^*, D)-design, and a $(K_{3\times 5}^*, D)$-design exist by Examples 13, 15, and 19, respectively, we have our desired (K_n^*, D)-design.

Case 2. $n \equiv 5 \pmod{10}$. Let $n = 10x + 5 = 5(2x + 1)$ for some positive integer x. Let $H_1, H_2, \ldots, H_{2x+1}$ be sets consisting of a single vertex each. By Theorem 7 a $\{K_3, K_5\}$-decomposition of K_{2x+1} exists. Therefore, by Lemma 10 a $\{K_{3\times5}, K_{5\times5}\}$-decomposition of $K_{(2x+1)\times5}$ exists. Let H'_i be the set obtained from H_i after blowing up each vertex in K_{2x+1} by 5. Now consider K^*_n to have vertex set $\bigcup_{i=1}^{2x+1} H'_i$ where each H'_i induces a K^*_5. Thus, K^*_n decomposes into copies of K^*_5, $K^*_{3\times5}$, and $K^*_{5\times5}$. Since a (K^*_5, D)-design, a $(K^*_{3\times5}, D)$-design, and a $(K^*_{5\times5}, D)$-design all exist by Examples 11, 19, and 20, respectively, we have our desired (K^*_n, D)-design. ■

Theorem 23. *Let* $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$. *If* $n \equiv 1 \pmod 5$ *with* $n \geq 6$, *then a* (K^*_n, D)*-design exists.*

Proof. Let $D \in \{\mathrm{D69}, \mathrm{D74}, \mathrm{D85}, \mathrm{D89}, \mathrm{D90}, \mathrm{D105}\}$ and let $n \equiv 1$ or $6 \pmod{10}$.

Case 1. $n \equiv 1 \pmod{10}$. Let $n = 10x+1 = 5(2x)+1$ for some positive integer x. When x is 1 or 2 the result follows from Examples 14 and 16, respectively, so we now consider when $x \geq 3$.

Subcase 1a. $x \equiv 0$ or $1 \pmod 3$. Here we can consider $V(K^*_n) = \left(\bigcup_{i=1}^{x} H'_i\right) \cup \{\infty\}$, where each H'_i is defined as in Subcase 1a of the proof of Theorem 22 with the modification that each $H'_i \cup \{\infty\}$ induces a K^*_{11}. Similarly to the proof of that Subcase 1a, the desired (K^*_n, D)-design can be constructed using (K^*_{11}, D)-designs—in place of (K^*_{10}, D)-designs—along with $(K^*_{3\times5}, D)$-designs, which exist by Examples 14 and 19, respectively.

Subcase 1b. $x \equiv 2 \pmod 3$. Here we can consider $V(K^*_n) = \left(\bigcup_{i=0}^{x-2} H'_i\right) \cup \{\infty\}$, where each H'_i is defined as in Subcase 1b of the proof of Theorem 22 with the modifications that $H'_0 \cup \{\infty\}$ induces a K^*_{21} and, for $1 \leq i \leq x-2$, each $H'_i \cup \{\infty\}$ induces a K^*_{11}. Similarly to the proof of that Subcase 1b, the desired (K^*_n, D)-design can be constructed using (K^*_{11}, D)-designs and a (K^*_{21}, D)-design—in place of (K^*_{10}, D)- and (K^*_{20}, D)-designs—along with $(K^*_{3\times5}, D)$-designs, which exist by Examples 14, 16, and 19, respectively.

Case 2. $n \equiv 6 \pmod{10}$. Here we can consider $V(K^*_n) = \left(\bigcup_{i=1}^{2x+1} H'_i\right) \cup \{\infty\}$, where each H'_i is defined as in Case 2 of the proof of Theorem 22 with the modification that $H'_i \cup \{\infty\}$ induces a K^*_6. Similarly to the proof of that Case 2, the desired (K^*_n, D)-design can be constructed by using (K^*_6, D)-designs—in place of (K^*_5, D)-designs—along with $(K^*_{3\times5}, D)$-designs and $(K^*_{5\times5}, D)$-designs, which exist by Examples 12, 19, and 20. ■

Our results from this section along with those in Theorem 6 are summarized in the following two main theorems.

Theorem 24. *Let $D \in \{\mathrm{D74}, \mathrm{D75}, \mathrm{D85}, \mathrm{D87}, \mathrm{D89}, \mathrm{D101}, \mathrm{D102}, \mathrm{D105}\}$. There exists a (K_n^*, D)-design if and only if $n \equiv 0$ or $1 \pmod 5$ and $n \geq 5$.*

Theorem 25. *Let $D \in \{\mathrm{D69}, \mathrm{D90}\}$. There exists a (K_n^*, D)-design if $n \equiv 1 \pmod 5$ and $n \geq 6$.*

Finally, we formally state our conjecture regarding the open results for the $\{\mathrm{D69}, \mathrm{D90}\}$ pair of digraphs.

Conjecture 26. *Let $D \in \{\mathrm{D69}, \mathrm{D90}\}$. There exists a (K_n^*, D)-design if $n \equiv 0 \pmod 5$ and $n \geq 25$.*

Acknowledgements

This research is supported by grant number A1359300 from the Division of Mathematical Sciences at the National Science Foundation. This work was done while the second, third, fifth, sixth and seventh authors were participants in *REU Site: Mathematics Research Experience for Pre-service and for In-service Teachers* at Illinois State University. The authors wish to thank Mariusz Meszka for finding the $(K_{25}^*, \mathrm{D69})$- and $(K_{30}^*, \mathrm{D69})$-designs that appear in Examples 17 and 18.

References

[1] R.J.R. Abel, F.E. Bennett and M. Greig, *PBD-Closure*, in: Handbook of Combinatorial Designs, 2nd Ed., C.J. Colbourn and J.H. Dinitz (Eds), (Chapman & Hall/CRC Press, Boca Raton, FL, 2007) 246–254.

[2] P. Adams, D. Bryant and M. Buchanan, *A survey on the existence of G-designs*, J. Combin. Des. **16** (2008) 373–410.
doi:10.1002/jcd.20170

[3] J.-C. Bermond, C. Huang, A. Rosa and D. Sotteau, *Decomposition of complete graphs into isomorphic subgraphs with five vertices*, Ars Combin. **10** (1980) 211–254.

[4] J.-C. Bermond and J. Schönheim, *G-decomposition of K_n, where G has four vertices or less*, Discrete Math. **19** (1977) 113–120.
doi:10.1016/0012-365X(77)90027-9

[5] D. Bryant and S. El-Zanati, *Graph decompositions*, in: Handbook of Combinatorial Designs, 2nd Ed., C.J. Colbourn and J.H. Dinitz (Eds), (Chapman & Hall/CRC Press, Boca Raton, FL, 2007) 477–486.

[6] R.C. Bunge, C.J. Cowan, L.J. Cross, S.I. El-Zanati, A.E. Hart, D.P. Roberts and A.M. Youngblood, *Decompositions of complete digraphs into small tripartite digraphs*, J. Combin. Math. Combin. Comput., to appear.

[7] R.C. Bunge, S.I. El-Zanati, H.J. Fry, K.S. Krauss, D.P. Roberts, C.A. Sullivan, A.A. Unsicker and N.E. Witt, *On the spectra of bipartite directed subgraphs of K_4^**, J. Combin. Math. Combin. Comput. **98** (2016) 375–390.

[8] C.J. Colbourn and J.H. Dinitz, Handbook of Combinatorial Designs, 2nd Ed., (Chapman & Hall/CRC Press, Boca Raton, FL, 2007).

[9] G. Ge, *Group divisible designs*, in: Handbook of Combinatorial Designs, 2nd Ed., C.J. Colbourn and J.H. Dinitz (Eds.), (Chapman & Hall/CRC Press, Boca Raton, FL, 2007) 255–260.

[10] G. Ge, S. Hu, E. Kolotoğlu and H. Wei, *A complete solution to spectrum problem for five-vertex graphs with application to traffic grooming in optical networks*, J. Combin. Des. **23** (2015) 233–273.
doi:10.1002/jcd.21405

[11] A. Hartman and E. Mendelsohn, *The last of the triple systems*, Ars Combin. **22** (1986) 25–41.

[12] D.G. Hoffman and K. Kirkpatrick, *G-designs of order n and index λ where G has* 5 *vertices or less*, Australas. J. Combin. **18** (1998) 13–37.

[13] R.C. Read and R.J. Wilson, An Atlas of Graphs, (Oxford University Press, Oxford, 1998).

[14] R. Wilson, *Construction and uses of pairwise balanced designs*, Math. Centre Tracts. **55** (1974) 18–41.

Received 26 June 2017
Revised 25 November 2017
Accepted 25 November 2017