Discussiones Mathematicae Graph Theory 40 (2020) 67–76 doi:10.7151/dmgt.2110

# ON THE METRIC DIMENSION OF DIRECTED AND UNDIRECTED CIRCULANT GRAPHS

Tomáš Vetrík

Department of Mathematics and Applied Mathematics University of the Free State Bloemfontein, South Africa

e-mail: vetrikt@ufs.ac.za

### Abstract

The undirected circulant graph  $C_n(\pm 1, \pm 2, \ldots, \pm t)$  consists of vertices  $v_0, v_1, \ldots, v_{n-1}$  and undirected edges  $v_i v_{i+j}$ , where  $0 \le i \le n-1, 1 \le j \le t$   $(2 \le t \le \frac{n}{2})$ , and the directed circulant graph  $C_n(1,t)$  consists of vertices  $v_0, v_1, \ldots, v_{n-1}$  and directed edges  $v_i v_{i+1}, v_i v_{i+t}$ , where  $0 \le i \le n-1$   $(2 \le t \le n-1)$ , the indices are taken modulo n. Results on the metric dimension of undirected circulant graphs  $C_n(\pm 1, \pm t)$  are available only for special values of t. We give a complete solution of this problem for directed graphs  $C_n(1,t)$  for every  $t \ge 2$  if  $n \ge 2t^2$ . Grigorious *et al.* [On the metric dimension of circulant and Harary graphs, Appl. Math. Comput. 248 (2014) 47–54] presented a conjecture saying that  $\dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) = t + p - 1$  for n = 2tk + t + p, where  $3 \le p \le t + 1$ . We disprove it by showing that  $\dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \le t + \frac{p+1}{2}$  for n = 2tk + t + p, where  $t \ge 4$  is even, p is odd,  $1 \le p \le t + 1$  and  $k \ge 1$ .

Keywords: metric dimension, resolving set, circulant graph, distance.

2010 Mathematics Subject Classification: 05C35, 05C12.

### 1. INTRODUCTION

Let V(G) be vertex set of a connected (undirected or directed) graph G. The distance d(u, v) between two vertices u, v in an undirected graph is the number of edges in a shortest path between u and v. In a directed graph G the distance d(u, v) from a vertex  $u \in V(G)$  to a vertex  $v \in V(G)$  is the length of a shortest directed path from u to v. A vertex w resolves two vertices u and v if  $d(u, w) \neq d(v, w)$ . For an ordered set of vertices  $W = \{w_1, w_2, \dots, w_z\}$ , the representation of distances of v with respect to W is the ordered z-tuple

$$r(v|W) = (d(v, w_1), d(v, w_2), \dots, d(v, w_z)).$$

A set  $W \subset V(G)$  is a resolving set of G if every two distinct vertices of G have different representations of distances with respect to W (if every two vertices of G are resolved by some vertex in W). The metric dimension of G is the number of vertices in a smallest resolving set and it is denoted by dim(G). The *i*-th coordinate in r(v|W) is 0 if and only if  $v = w_i$ . Thus in order to prove that Wis a resolving set of G, it suffices to show that  $r(u|W) \neq r(v|W)$  for every two different vertices  $u, v \in V(G) \setminus W$ .

The metric dimension is an invariant, which has applications in robot navigation [9], pharmaceutical chemistry [2], pattern recognition and image processing [10]. It has been extensively studied. For example, Imran [5] studied barycentric subdivisions of Cayley graphs and Saputro *et al.* [12] gave bounds on the metric dimension of the lexicographic product of graphs

Let n, m and  $a_1, a_2, \ldots, a_m$  be positive integers such that  $1 \leq a_1 < a_2 < \cdots < a_m \leq \frac{n}{2}$ . The undirected circulant graph  $C_n(\pm a_1, \pm a_2, \ldots, \pm a_m)$  consists of the vertices  $v_0, v_1, \ldots, v_{n-1}$  and undirected edges  $v_i v_{i+a_j}$ , where  $0 \leq i \leq n-1$ ,  $1 \leq j \leq m$ ; the indices are taken modulo n.

For generators  $a_1, a_2, \ldots, a_m$  such that  $1 \le a_1 < a_2 < \cdots < a_m \le n-1$ , the directed circulant graph  $C_n(a_1, a_2, \ldots, a_m)$  consists of the vertices  $v_0, v_1, \ldots, v_{n-1}$  and directed edges  $v_i v_{i+a_j}$ , where  $0 \le i \le n-1$ ,  $1 \le j \le m$ ; the indices are taken modulo n. The directed circulant graph  $C_n(-a_1, -a_2, \ldots, -a_m)$  contains the directed edges  $v_i v_{i-a_j}$ .

Circulant graphs form an important family of Cayley graphs. The metric dimension of undirected circulant graphs  $C_n(\pm 1, \pm t)$  was studied for special values of t. Javaid, Rahim and Ali [8] proved that if  $n \equiv 0, 2, 3 \pmod{4}$ , then  $dim(C_n(\pm 1, \pm 2)) = 3$ . Borchert and Gosselin [1] showed that if  $n \equiv 1 \pmod{4}$ , then  $dim(C_n(\pm 1, \pm 2)) = 4$ . The undirected circulant graphs  $C_n(\pm 1, \pm 3)$  were considered in [7] and the graphs  $C_n(\pm 1, \pm \frac{n}{2})$  for even n were investigated in [11]. We study the metric dimension for directed circulant graphs with 2 generators. We give a complete solution of this problem for directed graphs  $C_n(1, t)$  for every  $t \geq 2$  if  $n \geq 2t^2$ .

Exact values of the metric dimension of undirected graphs  $C_n(\pm 1, \pm 2, \pm 3)$ were given in [1] and [6]. Grigorious *et al.* [4] showed that t + 1 vertices  $v_0, v_1, \ldots, v_t$  resolve the graph  $C_n(\pm 1, \pm 2, \ldots, \pm t)$  if  $n \equiv r \pmod{2t}$ , where  $2 \leq r \leq t + 2$  and they gave the bound  $\dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \leq r - 1$ if  $n \equiv r \pmod{2t}$ , where  $t + 3 \leq r \leq 2t + 1$ . They presented a conjecture saying that  $\dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) = t + p - 1$  for n = 2tk + t + p, where  $3 \leq p \leq t+1$ . We disprove it for even  $t \geq 4$  and odd  $p \geq 5$  by showing that  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \leq t + \frac{p+1}{2}$  for n = 2tk+t+p where  $t \geq 4$  is even, p is odd,  $1 \leq p \leq t+1$  and  $k \geq 1$ . Note that Chau and Gosselin [3] recently proved that  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) = t+1$  if  $n \equiv 2 \pmod{2t}$  and  $n \equiv t+1 \pmod{2t}$ . They also showed that  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) = dim(C_{n+2t}(\pm 1, \pm 2, \ldots, \pm t))$  for large n, which implies that the metric dimension of the graphs  $C_n(\pm 1, \pm 2, \ldots, \pm t)$  is completely determined by the congruence class of n modulo 2t.

### 2. Directed Circulant Graphs

We study the metric dimension of directed circulant graphs  $C_n(1,t)$ . It is easy to see that the graph  $C_n(1,t)$  is isomorphic to the graph  $C_n(-1,-t)$  for  $2 \le t \le n-1$ . We present Theorems 1 and 2 for the graph  $C_n(-1,-t)$ , because it is easier to express distances from vertices in a graph to vertices in chosen resolving sets if we consider  $C_n(-1,-t)$  (especially in the proof of Theorem 2).

The distance from the vertex  $v_j$  to the vertex  $v_i$  in  $C_n(-1, -t)$ , where  $i, j \in \{0, 1, \ldots, n-1\}$ , is

(1)  
(1)  
(2)  

$$d(v_j, v_i) = \begin{cases} \left\lfloor \frac{j-i}{t} \right\rfloor + p, & p \equiv (j-i) \pmod{t}, & \text{if } j \ge i, \\ \left\lfloor \frac{n+j-i}{t} \right\rfloor + p, & p \equiv (n+j-i) \pmod{t}, & \text{if } j < i, \end{cases}$$

where  $0 \le p \le t - 1$ .

**Theorem 1.** Let  $t \ge 2$  and  $n \ge 2t^2$ . Then  $dim(C_n(-1, -t)) \ge t$ .

**Proof.** We prove the result by contradiction. Assume that  $dim(C_n(-1, -t)) \leq t - 1$ . Let  $W = \{v_{i_1}, v_{i_2}, \ldots, v_{i_{t-1}}\}$  be a resolving set of  $C_n(-1, -t)$ , where  $0 \leq i_1 \leq i_2 \leq \cdots \leq i_{t-1}$ . Since we have at most t - 1 different vertices in W and the graph  $C_n(-1, -t)$  has at least  $2t^2$  vertices,  $C_n(-1, -t)$  contains a set of 2t consecutive vertices  $V' = \{v_j, v_{j+1}, \ldots, v_{j+2t-1}\}$ , where  $0 \leq j \leq n - 1$ , such that no vertex of W is in V'. Without loss of generality we can assume that j = n - 2t, which means that  $V' = \{v_{n-2t}, v_{n-2t+1}, \ldots, v_{n-1}\}$  and  $i_{t-1} < n - 2t$ .

Since  $|W| \le t-1$ , there is a  $k \in \{0, 1, \ldots, t-1\}$ , such that no vertex  $v_i \in W$  satisfies  $i \equiv k \pmod{t}$ . So we can write any vertex of W in the form  $v_{tr+s}$ , where  $0 \le s \le t-1$ ,  $s \ne k$  and  $r \ge 0$ .

Let  $v_l$  be any vertex in the set of t vertices  $\{v_{n-2t}, v_{n-2t+1}, \ldots, v_{n-t-1}\}$ , such that  $l \equiv k \pmod{t}$ . Then we can write l = tx + k, where  $0 \leq k \leq t - 1$ . We show that the vertices  $v_{tx+k}, v_{tx+k+t-1} \in V'$  are not resolved by W. Note that

T. Vetrík

tx + k > tr + s. By (1) we have

$$d(v_{tx+k}, v_{tr+s}) = \begin{cases} \left\lfloor \frac{tx+k-(tr+s)}{t} \right\rfloor + k - s = x - r + \left\lfloor \frac{k-s}{t} \right\rfloor + k - s \\ = x - r + k - s & \text{if } k > s, \\ x - r + \left\lfloor \frac{k-s}{t} \right\rfloor + k - s + t \\ = x - r + k - s + t - 1 & \text{if } k < s, \end{cases}$$

$$d(v_{tx+k+t-1}, v_{tr+s}) = \begin{cases} \left\lfloor \frac{tx+k+t-1-(tr+s)}{t} \right\rfloor + k - 1 - s \\ = x - r + k - s & \text{if } k > s, \\ x + 1 - r + \left\lfloor \frac{k-1-s}{t} \right\rfloor + k - 1 - s + t \\ = x - r + k - s + t - 1 & \text{if } k < s. \end{cases}$$

Since  $d(v_{tx+k}, v_{tr+s}) = d(v_{tx+k+t-1}, v_{tr+s})$  for any vertex  $v_{tr+s} \in W$ , the graph  $C_n(-1, -t)$  is not resolved by W. A contradiction.

Let us present an upper bound on the metric dimension of directed circulant graphs with 2 generators.

**Theorem 2.** Let  $2 \le t < n$ . Then  $dim(C_n(-1, -t)) \le t$ .

**Proof.** We prove that  $W = \{v_0, v_1, \ldots, v_{t-1}\}$  is a resolving set of  $C_n(-1, -t)$ . First we find all vertices  $v_j$   $(1 \le j \le n-1)$  of  $C_n(-1, -t)$  such that  $d(v_j, v_0) = x$  for any  $x \ge 1$ . We can write j = tr + p where  $r \ge 0$  and  $0 \le p \le t - 1$ . Since by (1),  $d(v_{tr+p}, v_0) = r + p$ , we have r + p = x. Thus r = x - p ( $\ge 0$ ) and then  $v_{t(x-p)+p}$  for  $0 \le p \le t - 1$  and  $1 \le t(x-p) + p \le n - 1$  are the vertices of  $C_n(1,t)$  such that  $d(v_{t(x-p)+p}, v_0) = x$ .

It remains to show that these vertices are resolved by  $v_i$ , i = 1, 2, ..., t - 1. It suffices to consider only those vertices  $v_{t(x-p)+p}$  which are not in W, so we can assume that t(x-p) + p > i. For i = 1, 2, ..., t - 1, by (1),

(3) 
$$d(v_{t(x-p)+p}, v_i) = \begin{cases} x - p + \left\lfloor \frac{p-i}{t} \right\rfloor + p - i = x - i & \text{if } p \ge i, \\ x - p + \left\lfloor \frac{p-i}{t} \right\rfloor + p - i + t = x + t - 1 - i & \text{if } p < i. \end{cases}$$

We know that the first entry of  $r(v_{t(x-p)+p}|W)$  is x. From (3) it follows that the next p entries (where  $0 \le p \le t-1$ ) are x-i and the last t-1-p entries of  $r(v_{t(x-p)+p}|W)$  are x+t-1-i.

So if p = 0 (and if  $v_{tx}$  exists), the first entry of  $r(v_{tx}|W)$  is x and the other entries are x + t - 1 - i which means that  $r(v_{tx}|W) = (x, x + t - 2, x + t - 3, ..., x + t - 1 - (t - 1))$ . If p = 1, the first entry of  $r(v_{t(x-1)+1}|W)$  is x, the second entry is x - 1 and the other entries are x + t - 1 - i, so  $r(v_{t(x-1)+1}|W) = 0$ 

 $(x, x-1, x+t-3, x+t-4, \dots, x+t-1-(t-1)). \text{ Similarly } r(v_{t(x-2)+2}|W) = (x, x-1, x-2, x+t-4, \dots, x+t-1-(t-1)), \dots, r(v_{t(x-(t-1))+(t-1)}|W) = (x, x-1, x-2, \dots, x-(t-1)).$ 

Since all vertices  $v_j$ ,  $1 \le j \le n-1$ , such that  $d(v_j, v_0) = x$  are resolved by W, we have  $dim(C_n(-1, -t)) \le |W| = t$ .

From Theorems 1 and 2 we obtain Corollary 3.

Corollary 3. Let  $t \ge 2$  and  $n \ge 2t^2$ . Then  $dim(C_n(-1, -t)) = t$ .

Since the graphs  $C_n(-1, -t)$  and  $C_n(1, t)$  are isomorphic, we get the following corollary.

**Corollary 4.** Let  $t \ge 2$  and  $n \ge 2t^2$ . Then  $dim(C_n(1,t)) = t$ .

## 3. UNDIRECTED CIRCULANT GRAPHS

We give an upper bound on the metric dimension of undirected circulant graphs  $C_n(\pm 1, \pm 2, \ldots, \pm t)$  for  $n \equiv r \pmod{2t}$ , where r = 1 and  $r = t+1, t+3, \ldots, 2t-1$ .

The distance between two vertices  $v_i$  and  $v_j$  in  $C_n(\pm 1, \pm 2, \dots, \pm t)$ , where  $0 \le i < j < n$ , is

(4) 
$$d(v_i, v_j) = \min\left\{ \left\lceil \frac{j-i}{t} \right\rceil, \left\lceil \frac{n-(j-i)}{t} \right\rceil \right\}.$$

This equation can be simplified as

(5)  
(6)  

$$d(v_i, v_j) = \begin{cases} \left\lceil \frac{j-i}{t} \right\rceil & \text{if } 0 \le j-i \le \frac{n}{2}, \\ \left\lceil \frac{n-(j-i)}{t} \right\rceil & \text{if } \frac{n}{2} < j-i < n. \end{cases}$$

**Theorem 5.** Let n = 2tk + t + p where  $t \ge 4$  is even, p is odd,  $1 \le p \le t + 1$  and  $k \ge 1$ . Then

$$\dim \left( C_n(\pm 1, \pm 2, \dots, \pm t) \right) \le t + \frac{p+1}{2}.$$

**Proof.** Let n = 2tk + t + p where  $k \ge 1, t \ge 4$  is even and  $p = 1, 3, \ldots, t + 1$ . Let

$$W_1 = \{v_0, v_2, \dots, v_{t-2}\}, \quad W_2 = \{v_{t-1}, v_{t+1}, \dots, v_{2t-3}\}, \\ W_3 = \{v_{tk+t-1}, v_{tk+t+1}, \dots, v_{tk+t+p-2}\}.$$

We have  $|W_1| = |W_2| = \frac{t}{2}$  and  $|W_3| = \frac{p+1}{2}$ . Let us prove that  $W = W_1 \cup W_2 \cup W_3$  is a resolving set of the graph  $C_n(1, 2, \dots, t)$ .

We divide the vertex set of  $C_n(\pm 1, \pm 2, \dots, \pm t)$  into four disjoint sets:

$$V_1 = \{v_0, v_1, \dots, v_t\}, \qquad V_2 = \{v_{t+1}, v_{t+2}, \dots, v_{tk+t}\}, \\ V_3 = \{v_{tk+t+1}, v_{tk+t+2}, \dots, v_{tk+t+p-1}\}, \quad V_4 = \{v_{tk+t+p}, v_{tk+t+p+1}, \dots, v_{n-1}\}.$$

First we prove that any two vertices of  $V_2$  have different representations of distances with respect to W. For x = 1, 2, ..., k - 1; j = 1, 2, ..., t; i = 0, 2, ..., t - 2, we have  $v_i \in W_1$  and by (5),

$$d(v_{tx+j}, v_i) = x + \left\lceil \frac{j-i}{t} \right\rceil = \begin{cases} x+1 & \text{if } i < j, \\ x & \text{if } i \ge j, \end{cases}$$

and if x = k; j = 1, 2, ..., t, by (4), we get

$$d(v_{tk+j}, v_i) = \min\left\{ \left\lceil \frac{(tk+j) - i}{t} \right\rceil, \left\lceil \frac{n - [(tk+j) - i]}{t} \right\rceil \right\},\$$
$$= \min\left\{ k + \left\lceil \frac{j - i}{t} \right\rceil, k + 1 + \left\lceil \frac{p + i - j}{t} \right\rceil \right\} = \left\{ \begin{matrix} k+1 & \text{if } i < j, \\ k & \text{if } i \ge j. \end{matrix} \right\}$$

Since j (where  $1 \le j \le t$ ) is greater than  $\left\lceil \frac{j}{2} \right\rceil$  elements from the set  $\{0, 2, \ldots, t-2\}$ , the first  $\left\lceil \frac{j}{2} \right\rceil$  entries of  $r(v_{tx+j}|W_1)$  for  $x = 1, 2, \ldots, k$  are equal to x + 1 and the other  $\frac{t}{2} - \left\lceil \frac{j}{2} \right\rceil$  entries are x;  $r(v_{tx+j}|W_1) = (x + 1, \ldots, x + 1, x, \ldots, x)$ . Therefore the only vertices in  $V_2$  with the same representations of distances with respect to  $W_1$  are the pairs  $(v_{t+1}, v_{t+2}), (v_{t+3}, v_{t+4}), \ldots, (v_{tk+t-1}, v_{tk+t})$ . But since for  $x = 1, 2, \ldots, k$  and  $j = 1, 3, \ldots, t - 3$ , we obtain  $v_{t+j} \in W_2$  and by (5),

$$d(v_{tx+j}, v_{t+j}) = x - 1, \ d(v_{tx+j+1}, v_{t+j}) = x - 1 + \left\lceil \frac{1}{t} \right\rceil = x,$$

and for  $v_{t-1} \in W_2$ , we have

$$d(v_{tx+t-1}, v_{t-1}) = x, \ d(v_{tx+t}, v_{t-1}) = x + \left\lceil \frac{1}{t} \right\rceil = x+1,$$

vertices in  $W_2$  resolve the pairs  $(v_{t+1}, v_{t+2}), (v_{t+3}, v_{t+4}), \ldots, (v_{tk+t-1}, v_{tk+t})$ . Thus no two vertices in  $V_2$  have the same representations of distances with respect to W.

Let us study representations of distances of the vertices in  $V_4$ . For  $x = 1, 2, \ldots, k-1$ ;  $j = 0, 1, \ldots, t-1$ ;  $i = 0, 2, \ldots, t-2$ ; we have  $v_i \in W_1$  and by (6),

$$d(v_{n-tx+j}, v_i) = \left\lceil \frac{n - \left[ (n - tx + j) - i \right]}{t} \right\rceil = x + \left\lceil \frac{i - j}{t} \right\rceil = \begin{cases} x & \text{if } i \le j, \\ x + 1 & \text{if } i > j, \end{cases}$$

and if x = k, we get

$$d(v_{n-tk+j}, v_i) = \min\left\{ \left\lceil \frac{(n-tk+j)-i}{t} \right\rceil, \left\lceil \frac{n-\left[(n-tk+j)-i\right]}{t} \right\rceil \right\}$$
$$= \min\left\{ k+1 + \left\lceil \frac{p+j-i}{t} \right\rceil, k+\left\lceil \frac{i-j}{t} \right\rceil \right\} = \left\{ \begin{matrix} k & \text{if } i \le j, \\ k+1 & \text{if } i > j. \end{matrix} \right\}$$

Since j (where  $0 \le j \le t-1$ ) is greater than or equal to  $\lfloor \frac{j}{2} \rfloor + 1$  elements from the set  $\{0, 2, \ldots, t-2\}$ , the first  $\lfloor \frac{j}{2} \rfloor + 1$  entries of  $r(v_{n-tx+j}|W_1)$  (for  $x = 1, 2, \ldots, k$ ) are equal to x and the other entries are x + 1. The only vertices in  $V_4$  with the same representations of distances with respect to  $W_1$  are the pairs  $(v_{n-tk}, v_{n-tk+1}), (v_{n-tk+2}, v_{n-tk+3}), \ldots, (v_{n-2}, v_{n-1})$ . We show that most of these pairs are resolved by vertices in  $W_2$ . For  $x = 1, 2, \ldots, k-1$  and  $j = 1, 3, \ldots, t-3$ , we have  $v_{t+j} \in W_2$  and by (6),

$$d(v_{n-tx+j}, v_{t+j}) = x + 1, d(v_{n-tx+j-1}, v_{t+j}) = x + 1 + \left|\frac{1}{t}\right| = x + 2,$$

and for  $v_{t-1} \in W_2$ ,  $x = 1, 2, \dots, k$ , by (6),

$$d(v_{n-tx+t-1}, v_{t-1}) = x, \ d(v_{n-tx+t-2}, v_{t-1}) = x + \left\lceil \frac{1}{t} \right\rceil = x+1,$$

so vertices of  $W_2$  resolve all pairs of vertices  $(v_{n-tk+t-2}, v_{n-tk+t-1}), (v_{n-tk+t}, v_{n-tk+t+1}), \ldots, (v_{n-2}, v_{n-1}),$  which are the pairs  $(v_{tk+2t+p-2}, v_{tk+2t+p-1}), (v_{tk+2t+p}, v_{tk+2t+p+1}), \ldots, (v_{n-2}, v_{n-1}).$  It remains to resolve the pairs  $(v_{tk+t+p+1}, v_{tk+t+p+2}, v_{tk+t+p+3}), \ldots, (v_{tk+2t+p-4}, v_{tk+2t+p-3}).$ 

For 
$$j = 0, 2, ..., t - p - 3$$
, we have  $v_{t+p+j} \in W_2$  and by (5),

 $d(v_{tk+t+p+j}, v_{t+p+j}) = k, \ d(v_{tk+t+p+j+1}, v_{t+p+j}) = k + \lfloor \frac{1}{t} \rfloor = k+1,$ 

so the pairs  $(v_{tk+t+p}, v_{tk+t+p+1}), \ldots, (v_{tk+2t-3}, v_{tk+2t-2})$  are resolved by  $W_2$ .

For  $j = t - p - 1, t - p + 1, \dots, t - 4$ , we have  $v_{tk+p+j} \in W_3$  and by (5),

$$d(v_{tk+t+p+j}, v_{tk+p+j}) = 1, \ d(v_{tk+t+p+j+1}, v_{tk+p+j}) = 1 + \left\lceil \frac{1}{t} \right\rceil = 2,$$

so the pairs  $(v_{tk+2t-1}, v_{tk+2t}), \ldots, (v_{tk+2t+p-4}, v_{tk+2t+p-3})$  are resolved by  $W_3$ . Thus all pairs of vertices in  $V_4$  are resolved by W.

A vertex  $v \in V_2$  and a vertex in  $V_4$  can have the same representations of distances with respect to  $W_1$  only if all entries of  $r(v|W_1)$  are the same numbers. For  $x = 1, 2, \ldots, k$ , we have  $v_{tx+t-1}, v_{tx+t} \in V_2$  and  $r(v_{tx+t-1}|W_1) = r(v_{tx+t}|W_1) =$  $(x + 1, \ldots, x + 1)$ . For  $v_{n-tx+t-2}, v_{n-tx+t-1} \in V_4$  we have  $r(v_{n-tx+t-2}|W_1) =$  $r(v_{n-tx+t-1}|W_1) = (x, \ldots, x)$ , which implies that for  $x = 1, 2, \ldots, k-1$ , we obtain  $r(v_{tx+t-1}|W_1) = r(v_{tx+t}|W_1) = r(v_{n-tx-2}|W_1) = r(v_{n-tx-1}|W_1)$ . Since for  $v_{2t-3} \in W_2$ , by (5),

 $d(v_{tx+t-1}, v_{2t-3}) = x - 1 + \left\lceil \frac{2}{t} \right\rceil = x, \quad d(v_{tx+t}, v_{2t-3}) = x - 1 + \left\lceil \frac{3}{t} \right\rceil = x,$ 

T. Vetrík

and by (6),

$$d(v_{n-tx-2}, v_{2t-3}) = x+2+\left\lceil \frac{-1}{t} \right\rceil = x+2, \ d(v_{n-tx-1}, v_{2t-3}) = x+2+\left\lceil \frac{-2}{t} \right\rceil = x+2,$$
  
any vertex in  $V_2$  and any vertex in  $V_4$  have different representations of distances

with respect to W.

We study representations of the vertices in  $V_3$ . For j = 1, 2, ..., p - 1 and i = 0, 2, ..., t - 2, we have  $v_i \in W_1$  and by (4),

$$d(v_{tk+t+j}, v_i) = \min\left\{k+1 + \left\lceil \frac{j-i}{t} \right\rceil, k + \left\lceil \frac{p+i-j}{t} \right\rceil\right\} = k+1,$$

thus  $r(v_{tk+t+j}|W_1) = (k+1, \ldots, k+1)$ . The only vertices in  $V_2 \cup V_4$  with the same representations of distances with respect to  $W_1$  are  $v_{tk+t-1}$  and  $v_{tk+t}$ .

Let us prove that any two vertices in  $V_3 \cup \{v_{tk+t-1}, v_{tk+t}\}$  have different representations with respect to W. It suffices to consider the vertices in  $V' = (V_3 \cup \{v_{tk+t-1}, v_{tk+t}\}) \setminus W_3 = \{v_{tk+t}, v_{tk+t+2}, \ldots, v_{tk+t+p-1}\}$ . For  $j = 0, 2, \ldots, p-1$  and  $i = 1, 3, \ldots, t-3$ , we have  $v_{t+i} \in W_2$  and by (5)

$$d(v_{tk+t+j}, v_{t+i}) = k + \left\lceil \frac{j-i}{t} \right\rceil = \begin{cases} k & \text{if } i \ge j, \\ k+1 & \text{if } i < j. \end{cases}$$

Since j (for  $j \leq t-2$ ) is greater than  $\frac{j}{2}$  elements from the set  $\{1, 3, \ldots, t-3\}$ , the first  $\frac{j}{2}$  entries of  $r(v_{tk+t+j}|W'_2)$  where  $W'_2 = W_2 \setminus \{v_{t-1}\}$  are equal to k+1and the other  $\frac{t}{2} - \frac{j}{2} - 1$  entries are k. If p = t+1 and j = t, we obtain  $r(v_{tk+t+j}|W'_2) = r(v_{tk+2t}|W'_2) = (k+1,\ldots,k+1)$ . It follows that the only vertices of V' having the same representations of distances with respect to  $W'_2$  are  $v_{tk+2t}$ and  $v_{tk+2t-2}$  if p = t+1. These vertices are resolved by  $v_{tk+t-1} \in W_3$ , since by (5),  $d(v_{tk+2t}, v_{tk+t-1}) = 1 + \lfloor \frac{1}{t} \rfloor = 2$  and  $d(v_{tk+2t-2}, v_{tk+t-1}) = 1 + \lfloor \frac{-1}{t} \rfloor = 1$ . Thus all vertices of  $V_3$  are resolved by W.

We consider the vertices in  $V_1$ . For j = 1, 3, ..., t-1 and t; i = 0, 2, ..., t-2, we have  $v_i \in W_1$  and  $d(v_j, v_i) = \left\lceil \frac{|j-i|}{t} \right\rceil = 1$ , thus  $r(v_j|W_1) = (1, ..., 1)$  for  $v_j \in V_1 \setminus W_1$ . From the previous part of this proof we know that the only vertices in  $V_2 \cup V_3 \cup V_4$  having the representation with respect to  $W_1$  equal to (1, ..., 1)are  $v_{n-2}$  and  $v_{n-1}$ . Since  $v_{t-1} \in W_2$ , it remains to resolve all pairs of vertices in the set  $V'' = \{v_1, v_3, ..., v_{t-3}; v_t, v_{n-2}, v_{n-1}\}$ .

We study their representations with respect to  $W_2$ . For  $j = 1, 3, \ldots, t - 3$ and  $i = -1, 1, \ldots, t - 3$ , we have  $v_{t+i} \in W_2$  and by (5),

$$d(v_j, v_{t+i}) = 1 + \left\lceil \frac{i-j}{t} \right\rceil = \begin{cases} 1 & \text{if } i \le j, \\ 2 & \text{if } i > j. \end{cases}$$

Since j is greater than or equal to  $\frac{j+3}{2}$  elements from the set  $\{-1, 1, \ldots, t-3\}$ , the first  $\frac{j+3}{2}$  entries of  $r(v_j|W_2)$  are equal to 1 and the other  $\frac{t}{2} - \frac{j+3}{2}$  entries

74

are 2. The first two entries of  $r(v_j|W_3)$  are always 1. For  $v_t$  and any  $v_{t+i} \in W_2$ ,  $d(v_t, v_{t+i}) = \left\lceil \frac{|i|}{t} \right\rceil = 1$ , therefore  $r(v_t|W_2) = (1, \dots, 1)$ . For  $i = -1, 1, \dots, t-3$ , by (6),

$$d(v_{n-1}, v_{t+i}) = 1 + \left\lceil \frac{i+1}{t} \right\rceil = \begin{cases} 1 & \text{if } i = -1, \\ 2 & \text{if } i \ge 1, \end{cases}$$

so  $r(v_{n-1}|W_2) = (1, 2, ..., 2)$ . We have  $d(v_{n-2}, v_{t+i}) = 1 + \lfloor \frac{i+2}{t} \rfloor = 2$ , thus  $r(v_{n-2}|W_2) = (2, ..., 2)$ .

The only pair of vertices in V'' having the same representations with respect to  $W_2$  is  $(v_{t-3}, v_t)$ , which is resolved by  $v_{tk+t-1} \in W_3$ , since by (5) we have  $d(v_{t-3}, v_{tk+t-1}) = k + \lfloor \frac{2}{t} \rfloor = k + 1$  and  $d(v_t, v_{tk+t-1}) = k + \lfloor \frac{-1}{t} \rfloor = k$ .

Every two distinct vertices of the graph  $C_n(\pm 1, \pm 2, \dots, \pm t)$  have different representations of distances with respect to W, thus W is a resolving set of  $C_n(\pm 1, \pm 2, \dots, \pm t)$ . Hence  $dim(C_n(\pm 1, \pm 2, \dots, \pm t)) \leq |W| = t + \frac{p+1}{2}$ .

### 4. Conclusion

We studied the metric dimension of undirected and directed circulant graphs. Results on the metric dimension of undirected circulant graphs  $C_n(\pm 1, \pm t)$  are available only for special values of t. In Section 2 we found exact values of the metric dimension for directed circulant graphs  $C_n(1,t)$  by showing that if  $t \ge 2$  and  $n \ge 2t^2$ , then  $dim(C_n(1,t)) = t$ .

In Section 3 we presented a bound on the metric dimension of undirected circulant graphs. We proved that for n = 2tk + t + p, where  $t \ge 4$  is even, p is odd,  $1 \le p \le t + 1$  and  $k \ge 1$ ,  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \le t + \frac{p+1}{2}$ . Note that by [13],  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \le t + \frac{p}{2}$  if t and p are even,  $2 \le p \le t$ , thus we have  $dim(C_n(\pm 1, \pm 2, \ldots, \pm t)) \le t + \lceil \frac{p}{2} \rceil$  for n = 2tk + t + p, where  $t \ge 4$  is even,  $1 \le p \le t + 1$  and  $k \ge 1$ ,

### Acknowledgments

This work has been supported by the National Research Foundation of South Africa; grant numbers: 112122, 90793.

#### References

- A. Borchert and S. Gosselin, The metric dimension of circulant graphs and Cayley hypergraphs, Util. Math. 106 (2018) 125–147.
- [2] G. Chartrand, L. Eroh, M.A. Johnson and O.R. Oellermann, Resolvability in graphs and the metric dimension of a graph, Discrete Appl. Math. 105 (2000) 99–113. doi:10.1016/S0166-218X(00)00198-0

- K. Chau and S. Gosselin, The metric dimension of circulant graphs and their Cartesian products, Opuscula Math. 37 (2017) 509–534. doi:10.7494/OpMath.2017.37.4.509
- [4] C. Grigorious, P. Manuel, M. Miller, B. Rajan and S. Stephen, On the metric dimension of circulant and Harary graphs, Appl. Math. Comput. 248 (2014) 47–54. doi:10.1016/j.amc.2014.09.045
- M. Imran, On the metric dimension of barycentric subdivision of Cayley graphs, Acta Math. Appl. Sin. Eng. Ser. 32 (2016) 1067–1072. doi:10.1007/s10255-016-0627-0
- M. Imran, A.Q. Baig, S.A. Ul. Hag Bokhary and I. Javaid, On the metric dimension of circulant graphs, Appl. Math. Lett. 25 (2012) 320–325. doi:10.1016/j.aml.2011.09.008
- [7] I. Javaid, M.N. Azhar and M. Salman, Metric dimension and determining number of Cayley graphs, World Appl. Sci. J. 18 (2012) 1800–1812.
- [8] I. Javaid, M.T. Rahim and K. Ali, Families of regular graphs with constant metric dimension, Util. Math. 75 (2008) 21–33.
- S. Khuller, B. Raghavachari and A. Rosenfeld, *Landmarks in graphs*, Discrete Appl. Math. **70** (1996) 217–229. doi:10.1016/0166-218X(95)00106-2
- R.A. Melter and I. Tomescu, Metric bases in digital geometry, Comput. Vision Graphics Image Process. 25 (1984) 113–121. doi:10.1016/0734-189X(84)90051-3
- [11] M. Salman, I. Javaid and M.A. Chaudhry, Resolvability in circulant graphs, Acta Math. Sin. Eng. Ser. 28 (2012) 1851–1864. doi:10.1007/s10114-012-0417-4
- [12] S.W. Saputro, R. Simanjuntak, S. Uttunggadewa, H. Assiyatun, E.T. Baskoro, A.N.M. Salman and M. Bača, *The metric dimension of the lexicographic product* of graphs, Discrete Math. **313** (2013) 1045–1051. doi:10.1016/j.disc.2013.01.021
- [13] T. Vetrík, The metric dimension of circulant graphs, Canad. Math. Bull. 60 (2017) 206–216. doi:10.4153/CMB-2016-048-1

Received 25 May 2017 Revised 23 January 2018 Accepted 23 January 2018