Discussiones Mathematicae Graph Theory 40 (2020) 985–994 doi:10.7151/dmgt.2155

LIGHT MINOR 5-STARS IN 3-POLYTOPES WITH MINIMUM DEGREE 5 AND NO 6-VERTICES¹

Oleg V. Borodin, Anna O. Ivanova

AND

Ekaterina I. Vasil'eva

Institute of Mathematics Siberian Branch Russian Academy of Sciences Novosibirsk, 630090, Russia

e-mail: brdnoleg@math.nsc.ru shmgnanna@mail.ru ekaterinavasilyeva93@gmail.com

Abstract

In 1940, Lebesgue gave an approximate description of the neighborhoods of 5-vertices in the class $\mathbf{P_5}$ of 3-polytopes with minimum degree 5.

Given a 3-polytope P, by w(P) denote the minimum of the degree-sum (weight) of the neighborhoods of 5-vertices (minor 5-stars) in P.

In 1996, Jendrol' and Madaras showed that if a polytope P in $\mathbf{P_5}$ is allowed to have a 5-vertex adjacent to four 5-vertices, then w(P) can be arbitrarily large.

For each P in $\mathbf{P_5}$ without vertices of degree 6 and 5-vertices adjacent to four 5-vertices, it follows from Lebesgue's Theorem that $w(P) \leq 68$. Recently, this bound was lowered to $w(P) \leq 55$ by Borodin, Ivanova, and Jensen and then to $w(P) \leq 51$ by Borodin and Ivanova.

In this note, we prove that every such polytope P satisfies $w(P) \leq 44$, which bound is sharp.

Keywords: planar map, planar graph, 3-polytope, structural properties, 5-star, weight, height.

2010 Mathematics Subject Classification: 05C15.

¹The work was funded by the Russian Science Foundation, grant 16-11-10054.

1. INTRODUCTION

The degree of a vertex or face x in a convex finite 3-dimensional polytope (called a 3-polytope) is denoted by d(x). A k-vertex is a vertex v with d(v) = k. A k^+ -vertex (k^- -vertex) is one of degree at least k (at most k). Similar notation is used for the faces. A 3-polytope with minimum degree 5 is denoted by P_5 , and the set of such 3-polytopes is \mathbf{P}_5 .

The weight of a subgraph S of P_5 is the degree sum of the vertices of S in P_5 , and the *height* of S is the maximum degree of the vertices of S in P_5 . A k-star, a star with k rays, is *minor* if its center v has degree at most 5. In particular, the neighborhoods of 5-vertices are minor 5-stars and vice versa. All stars considered in this note are minor.

By $w(S_k)$ and $h(S_k)$ we denote the minimum weight and height, respectively, of minor k-stars in a given 3-polytope P_5 .

In 1904, Wernicke [13] proved that every P_5 has a 5-vertex adjacent to a 6⁻-vertex. This result was strengthened by Franklin [9] in 1922 to the existence of a 5-vertex with two 6⁻-neighbors. In 1940, in attempts to solve the Four Color Problem, Lebesgue [12, p. 36] gave an approximate description of the neighborhoods of 5-vertices in P_5 s. In particular, this description implies the results in [9,13] and shows that there is a 5-vertex with three 7⁻-neighbors.

The bounds $w(S_1) \leq 11$ (Wernicke [13]) and $w(S_2) \leq 17$ (Franklin [9]) are tight. It was proved by Lebesgue [12] that $w(S_3) \leq 24$, which was improved in 1996 by Jendrol' and Madaras [10] to the sharp bound $w(S_3) \leq 23$. Furthermore, Jendrol' and Madaras [10] gave a precise description of minor 3-stars in P_5 s.

Lebesgue [12] proved $w(S_4) \leq 31$, which was strengthened by Borodin and Woodall [8] to the tight bound $w(S_4) \leq 30$. Note that $w(S_3) \leq 23$ easily implies $w(S_2) \leq 17$ and immediately follows from $w(S_4) \leq 30$ (in both cases, it suffices to delete a vertex of maximum degree from a minor star of minimum weight). Recently, Borodin and Ivanova [1] obtained a precise description of 4-stars in P_5 s.

The more general problem of precisely describing 5-stars at 5-vertices in P_{5} s inspired by Lebesgue's Theorem is still widely open.

Jendrol' and Madaras [10] showed that if a polytope P_5 has a 5-vertex adjacent to four 5-vertices, called a *minor* $(5, 5, 5, 5, \infty)$ -*star*, then $h(S_5)$ and hence $w(S_5)$ can be arbitrarily large. Therefore, in what follows we consider P_5 s without minor $(5, 5, 5, 5, \infty)$ -stars.

Recently, precise upper bounds for the height and weight of minor 5-stars have been obtained for some restricted subclasses in \mathbf{P}_5 . A lot of earlier results on the structure of stars in 3-polytopes can be found in [11].

For every P_5 having no vertices of degree from 6 to 9, Lebegue's bounds $h(S_5) \leq 14$ and $w(S_5) \leq 44$ were improved by Borodin and Ivanova [3] to the sharp bounds $h(S_5) \leq 12$ and $w(S_5) \leq 42$.

LIGHT MINOR 5-STARS IN 3-POLYTOPES WITH MINIMUM DEGREE 5 AND ...987

For each P_5 with no 6- to 8-vertices, it follows from Lebesgue's Theorem that $h(S_5) \leq 17$ and $w(S_5) \leq 46$, which bounds were improved in Borodin, Ivanova and Nikiforov [7] to the best possible bounds $h(S_5) \leq 12$ and $w(S_5) \leq 42$.

Under the absence of 6- and 7-vertices, Lebegue's bound $h(S_5) \leq 23$ was improved by Borodin *et al.* [5] to the sharp bound $h(S_5) \leq 14$.

For each P_5 with no 6-vertices, it follows from Lebesgue's Theorem that $h(S_5) \leq 41$. This bound was lowered to $h(S_5) \leq 28$ by Borodin, Ivanova, and Jensen [4], then to $h(S_5) \leq 23$ in Borodin-Ivanova [2], and finally to the tight bound $h(S_5) \leq 17$ by Borodin, Ivanova, and Nikiforov [6].

As for the minimum weight of minor 5-stars in P_5 s under the absence of 6-vertices, Lebesgue's bound $w(S_5) \leq 68$ was lowered to $w(S_5) \leq 55$ by Borodin, Ivanova, and Jensen [4] and then to $w(S_5) \leq 51$ in Borodin-Ivanova [2]. The purpose of this paper is to prove the following fact.

Theorem 1. Every 3-polytope with minimum degree 5 and neither 6-vertices nor minor $(5, 5, 5, 5, \infty)$ -stars has a minor 5-star with weight at most 44, which bound is best possible.

We note that a light minor 5-star ensured by Theorem 1 has height at most $44 - 4 \times 5 - 7 = 17$. The tightness of the bounds 44 and 17 is confirmed by a construction in [6].

2. Proof of Theorem 1

Discharging.

Suppose that a 3-polytope P'_5 is a counterexample to the main statement of Theorem 1. Thus each minor 5-star in P'_5 has weight at least 45 and at most three 5-vertices.

Let P_5 be a counterexample with the maximum number of edges on the same set of vertices as P'_5 .

Remark 2. P_5 has no 4⁺-face with two nonconsecutive 7⁺-vertices along the boundary, for otherwise adding a diagonal between these vertices would result in a counterexample with greater number of edges.

Let V, E, and F be the sets of vertices, edges, and faces of P_5 . Euler's formula |V| - |E| + |F| = 2 implies

(1)
$$\sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2d(f) - 6) = -12.$$

We assign an *initial charge* $\mu(v) = d(v) - 6$ to each $v \in V$ and $\mu(f) = 2d(f) - 6$ to each $f \in F$, so that only 5-vertices have a negative initial charge.

Using the properties of P_5 as a counterexample to Theorem 1, we define a local redistribution of charges, preserving their sum such that the final charge $\mu(x)$ is non-negative for all $x \in V \cup F$. This will contradict the fact that the sum of the final charges is, by (1), equal to -12.

The final charge $\mu'(x)$ whenever $x \in V \cup F$ is defined by applying the rules R1–R11 below (see Figure 1).

For a vertex v, let $v_1, \ldots, v_{d(v)}$ be the vertices adjacent to v in a fixed cyclic order. If f is a face, then $v_1, \ldots, v_{d(f)}$ are the vertices incident with f in the same cyclic order.

If d is an integer with $8 \le d \le 15$, then we put $\xi_d = \frac{d-6}{d}$.

A vertex is simplicial if it is completely surrounded by 3-faces. A simplicial 5-vertex v is helpful if $d(v_1) \ge 12$, $d(v_2) = d(v_4) = 5$, $d(v_3) = 7$, and $d(v_5) \ge 12$ (see Figure 1, R10). A simplicial 5-vertex v is strong if $d(v_1) = d(v_2) = 5$, $7 \le d(v_3) \le 11$, and $7 \le d(v_5) \le 11$ (so $d(v_4) \ge 45 - 2 \times 11 - 3 \times 5 \ge 8$) (see Figure 1, R11).

R1. Each 4^+ -face gives $\frac{1}{2}$ to each incident 5-vertex.

R2. If a 5-vertex v is incident with precisely one 4^+ -face, then v receives $\frac{1}{2}$ from each adjacent 16^+ -vertex.

R3. A simplicial 5-vertex v with at least two 12⁺-neighbors receives $\frac{1}{2}$ from each adjacent 16⁺-vertex.

R4. A simplicial 5-vertex v with $d(v_4) \neq 5$, $d(v_5) \geq 16$, and no other 12^+ -neighbors receives the following charge from v_5 :

- (a) if $d(v_1) \neq 5$, then 1, and
- (b) if $d(v_1) = 5$, then $\frac{3}{4}$ provided that $d(v_5) \le 17$ or $\frac{5}{6}$ otherwise.

R5. A simplicial 5-vertex v with $d(v_5) \ge 18$, $d(v_1) = d(v_4) = 5$, and $7 \le d(v_2) \le d(v_3) \le 11$ receives $\frac{2}{3}$ from v_5 .

R6. A simplicial 5-vertex v with $16 \le d(v_5) \le 17$, $d(v_1) = d(v_4) = 5$, and $7 \le d(v_2) \le d(v_3) \le 11$ receives from v_5 :

- (n) $\frac{5}{8}$ if neither v_2 nor v_3 is a 7-vertex having six simplicial 5-neighbors ("normally"), and
- (e) $\frac{2}{3}$ otherwise ("as an exception").

R7. A simplicial 5-vertex v with $d(v_5) \ge 16$, $d(v_1) = d(v_2) = d(v_4) = 5$, and $7 \le d(v_3) \le 11$ receives the following charge from v_5 .

- (n) If v_1 is not simplicial or v_2 is not strong (that is "normal"), then $\frac{3}{4}$ if $d(v_5) \le 17$ or $\frac{5}{6}$ otherwise.
- (e) If v_1 is simplicial and v_2 is strong (which is "an exception"), then $\frac{5}{8}$ if $d(v_5) \le 17$ or $\frac{2}{3}$ otherwise.

Light Minor 5-Stars in 3-Polytopes with Minimum Degree 5 and $\dots 989$

R8. A *d*-vertex *v* with $8 \le d(v) \le 15$ gives its 5-neighbor v_2 :

- (a) ξ_d if $d(v_1) = d(v_3) = 5$,
- (b) $\frac{3\xi_d}{2}$ if $d(v_1) = 5$ and $d(v_3) \neq 5$, and
- (c) $2\xi_d$ if $d(v_1) \neq 5$ and $d(v_3) \neq 5$.

R9. A 7-vertex v gives each adjacent simplicial 5-vertex:

- (n) $\frac{1}{5}$ "as a norm", that is if v has at most five simplicial 5-neighbors, or
- (e) $\frac{1}{6}$ "as an exception".

R10. A 7-vertex v receives $\frac{1}{6}$ from each helpful 5-neighbor.

R11. A strong 5-vertex gives $\frac{1}{6}$ to each 5-neighbor.



Figure 1. Rules of discharging.

Checking $\mu'(x) \ge 0$ whenever $x \in V \cup F$. If f is a 4⁺-face, then $\mu'(f) \ge 2d(f) - 6 - d(f) \times \frac{1}{2} = \frac{3(d(f)-4)}{2} \ge 0$ by R1. Now suppose $v \in V$.

Case 1. $d(v) \ge 18$. We know that v gives one of the charges in $\left\{\frac{1}{2}, \frac{2}{3}, \frac{5}{6}, 1\right\}$ to each adjacent 5-vertex incident with at least four 3-faces by R2–R7. Since $d(v) - 6 \ge \frac{2d(v)}{3}$, it suffices to average these donations so that each 5⁺-neighbor will receive at most $\frac{2}{3}$ from v.

To this end, we first switch $\frac{1}{6}$ from 1 given to a 5-neighbor v_k by R4(a) to each of the 7⁺-neighbors v_{k-1} and v_{k+1} (hereafter, addition modulo d(v)). As a result, the averaged donation from v to v_k becomes $1 - 2 \times \frac{1}{6} = \frac{2}{3}$.

Next, if $\frac{5}{6}$ is given to a 5-neighbor v_k by R4(b), then we switch $\frac{1}{6}$ to its common 7⁺-neighbor with v.

Finally, the donation of $\frac{5}{6}$ by R7(n) happens to a simplicial 5-neighbor v_k of v having cyclic neighbors $v_{k-1}, x_k, y_k, v_{k+1}$ with $7 \le d(x_k) \le 11$ and 5-neighbors v_{k-1}, y_k, v_{k+1} , where either v_{k+1} is not simplicial or y_k is not strong.

If v_{k+1} is not simplicial, then we switch $\frac{1}{6}$ from v_k to v_{k+1} and note that the latter receives at most $\frac{1}{2}$ from v by R2.

From now on suppose that v_{k+1} is simplicial, and let z_k be the vertex conjugated with v_k with respect to the edge $y_k v_{k+1}$. Since v_k receives $\frac{5}{6}$ by R7(n) by our assumption, it follows that $d(z_k) \notin \{7, \ldots, 11\}$, for otherwise y_k is strong since it has the fifth neighbor of degree at least $w(S_5) - 3 \times 5 - 2 \times 11 = 8$ and is simplicial in view of Remark 2.

If $d(z_k) \ge 12$, then we switch $\frac{1}{6}$ from v_k to v_{k+1} , where v_{k+1} this time receives $\frac{1}{2}$ by R3. Note that v_{k+2} receives $\frac{1}{2}$ by R2 or R3, which implies that $\frac{1}{6}$ is switched to v_{k+1} only once.

It remains to assume that $d(z_k) = 5$. This implies that $d(v_{k+2}) \ge 7$ since v_{k+1} cannot have four 5-neighbors. Here, we switch $\frac{1}{6}$ from v_k to v_{k+2} . (Of course, v_{k+1} also switches $\frac{1}{6}$ from its $\frac{5}{6}$ obtained by R4(b) to v_{k+2} , as said above.)

It is not hard to see that no 5-vertex v_{k+1} can receive $\frac{1}{6}$ in the course of our averaging both from v_k and v_{k+2} since then v_{k+1} would have four 5-neighbors, which is impossible.

As a result, the averaged donation of v to each 5-neighbor becomes at most $1-2\times\frac{1}{6}=\frac{5}{6}-\frac{1}{6}=\frac{1}{2}+\frac{1}{6}=\frac{2}{3}$ and that to each 7⁺-neighbor is at most $0+4\times\frac{1}{6}=\frac{2}{3}$, as desired.

Case 2. $16 \le d(v) \le 17$. We now show that the neighbors of v receive from v by R2–R7 at most $\frac{5}{8}$ on the average, which implies that $\mu'(v) \ge d(v) - 6 - \frac{5d(v)}{8} = \frac{3(d(v)-16)}{8} \ge 0$. We proceed similarly to Case 1 with a 5-vertex v_k getting more than $\frac{5}{8}$ from v by R4, R6(e) or R7(n).

If v_k is as in R4(a), then we shift $\frac{1}{4}$ from 1 obtained by v_k to each of the 7⁺-vertices v_{k-1} and v_{k+1} . In R4(b), we shift $\frac{1}{8}$ from $\frac{3}{4}$ to a unique 7⁺-vertex in $\{v_{k-1}, v_{k+1}\}$.

990

LIGHT MINOR 5-STARS IN 3-POLYTOPES WITH MINIMUM DEGREE 5 AND ...991

Now consider R6(e), which has no analogues in Case 1. By symmetry, we can assume that v_{k+1} lies in a common 3-face with v_k and a 7-vertex having six simplicial 5-neighbors. Now $d(v_{k+2}) \ge 7$ as v_{k+1} cannot have three 5-neighbors in addition to a 7-neighbor and a 17⁻-neighbor since $w(S_5) \ge 45$ by assumption. Recall that v_{k+1} receives at most $\frac{3}{4}$ by R4(b), R3, or R7(n) and that $\frac{1}{8}$ was already switched from v_{k+1} to v_{k+2} in the previous paragraph. Here, we also switch $\frac{1}{8}$ from $\frac{2}{3}$ received by v_k to v_{k+2} .

In the situation of R7(n), let v_{k+1} lie in a 3-face incident with three 5-vertices. Arguing as in Case 1, we see that either v_{k+1} receives $\frac{1}{2}$ from v, in which case we switch $\frac{1}{8}$ from v_k to v_{k+1} , or we have $d(v_{k+2}) \ge 7$, in which case we switch $\frac{1}{8}$ from v_k to v_{k+2} .

As a result of this averaging, each 5-neighbor of v receives at most $1-2\times\frac{1}{4}<\frac{3}{4}-\frac{1}{8}=\frac{1}{2}+\frac{1}{8}=\frac{5}{8}$, while each 7⁺-neighbor receives at most $4\times\frac{1}{8}=\frac{1}{4}+2\times\frac{1}{8}=2\times\frac{1}{4}<\frac{5}{8}$ from v, as desired.

Case 3. $8 \leq d(v) \leq 15$. To satisfy R8, we first send $\xi_{d(v)}$ to each neighbor v_k , and then each 7⁺-neighbor v_k transfers $\frac{\xi_{d(v)}}{2}$ to each 5-vertex in $\{v_{k-1}, v_{k+1}\}$. This shows that $\mu'(v) \geq d(v) - 6 - d(v) \times \xi_{d(v)} = 0$.

Case 4. d(v) = 7. If v has at most five simplicial 5-neighbors, then $\mu'(v) \ge 7 - 6 - 5 \times \frac{1}{5} = 0$ by R9(n). If v has precisely six simplicial 5-neighbors, then $\mu'(v) \ge 1 - 6 \times \frac{1}{6} = 0$ by R9(e).

Finally, suppose v is completely surrounded by simplicial 5-vertices. This implies that there is a 7-cycle $C_7 = w_1 \cdots w_7$ avoiding v, where each v_k lies in a 3-face $w_k v_k w_{k+1}$ (addition modulo 7). Note that $d(w_k)+d(w_{k+1}) \ge 45-3\times 5-7 = 23$ whenever $1 \le k \le 7$. By the oddness of 7, v has a helpful neighbor, which gives $\frac{1}{6}$ to v by R10. As a result, we have $\mu'(v) \ge 1 + \frac{1}{6} - 7 \times \frac{1}{6} = 0$ in view of R9(e), as required.

Case 5. d(v) = 5. If v is incident with at least two 4⁺-faces, then $\mu'(v) \ge 5 - 6 + 2 \times \frac{1}{2} = 0$ by R1.

If v is incident with precisely one 4⁺-face, then we are done when v has a 12^+ -neighbor since v receives $\frac{1}{2}$ by R1 and at least $\frac{1}{2}$ by R2 or R8.

So suppose otherwise. Note that v then has two 8⁺-neighbors, for otherwise v would have an 11⁻-neighbor and four 7⁻-neighbors, which implies $w(S_5) \leq 5 + 4 \times 7 + 11 < 45$, a contradiction. Thus $\mu'(v) \geq -1 + \frac{1}{2} + 2 \times \frac{1}{4} = 0$ by R1 and R8.

From now on we can assume that v is simplicial.

Subcase 5.1. v is helpful, with $d(v_1) \ge 12$, $d(v_2) = d(v_4) = 5$, $d(v_3) = 7$, and $d(v_5) \ge 12$. Now v receives $\frac{1}{2}$ from each of v_1, v_5 by R3 and/or R8. Also v receives at least $\frac{1}{6}$ from v_3 by R9 and returns $\frac{1}{6}$ to v_3 by R10. This implies $\mu'(v) \ge -1 + 2 \times \frac{1}{2} + \frac{1}{6} - \frac{1}{6} = 0$, as desired. Subcase 5.2. v is strong, with $d(v_1) = d(v_2) = 5$, $7 \le d(v_3) \le d(v_5) \le 11$, and $d(v_4) \ge 45 - 3 \times 5 - 2 \times 11 = 8$. Now v must collect the total of at least $\frac{4}{3}$ from v_3, v_4, v_5 in order to be able to give $2 \times \frac{1}{6}$ to v_1, v_2 according to R11 (and leave 1 for itself).

We are easily done if $d(v_4) \ge 12$, for then v_4 gives v at least 1 by R4(a) or R8(c) while each of v_3, v_5 gives at least $\frac{1}{6}$ by R8 and R9.

So suppose $d(v_4) \leq 11$. Since $d(v_3) + d(v_4) + d(v_5) \geq w(S_5) - 3 \times 5 = 30$, this implies that v has no neighbors of degree less than $30 - 2 \times 11 = 8$. If $d(v_4) = 8$, then $d(v_3) = d(v_5) = 11$, which implies that v receives $\frac{1}{2}$ from v_4 by R8(c) and $2 \times \frac{15}{22}$ from v_3 , v_5 by R8(b), as desired. If $d(v_4) \geq 9$, then v receives at least $\frac{2}{3}$ from v_4 by R8(c) and at least $2 \times \frac{3}{8}$ from v_3 , v_5 by R8(b), and we are done.

Subcase 5.3. v does not give charge away by R10 and R11. So we must check that v collects the total of at least 1 from its neighbors by R3–R9. If v has at least two 12⁺-neighbors, then $\mu'(v) \ge -1 + 2 \times \frac{1}{2} = 0$ by R3 or R8(a),(b). So in what follows we assume that v has at most one 12⁺-neighbor, which means that R3 is not applied to v.

Subcase 5.3.1. v has at most one 5-neighbor. Here, v receives at least $\frac{3}{8}$ from an 8-neighbor and at least $\frac{1}{2}$ from a 9⁺-neighbor by R4–R8. This implies, in view of R9, that $\mu'(v) \ge -1+3 \times \frac{1}{6} + \frac{1}{2} = 0$ in the presence of a 9⁺-neighbor or $\mu'(v) \ge$ $-1+2 \times \frac{1}{6} + 2 \times \frac{3}{8} > 0$ when v has at least two 8-neighbors. However, one of this situations is inevitable, since otherwise we would have $w(S_5) \le 5+4 \times 7+8 < 45$, which is impossible.

Subcase 5.3.2. v has precisely two 5-neighbors. Note that the total degree of the three 7⁺-neighbors of v is at least $45 - 3 \times 5 = 30$.

Suppose v has no 7-neighbor. Each 8⁺-neighbor v_2 gives v by R4–R8 at least $\frac{1}{4}$ if $d(v_1) = d(v_3) = 5$ and at least $\frac{3}{8}$ if $d(v_1) \neq 5$, so $\mu'(v) \geq -1 + \frac{1}{4} + 2 \times \frac{3}{8} = 0$, and we are done.

Next suppose v has at least one 7-neighbor. Now the other two 7⁺-neighbors have the total degree at least 30 - 7 = 23, so there is a 12^+ -neighbor, say v_2 , among them.

If v_2 gives v at least $\frac{3}{4}$ to v by R4 or R8, then $\mu'(v) > 0$, since the other two 7⁺-neighbors give at least $2 \times \frac{1}{6}$ by R4–R9.

So suppose $d(v_1) = d(v_3) = 5$ and $7 = d(v_4) \le d(v_5)$. Now if $d(v_2) \ge 18$, then we have $\mu'(v) \ge -1 + \frac{2}{3} + 2 \times \frac{1}{6} = 0$ by R4–R9.

For $16 \leq d(v_2) \leq 17$ we are similarly done if v_2 gives $\frac{2}{3}$ by R6(e), so suppose R6(n) is applied to v_2 rather than R6(e). If $d(v_5) \geq 8$, then $\mu'(v) \geq -1 + \frac{5}{8} + \frac{1}{6} + \frac{3}{8} > 0$. It remains to assume that $d(v_4) = d(v_5) = 7$ and neither of v_4, v_5 has six simplicial 5-neighbors (as if we apply R9(e) to v_4 or v_5 it would mean we should apply R6(e) to v, and then $\mu'(v) \geq -1 + \frac{2}{3} + 2 \times \frac{1}{6} = 0$). This means that $\mu'(v) \geq -1 + \frac{5}{8} + 2 \times \frac{1}{5} = \frac{1}{40}$ by R6(n) and R9(n).

Finally, suppose $12 \le d(v_2) \le 15$. Now $d(v_5) \ge w(S_5) - 3 \times 5 - 7 - 15 = 8$, and it suffices to observe that v receives at least $\frac{1}{2}, \frac{1}{6}, \frac{3}{8}$ from v_2, v_4, v_5 , respectively, which makes $\mu'(v) > 0$, as desired.

Subcase 5.3.3. v has precisely three 5-neighbors. Note that the total degree of the two 7⁺-neighbors of v is at least $45 - 4 \times 5 = 25$.

First suppose $7 \le d(v_1) \le d(v_2)$. By the above assumption that R3 is not applied, we have $d(v_1) \le 11$, which implies that v has a 14⁺-neighbor. Note that v_2 gives v at least $\frac{3}{4}$ by R4(b) or R8, while v_1 gives v at least $\frac{3}{8}$ by R8 if $d(v_1) \ge 8$, and then we have $\mu'(v) \ge 0$. But if $d(v_1) = 7$, then $d(v_2) \ge 25 - 7 = 18$, and $\mu'(v) \ge -1 + \frac{5}{6} + \frac{1}{6} = 0$ by R4(b) combined with R9.

Thus from now on we can assume that $7 \le d(v_1) \le 11$ and $d(v_3) \ge 14$. If $d(v_3) \le 15$, then v receives from v_1 and v_3 at least $1 = \frac{2}{5} + \frac{3}{5} = \xi_{10} + \xi_{15} < \xi_{11} + \xi_{14}$ by R8(a), as desired.

Next suppose $16 \le d(v_3) \le 17$, which implies that $d(v_1) \ge 8$. Since v_1 gives v at least $\frac{1}{4}$ by R8(a) while v_3 gives either $\frac{3}{4}$ or $\frac{5}{8}$ by R7, we are done unless v_3 gives $\frac{5}{8}$ by R7(e). The latter happens when v_5 is strong, in which case v receives $\frac{1}{6}$ from v_5 by R11, which yields $\mu'(v) \ge -1 + \frac{1}{4} + \frac{1}{6} + \frac{5}{8} \ge 0$.

Finally, suppose $d(v_3) \ge 18$. Now v_1 gives v at least $\frac{1}{6}$ by R9 while v_3 gives either $\frac{5}{6}$ or $\frac{2}{3}$ by R7. Since the donation of $\frac{2}{3}$ by R7(e) to v is accompanied by receiving $\frac{1}{6}$ by R11 from a strong vertex v_5 , we have $\mu'(v) \ge -1 + 2 \times \frac{1}{6} + \frac{2}{3} = -1 + \frac{1}{6} + \frac{5}{6} = 0$.

Thus we have proved $\mu'(x) \ge 0$ whenever $x \in V \cup F$, which contradicts (1) and completes the proof of Theorem 1.

References

- O.V. Borodin and A.O. Ivanova, Describing 4-stars at 5-vertices in normal plane maps with minimum degree 5, Discrete Math. **313** (2013) 1710–1714. doi:10.1016/j.disc.2013.04.025
- O.V. Borodin and A.O. Ivanova, Light and low 5-stars in normal plane maps with minimum degree 5, Sib. Math. J. 57 (2016) 470-475. doi:10.1134/S0037446616030071
- [3] O.V. Borodin and A.O. Ivanova, Light neighborhoods of 5-vertices in 3-polytopes with minimum degree 5, Sib. Elektron. Mat. Izv. 13 (2016) 584–591. doi:10.17377/semi.2016.13.045
- [4] O.V. Borodin, A.O. Ivanova and T.R. Jensen, 5-stars of low weight in normal plane maps with minimum degree 5, Discuss. Math. Graph Theory 34 (2014) 539–546. doi:10.7151/dmgt.1748
- [5] O.V. Borodin, A.O. Ivanova, O.N. Kazak and E.I. Vasil'eva, Heights of minor 5-stars in 3-polytopes with minimum degree 5 and no vertices of degree 6 and 7, Discrete Math. 341 (2018) 825–829. doi:10.1016/j.disc.2017.11.021

- [6] O.V. Borodin, A.O. Ivanova and D.V. Nikiforov, Low minor 5-stars in 3-polytopes with minimum degree 5 and no 6-vertices, Discrete Math. 340 (2017) 1612–1616. doi:10.1016/j.disc.2017.03.002
- [7] O.V. Borodin, A.O. Ivanova and D.V. Nikiforov, Low and light minor 5-stars in 3polytopes with minimum degree 5 and restrictions on the degrees of major vertices, Sib. Math. J. 58 (2017) 600–605. doi:10.1134/S003744661704005X
- [8] O.V. Borodin and D.R. Woodall, Short cycles of low weight in normal plane maps with minimum degree 5, Discuss. Math. Graph Theory 18 (1998) 159–164. doi:10.7151/dmgt.1071
- [9] P. Franklin, The four colour problem, Amer. J. Math. 44 (1922) 225–236. doi:10.2307/2370527
- [10] S. Jendrol' and T. Madaras, On light subgraphs in plane graphs of minimum degree five, Discuss. Math. Graph Theory 16 (1996) 207–217. doi:10.7151/dmgt.1035
- [11] S. Jendrol' and H.-J. Voss, Light subgraphs of graphs embedded in the plane—A survey, Discrete Math. **313** (2013) 406–421. doi:10.1016/j.disc.2012.11.007
- [12] H. Lebesgue, Quelques conséquences simples de la formule d'Euler, J. Math. Pures Appl. 19 (1940) 27–43.
- P. Wernicke, Über den kartographischen Vierfarbensatz, Math. Ann. 58 (1904) 413–426. doi:10.1007/BF01444968

Received 30 November 2017 Revised 11 June 2018 Accepted 12 June 2018