

MINIMAL GRAPHS WITH RESPECT TO GEOMETRIC DISTANCE REALIZABILITY

TOMÁŠ MADARAS

AND

PAVOL ŠIROCZKI

Institute of Mathematics
P.J. Šafárik University in Košice
Jesenná 5, 04001 Košice, Slovakia

e-mail: tomas.madaras@upjs.sk
siroczi@gmail.com

Abstract

A graph G is minimal non-unit-distance graph if there is no drawing of G in Euclidean plane having all edges of unit length, but, for each edge e of G , $G - e$ has such a drawing. We prove that, for infinitely many n , the number of non-isomorphic n -vertex minimal non-unit-distance graphs is at least exponential in n .

Keywords: unit-distance graph, odd-distance graph, Euclidean plane.

2010 Mathematics Subject Classification: 05C62.

1. INTRODUCTION

Throughout this paper, we consider connected graphs without loops or multiple edges, and their drawings in the plane. By *drawing* $D = D(G)$ of a graph $G = (V, E)$ in the plane, we mean a function ϕ defined on $V \cup E$ which assigns each vertex $v \in V$ a point $\phi(v) \in \mathbb{R}^2$, and, each edge $uv \in E$ is mapped to a simple curve $\phi(uv) \subset \mathbb{R}^2$ with endpoints $\phi(u), \phi(v)$. It is usually assumed that ϕ is injective on V and, for each edge $uv \in E$ and each $w \in V, w \neq u, v$, $\phi(w) \notin \phi(uv)$ holds; a drawing which violates some of these two properties is called *degenerate*. When mentioning distances between points $\phi(x), \phi(y)$ (which correspond to vertices x, y of G) in the plane, we refer to the Euclidean distance

(here denoted as $\text{dist}(x, y)$) unless specified otherwise. All graph-theoretic terms which are not defined here are used in accordance with [11].

To describe properties of graphs of a graph family $\mathcal{G}$, one aims to obtain a complete characterization of members of $\mathcal{G}$, usually in terms of forbidden subgraphs, induced subgraphs, minors or topological minors. A related source of information on properties of $\mathcal{G}$ is the set $\mathcal{M}(\mathcal{G})$ of *minimal non- $\mathcal{G}$ -graphs* (that is, the graphs which do not belong to $\mathcal{G}$, but each of their proper subgraphs is in $\mathcal{G}$) and the function $f(\mathcal{G}, n)$ whose value is the number of non-isomorphic n -vertex graphs from $\mathcal{M}(\mathcal{G})$. The asymptotic character of $f(\mathcal{G}, n)$ may serve as an indirect indicator for 'tractability' of the family $\mathcal{G}$ in terms of efficient algorithmic recognition of good characterization. For example, Kuratowski's theorem [6] yields that the minimal non-planar graphs are exactly the subdivisions of K_5 or $K_{3,3}$, and the number of such graphs on n vertices is certainly polynomial in n (because it is bounded from above by the number of ways how to redistribute 2-valent vertices on the edges of K_5 or $K_{3,3}$). On the other hand, for 1-planar graphs (for which there exists a drawing in the plane such that each its edge is crossed at most once), it was shown in [5] that, for each $n \geq 63$, there exist at least $2^{\frac{n-54}{4}}$ non-isomorphic minimal non-1-planar graphs. Note that $f(\mathcal{G}, n)$ is not related to the number of distinct n -vertex graphs of $\mathcal{G}$, as this number is exponential for both planar and 1-planar graphs.

In this paper, we study the function $f(\mathcal{UD}, n)$ for the family $\mathcal{UD}$ of graphs defined by the property that, for each $G \in \mathcal{UD}$, there exists a non-degenerate drawing D of G in the Euclidean plane such that all edges of D are unit segments (D is further referred as *unit-distance drawing* of G). The family $\mathcal{UD}$ is a part of larger family of *unit-distance graphs* in $\mathbb{R}^2$: here the vertex set V is a subset of $\mathbb{R}^2$ and the edge set E is a subset of all pairs $\{x, y\}$, $x, y \in V$ with $\text{dist}(x, y) = 1$ (see [1]). The latter family was widely studied mainly in connection with the (Hadwiger)-Nelson problem of determining the chromatic number of the plane (for its history and connections to other areas of mathematics, see the excellent monograph [9]). The problem of characterization of minimal non-unit-distance graphs was first mentioned in [2] and seems to be still open. It is easy to see that the family $\mathcal{UD}$ is not closed under taking minors, since any subdivision of $K_{2,3}$ (which is not a unit-distance graph) has an unit-distance drawing. In [4] and [10], it is proven that the problem of recognition of unit-distance graphs (and, more generally, the problem of determining the minimum dimension of Euclidean space in which a graph has unit-distance drawing) is NP-hard.

We prove the following result.

Theorem 1. *For infinitely many integers n , there exist at least exponentially many non-isomorphic n -vertex graphs from $\mathcal{M}(\mathcal{UD})$.*

2. THE PROOF

For the purpose of the subsequent proof, we briefly recall the equidistribution theorem.

Theorem 2 (Equidistribution theorem, [12]). *Let a be an irrational number. Then the sequence $\{n \cdot a - \lfloor n \cdot a \rfloor\}_{n=1}^{\infty}$ is uniformly distributed on the unit interval $(0, 1)$.*

Claim 3. *The inequality*

$$(*) \quad n - \sqrt{3} \cdot \left\lfloor \frac{n}{\sqrt{3}} \right\rfloor < \frac{2 - \sqrt{3}}{2}$$

holds for infinitely many positive integers n .

Proof. The equidistribution theorem for $a = \frac{1}{\sqrt{3}}$ yields that the sequence $\left\{ \frac{n}{\sqrt{3}} - \left\lfloor \frac{n}{\sqrt{3}} \right\rfloor \right\}_{n=1}^{\infty}$ is uniformly distributed on the interval $(0, 1)$. Thus, the sequence $s = \left\{ n - \sqrt{3} \cdot \left\lfloor \frac{n}{\sqrt{3}} \right\rfloor \right\}_{n=1}^{\infty}$ is uniformly distributed on the interval $(0, \sqrt{3})$ as the elements of this sequence are exactly the elements of the previous sequence multiplied by $\sqrt{3}$. Our claim now follows from the fact that the terms on the right side of $(*)$ are positive and the uniform distribution of s . ■



From now on we will denote the set of all positive integers satisfying $(*)$ by $\mathcal{S}$ and, for a positive integer $n \in \mathcal{S}$, we define $N = \left\lfloor \frac{n}{\sqrt{3}} \right\rfloor$; note that $\min \mathcal{S} = 7$.

We continue by defining, for any $n \in \mathcal{S}$, the set $\mathcal{G}_n$ of graphs $G_{n,S}$ in the following way: start with a 'snake' S_n consisting of $4n - 1$ triangles (see Figure 1), denote its 2-valent vertices (the left and the right one, respectively) as v_0 and v_{2n} . Note that S_n has (up to mirror symmetry) a unique unit-distance drawing.

Let B_1 and B_2 be the graphs in Figure 2. We define the sequence $S = \{D_i\}_{i=1}^{2N}$, where each D_i is either a copy of B_1 or B_2 , and the number of occurrences of B_1 and B_2 in S is equal. Concatenate the graphs of S by identifying vertices $y_i \in D_i$ and $x_{i+1} \in D_{i+1}$, thus forming a 'chain'. Finally, identify vertex $x_1 \in D_1$ with vertex $v_0 \in S_n$ and $y_{2N} \in D_{2N}$ with $v_{2n} \in S_n$ obtaining the graph $G_{n,S}$ (the resulting graph clearly depends on the choice of S).

Claim 4. *The number of non-isomorphic graphs in $\mathcal{G}_n$ is at least exponential in terms of number of their vertices.*

Proof. We have $\frac{1}{2} \binom{2N}{N}$ possible orderings of terms in S (the term $\frac{1}{2}$ prevents us from including mirror symmetry), which is at least $\frac{4^N}{2 \cdot (2N+1)} > 2^N$ for $N \geq 5$. On the other hand, the number of vertices of any graph $G_{n,S} \in \mathcal{G}_n$ is $4n + 1 +$

Figure 1. The 'snake' S_n .Figure 2. The blocks B_1 and B_2 .

$8N$, which is linear in N . Thus the number of non-isomorphic graphs in $\mathcal{G}_n$ is exponential in the size of their vertex set (it is easy to see that for two sequences $S_1 = \{D_i\}_{i=1}^{2N}$ and $S_2 \neq S_1$, the graphs G_{n,S_1} and G_{n,S_2} are isomorphic only if $S_2 = \{D_{2N-i}\}_{i=1}^{2N}$, but then their drawings possess mirror symmetry). ■

Claim 5. *Every graph $G \in \mathcal{G}_n$ is not a unit-distance graph.*


Proof. Assume that a unit distance drawing of G exists. There is a unique (up to mirror symmetry) unit-distance realization of the graph S_n . The distance between vertices v_0 and v_{2n} in this drawing is necessarily $2n$. Similarly, for $B_i, i \in \{1, 2\}$, the unit-distance realization of B_i is also unique (up to mirror symmetry) and the distance between vertices x and y in this drawing is exactly $\sqrt{3}$. Thus, the distance between vertices x_1 and y_{2N} is at most $2N\sqrt{3} = 2\sqrt{3} \cdot \left\lfloor \frac{n}{\sqrt{3}} \right\rfloor < 2n$, the last inequality follows from the irrationality of $\sqrt{3}$. Notice that in G vertex x_1 is identified with vertex v_0 and vertex v_{2n} is identified with y_{2N} , but the distance between these vertices is either $2n$ (from the realisation of S_n) and, on the other hand, strictly less than $2n$ (from the properties of the drawings of blocks $B_i, i \in [1, 2N]$). This contradiction shows that G is not a unit-distance graph. ■

Claim 6. *Every graph $G \in \mathcal{G}_n$ belongs to $\mathcal{M}(UD)$.*


Proof. We need to show that for every edge $e \in E(G)$, the graph $G - e$ belongs to UD . This is done by case analysis.

First, assume that the removed edge e belongs to block D_i for some $i \in [1, 2N]$ and this block is isomorphic with B_1 . Here we need to distinguish two cases (up

to mirror symmetry) labeled with e_1, e_2 in Figure 2. In each of these cases the drawing of this block can be deformed in such a way that the distance between vertices x_i and y_i in this drawing of $D_i - e$ is arbitrarily close to 2. The drawings for each case are illustrated in Figure 3. For each $j \in [1, 2N], j \neq i$, the distance between vertices x_j and y_j is exactly $\sqrt{3}$. Thus the sum $\sum_{i=1}^{2N} \text{dist}(x_j, y_j)$ is arbitrary close to $(2N - 1)\sqrt{3} + 2$. On the other hand, from (*) it follows that $2n < (2N - 1)\sqrt{3} + 2$, so there exists such a drawing that the considered sum is bigger than $2n$.

Figure 3. Deformation of B_1 without edge.

Next, assume that the removed edge e belongs to block D_i for some $i \in [1, 2N]$ and this block is isomorphic with B_2 . In this case we need to distinguish 5 subcases (again, up to mirror symmetry) labelled $e_i, i \in [1, 5]$, in Figure 2. Again, in each of the subcases, the drawing of this block can be deformed in such a way that the distance between vertices x_i and y_i in this drawing of $D_i - e$ is arbitrarily close to 2. The deformation for each case is illustrated in Figure 4. Again, similarly to the previous case, it follows that $\sum_{i=1}^{2N} \text{dist}(x_j, y_j) > 2n$.

Figure 4. Deformation of B_2 without edge.

Finally, assume that the edge e belongs to S_n . Figure 5 shows all five cases how to deform $S_n - e$ to obtain a non-degenerate unit-distance drawing in which the Euclidean distance between v_0 and v_{2n} will be less than $2n$. The difference between $2n$ and $2N$ is less than $2 - \sqrt{3}$, so it suffices to decrease the distance between vertices v_0 and v_{2n} by any value bigger than $2 - \sqrt{3}$. This is obviously possible for the first case (vertex v_0 arbitrary close to v_2) and the second case (v_0 arbitrary close to v_1). In the third case, the worst possibility (in terms of distance between v_0 and v_{2n}) is to remove the edge u_1u_2 , but v_0 could be rotated arbitrary close to vertex v_2 , thus the distance between v_0 and v_{2n} would be arbitrary close to $2(n-1)$.




Figure 5. Deformation of S_n without edge.

After deforming the considered graph as illustrated for the fourth case, the distance between vertices v_0 and v_{2n} is arbitrary close to $\left((2n - \frac{1}{2})^2 + (\frac{\sqrt{3}}{2})^2\right)^{\frac{1}{2}}$ (consider the right-angled triangle where the line segment connecting v_0 and v_{2n} would form the hypotenuse), which converges to $2n - \frac{1}{2}$ as n increases (even for the smallest possible $n = 7$ this distance is 13.52775 which is quite close to 13.5). It follows from (*) that $2n - \frac{1}{2} < 2N\sqrt{3} + 2 - \sqrt{3} - \frac{1}{2} < 2N\sqrt{3}$.

In fifth case, the vertex v_0 could be rotated around the vertex u_{i+1} to be close to the line segment connecting vertices v_{i+1} and v_{2n} . This time the worst case would be to remove the edge v_1v_2 , but the vertex v_0 would be close to vertex v_3 and the distance between v_0 and v_{2n} in this case would be close to $2n - 3$.

In all five described cases, the distance between vertices v_0 and v_{2n} in a unit-distance drawing of $S_n - e$ can be decreased at least by a factor arbitrary close to $\frac{1}{2}$, which is bigger than $2 - \sqrt{3}$.

Now, it suffices to show that the chain of graphs corresponding to the sequence S can be attached to v_0 and v_{2n} to obtain a non-degenerate unit-distance drawing of G . This is implied by the following general auxiliary result.

Lemma 7. *Let $\{H_i\}_{i=1}^k$ be a sequence of graphs such that, for each $i \in [1, k]$ there exists a non-degenerate unit-distance drawing D_i of H_i containing two vertices $x_{i-1}^i, x_i^i \in V(H_i)$ having Euclidean distance $d_i > 1$. Further, let D be a non-degenerate unit-distance drawing of a graph H containing vertices u, v with their Euclidean distance less than $\sum_{i=1}^k d_i$. Consider the graph G' obtained by identification of each vertex $x_i^i \in V(H_i)$ with vertex $x_{i+1}^{i+1} \in V(H_{i+1})$ and subsequent identification of u with x_0^1 and v with vertex x_k^k . Then there exists a non-degenerate unit-distance drawing of G' .*

Proof. For $i = 1, \dots, k - 1$, let x_i be the vertex resulting from the identification of x_i^i with x_{i+1}^{i+1} , and let $x_0 = u, x_k = v$. Start with the placement of D_1 after identifying u with x_0^1 . If we place the unit-distance drawing D_1 of H_1 in such a way that x_1 in H_1 lies on the line segment connecting points x_0 and x_k , then the Euclidean distance between x_1 and x_k will be strictly less than $\sum_{i=2}^k d_i$. Then there exists, by continuity of Euclidean distance, $\varepsilon \in \mathbb{R}$ such that when rotating the drawing of H_i by an angle $\alpha \in (-\varepsilon, \varepsilon)$ around the point x_0 , the Euclidean distance between x_1 and x_k is still strictly less than $\sum_{i=2}^k d_i$.

We have to show that the placement of H_1 can be arranged in a way that the unit-distance drawing obtained is not degenerate. First, let us consider the case when a vertex $w \in V(D_1)$ lies on an edge $e \in E(D)$. The feasible positions for w are determined by the angle $\alpha \in (-\varepsilon, \varepsilon)$ and form a circular arc. An edge $e \in E(D)$ can intersect this arc in at most two points, so there are a finite number of forbidden positions for w . Now, consider the case when $w \in V(D)$ and $e \in E(D_1)$. We argue in the same way, but this time we fix the unit-distance

drawing of H_1 and rotate the drawing D by angle α around the vertex x_0 . As H_1 and H are finite graphs, an appropriate α can be chosen such that all vertices of $V(D_1) \cup V(D)$ are distinct and none of them appears in the interior of a line segment of $D_1 \cup D$.

The above described step can be repeated for all $i \in [1, k-3]$, however, when placing the drawing D_{k-2} of H_{k-2} , one has to be more careful, because the position of x_{k-2} determines the position of x_{k-1} . When preparing to place D_{k-2} the situation is that we have a point x_{k-3} whose distance from x_k is less than $d_{k-2} + d_{k-1} + d_k$. Again, if we place D_{k-2} such that x_{k-2} lies on the line segment determined by points x_{k-3} and x_k , the Euclidean distance of x_{k-2} and x_k is less than $d_{k-1} + d_k$ and there exists $\varepsilon \in \mathbb{R}$ such that when rotating the drawing D_{k-2} by an angle $\alpha \in (-\varepsilon, \varepsilon)$ around x_{k-3} , the Euclidean distance between x_{k-2} and x_k remains smaller than $d_{k-1} + d_k$. The possible positions for x_{k-1} form a circular arc (intersection of the circle with unit diameter centered in x_k and the disk centered in x_{k-3} with radius d_{k-2}). Thus, there are infinitely many possible positions for x_{k-2} and x_{k-1} and now we can again apply the argument as in the previous cases to avoid a degeneracy. $\square$

Now, that Lemma 7 is proved, the proof of Claim 6 is complete. $\blacksquare$

3. CONCLUDING REMARKS

Since our result covers only a subset of the set of positive integers, it would be desirable to prove an exponential lower bound for $f(\mathcal{UD}, n)$ for all n .

Note that, in unit-distance drawings of graphs from $\mathcal{UD}$, we do not require the condition $\text{dist}(x, y) \neq 1$ for nonadjacent vertices x, y (unit-distance drawings which satisfy this condition are called *faithful*). The recent paper [1] considers labelled unit-distance and faithful unit-distance graphs in general Euclidean spaces $\mathbb{R}^d$; it is proved that, for prescribed d and given number of vertices, there is far more unit-distance graphs than the faithful ones. Concerning the structure of minimal non-faithful unit-distance graphs (for fixed d), the authors address the problem of minimum number of edges of such graphs and provide lower and upper bounds for bipartite case.

Along with unit-distance graphs, there are studied also *odd distance graphs* for which there exist drawings with all edges represented by line segments of odd lengths. It is known (see [3]) that the complete graph K_4 is not an odd distance graph, and recently in [8] it was proved that this is also not the case for the wheel graph W_6 (observe that both these graphs also belong to $\mathcal{M}(\mathcal{UD})$). Hence, the related problem would be to characterize all minimal non-odd-distance graphs; note that, by the result of Piepmeyer (see [7]), these graphs have chromatic number at least 4.

Acknowledgement

This work was supported by the Slovak Research and Development Agency under the Contract No. APVV-15-0116.

REFERENCES

- [1] N. Alon and A. Kupavskii, *Two notions of unit distance graphs*, J. Combin. Theory Ser. A **125** (2014) 1–17.
doi:10.1016/j.jcta.2014.02.006
- [2] P. Erdős, F. Harary and W.T. Tutte, *On the dimension of a graph*, Mathematika **12** (1965) 118–122.
doi:10.1112/S0025579300005222
- [3] R.L. Graham, B.L. Rothschild and E.G. Strauss, *Are there $n + 2$ points in $\mathbb{E}^n$ with odd integral distances?*, Amer. Math. Monthly **81** (1974) 21–25.
doi:10.1080/00029890.1974.11993491
- [4] B. Horvat, J. Kratochvíl and T. Pisanski, *On the computational complexity of degenerate unit distance representations of graphs*, Lecture Notes in Comput. Sci. **6460** (2011) 274–285.
doi:10.1007/978-3-642-19222-7_28
- [5] V.P. Korzhik and B. Mohar, *Minimal obstructions for 1-immersions and hardness of 1-planarity testing*, J. Graph Theory **72** (2013) 30–71.
doi:10.1002/jgt.21630
- [6] K. Kuratowski, *Sur le problème des courbes gauches en topologie*, Fund. Math. **15** (1930) 271–283.
doi:10.4064/fm-15-1-271-283
- [7] L. Piepmeyer, *The maximum number of odd integral distances between points in the plane*, Discrete Comput. Geom. **16** (1996) 113–115.
doi:10.1007/BF02711135
- [8] M. Rosenfeld and N.L. Tiên, *Forbidden subgraphs of the odd-distance graph*, J. Graph Theory **75** (2014) 323–330.
doi:10.1002/jgt.21738
- [9] A. Soifer, *The Mathematical Coloring Book* (Springer-Verlag, New York, 2009).
doi:10.1007/978-0-387-74642-5
- [10] M. Tikhomirov, *On computational complexity of length embeddability of graphs*, Discrete Math. **339** (2016) 2605–2612.
doi:10.1016/j.disc.2016.05.011
- [11] D.B. West, *Introduction to Graph Theory* (Prentice Hall, 1996).
- [12] H. Weyl, *Über die Gibbs'sche Erscheinung und verwandte Konvergenzphänomene*, Rend. Circ. Mat. **30** (1910) 377–407.
doi:10.1007/BF03014883

Received 9 April 2018

Revised 6 August 2018

Accepted 6 August 2018