

## CHANGING AND UNCHANGING OF THE DOMINATION NUMBER OF A GRAPH: PATH ADDITION NUMBERS

VLADIMIR SAMODIVKIN

*Department of Mathematics  
University of Architecture, Civil Engineering and Geodesy  
Sofia 1164, Bulgaria*

**e-mail:** vl.samodivkin@gmail.com

### Abstract

Given a graph  $G = (V, E)$  and two its distinct vertices  $u$  and  $v$ , the  $(u, v)$ - $P_k$ -addition graph of  $G$  is the graph  $G_{u,v,k-2}$  obtained from disjoint union of  $G$  and a path  $P_k : x_0, x_1, \dots, x_{k-1}$ ,  $k \geq 2$ , by identifying the vertices  $u$  and  $x_0$ , and identifying the vertices  $v$  and  $x_{k-1}$ . We prove that  $\gamma(G) - 1 \leq \gamma(G_{u,v,k})$  for all  $k \geq 1$ , and  $\gamma(G_{u,v,k}) > \gamma(G)$  when  $k \geq 5$ . We also provide necessary and sufficient conditions for the equality  $\gamma(G_{u,v,k}) = \gamma(G)$  to be valid for each pair  $u, v \in V(G)$ . In addition, we establish sharp upper and lower bounds for the minimum, respectively maximum,  $k$  in a graph  $G$  over all pairs of vertices  $u$  and  $v$  in  $G$  such that the  $(u, v)$ - $P_k$ -addition graph of  $G$  has a larger domination number than  $G$ , which we consider separately for adjacent and non-adjacent pairs of vertices.

**Keywords:** domination number, path addition.

**2010 Mathematics Subject Classification:** 05C69.

### 1. INTRODUCTION

For basic notation and graph theory terminology not explicitly defined here, we in general follow Haynes *et al.* [8]. We denote the vertex set and the edge set of a graph  $G$  by  $V(G)$  and  $E(G)$ , respectively. The complement  $\overline{G}$  of  $G$  is the graph whose vertex set is  $V(G)$  and whose edges are the pairs of nonadjacent vertices of  $G$ . We write  $K_n$  for the *complete graph* of order  $n$ ,  $K_{m,n}$  for the *complete bipartite graph* with partite sets of order  $m$  and  $n$ , and  $P_n$  for the *path* on  $n$  vertices. Let  $C_m$  denote the *cycle* of length  $m$ . For any vertex  $x$  of a graph  $G$ ,  $N_G(x)$  denotes the set of all neighbors of  $x$  in  $G$ ,  $N_G[x] = N_G(x) \cup \{x\}$  and the

degree of  $x$  is  $\deg(x, G) = |N_G(x)|$ . The *minimum* and *maximum* degrees of a graph  $G$  are denoted by  $\delta(G)$  and  $\Delta(G)$ , respectively. For a subset  $A \subseteq V(G)$ , let  $N_G(A) = \bigcup_{x \in A} N_G(x)$  and  $N_G[A] = N_G(A) \cup A$ . A *vertex cover* of a graph is a set of vertices such that each edge of the graph is incident to at least one vertex of the set. Let  $G$  be a graph and  $uv$  be an edge of  $G$ . By subdividing the edge  $uv$  we mean forming a graph  $H$  from  $G$  by adding a new vertex  $w$  and replacing the edge  $uv$  by  $uw$  and  $wv$ . Formally,  $V(H) = V(G) \cup \{w\}$  and  $E(H) = (E(G) \setminus \{uv\}) \cup \{uw, wv\}$ . For a graph  $G$ , let  $x \in S \subseteq V(G)$ . A vertex  $y \in V(G)$  is a *S-private neighbor* of  $x$  if  $N_G[y] \cap S = \{x\}$ . The set of all *S-private* neighbors of  $x$  is denoted by  $pn_G[x, S]$ .

The study of domination and related subset problems is one of the fastest growing areas in graph theory. For a comprehensive introduction to the theory of domination in graphs we refer the reader to Haynes *et al.* [8]. A *dominating set* for a graph  $G$  is a subset  $D \subseteq V(G)$  of vertices such that every vertex not in  $D$  is adjacent to at least one vertex in  $D$ . The *domination number* of  $G$ , denoted by  $\gamma(G)$ , is the smallest cardinality of a dominating set of  $G$ . A dominating set of  $G$  with cardinality  $\gamma(G)$  is called a  $\gamma$ -*set* of  $G$ . The concept of  $\gamma$ -bad/good vertices in graphs was introduced by Fricke *et al.* in [5]. A vertex  $v$  of a graph  $G$  is called

- (i) [5]  $\gamma$ -*good*, if  $v$  belongs to some  $\gamma$ -set of  $G$ , and
- (ii) [5]  $\gamma$ -*bad*, if  $v$  belongs to no  $\gamma$ -set of  $G$ .

A graph  $G$  is said to be  $\gamma$ -*excellent* whenever all its vertices are  $\gamma$ -good [5]. Brigham *et al.* [3] defined a vertex  $v$  of a graph  $G$  to be  $\gamma$ -*critical* if  $\gamma(G - v) < \gamma(G)$ , and  $G$  to be *vertex domination-critical* (from now on called *vc-graph*) if each vertex of  $G$  is  $\gamma$ -critical. For a graph  $G$  we define  $V^-(G) = \{x \in V(G) \mid \gamma(G - x) < \gamma(G)\}$ .

It is often of interest to know how the value of a graph parameter  $\mu$  is affected when a change is made in a graph, for instance vertex or edge removal, edge addition, edge subdivision and edge contraction. In this connection, here we consider this question in the case  $\mu = \gamma$  when a path is added to a graph.

Path-addition is an operation that takes a graph and adds an internally vertex-disjoint path between two vertices together with a set of supplementary edges. This operation can be considered as a natural generalization of the edge addition. Formally, let  $u$  and  $v$  be distinct vertices of a graph  $G$ . The  $(u, v)$ - $P_k$ -*addition graph* of  $G$  is the graph  $G_{u,v,k-2}$  obtained from disjoint union of  $G$  and a path  $P_k : x_0, x_1, \dots, x_{k-1}$ ,  $k \geq 2$ , by identifying the vertices  $u$  and  $x_0$ , and identifying the vertices  $v$  and  $x_{k-1}$ . When  $k \geq 3$  we call  $x_1, x_2, \dots, x_{k-2}$  *path-addition vertices*. By  $pa_\gamma(u, v)$  we denote the minimum number  $k$  such that  $\gamma(G) < \gamma(G_{u,v,k})$ . For every graph  $G$  with at least 2 vertices we define

- ▷ the *e-path addition* ( $\bar{e}$ -*path addition*) *number with respect to domination*, de-

noted  $epa_\gamma(G)$  ( $\bar{epa}_\gamma(G)$ , respectively), to be

- $epa_\gamma(G) = \min\{pa_\gamma(u, v) \mid u, v \in V(G), uv \in E(G)\}$ ,
  - $\bar{epa}_\gamma(G) = \min\{pa_\gamma(u, v) \mid u, v \in V(G), uv \notin E(G)\}$ , and
- ▷ the *upper e-path addition* (*upper  $\bar{e}$ -path addition*) *number with respect to domination*, denoted  $Epa_\gamma(G)$  ( $\bar{Epa}_\gamma(G)$ , respectively), to be
- $Epa_\gamma(G) = \max\{pa_\gamma(u, v) \mid u, v \in V(G), uv \in E(G)\}$ ,
  - $\bar{Epa}_\gamma(G) = \max\{pa_\gamma(u, v) \mid u, v \in V(G), uv \notin E(G)\}$ .

If  $G$  is complete, then we write  $\bar{Epa}_\gamma(G) = \bar{epa}_\gamma(G) = \infty$ , and if  $G$  is edgeless then  $epa_\gamma(G) = Epa_\gamma(G) = \infty$ . In what follows the subscript  $\gamma$  will be omitted from the notation.

The remainder of this paper is organized as follows. In Section 2, we prove that  $1 \leq epa(G) \leq 3$  and  $2 \leq Epa(G) \leq 3$ , and we present necessary and sufficient conditions for  $pa(u, v) = i$ ,  $i = 1, 2, 3$ , where  $uv \in E(G)$ . In Section 3, we show that  $1 \leq \bar{epa}(G) \leq \bar{Epa}(G) \leq 5$ , and we give necessary and sufficient conditions for  $\bar{epa}(G) = \bar{Epa}(G) = j$ ,  $1 \leq j \leq 5$ . We conclude in Section 4 with open problems.

We end this section with some known results which will be useful in proving our main results.

**Lemma 1** [2]. *If  $G$  is a graph and  $H$  is any graph obtained from  $G$  by subdividing some edges of  $G$ , then  $\gamma(H) \geq \gamma(G)$ .*

**Lemma 2.** *Let  $G$  be a graph and  $v \in V(G)$ .*

- (i) [5] *If  $v$  is  $\gamma$ -bad, then  $\gamma(G - v) = \gamma(G)$ .*
- (ii) [3]  *$v$  is  $\gamma$ -critical if and only if  $\gamma(G - v) = \gamma(G) - 1$ .*
- (iii) [5] *If  $v$  is  $\gamma$ -critical, then all its neighbors are  $\gamma$ -bad vertices of  $G - v$ .*
- (iv) [11] *If  $e \in E(\bar{G})$ , then  $\gamma(G) - 1 \leq \gamma(G + e) \leq \gamma(G)$ .*

In most cases, Lemma 2 will be used in the sequel without specific reference.

## 2. THE ADJACENT CASE

The aim of this section is to prove that  $1 \leq pa(u, v) \leq 3$  and to find necessary and sufficient conditions for  $pa(u, v) = i$ ,  $i = 1, 2, 3$ , where  $uv \in E(G)$ .

**Observation 3.** *If  $u$  and  $v$  are adjacent vertices of a graph  $G$ , then  $\gamma(G) = \gamma(G_{u,v,0}) \leq \gamma(G_{u,v,k}) \leq \gamma(G_{u,v,k+1})$  for  $k \geq 1$ .*

**Proof.** The equality  $\gamma(G) = \gamma(G_{u,v,0})$  is obvious. For any  $\gamma$ -set  $M$  of  $G_{u,v,1}$  both  $M_u = (M \setminus \{x_1\}) \cup \{u\}$  and  $M_v = (M \setminus \{x_1\}) \cup \{v\}$  are dominating sets of  $G$ , and at

least one of them is a  $\gamma$ -set of  $G_{u,v,1}$ . Hence  $\gamma(G) \leq \min\{|M_u|, |M_v|\} = \gamma(G_{u,v,1})$ . The rest follows by Lemma 1. ■

**Theorem 4.** *Let  $u$  and  $v$  be adjacent vertices of a graph  $G$ . Then  $\gamma(G) \leq \gamma(G_{u,v,1}) \leq \gamma(G) + 1$  and the following is true.*

- (i)  $\gamma(G) = \gamma(G_{u,v,1})$  if and only if at least one of  $u$  and  $v$  is a  $\gamma$ -good vertex of  $G$ .
- (ii)  $\gamma(G_{u,v,1}) = \gamma(G) + 1$  if and only if both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ .

**Proof.** The left side inequality follows by Observation 3. If  $D$  is a  $\gamma$ -set of  $G$ , then  $D \cup \{x_1\}$  is a dominating set of  $G_{u,v,1}$ , which implies  $\gamma(G_{u,v,1}) \leq \gamma(G) + 1$ .

If at least one of  $u$  and  $v$  belongs to some  $\gamma$ -set  $D_1$  of  $G$ , then  $D_1$  is a dominating set of  $G_{u,v,1}$ . This clearly implies  $\gamma(G) = \gamma(G_{u,v,1})$ .

Let now both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ , and suppose that  $\gamma(G_{u,v,1}) = \gamma(G)$ . In this case for any  $\gamma$ -set  $M$  of  $G_{u,v,1}$  is fulfilled  $u, v \notin M$  and  $x_1 \in M$ . But then  $(M \setminus \{x_1\}) \cup \{u\}$  is a  $\gamma$ -set for both  $G$  and  $G_{u,v,1}$ , a contradiction. ■

**Corollary 5.** *Let  $G$  be a graph with edges. Then  $\text{Epa}(G) \geq 2$  and  $\text{epa}(G) = 1$  if and only if the set of all  $\gamma$ -bad vertices of  $G$  is neither empty nor independent.*

**Theorem 6.** *Let  $u$  and  $v$  be adjacent vertices of a graph  $G$ . Then  $\gamma(G) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ . Moreover,*

- (A)  $\gamma(G_{u,v,2}) = \gamma(G) + 1$  if and only if at least one of the following holds:
  - (i) both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ ,
  - (ii) at least one of  $u$  and  $v$  is  $\gamma$ -good,  $u, v \notin V^-(G)$  and each  $\gamma$ -set of  $G$  contains at most one of  $u$  and  $v$ .
- (B)  $\gamma(G_{u,v,2}) = \gamma(G)$  if and only if at least one of the following is true:
  - (iii) there exists a  $\gamma$ -set of  $G$  which contains both  $u$  and  $v$ ,
  - (iv) at least one of  $u$  and  $v$  is in  $V^-(G)$ .

**Proof.** The left side inequality follows by Observation 3. If  $D$  is an arbitrary  $\gamma$ -set of  $G$ , then  $D \cup \{x_1\}$  is a dominating set of  $G_{u,v,2}$ . Hence  $\gamma(G_{u,v,2}) \leq \gamma(G) + 1$ .

(A)  $\Rightarrow$  Assume that the equality  $\gamma(G_{u,v,2}) = \gamma(G) + 1$  holds. By Theorem 4 we know that  $\gamma(G_{u,v,1}) \in \{\gamma(G), \gamma(G) + 1\}$ . If  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ , then again by Theorem 4, both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ . So let  $\gamma(G) = \gamma(G_{u,v,1})$ . Then at least one of  $u$  and  $v$  is a  $\gamma$ -good vertex of  $G$  (Theorem 4). Clearly there is no  $\gamma$ -set of  $G$  which contains both  $u$  and  $v$ . If  $u \in V^-(G)$  and  $U$  is a  $\gamma$ -set of  $G - u$ , then  $U \cup \{x_1\}$  is a dominating set of  $G_{u,v,2}$  and  $|U \cup \{x_1\}| = \gamma(G)$ , a contradiction. Thus  $u, v \notin V^-(G)$ .

(A)  $\Leftarrow$  If both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ , then  $\gamma(G_{u,v,1}) = \gamma(G) + 1$  (Theorem 4). But we know that  $\gamma(G_{u,v,1}) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ ; hence

$\gamma(G_{u,v,2}) = \gamma(G) + 1$ . Finally let (ii) hold and  $M$  be a  $\gamma$ -set of  $G_{u,v,2}$ . If  $x_1, x_2 \notin M$ , then  $u, v \in M$  which leads to  $\gamma(G_{u,v,2}) > \gamma(G)$ . If  $x_1, x_2 \in M$ , then  $(M \setminus \{x_1, x_2\}) \cup \{u, v\}$  is a dominating set of  $G$  of cardinality more than  $\gamma(G)$ . Now let without loss of generality  $x_1 \in M$  and  $x_2 \notin M$ . If  $M \setminus \{x_1\}$  is a dominating set of  $G$ , then  $\gamma(G) + 1 \leq |M| = \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ . So, let  $M \setminus \{x_1\}$  be no dominating set of  $G$ . Hence  $M \setminus \{x_1\}$  is a dominating set of  $G - u$ . Since  $u \notin V^-(G)$ ,  $\gamma(G) \leq \gamma(G - u) \leq |M \setminus \{x_1\}| < \gamma(G_{u,v,2})$ .

(B)  $\Rightarrow$  Let  $\gamma(G_{u,v,2}) = \gamma(G)$ . Suppose that neither (iii) nor (iv) is valid. Hence  $u, v \notin V^-(G)$  and no  $\gamma$ -set of  $G$  contains both  $u$  and  $v$ . But then at least one of (i) and (ii) holds, and from (A) we conclude that  $\gamma(G_{u,v,2}) = \gamma(G) + 1$ , a contradiction.

(B)  $\Leftarrow$  Let at least one of (iii) and (iv) be hold. Then neither (i) nor (ii) is fulfilled. Now by (A) we have  $\gamma(G_{u,v,2}) \neq \gamma(G) + 1$ . Since  $\gamma(G) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ , we obtain  $\gamma(G) = \gamma(G_{u,v,2})$ . ■

The *independent domination number* of a graph  $G$ , denoted by  $i(G)$ , is the minimum size of an independent dominating set of  $G$ . It is obvious that  $i(G) \geq \gamma(G)$ . In a graph  $G$ ,  $i(G)$  is *strongly equal* to  $\gamma(G)$ , written  $i(G) \equiv \gamma(G)$ , if each  $\gamma$ -set of  $G$  is independent. It remains an open problem to characterize the graphs  $G$  with  $i(G) \equiv \gamma(G)$  [7].

**Corollary 7.** *Let  $G$  be a graph with edges. Then (a)  $\text{epa}(G) \geq 2$  if and only if the set of all  $\gamma$ -bad vertices is either empty or independent, and (b)  $\text{Epa}(G) = 2$  if and only if  $i(G) \equiv \gamma(G)$ .*

**Proof.** (a) Immediately by Corollary 5.

(b)  $\Rightarrow$  Let  $\text{Epa}(G) = 2$ . If  $D$  is a  $\gamma$ -set of  $G$  and  $u, v \in D$  are adjacent, then  $D$  is a dominating set of  $G_{u,v,2}$ , a contradiction.

(b)  $\Leftarrow$  Let all  $\gamma$ -sets of  $G$  be independent. Suppose  $u \in V^-(G)$  and  $D$  is a  $\gamma$ -set of  $G - u$ . Then  $D_1 = D \cup \{v\}$  is a  $\gamma$ -set of  $G$ , where  $v$  is any neighbor of  $u$ . But  $D_1$  is not independent. Hence  $V^-(G)$  is empty. Thus, for any 2 adjacent vertices  $u$  and  $v$  of  $G$  is fulfilled either (A)(i) or (A)(ii) of Theorem 6. Therefore  $\text{Epa}(G) \leq 2$ . The result now follows by Corollary 5. ■

Denote by  $\mathbb{Z}_n = \{0, 1, \dots, n-1\}$  the additive group of order  $n$ . Let  $S$  be a subset of  $\mathbb{Z}_n$  such that  $0 \notin S$  and  $x \in S$  implies  $-x \in S$ . The *circulant graph* with distance set  $S$  is the graph  $C(n; S)$  with vertex set  $\mathbb{Z}_n$  and vertex  $x$  adjacent to vertex  $y$  if and only if  $x - y \in S$ .

Let  $n \geq 3$  and  $k \in \mathbb{Z}_n \setminus \{0\}$ . The *generalized Petersen graph*  $P(n, k)$  is the graph on the vertex-set  $\{x_i, y_i \mid i \in \mathbb{Z}_n\}$  with adjacencies  $x_i x_{i+1}$ ,  $x_i y_i$ , and  $y_i y_{i+k}$  for all  $i$ .

**Example 8.** A special case of graphs  $G$  with  $\text{Epa}(T) = 2$  are graphs for which each  $\gamma$ -set is efficient dominating (an efficient dominating set in a graph  $G$  is a

set  $S$  such that  $\{N[s] \mid s \in S\}$  is a partition of  $V(G)$ ). We list several examples of such graphs [10].

- (a) A *crown graph*  $H_{n,n}$ ,  $n \geq 3$ , which is obtained from the complete bipartite graph  $K_{n,n}$  by removing a perfect matching.
- (b) Circulant graphs  $G = C(n = (2k+1)t; \{1, \dots, k\} \cup \{n-1, \dots, n-k\})$ , where  $k, t \geq 1$ .
- (c) Circulant graphs  $G = C(n; \{\pm 1, \pm s\})$ , where  $2 \leq s \leq n-2$ ,  $s \neq n/2$ ,  $5 \mid n$  and  $s \equiv \pm 2 \pmod{5}$ .
- (d) The *generalized Petersen graph*  $P(n, k)$ , where  $n \equiv 0 \pmod{4}$  and  $k$  is odd.

**Theorem 9.** *If  $u$  and  $v$  are adjacent vertices of a graph  $G$ , then  $\gamma(G_{u,v,3}) = \gamma(G) + 1$ .*

**Proof.** If  $D$  is a  $\gamma$ -set of  $G$ , then  $D \cup \{x_2\}$  is a dominating set of  $G$ . Hence  $\gamma(G_{u,v,3}) \leq \gamma(G) + 1$ .

Let  $M$  be a  $\gamma$ -set of  $G_{u,v,3}$ . Then at least one of  $x_1, x_2$  and  $x_3$  is in  $M$ . If  $x_2 \in M$ , then clearly  $\gamma(G_{u,v,3}) = \gamma(G) + 1$ . If  $x_2 \notin M$  and  $x_1, x_3 \in M$ , then  $(M \setminus \{x_1, x_3\}) \cup \{u\}$  is a dominating set of  $G$ . If  $x_2, x_3 \notin M$  and  $x_1 \in M$ , then  $v \in M$  and  $M \setminus \{x_1\}$  is a dominating set of  $G$ . All this leads to  $\gamma(G_{u,v,3}) = \gamma(G) + 1$ . ■

**Corollary 10.** *Let  $G$  be a graph with edges. Then  $\text{epa}(G) \leq \text{Epa}(G) \leq 3$ . Moreover,  $\text{Epa}(G) = 3$  if and only if  $G$  has a  $\gamma$ -set that is not independent, and  $\text{epa}(G) = 3$  if and only if for each pair of adjacent vertices  $u$  and  $v$  at least one of the following is valid.*

- (i) *There exists a  $\gamma$ -set of  $G$  which contains both  $u$  and  $v$ .*
- (ii) *At least one of  $u$  and  $v$  is in  $V^-(G)$ .*

**Proof.** By Corollary 5 and Theorem 9 we have  $1 \leq \text{epa}(G) \leq \text{Epa}(G) \leq 3$  and  $2 \leq \text{Epa}(G)$ . Since  $\text{Epa}(G) = 2$  if and only if  $i(G) \equiv \gamma(G)$  (by Corollary 7),  $\text{Epa}(G) = 3$  if and only if  $G$  has a  $\gamma$ -set that is not independent.

Clearly  $\text{epa}(G) = 3$  if and only if  $\gamma(G_{u,v,2}) = \gamma(G)$  for each pair of adjacent vertices  $u$  and  $v$  of  $G$ . Then because of Theorem 6(B), we have that  $\text{epa}(G) = 3$  if and only if for each pair of adjacent vertices  $u$  and  $v$  of  $G$  at least one of (i) and (ii) holds. ■

**Corollary 11.** *Let  $G$  be a graph with edges. If  $V^-(G)$  has a subset which is a vertex cover of  $G$ , then  $\text{epa}(G) = 3$ . In particular, if  $G$  is a vc-graph then  $\text{epa}(G) = 3$ .*

We define the following classes of graphs  $G$  with  $\Delta(G) \geq 1$ .

- $\mathcal{A} = \{G \mid \text{epa}(G) = 3\}$ ,

- $\mathcal{A}_1 = \{G \mid V^-(G) \text{ is a vertex cover of } G\}$ ,
- $\mathcal{A}_2 = \{G \mid \text{each two adjacent vertices belongs to some } \gamma\text{-set of } G\}$ ,
- $\mathcal{A}_3 = \{G \mid G \text{ is a vc-graph}\}$ .

Clearly,  $\mathcal{A}_3 \subseteq \mathcal{A}_1$  and by Corollaries 10 and 11,  $\mathcal{A}_1 \cup \mathcal{A}_2 \subseteq \mathcal{A}$ . These relationships are illustrated in the Venn diagram in Figure 1(left). To continue we relabel this diagram in six regions  $\mathbf{R}_0$ – $\mathbf{R}_5$  as shown in Figure 1(right). In what follows in this section we show that none of  $\mathbf{R}_0$ – $\mathbf{R}_5$  is empty. The *corona* of a graph  $H$  is the graph  $G = H \circ K_1$  obtained from  $H$  by adding a degree-one neighbor to every vertex of  $H$ . If  $F$  and  $H$  are disjoint graphs,  $v_F \in V(F)$  and  $v_H \in V(H)$ , then the *coalescence*  $(F \cdot H)(v_F, v_H : v)$  of  $F$  and  $H$  via  $v_F$  and  $v_H$ , is the graph obtained from the union of  $F$  and  $H$  by identifying  $v_F$  and  $v_H$  in a vertex labeled  $v$ .




Figure 1. Left: Classes of graphs with  $\text{epa} = 3$ . Right: Regions of Venn diagram.

**Remark 12.** It is easy to see that all the following hold.

- If  $H$  is a connected graph of order  $n \geq 2$ , then  $G = H \circ K_1 \in \mathbf{R}_0$ .
- Let  $G_k^1$  be a graph obtained from the cycle  $C_{3k+1} : x_0, x_1, x_2, \dots, x_{3k}, x_0$ ,  $k \geq 2$ , by adding a vertex  $y$  and edges  $yx_0, yx_2$ . Then  $\gamma(G_k^1) = k + 1$ ,  $G_k^1$  is  $\gamma$ -excellent,  $V^-(G_k^1) = \{x_0, x_2\} \cup \bigcup_{r=1}^{k-1} \{x_{3r+1}, x_{3r+2}\}$  is a vertex cover of  $G$ , and there is no  $\gamma$ -set of  $G_k^1$  that contains both  $x_{3r+1}$  and  $x_{3r+2}$ . Thus  $G_k^1$  is in  $\mathbf{R}_1$ .
- The graph  $H_{10}$  depicted in Figure 2 is in  $\mathcal{A}_3$  and  $\gamma(H_{10}) = 3$  [1]. It is obvious that no  $\gamma$ -set of  $H_{10}$  contains both  $u$  and  $v$ . Hence  $H_{10} \in \mathbf{R}_2$ . Consider now the graph  $G_k^2 = (C_{3k+1} \cdot H_{10})(x_0, w : z)$ , where  $C_{3k+1} : x_0, x_1, x_2, \dots, x_{3k}, x_0$ ,  $k \geq 2$ , is a cycle on  $3k + 1$  vertices and  $w$  is any of the two common neighbors of  $u$  and  $v$  in  $H_{10}$ . Since both  $C_{3k+1}$  and  $H_{10}$  are vc-graphs, by [4] we have that  $G_k^2$  is vc-graph and  $\gamma(G_k^2) = \gamma(C_{3k+1}) + \gamma(H_{10}) - 1$ . Let  $D$  be an arbitrary  $\gamma$ -set of  $G_k^2$ ,  $D_1 = D \cap V(H_{10})$  and  $D_2 = D \cap V(C_{3k+1})$ . Then exactly one of the following holds.
  - $z \in D$ ,  $D_1$  is a  $\gamma$ -set of  $H_{10}$  and  $D_2$  is a  $\gamma$ -set of  $C_{3k+1}$ .
  - $z \notin D$ ,  $D_1$  is a  $\gamma$ -set of  $H_{10}$  and  $D_2 \cup \{x_0\}$  is a  $\gamma$ -set of  $C_{3k+1}$ .

(c)  $z \notin D$ ,  $D_1 \cup \{w\}$  is a  $\gamma$ -set of  $H_{10}$  and  $D_2$  is a  $\gamma$ -set of  $C_{3k+1}$ .

Since no  $\gamma$ -set of  $H_{10}$  contains both  $u$  and  $v$ , by (a), (b) and (c) we conclude that at most one of  $u$  and  $v$  is in  $D$ . Thus  $G_k^2 \in \mathbf{R}_2$ .

(iv)  $C_{3k+1} \in \mathbf{R}_3$  for all  $k \geq 1$ .

(v)  $K_{2,n} \in \mathbf{R}_4$  for all  $n \geq 3$ .

(vi)  $K_{n,n} \in \mathbf{R}_5$  for all  $n \geq 3$ .

Thus all regions  $\mathbf{R}_0, \mathbf{R}_1, \mathbf{R}_2, \mathbf{R}_3, \mathbf{R}_4, \mathbf{R}_5$  are nonempty.




Figure 2. Graph  $H_{10}$  is in  $\mathbf{R}_2$ .

### 3. THE NONADJACENT CASE

In this section we show that  $1 \leq \bar{epa}(G) \leq \overline{Epa}(G) \leq 5$  and we obtain necessary and sufficient conditions for  $\bar{epa}(G) = \overline{Epa}(G) = j$ ,  $1 \leq j \leq 5$ .

We begin with an easy observation which is an immediate consequence by Lemma 2(iv) and Lemma 1.

**Observation 13.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . Then  $\gamma(G) - 1 \leq \gamma(G_{u,v,0}) \leq \gamma(G)$  and  $\gamma(G_{u,v,k}) \leq \gamma(G_{u,v,k+1})$  for  $k \geq 0$ .*

**Theorem 14.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . Then  $\gamma(G) - 1 \leq \gamma(G_{u,v,1}) \leq \gamma(G) + 1$ . Moreover,*

- (i)  $\gamma(G) - 1 = \gamma(G_{u,v,1})$  if and only if  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ .
- (ii)  $\gamma(G_{u,v,1}) = \gamma(G) + 1$  if and only if both  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ ,  $u \notin V^-(G - v)$  and  $v \notin V^-(G - u)$ . If  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ , then  $x_1 \in V^-(G_{u,v,1})$ .

**Proof.** The left side inequality follows by Observation 13.

(i)  $\Rightarrow$  Assume the equality  $\gamma(G) - 1 = \gamma(G_{u,v,1})$  holds and let  $M$  be any  $\gamma$ -set of  $G_{u,v,1}$ . Then at least one and not more than two of  $x_1, u$  and  $v$  must be in  $M$ . Hence  $M_1 = (M \setminus \{x_1\}) \cup \{u, v\}$  is a dominating set of  $G$  and  $\gamma(G) \leq |M_1| \leq |M| + 1 = \gamma(G_{u,v,1}) + 1 = \gamma(G)$ . This immediately implies that  $M_1$  is a  $\gamma$ -set of  $G$ . Hence  $x_1 \in M$  and  $pn[x_1, M] = \{x_1, u, v\}$ . Since  $M_1 \setminus \{u, v\}$  is a dominating set of  $G - \{u, v\}$ , we have  $\gamma(G) - 2 \leq \gamma(G - \{u, v\}) \leq |M_1 \setminus \{u, v\}| = \gamma(G) - 2$ .



(i)  $\Leftarrow$  Suppose now  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ . Then for any  $\gamma$ -set  $U$  of  $G - \{u, v\}$ , the set  $U \cup \{x_1\}$  is a dominating set of  $G_{u,v,1}$ . This leads to  $\gamma(G_{u,v,1}) \leq |U \cup \{x_1\}| = \gamma(G) - 1 \leq \gamma(G_{u,v,1})$ .

Now we will prove the right side inequality. Let  $D$  be any  $\gamma$ -set of  $G$ . If at least one of  $u$  and  $v$  is in  $D$ , then  $D$  is a dominating set  $G_{u,v,1}$  and  $\gamma(G_{u,v,1}) \leq \gamma(G)$ . So, let neither  $u$  nor  $v$  belong to some  $\gamma$ -set of  $G$ . Then  $D \cup \{x_1\}$  is a dominating set of  $G_{u,v,1}$  and  $\gamma(G_{u,v,1}) \leq \gamma(G) + 1$ .

(ii)  $\Rightarrow$  Assume that  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ . Then  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$  and for any  $\gamma$ -set  $D$  of  $G$ ,  $D \cup \{x_1\}$  is a  $\gamma$ -set of  $G_{u,v,1}$ . Hence  $x_1 \in V^-(G_{u,v,1})$ . Suppose  $u \in V^-(G - v)$  and let  $U$  be a  $\gamma$ -set of  $G - \{u, v\}$ . Then  $U_1 = U \cup \{x_1\}$  is a dominating set of  $G_{u,v,1}$  and  $\gamma(G) + 1 = \gamma(G_{u,v,1}) \leq |U_1| = 1 + \gamma((G - v) - u) = \gamma(G - v) = \gamma(G)$ , a contradiction. Thus  $u \notin V^-(G - v)$  and by symmetry,  $v \notin V^-(G - u)$ .

(ii)  $\Leftarrow$  Let both  $u$  and  $v$  be  $\gamma$ -bad vertices of  $G$ ,  $u \notin V^-(G - v)$  and  $v \notin V^-(G - u)$ . Hence  $\gamma(G - \{u, v\}) \geq \gamma(G)$ . Consider any  $\gamma$ -set  $M$  of  $G_{u,v,1}$ . If one of  $u$  and  $v$  belongs to  $M$ , then  $\gamma(G) + 1 = \gamma(G_{u,v,1})$ . So, let  $x_1$  is in each  $\gamma$ -set of  $G_{u,v,1}$ . But then  $pn[x_1, M] = \{x_1, u, v\}$ . Hence  $\gamma(G_{u,v,1}) - 1 = \gamma(G - \{u, v\}) \geq \gamma(G) \geq \gamma(G_{u,v,1}) - 1$ . ■

**Corollary 15.** *Let  $G$  be a noncomplete graph. Then  $1 \leq \bar{epa}(G) \leq \bar{Epa}(G)$  and the following assertions hold.*

- (i)  $\bar{epa}(G) = 1$  if and only if there are nonadjacent  $\gamma$ -bad vertices  $u$  and  $v$  of  $G$  such that  $u \notin V^-(G - v)$  and  $v \notin V^-(G - u)$ .
- (ii)  $\bar{Epa}(G) = 1$  if and only if  $\gamma(G) = 1$ .

**Proof.** Observation 13 implies  $1 \leq \bar{epa}(G)$ .

(i) Immediately by Theorem 14.

(ii) If  $\gamma(G) = 1$ , then clearly  $\bar{Epa}(G) = 1$ . If  $\gamma(G) \geq 2$ , then  $G$  has 2 non-adjacent vertices at least one of which is  $\gamma$ -good. By Theorem 14,  $\bar{Epa}(G) \geq 2$ . ■

**Theorem 16.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . Then  $\gamma(G) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ . Moreover,*

- (C)  $\gamma(G_{u,v,2}) = \gamma(G)$  if and only if one of the following holds.
  - (i) There is a  $\gamma$ -set of  $G$  which contains both  $u$  and  $v$ .
  - (ii) At least one of  $u$  and  $v$  is in  $V^-(G)$ .
- (D)  $\gamma(G_{u,v,2}) = \gamma(G) + 1$  if and only if  $u, v \notin V^-(G)$  and any  $\gamma$ -set of  $G$  contains at most one of  $u$  and  $v$ .

**Proof.** For any  $\gamma$ -set  $D$  of  $G$ ,  $D \cup \{x_2\}$  is a dominating set of  $G_{u,v,2}$ . Hence  $\gamma(G_{u,v,2}) \leq \gamma(G) + 1$ . Suppose  $\gamma(G_{u,v,2}) \leq \gamma(G) - 1$  and let  $M$  be a  $\gamma$ -set of  $G_{u,v,2}$ . Then at least one of  $x_1$  and  $x_2$  is in  $M$ . If  $x_1, x_2 \in M$ , then  $M_1 = (M \setminus \{x_1, x_2\}) \cup$

$\{u, v\}$  is a dominating set of  $G$  and  $|M_1| \leq \gamma(G_{u,v,2})$ , a contradiction. So let without loss of generality,  $x_1 \in M$  and  $x_2 \notin M$ . If  $u \in M$  or  $v \in M$ , then again  $M_1$  is a dominating set of  $G$  and  $|M_1| \leq \gamma(G_{u,v,2})$ , a contradiction. Thus  $x_1 \in M$  and  $u, v \notin M$ . But then  $(M \setminus \{x_1\}) \cup \{u\}$  is a dominating set of  $G$ , contradicting  $\gamma(G_{u,v,2}) < \gamma(G)$ . Thus  $\gamma(G) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ .

(C)  $\Rightarrow$  Let  $\gamma(G_{u,v,2}) = \gamma(G)$ . Assume that neither (i) nor (ii) hold. Let  $M$  be a  $\gamma$ -set of  $G_{u,v,2}$ . If  $x_1, x_2 \in M$ , then  $M_1 = (M \setminus \{x_1, x_2\}) \cup \{u, v\}$  is a dominating set of  $G$  of cardinality not more than  $\gamma(G)$  and  $u, v \in M_1$ , a contradiction. Let without loss of generality  $x_1 \in M$  and  $x_2 \notin M$ . Since  $M \setminus \{x_1\}$  is no dominating set of  $G$ ,  $u \in pn[x_1, M]$ . But then  $M_3 = (M \setminus \{x_1\}) \cup \{u\}$  is a  $\gamma$ -set of  $G$  and  $u \in V^-(G)$ , a contradiction. Thus at least one of (i) and (ii) is valid.

(C)  $\Leftarrow$  If both  $u$  and  $v$  belong to some  $\gamma$ -set  $D$  of  $G$ , then  $D$  is a dominating set of  $G_{u,v,2}$ . Hence  $\gamma(G_{u,v,2}) = \gamma(G)$ . Finally let  $u \in V^-(G)$  and  $D$  a  $\gamma$ -set of  $G - u$ . Then  $D \cup \{x_1\}$  is a dominating set of  $G_{u,v,2}$  of cardinality  $\gamma(G)$ . Thus  $\gamma(G_{u,v,2}) = \gamma(G)$ .

(D) Immediately by (C) and  $\gamma(G) \leq \gamma(G_{u,v,2}) \leq \gamma(G) + 1$ .  $\blacksquare$

**Corollary 17.** *Let  $G$  be a noncomplete graph. Then the following assertions hold.*

- (i)  $\bar{epa}(G) \leq 2$  if and only if there are nonadjacent vertices  $u, v \in V(G) \setminus V^-(G)$  such that any  $\gamma$ -set of  $G$  contains at most one of them.
- (ii)  $\bar{Epa}(G) = 2$  if and only if  $\gamma(G) \geq 2$  and each  $\gamma$ -set of  $G$  is a clique.

**Proof.** (i) Immediately by Theorem 16.

(ii)  $\Rightarrow$  Let  $\bar{Epa}(G) = 2$ . By Corollary 15,  $\gamma(G) \geq 2$ . Suppose  $G$  has a  $\gamma$ -set, say  $D$ , which is not a clique. Then there are nonadjacent  $u, v \in D$ . By Theorem 16(C),  $\gamma(G_{u,v,2}) = \gamma(G)$ , which contradict  $\bar{Epa}(G) = 2$ . Thus, each  $\gamma$ -set of  $G$  is a clique.

(ii)  $\Leftarrow$  Let  $\gamma(G) \geq 2$  and let each  $\gamma$ -set of  $G$  be a clique. If  $G$  has a vertex  $z \in V^-(G)$  and  $M_z$  is a  $\gamma$ -set of  $G - z$ , then  $M = M_z \cup \{z\}$  is a  $\gamma$ -set of  $G$  and  $z$  is an isolated vertex of the graph induced by  $M$ , a contradiction. Thus  $V^-(G)$  is empty. Now by Theorem 16(D),  $\bar{Epa}(G) = 2$ .  $\blacksquare$

**Example 18.** The join of two graphs  $G_1$  and  $G_2$  with disjoint vertex sets is the graph, denoted by  $G_1 + G_2$ , with the vertex set  $V(G_1) \cup V(G_2)$  and edge set  $E(G_1) \cup E(G_2) \cup \{uv \mid u \in V(G_1), v \in V(G_2)\}$ . Let  $\gamma(G_i) \geq 3$ ,  $i = 1, 2$ . Then  $\gamma(G_1 + G_2) = 2$  and each  $\gamma$ -set of  $G_1 + G_2$  contains exactly one vertex of  $G_i$ ,  $i = 1, 2$ . Hence  $\bar{Epa}(G_1 + G_2) = 2$ . In particular,  $\bar{Epa}(K_{m,n}) = 2$  when  $m, n \geq 3$ .

**Theorem 19.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . Then  $\gamma(G) \leq \gamma(G_{u,v,3}) \leq \gamma(G) + 1$ . Moreover,  $\gamma(G_{u,v,3}) = \gamma(G)$  if and only if at least one of the following holds.*

- (i)  $u \in V^-(G)$  and  $v$  is a  $\gamma$ -good vertex of  $G - u$ ,
- (ii)  $v \in V^-(G)$  and  $u$  is a  $\gamma$ -good vertex of  $G - v$ .

**Proof.** If  $D$  is a dominating set of  $G$ , then  $D \cup \{x_2\}$  is a dominating set of  $G_{u,v,3}$ . Hence  $\gamma(G_{u,v,3}) \leq \gamma(G) + 1$ . We already know that  $\gamma(G) \leq \gamma(G_{u,v,2})$  and  $\gamma(G_{u,v,2}) \leq \gamma(G_{u,v,3})$ . But then  $\gamma(G) \leq \gamma(G_{u,v,3})$ .

$\Rightarrow$  Let  $\gamma(G_{u,v,3}) = \gamma(G)$  and let  $M$  be a  $\gamma$ -set of  $G_{u,v,3}$  such that  $Q = M \cap \{x_1, x_2, x_3\}$  has minimum cardinality. Clearly  $|Q| = 1$ . If  $\{x_2\} = Q$ , then  $M \setminus \{x_2\}$  is a dominating set of  $G$ , contradicting  $\gamma(G_{u,v,3}) = \gamma(G)$ . Let without loss of generality  $\{x_1\} = Q$ . This implies  $v \in M$ ,  $x_3 \in pn[v, M]$  and  $pn[x_1, M] = \{u, x_1, x_2\}$ . Then  $M_2 = (M \setminus \{x_1\}) \cup \{u\}$  is a  $\gamma$ -set of  $G$ ,  $pn[u, M_2] = \{u\}$  and  $v \in M_2$ ; hence (i) holds.

$\Leftarrow$  Let without loss of generality (i) is true. Then there is a  $\gamma$ -set  $D$  of  $G$  such that  $u, v \in D$  and  $D \setminus \{u\}$  is a  $\gamma$ -set of  $G - u$ . But then  $(D \setminus \{u\}) \cup \{x_1\}$  is a dominating set of  $G_{u,v,3}$ , which implies  $\gamma(G) \geq \gamma(G_{u,v,3})$ . ■

**Corollary 20.** Let  $G$  be a noncomplete graph. Then the following holds.

- (E)  $\bar{epa}(G) \leq 3$  if and only if there is a pair of nonadjacent vertices  $u$  and  $v$  such that neither (i) nor (ii) is valid, where
  - (i)  $u \in V^-(G)$  and  $v$  is a  $\gamma$ -good vertex of  $G - u$ ,
  - (ii)  $v \in V^-(G)$  and  $u$  is a  $\gamma$ -good vertex of  $G - v$ .
- (F)  $\bar{epa}(G) = \bar{Epa}(G) = 3$  if and only if all vertices of  $G$  are  $\gamma$ -good,  $V^-(G)$  is empty and for every 2 nonadjacent vertices  $u$  and  $v$  of  $G$  there is a  $\gamma$ -set of  $G$  which contains them both.

**Proof.** (F)  $\Rightarrow$  Let  $\bar{epa}(G) = \bar{Epa}(G) = 3$ . If  $u \in V^-(G)$  and  $D$  is a  $\gamma$ -set of  $G - u$ , then for  $u$  and each  $v \in D$  is fulfilled (i) of Theorem 19. But then  $\bar{Epa}(G) \neq 3$ , a contradiction. So,  $V^-(G)$  is empty. Suppose that  $G$  has  $\gamma$ -bad vertices. Then there is a  $\gamma$ -bad vertex which is nonadjacent to some other vertex of  $G$ . But Theorem 16(D) implies  $\bar{epa}(G) < 3$ , a contradiction. Thus all vertices of  $G$  are  $\gamma$ -good. Now let  $u, v \in V(G)$  be nonadjacent. If there is no  $\gamma$ -set of  $G$  which contains both  $u$  and  $v$ , then by Theorem 16(D) we have  $\gamma(G_{u,v,2}) = \gamma(G) + 1$ , a contradiction.

(F)  $\Leftarrow$  Let  $V^-(G)$  be empty and for each pair  $u, v$  of nonadjacent vertices of  $G$  there is a  $\gamma$ -set  $D_{uv}$  of  $G$  with  $u, v \in D_{uv}$ . By Theorem 19,  $\gamma(G_{u,v,3}) = \gamma(G) + 1$ , and by Theorem 16,  $\gamma(G_{u,v,2}) = \gamma(G)$ . Hence  $pa(u, v) = 3$ . ■

**Example 21.** Denote by  $\mathcal{U}$  the class of all graphs  $G$  with  $\bar{epa}(G) = \bar{Epa}(G) = 3$ . Then all the following holds. (a) Circulant graphs  $C(2k + 1; \{\pm 1, \pm 2, \dots, \pm(k - 1)\}) \in \mathcal{U}$  for all  $k \geq 1$ . (b) Let  $G$  be a disconnected graph. Then  $G \in \mathcal{U}$  if and only if  $G$  has no isolated vertices and each its component is either in  $\mathcal{U}$  or is complete.

**Theorem 22.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . Then  $\gamma(G) \leq \gamma(G_{u,v,4}) \leq \gamma(G) + 2$ . Moreover, the following assertions are valid.*

- (G)  $\gamma(G_{u,v,4}) = \gamma(G) + 2$  if and only if  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ .
- (H) If  $\gamma(G_{u,v,1}) = \gamma(G)$  and  $\gamma(G_{u,v,i}) = \gamma(G) + 1$  for some  $i \in \{2, 3\}$ , then  $\gamma(G_{u,v,4}) = \gamma(G) + 1$ .
- (I) Let  $\gamma(G_{u,v,3}) = \gamma(G)$ . Then  $\gamma(G_{u,v,4}) \leq \gamma(G) + 1$  and the equality holds if and only if  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$ .
- (J)  $\gamma(G_{u,v,4}) = \gamma(G)$  if and only if  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ .

**Proof.** Since  $\gamma(G) \leq \gamma(G_{u,v,3})$  (by Theorem 19) and  $\gamma(G_{u,v,3}) \leq \gamma(G_{u,v,4})$  (by Observation 13), we have  $\gamma(G) \leq \gamma(G_{u,v,4})$ . Let  $S$  be a  $\gamma$ -set of  $G$ . Then  $S \cup \{x_1, x_4\}$  is a dominating set of  $G_{u,v,4}$ , which leads to  $\gamma(G_{u,v,4}) \leq \gamma(G) + 2$ .

**Claim 1.** *If  $\gamma(G_{u,v,1}) \leq \gamma(G)$ , then  $\gamma(G_{u,v,4}) \leq \gamma(G) + 1$ .*

**Proof.** Assume that  $v$  is a  $\gamma$ -bad vertex of  $G$ ,  $u \in V^-(G - v)$  and  $R$  a  $\gamma$ -set of  $G - \{u, v\}$ . Then  $|R| = \gamma((G - v) - u) = \gamma(G - v) - 1 = \gamma(G) - 1$  and  $R \cup \{x_1, x_4\}$  is a dominating set of  $G_{u,v,4}$ . Hence  $\gamma(G_{u,v,4}) \leq |R| + 2 = \gamma(G) + 1$ .

Assume now that  $D$  is a  $\gamma$ -set of  $G$  with  $u \in D$ . Then  $D \cup \{x_3\}$  is a dominating set of  $G_{u,v,4}$ . Hence again  $\gamma(G_{u,v,4}) \leq \gamma(G) + 1$ . Now by Theorem 14 we immediately obtain the required.  $\square$

(G) Let  $\gamma(G_{u,v,4}) = \gamma(G) + 2$ . By Claim 1,  $\gamma(G_{u,v,1}) > \gamma(G)$  and by Theorem 14,  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ .

Let now  $\gamma(G_{u,v,1}) = \gamma(G) + 1$ . By Theorem 14,  $u$  and  $v$  are  $\gamma$ -bad vertices of  $G$ ,  $u \notin V^-(G - v)$  and  $v \notin V^-(G - u)$ . Let  $M$  be a  $\gamma$ -set of  $G_{u,v,4}$  such that  $R = M \cap \{x_1, x_2, x_3, x_4\}$  has minimum cardinality. Clearly  $|R| \in \{1, 2\}$ . Assume first  $|R| = 1$  and without loss of generality  $\{x_2\} = M$ . Then  $M \setminus \{x_2\}$  is a dominating set of  $G$  with  $v \in M \setminus \{x_2\}$ . Since  $v$  is a  $\gamma$ -bad vertex of  $G$ ,  $|M \setminus \{x_2\}| > \gamma(G)$  and then  $\gamma(G_{u,v,4}) = |M| > \gamma(G) + 1$ . Let now  $|R| = 2$  and without loss of generality  $x_1, x_4 \in M$ . Since  $|M \cap \{x_1, x_2, x_3, x_4\}|$  is minimum,  $u, v \notin M$  and  $M \setminus \{x_1, x_4\}$  is a dominating set of  $G - \{u, v\}$ . But then  $\gamma(G_{u,v,4}) = 2 + |M \setminus \{x_1, x_4\}| \geq 2 + \gamma((G - u) - v) \geq 2 + \gamma(G - u) = 2 + \gamma(G)$ .

(H) Let  $\gamma(G_{u,v,1}) = \gamma(G)$ . By Claim 1,  $\gamma(G_{u,v,4}) \leq \gamma(G) + 1$ . If  $\gamma(G_{u,v,i}) = \gamma(G) + 1$  for some  $i \in \{1, 2\}$ , then since  $\gamma(G_{u,v,4}) \geq \gamma(G_{u,v,i})$ , we obtain  $\gamma(G_{u,v,4}) = \gamma(G) + 1$ .

(I) Let  $\gamma(G_{u,v,3}) = \gamma(G)$ . Hence at least one of (i) and (ii) of Theorem 19 holds, and by (E),  $\gamma(G_{u,v,4}) \leq \gamma(G) + 1$ .

Assume that the equality holds. If  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ , then for any  $\gamma$ -set  $U$  of  $G - \{u, v\}$ ,  $U \cup \{x_1, x_4\}$  is a dominating set of  $G_{u,v,4}$ . Hence  $\gamma(G_{u,v,4}) = \gamma(G)$ , a contradiction.

Let now  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$  and without loss of generality condition (i) of Theorem 19 be satisfied. Suppose  $\gamma(G_{u,v,4}) = \gamma(G)$ . Hence for each  $\gamma$ -set  $M$  of  $G_{u,v,4}$  are fulfilled:  $x_1, x_4 \in M$ ,  $x_2, x_3, u, v \notin M$ ,  $pn[x_1, M] = \{x_1, x_2, u\}$  and  $pn[x_4, M] = \{x_3, x_4, v\}$ . But then  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ , a contradiction. Thus  $\gamma(G_{u,v,4}) = \gamma(G) + 1$ .

(J) If  $\gamma(G_{u,v,4}) = \gamma(G)$ , then  $\gamma(G_{u,v,3}) = \gamma(G)$  and by (G),  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ .

Now let  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ . But then for each  $\gamma$ -set  $D$  of  $G - \{u, v\}$ , the set  $D \cup \{x_1, x_4\}$  is a dominating set of  $G_{u,v,4}$ . Thus  $\gamma(G_{u,v,4}) = \gamma(G)$ . ■

**Theorem 23.** *Let  $u$  and  $v$  be nonadjacent vertices of a graph  $G$ . If  $\gamma(G_{u,v,k}) = \gamma(G)$ , then  $k \leq 4$ . If  $k \geq 5$ , then  $\gamma(G_{u,v,k}) > \gamma(G)$ . If  $\gamma(G_{u,v,4}) = \gamma(G)$ , then  $\gamma(G_{u,v,5}) = \gamma(G) + 1$ .*

**Proof.** By Theorem 22,  $\gamma(G) \leq \gamma(G_{u,v,4}) \leq \gamma(G) + 2$ . If  $\gamma(G_{u,v,4}) > \gamma(G)$ , then  $\gamma(G_{u,v,k}) > \gamma(G)$  for all  $k \geq 5$  because of Observation 13. So, let  $\gamma(G_{u,v,4}) = \gamma(G)$ . By Theorem 22(H),  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ . But then for each  $\gamma$ -set  $D$  of  $G - \{u, v\}$ , the set  $D \cup \{x_1, x_3, x_5\}$  is a dominating set of  $G_{u,v,5}$ . Hence  $\gamma(G_{u,v,5}) \leq \gamma(G) + 1$ . Let now  $M$  be a  $\gamma$ -set of  $G_{u,v,5}$ . Then at least one of  $x_2, x_3, x_4$  is in  $M$  and hence  $\gamma(G_{u,v,5}) = |M| \geq \gamma(G) + 1$ . Thus  $\gamma(G_{u,v,5}) = \gamma(G) + 1$ . Now using again Observation 13 we conclude that  $\gamma(G_{u,v,k}) > \gamma(G)$  for all  $k \geq 5$ . ■

**Corollary 24.** *Let  $G$  be a noncomplete graph. Then  $\bar{epa}(G) \leq \bar{Epa}(G) \leq 5$ . Moreover, the following holds.*

- (i)  $\bar{Epa}(G) = 5$  if and only if there are nonadjacent vertices  $u$  and  $v$  of  $G$  with  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ .
- (ii)  $\bar{epa}(G) = 5$  if and only if  $G$  is edgeless.
- (iii)  $\bar{epa}(G) = \bar{Epa}(G) = 4$  if and only if for each pair  $u, v$  of nonadjacent vertices of  $G$ ,  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$  and at least one of the following holds:
  - (a)  $u \in V^-(G)$  and  $v$  is a  $\gamma$ -good vertex of  $G - u$ ,
  - (b)  $v \in V^-(G)$  and  $u$  is a  $\gamma$ -good vertex of  $G - v$ .

**Proof.** By Theorem 23,  $\bar{Epa}(G) \leq 5$ .

(i)  $\Rightarrow$  Let  $\bar{Epa}(G) = 5$ . Then there is a pair  $u, v$  of nonadjacent vertices of  $G$  such that  $\gamma(G_{u,v,4}) = \gamma(G)$ . Now by Theorem 22(H),  $\gamma(G - \{u, v\}) = \gamma(G) - 2$ .

(i)  $\Leftarrow$  Let  $\gamma(G - \{u, v\}) = \gamma(G) - 2$  and  $D$  be a  $\gamma$ -set of  $G - \{u, v\}$ , where  $u$  and  $v$  are nonadjacent vertices of  $G$ . Hence  $D_1 = D \cup \{x_1, x_4\}$  is a dominating set of  $G_{u,v,4}$  and  $|D_1| = \gamma(G)$ . This implies  $\gamma(G_{u,v,4}) = \gamma(G)$ . The result now follows by Theorem 23.

(ii) If  $G$  has no edges, then the result is obvious. So let  $G$  have edges and  $\bar{epa}(G) = 5$ . Then for any 2 nonadjacent vertices  $u$  and  $v$  of  $G$  is satisfied

$\gamma(G - \{u, v\}) = \gamma(G) - 2$  (by (i)). Hence we can choose  $u$  and  $v$  so that they have a neighbor in common, say  $w$ . But then  $w$  is a  $\gamma$ -bad vertex of  $G - u$  which implies  $v \notin V^-(G - u)$ . This leads to  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$ , a contradiction.

(iii)  $\Rightarrow$  Let  $\bar{epa}(G) = \bar{Epa}(G) = 4$ . Then for each two nonadjacent  $u, v \in V(G)$  we have  $\gamma(G) = \gamma(G_{u,v,3}) < \gamma(G_{u,v,4})$ . Now by Theorem 22( $\mathbb{G}$ ),  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$  and by Theorem 19, at least one of (a) and (b) is valid.

(iii)  $\Leftarrow$  Consider any two nonadjacent vertices  $u, v$  of  $G$ . Then  $\gamma(G - \{u, v\}) \geq \gamma(G) - 1$  and at least one of (a) and (b) is valid. Theorem 19 now implies  $\gamma(G) = \gamma(G_{u,v,3})$ , and by Theorem 22,  $pa(u, v) = 4$ . ■

**Example 25.** Let  $G_n$  be the Cartesian product of two copies of  $K_n$ ,  $n \geq 2$ . We consider  $G_n$  as an  $n \times n$  array of vertices  $\{x_{i,j} \mid 1 \leq i \leq j \leq n\}$ , where the closed neighborhood of  $x_{i,j}$  is the union of the sets  $\{x_{1,j}, x_{2,j}, \dots, x_{n,j}\}$  and  $\{x_{i,1}, x_{i,2}, \dots, x_{i,n}\}$ . Note that  $V(G_n) = V^-(G_n)$  and  $\gamma(G_n) = n$  [6]. It is easy to see that the following sets are  $\gamma$ -sets of  $G_n - x_{1,1}$ :  $D_i = \{x_{2,i}, x_{3,i+1}, \dots, x_{n,n+i-2}\}$ ,  $i = 2, 3, \dots, n$ , where  $x_{k,j} = x_{k,j-n+1}$  for  $j > n$  and  $2 \leq k \leq n$ . Since  $D = \bigcup_{i=2}^n D_i = V(G_n) \setminus N[x_{1,1}]$ , all  $\gamma$ -bad vertices of  $G_n - x_{1,1}$  are the neighbors of  $x_{1,1}$  in  $G_n$ . Since each vertex of  $D$  is adjacent to some neighbor of  $x_{1,1}$ ,  $V^-(G_n - x_{1,1})$  is empty. Now by Theorem 19 we have  $pa(x_{1,1}, y) \geq 4$ , and by Theorem 22( $\mathbb{H}$ ),  $pa(x_{1,1}, y) < 5$ . Thus  $pa(x_{1,1}, y) = 4$ . By reason of symmetry, we obtain  $\bar{epa}(G_n) = \bar{Epa}(G_n) = 4$ .

#### 4. OBSERVATIONS AND OPEN PROBLEMS

A constructive characterization of the trees  $T$  with  $i(T) \equiv \gamma(T)$ , and therefore a constructive characterization of the trees  $T$  with  $Epa(T) = 2$  (by Corollary 7), was provided in [9].

**Problem 26.** Characterize all unicyclic graphs  $G$  with  $Epa(G) = 2$ .

**Problem 27.** Find results on  $\gamma$ -excellent graphs  $G$  with  $\bar{Epa}(G) = 2$ .

**Problem 28.** Characterize all graphs  $G$  with  $\bar{epa}(G) = \bar{Epa}(G) = 4$ .

**Corollary 29.** Let  $G$  be a connected noncomplete graph with edges. Then

- (i)  $2 \leq epa(G) + \bar{Epa}(G) \leq 8$ ,
- (ii)  $2 \leq epa(G) + \bar{epa}(G) \leq 7$ ,
- (iii)  $3 \leq Epa(G) + \bar{Epa}(G) \leq 8$ ,
- (iv)  $3 \leq Epa(G) + \bar{epa}(G) \leq 7$ .

**Proof.** (i)–(iv) The left-side inequalities immediately follow by Corollary 5 and Corollary 15. The right-side inequalities hold because of Corollary 10 and Corollary 24. ■

Note that all bounds stated in Corollary 29 are attainable. We leave finding examples demonstrating this to the reader.

**Problem 30.** Characterize all graphs  $G$  that attain the bounds in Corollary 29.

### Acknowledgments

The author wishes to express his sincere thanks to the reviewers for their very helpful comments and remarks.

### REFERENCES

- [1] N. Ananchuen, W. Ananchuen and R.E.L. Aldred, *Maximal 3-gamma-vertex-critical graphs*, Util. Math. **88** (2012) 75–90.
- [2] A. Bhattacharya and G.R. Vijayakumar, *Effect of edge-subdivision on vertex-domination in a graph*, Discuss. Math. Graph Theory **22** (2002) 335–347.  
doi:10.7151/dmgt.1179
- [3] R.C. Brigham, P.Z. Chinn and R.D. Dutton, *A Study of Vertex Domination Critical Graphs* (Technical Report, University of Central Florida, 1984).
- [4] R.C. Brigham, P.Z. Chinn and R.D. Dutton, *Vertex domination-critical graphs*, Networks **18** (1988) 173–179.  
doi:10.1002/net.3230180304
- [5] G.H. Fricke, T.W. Haynes, S.M. Hedetniemi, S.T. Hedetniemi and R.C. Laskar, *Excellent trees*, Bull. Inst. Combin. Appl. **34** (2002) 27–38.
- [6] B. Hartnell and D. Rall, *Bounds on the bondage number of a graph*, Discrete Math. **128** (1994) 173–177.  
doi:10.1016/0012-365X(94)90111-2
- [7] W. Goddard and M.A. Henning, *Independent domination in graphs: A survey and recent results*, Discrete Math. **313** (2013) 839–854.  
doi:10.1016/j.disc.2012.11.031
- [8] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of domination in graphs* (Marcel Dekker Inc., New York, 1998).
- [9] T.W. Haynes, M.A. Henning and P.J. Slater, *Strong equality of domination parameters in trees*, Discrete Math. **260** (2003) 77–87.  
doi:10.1016/S0012-365X(02)00451-X
- [10] V. Samodivkin, *Common extremal graphs for three inequalities involving domination parameters*, Trans. Comb. **6** (2017) 1–9.  
doi:10.22108/TOC.2017.21464
- [11] H.B. Walikar and B.D. Acharya, *Domination critical graphs*, Nat. Acad. Sci. Lett. **2** (1979) 70–72.

Received 18 June 2018

Revised 2 November 2018

Accepted 3 November 2018