

T -COLORINGS, DIVISIBILITY AND THE CIRCULAR CHROMATIC NUMBER

ROBERT JANCZEWSKI

Department of Algorithms and Systems Modelling
Gdańsk University of Technology
Narutowicza 11/12, Gdańsk, Poland
e-mail: skalar@eti.pg.gda.pl

ANNA MARIA TRZASKOWSKA

Department of Applied Informatics in Management
Gdańsk University of Technology
Narutowicza 11/12, Gdańsk, Poland
e-mail: anna.trzaskowska@pg.edu.pl

AND

KRZYSZTOF TUROWSKI

Center for Science of Information
Purdue University West Lafayette, Indiana, USA
e-mail: krzysztof.szymon.turowski@gmail.com

Abstract

Let T be a T -set, i.e., a finite set of nonnegative integers satisfying $0 \in T$, and G be a graph. In the paper we study relations between the T -edge spans $\text{esp}_T(G)$ and $\text{esp}_{d \odot T}(G)$, where d is a positive integer and

$$d \odot T = \{0 \leq t \leq d(\max T + 1) : d \mid t \Rightarrow t/d \in T\}.$$

We show that $\text{esp}_{d \odot T}(G) = d \text{esp}_T(G) - r$, where r , $0 \leq r \leq d - 1$, is an integer that depends on T and G . Next we focus on the case $T = \{0\}$ and show that

$$\text{esp}_{d \odot \{0\}}(G) = \lceil d(\chi_c(G) - 1) \rceil,$$

where $\chi_c(G)$ is the circular chromatic number of G . This result allows us to formulate several interesting conclusions that include a new formula for the circular chromatic number

$$\chi_c(G) = 1 + \inf \{ \text{esp}_{d \odot \{0\}}(G)/d : d \geq 1 \}$$

and a proof that the formula for the T -edge span of powers of cycles, stated as conjecture in [Y. Zhao, W. He and R. Cao, *The edge span of T -coloring on graph C_n^d* , Appl. Math. Lett. 19 (2006) 647–651], is true.

Keywords: T -coloring, circular chromatic number.

2010 Mathematics Subject Classification: 05C15.

1. INTRODUCTION

In the paper we study relations between two different generalizations of ordinary vertex colorings: T -colorings and (k, d) -colorings. Let G be a graph with n -vertex set V and edge set E . Given integers $1 \leq d \leq k$, by a (k, d) -coloring of G we mean any function $c: V \rightarrow [0, k-1]$ ($[a, b] := \{a, a+1, \dots, b\}$ for any integers $a \leq b$) such that

$$d \leq |c(u) - c(v)| \leq k - d$$

whenever $uv \in E$. This notion may be viewed as a generalization of a k -coloring since (k, d) -colorings of G are k -colorings of G and $(k, 1)$ -colorings are the same as k -colorings that use colors from the interval $[0, k-1]$. The *circular chromatic number*, introduced by Vince [12] as a generalization of the chromatic number, is defined by the formula

$$\chi_c(G) = \inf \{k/d: G \text{ has a } (k, d)\text{-coloring}\}.$$

The circular chromatic number was studied by many authors, see [14, 15] for a survey of results. It was shown for example [12] that the distance between the circular and ordinary chromatic number does not exceed 1, i.e.

$$\chi(G) - 1 < \chi_c(G) \leq \chi(G).$$

In the same paper Vince proved two useful facts: (1) G has a (k, d) -coloring if and only if $\chi_c(G) \leq k/d$; (2) $\chi_c(G)$ is a rational number which has a form k/d , where $k \leq n$. We will use these observations to show that there is a relation between $\chi_c(G)$ and $\text{esp}_T(G)$ the T -edge span defined below. Given a T -set T , i.e., a finite set that consists of nonnegative integers and satisfies $0 \in T$, by a T -coloring of G we mean any function $c: V \rightarrow \mathbb{Z}$ such that

$$|c(u) - c(v)| \notin T$$

whenever $uv \in E$. T -colorings were introduced as a model for the frequency assignment problem in [5]. This notion also may be viewed as a generalization of ordinary vertex colorings since T -colorings are vertex colorings and vertex

colorings are $\{0\}$ -colorings. The T -edge span, introduced by Cozzens and Roberts [1], is defined as

$$\text{esp}_T(G) = \min\{\text{esp}(c) : c \text{ is a } T\text{-coloring of } G\},$$

where $\text{esp}(c) = \max\{|c(u) - c(v)| : uv \in E\}$ is the *edge span* of c (if G is an empty graph then $\text{esp}(c) = 0$). If we replace $\text{esp}(c)$ by $\text{sp}(c)$ (the *span* of c , i.e., $\max\{|c(u) - c(v)| : u, v \in V\}$) we will receive the T -span of G . Both parameters were studied by many authors, there are results concerning computational complexity of the problem of computing $\text{sp}_T(G)$ [2, 3], the behaviour of the greedy algorithm [7] and formulas describing $\text{sp}_T(G)$ and $\text{esp}_T(G)$ for some T -sets T and some graphs G [8, 9, 13].

The remainder of the paper is organized as follows. In Section 2 we study relations between $\text{esp}_T(G)$ and $\text{esp}_{d \odot T}(G)$, where d is a positive integer and $d \odot T = \{0 \leq t \leq d(\max T + 1) : d \mid t \Rightarrow t/d \in T\}$. We show that $\text{esp}_{d \odot T}(G) = d \text{esp}_T(G) - r$, where r , $0 \leq r \leq d - 1$, is an integer that depends on T and G . In Section 3 we study the distance between the T -span and T -edge span and show that it cannot exceed $\max T$. We also give examples that prove that this bound is tight. Section 4 contains our main results. We show that if T is an interval, i.e., $T = [0, d - 1]$ (or equivalently $T = d \odot \{0\}$), then (k, d) -colorings ($k \geq d$) are nonnegative T -colorings with span bounded by $k - 1$ and edge span bounded by $k - d$. We use this relation to show that

$$\text{esp}_{d \odot \{0\}}(G) = \lceil d(\chi_c(G) - 1) \rceil.$$

We also discuss whether it is possible to extend this relation to all T -sets. Using the above formula we show that

$$\chi_c(G) = 1 + \inf \{ \text{esp}_{d \odot \{0\}}(G)/d : d \geq 1 \}$$

and discuss how these formulas allow us to move known results from the world of the T -edge span to the world of the circular chromatic number and vice versa. The last section is devoted to the powers of cycles investigated in [13]. The authors conjectured and partially proved that

$$\text{esp}_{d \odot \{0\}}(C_n^p) = pd + \lceil rd/q \rceil,$$

where $q \geq 2$ and r are the quotient and the remainder of the division of n by $p + 1$, respectively. We show that it is true in general.

2. T -EDGE SPAN AND $d \odot T$ -EDGE SPAN

The operation \odot was introduced in [6], where it was shown that $\text{sp}_{d \odot T}(G) = d \text{sp}_T(G)$. Below we prove a similar formula for the T -edge span, but before we proceed we need to recall the following result.

Lemma 1 (Lemma 2.2(i) of [6]). *If a and b are real numbers, then $\lfloor |a - b| \rfloor \leq \lfloor |a| - |b| \rfloor \leq \lceil |a - b| \rceil$.*

Lemma 2. *Let G be a graph, T be a T -set and d be a positive integer.*

- (1) *If c is a T -coloring of G , then dc is a $d \odot T$ -coloring of G .*
- (2) *If c is a $d \odot T$ -coloring of G , then $\lfloor c/d \rfloor$ is a T -coloring of G .*

Proof. Let uv be an edge of G (if G is empty, then our claim is obvious).

(1) If $|c(u) - c(v)| \geq \max T + 1$, then $|dc(u) - dc(v)| \geq d(\max T + 1) = \max d \odot T + 1$. If $|c(u) - c(v)| < \max T + 1$ and $|dc(u) - dc(v)| \in d \odot T$, then the definition of $d \odot T$ gives $|c(u) - c(v)| \in T$, a contradiction. Hence $|dc(u) - dc(v)| \notin d \odot T$ in both cases.

(2) If $|c(u) - c(v)| \geq \max d \odot T + 1 = d(\max T + 1)$, then $|\lfloor c(u)/d \rfloor - \lfloor c(v)/d \rfloor| \geq \lfloor |c(u) - c(v)|/d \rfloor \geq \max T + 1$ by Lemma 1. If $|c(u) - c(v)| < \max d \odot T + 1$, then the definition of $d \odot T$ gives $d \mid |c(u) - c(v)|$ and, by Lemma 1, $|\lfloor c(u)/d \rfloor - \lfloor c(v)/d \rfloor| = |c(u) - c(v)|/d \notin T$. Hence $|\lfloor c(u)/d \rfloor - \lfloor c(v)/d \rfloor| \notin T$ in both cases. ■

Theorem 3. *Let G be a graph, T be a T -set and d be a positive integer. There is an integer $0 \leq r \leq d - 1$ such that $\text{esp}_{d \odot T}(G) = d \text{esp}_T(G) - r$.*

Proof. Let c be a T -coloring of G such that $\text{esp}(c) = \text{esp}_T(G)$. By Lemma 2, dc is a $d \odot T$ -coloring of G . Hence

$$(1) \quad \text{esp}_{d \odot T}(G) \leq \text{esp}(dc) = d \text{esp}(c) = d \text{esp}_T(G).$$

Let c' be a $d \odot T$ -coloring of G such that $\text{esp}(c') = \text{esp}_{d \odot T}(G)$. By Lemma 2, $\lfloor c'/d \rfloor$ is a T -coloring of G . Let uv be an edge of G such that $\text{esp}(\lfloor c'/d \rfloor) = |\lfloor c'(u)/d \rfloor - \lfloor c'(v)/d \rfloor|$ (if G is empty our claim is obvious). Then

$$\begin{aligned} d \text{esp}_T(G) - d &\leq d \text{esp}(\lfloor c'/d \rfloor) - d = d |\lfloor c'(u)/d \rfloor - \lfloor c'(v)/d \rfloor| - d \\ (2) \quad &\leq d \lceil |c'(u) - c'(v)|/d \rceil - d \leq d \lceil \text{esp}(c')/d \rceil - d \\ &= d \lceil \text{esp}_{d \odot T}(G)/d \rceil - d < \text{esp}_{d \odot T}(G). \end{aligned}$$

To complete the proof it suffices to combine (1) with (2). ■

The open problem is a formula for r . Later we will show how to compute r provided that $T = \{0\}$ and that r can be any integer from $[0, d - 1]$.

Corollary 4. *Let G be a graph, T be a T -set and d be a positive integer. Then $\text{esp}_T(G) = \lceil \text{esp}_{d \odot T}(G)/d \rceil$.*

3. THE DISTANCE BETWEEN THE T -SPAN AND T -EDGE SPAN

It is known [1] that $\text{esp}_T(G) \leq \text{sp}_T(G)$. We are going to show that $\text{sp}_T(G) \leq \text{esp}_T(G) + \max T$ and give examples in which the difference $\text{sp}_T(G) - \text{esp}_T(G)$ equals $\max T$.

Lemma 5. *Let G be a graph and T be a T -set. If $c' : V \rightarrow \mathbb{Z}$ is a T -coloring of G and $c : V \rightarrow \mathbb{Z}$ is the remainder of the division of c' by $\text{esp}(c') + \max T + 1$, i.e., $c(v) = c'(v) \bmod (\text{esp}(c') + \max T + 1)$ for $v \in V$, then*

- (1) c is a T -coloring of G .
- (2) $\text{sp}(c) \leq \text{esp}(c') + \max T$.
- (3) $\text{esp}(c) \leq \text{esp}(c') + \max T + 1 - \min(\mathbb{N} \setminus T)$.

Proof. Observe that (2) follows immediately from the definition of c . To prove (1) and (3), take an edge uv of G (if G is empty, our claim is obvious). Let q be the quotient of the division of c' by $\text{esp}(c') + \max T + 1$. Without loss of generality we may assume that $q(u) \geq q(v)$. It is easy to see that $q(u) \leq q(v) + 1$ since otherwise

$$\begin{aligned} \text{esp}(c') &\geq |c'(u) - c'(v)| \\ &= |(\text{esp}(c') + \max T + 1)(q(u) - q(v)) + c(u) - c(v)| \\ &\geq (\text{esp}(c') + \max T + 1)|q(u) - q(v)| - |c(u) - c(v)| \\ &\geq 2(\text{esp}(c') + \max T + 1) - \text{esp}(c') - \max T \\ &= \text{esp}(c') + \max T + 2 > \text{esp}(c'). \end{aligned}$$

Hence there are two cases to consider.

(a) $q(u) = q(v) + 1$. Then $|c'(u) - c'(v)| = |(\text{esp}(c') + \max T + 1)(q(u) - q(v)) + (c(u) - c(v))| = |\text{esp}(c') + \max T + 1 + (c(u) - c(v))|$. Since $\text{esp}(c') + \max T + 1 > \text{esp}(c') \geq |c'(u) - c'(v)|$ and $|c(u) - c(v)| \leq \text{esp}(c') + \max T$, we have $|c(u) - c(v)| = \text{esp}(c') + \max T + 1 - |c'(u) - c'(v)|$. This gives $|c(u) - c(v)| \geq \max T + 1$ and $|c(u) - c(v)| \leq \text{esp}(c') + \max T + 1 - \min(\mathbb{N} \setminus T)$ since $|c'(u) - c'(v)| \notin T$ implies $|c'(u) - c'(v)| \geq \min(\mathbb{N} \setminus T)$.

(b) $q(u) = q(v)$. Then $|c'(u) - c'(v)| = |c(u) - c(v)|$, which gives $|c(u) - c(v)| \notin T$ and $|c(u) - c(v)| \leq \text{esp}(c') \leq \text{esp}(c') + \max T + 1 - \min(\mathbb{N} \setminus T)$. ■

Corollary 6. *Let G be a graph and T be a T -set. Then*

- (1) *There is a T -coloring c of G such that $\text{sp}(c) \leq \text{esp}_T(G) + \max T$ and $\text{esp}(c) \leq \text{esp}_T(G) + \max T + 1 - \min(\mathbb{N} \setminus T)$.*
- (2) *If T is an interval, then there is a T -coloring c of G such that $\text{esp}(c) = \text{esp}_T(G)$ and $\text{sp}(c) \leq \text{esp}_T(G) + \max T$.*
- (3) $\text{esp}_T(G) \leq \text{sp}_T(G) \leq \text{esp}_T(G) + \max T$.

Proof. (1) Let c' be a T -coloring of G satisfying $\text{esp}(c') = \text{esp}_T(G)$ and c be the remainder of the division of c' by $\text{esp}_T(G) + \max T + 1$. The claim follows from Lemma 5.

(2) Follows from (1) since $\min(\mathbb{N} \setminus T) = \max T + 1$ if T is an interval.

(3) Follows from (1) and the definition of the T -span. ■

The above inequalities are tight. It is known [1] that $\text{esp}_T(G) = \text{sp}_T(G)$ for all weakly perfect graphs and all T -sets T . It is also easy to see that if T is an interval, then $\text{sp}_T(C_{2n+1}) = 2 \max T + 2$ ($\text{sp}_T(G) = (\max T + 1)(\chi(G) - 1)$ if T is an interval, see [1]) and $\text{esp}_T(C_{2n+1}) = \lceil (\max T + 1)(1 + 1/n) \rceil$ (see Theorem 8) which gives $\text{sp}_T(C_{2n+1}) = \text{esp}_T(C_{2n+1}) + \max T$ provided that $n \geq \max T + 1$.

4. THE RELATION BETWEEN (k, d) -COLORINGS AND T -COLORINGS

Now we are ready to prove that there is a relation between (k, d) -colorings and T -colorings provided that T is an interval.

Lemma 7. *Let G be a graph and d be a positive integer. If $T = [0, d - 1]$, then for every function $c: V \rightarrow \mathbb{Z}$ and every integer $k \geq d$ the following conditions are equivalent:*

- (1) c is a T -coloring of G such that $\text{sp}(c) \leq k - 1$ and $\text{esp}(c) \leq k - d$.
- (2) $c - \min c(V)$ is a (k, d) -coloring of G .

Proof. Let uv be an edge of G (our claim is obvious if G is empty) and $c' = c - \min c(V)$.

(\Rightarrow) c is a T -coloring of G and T is an interval, so $|c'(u) - c'(v)| = |c(u) - c(v)| \geq d$. Moreover, $|c'(u) - c'(v)| = |c(u) - c(v)| \leq \text{esp}(c) \leq k - d$ and $c'(V) \subseteq [0, \text{sp}(c)] \subseteq [0, k - 1]$.

(\Leftarrow) c' is a (k, d) -coloring of G , so $|c(u) - c(v)| = |c'(u) - c'(v)| \geq d$ and $|c(u) - c(v)| = |c'(u) - c'(v)| \leq k - d$. This proves that c is a T -coloring and gives $\text{esp}(c) \leq k - d$. To complete the proof it suffices to observe that $c'(V) \subseteq [0, k - 1]$ implies $\text{sp}(c) = \text{sp}(c') \leq k - 1$. ■

Theorem 8. *Let G be a graph and d be a positive integer. If $T = [0, d - 1]$, then*

$$\text{esp}_T(G) = \lceil d(\chi_c(G) - 1) \rceil.$$

Proof. Without loss of generality we assume that G is not empty. Then $k = \lceil d\chi_c(G) \rceil - 1 \geq d$. If $\text{esp}_T(G) \leq k - d$, then, by Corollary 6, there is a T -coloring c of G such that $\text{esp}(c) = \text{esp}_T(G) \leq k - d$ and $\text{sp}(c) \leq \text{esp}_T(G) + d - 1 \leq k - 1$.

Lemma 7 implies now that $c - \min c(V)$ is a (k, d) -coloring, which finally gives $d\chi_c(G) \leq k$, a contradiction. Hence

$$\text{esp}_T(G) \geq k - d + 1.$$

On the other hand, $(k+1)/d \geq \chi_c(G)$ so there exists a $(k+1, d)$ -coloring c of G . Without loss of generality we assume that $\min c(V) = 0$. By Lemma 7, c has to be a T -coloring of G with $\text{esp}(c) \leq k - d + 1$. This gives

$$\text{esp}_T(G) \leq k - d + 1.$$

Combining these inequalities together, we get $\text{esp}_T(G) = k - d + 1 = \lceil d\chi_c(G) \rceil - d = \lceil d(\chi_c(G) - 1) \rceil$. ■

Since T is an interval, we know that $|T| = \max T + 1$ and the above formula may be expressed as

$$\text{esp}_T(G) = \lceil |T|(\chi_c(G) - 1) \rceil.$$

This resembles Tesman's inequality $\text{sp}_T(G) \leq |T|(\chi(G) - 1)$ which holds for all T -sets T and all graphs G [11], so it is interesting to ask the following question.

Does $\text{esp}_T(G) \leq \lceil |T|(\chi_c(G) - 1) \rceil$ for all T -sets T and all graphs G ?

Unfortunately, the answer is negative even for odd cycles. To show this, let us consider integers $1 \leq k \leq n-1$ and set $T = \{0, 2, \dots, 2k\}$ and $G = C_{2n+1}$. Then $\lceil |T|(\chi_c(G) - 1) \rceil = \lceil (k+1)(1 + 1/n) \rceil = k+2$ and $\text{esp}_T(C_{2n+1}) \geq 2k+2$ since otherwise the differences of colors assigned to adjacent vertices of G in any T -coloring of G with minimal edge span would be odd and their sum would not be 0, a contradiction.

Theorem 8 shows also that the value of integer r of Theorem 3 can be arbitrary. Indeed, if we take $0 \leq r \leq d-1$ and a planar graph G such that $\chi_c(G) = 3 - r/d$ (which exists by [10]), then $\chi(G) = 3$ and $\text{esp}_{d\odot\{0\}}(G) = \lceil d(\chi_c(G) - 1) \rceil = \lceil d(2 - r/d) \rceil = 2d - r = d(\chi(G) - 1) - r = d\text{esp}_{\{0\}}(G) - r$. The open question is if this is true for all T -sets T .

Theorem 9. *Let G be a graph. Then*

$$\chi_c(G) = 1 + \inf \{ \text{esp}_{d\odot\{0\}}(G)/d : d \geq 1 \}.$$

Moreover, if $\chi_c(G) = k/d$ ($1 \leq d \leq k$), then $\chi_c(G) = 1 + \text{esp}_{d\odot\{0\}}(G)/d$.

Proof. $\chi_c(G) - 1 \leq \text{esp}_{d\odot\{0\}}(G)/d$ by Theorem 8. To complete the proof it suffices to observe that if $\chi_c(G) = k/d$, then the same theorem gives $\chi_c(G) - 1 = \text{esp}_{d\odot\{0\}}(G)/d$. ■

Theorems 8 and 9 have two important consequences. Firstly, if we know a formula for $\chi_c(G)$, then we can easily obtain a formula for $\text{esp}_T(G)$ for all T -sets T that are intervals. For example, Fan [4] proved that $\chi_c(G) = \chi(G)$ if the complement of G is non-Hamiltonian, which gives

Corollary 10. *If G is a graph whose complement is non-Hamiltonian, then*

$$\text{esp}_{d \odot \{0\}}(G) = d(\chi(G) - 1) = \text{sp}_{d \odot \{0\}}(G)$$

for every $d \geq 1$.

Secondly, if the problem of computing $\chi_c(G)$ for graphs G from a certain class \mathcal{G} is polynomially solvable, then we can compute $\text{esp}_T(G)$ for $G \in \mathcal{G}$ and any interval T in a polynomial time, too.

5. POWERS OF CYCLES

Let $p \geq 1$ and $n \geq 2p + 2$ be integers. Let q and r are the quotient and the remainder of the division of n by $p + 1$, respectively.

Zhao *et al.* in [13] proved the following theorem.

Theorem 11. *If $q = pl + t$ for $l \geq 0$, $0 \leq t \leq p - 1$ such that $p \geq td$, then*

$$\text{esp}_{d \odot \{0\}}(C_n^p) = pd + \lceil rd/q \rceil.$$

Moreover, they conjectured that this equality holds for any $n \geq 2p + 2$, not only when $p \geq td$. We will show that it is true. Recall that it is known that if G is a n -vertex graph, then $\chi_c(G) \geq n/\alpha(G)$, where $\alpha(G)$ is the independence number of G .

Theorem 12. $\chi_c(C_n^p) = n/q$.

Proof. Let v_0, v_1, \dots, v_{n-1} be a cyclic ordering of vertices of C_n^p . We claim that a function given by

$$c(v_i) = (iq) \bmod n$$

is a (n, q) -coloring of C_n^p . Indeed, the definition of c gives $0 \leq c \leq n - 1$ and, if $v_i v_j$ ($i > j$) is an edge of C_n^p , then either $1 \leq i - j \leq p$ and $|c(v_i) - c(v_j)| = (i - j)q$ or $1 \leq n + j - i \leq p$ and $|c(v_i) - c(v_j)| = (n - i + j)q$. In both cases it is easy to verify that $q \leq |c(v_i) - c(v_j)| \leq qp \leq n - q$.

To complete the proof it suffices to observe that $\alpha(C_n^p) \leq q$ and use inequality $\chi_c(G) \geq n/\alpha(G)$. ■

Theorem 13. $\text{esp}_{d \odot \{0\}}(C_n^p) = pd + \lceil rd/q \rceil$.

Proof. Follows immediately from Theorems 8 and 12. ■

6. CONCLUSION

We proved the general relation between the circular chromatic number and T -edge span for $T = d \odot \{0\}$. Moreover, we applied it to solve an open conjecture concerning the T -edge span for powers of cycles C_n^p .

Possible further fields of research include for example finding the necessary conditions for $\text{esp}_T(G) \leq \lceil |T|(\chi_c(G) - 1) \rceil$, or analyzing dependence between $\text{esp}_T(G)$ and $\chi_c(G)$ on the structure of a set T .

REFERENCES

- [1] M.B. Cozzens and F.S. Roberts, *T-colorings of graphs and the channel assignment problem*, Congr. Numer. **35** (1982) 191–208.
- [2] K. Giaro, R. Janczewski and M. Małafiejski, *A polynomial algorithm for finding T-span of generalized cacti*, Discrete Appl. Math. **129** (2003) 371–382.
doi:10.1016/S0166-218X(02)00575-9
- [3] K. Giaro, R. Janczewski and M. Małafiejski, *The complexity of the T-coloring problem for graphs with small degree*, Discrete Appl. Math. **129** (2003) 361–369.
doi:10.1016/S0166-218X(02)00576-0
- [4] G. Fan, *Circular chromatic number and Mycielski graphs*, Combinatorica **24** (2004) 127–135.
doi:10.1007/s00493-004-0008-9
- [5] W.K. Hale, *Frequency assignment: theory and applications*, Proc. IEEE **68** (1980) 1497–1514.
doi:10.1109/PROC.1980.11899
- [6] R. Janczewski, *Divisibility and T-span of graphs*, Discrete Math. **234** (2001) 171–179.
doi:10.1016/S0012-365X(00)00378-2
- [7] R. Janczewski, *Greedy T-colorings of graphs*, Discrete Math. **309** (2009) 1685–1690.
doi:10.1016/j.disc.2008.01.049
- [8] J.S.-T. Juan, I. Sun and P.-X. Wu, *T-coloring on folded hypercubes*, Taiwanese J. Math. **13** (2009) 1331–1341.
doi:10.11650/twjm/1500405511
- [9] A. Raychaudhuri, *Further results on T-coloring and frequency assignment problems*, SIAM J. Discrete Math. **7** (1994) 605–613.
doi:10.1137/S0895480189171746
- [10] D. Moser, *The star-chromatic number of planar graphs*, J. Graph Theory **24** (1997) 33–43.
doi:10.1002/(SICI)1097-0118(199701)24:1<33::AID-JGT5>3.0.CO;2-K
- [11] B. Tesman, *Applications of forbidden difference graphs to T-coloring*, Congr. Numer. **74** (1990) 15–24.

- [12] A. Vince, *Star chromatic number*, J. Graph Theory **12** (1988) 551–559.
doi:10.1002/jgt.3190120411
- [13] Y. Zhao, W. He and R. Cao, *The edge span of T -coloring on graph C_n^d* , Appl. Math. Lett. **19** (2006) 647–651.
doi:10.1016/j.aml.2005.08.016
- [14] X. Zhu, *Circular chromatic number: a survey*, Discrete Math. **229** (2001) 371–410.
doi:10.1016/S0012-365X(00)00217-X
- [15] X. Zhu, *Recent developments in circular colouring of graphs*, Algorithms Combin. **26** (2006) 497–550.
doi:10.1007/3-540-33700-8_25

Received 19 March 2018

Revised 16 October 2018

Accepted 7 December 2018