Discussiones Mathematicae Graph Theory 39 (2019) 731–740 doi:10.7151/dmgt.2214

# HAMILTONIAN NORMAL CAYLEY GRAPHS

JUAN JOSÉ MONTELLANO-BALLESTEROS<sup>1</sup>

AND

ANAHY SANTIAGO ARGUELLO

Instituto de Matemáticas Universidad Nacional Autónoma de México Ciudad Universitaria, México, D.F., C.P. 04510, México

> e-mail: juancho@im.unam.mx jpscw@hotmail.com

## Abstract

A variant of the Lovász Conjecture on hamiltonian paths states that every finite connected Cayley graph contains a hamiltonian cycle. Given a finite group G and a connection set S, the Cayley graph Cay(G, S) will be called normal if for every  $g \in G$  we have that  $g^{-1}Sg = S$ . In this paper we present some conditions on the connection set of a normal Cayley graph which imply the existence of a hamiltonian cycle in the graph.

 ${\bf Keywords:} \ {\rm Cayley \ graph, \ hamiltonian \ cycle, \ normal \ connection \ set.}$ 

2010 Mathematics Subject Classification: 05C45, 05C99.

### 1. INTRODUCTION

Let G be a finite group. A subset  $S \subseteq G$  will be called symmetric if  $S = S^{-1}$ . Given a symmetric subset  $S \subseteq G \setminus \{e\}$  (with e the identity of G), the Cayley graph Cay(G, S) is the graph with vertex set G and a pair  $\{\alpha, \beta\}$  is an edge of Cay(G, S) if and only if there is  $s \in S$  such that  $\alpha = \beta s$  (since S is symmetric, observe that  $s^{-1} \in S$  and  $\beta = \alpha s^{-1}$ ). A Cayley graph Cay(G, S) will be called *normal* if for every  $\alpha \in G$ ,  $\alpha^{-1}S\alpha = S$ . In the literature there is another definition of normal Cayley graph, which is different from the one used in this paper, that said that a Cayley graph on a group G is normal if the right regular representation

<sup>&</sup>lt;sup>1</sup>Research supported by PAPIIT- México under project IN107218.

of the group G is normal in the full automorphism group of the graph (see, for instance [15, 19]).

The problem of finding hamiltonian cycles in graphs is a difficult problem, and since 1969 has received a great attention by the Lovász Conjecture which states that every vertex-transitive graph has a hamiltonian path. A variant of the Lovász Conjecture on hamiltonian paths states that every finite connected Cayley graph contains a hamiltonian cycle (see, for instance [1,3,14,18]). In particular, there are several works on the existence of hamiltonian cycles in Cayley graphs generated by two elements (see, for instance [6-10,12,20]).

In this paper we present the following results.

**Theorem 1.** Let G be a finite non-abelian simple group such that  $\langle \delta_1, \delta_2 \rangle = G$ . If Cay(G, S) is a normal Cayley graph with  $\{\delta_1, \delta_2\} \subseteq S$ , then Cay(G, S) contains a hamiltonian cycle.

**Theorem 2.** Let G be a finite group,  $G = G_0 \supseteq G_1, \ldots, G_{l-1} \supseteq G_l$  be a composition series of G and let  $\{\delta_0, \ldots, \delta_{l+1}\} \subseteq G$  such that, for each  $0 \le i \le l$ ,  $G_i/G_{i+1} = \langle \delta_i G_{i+1}, \delta_{i+1} G_{i+1} \rangle$ . If Cay(G, S) is a normal Cayley graph with  $\{\delta_0, \ldots, \delta_{l+1}\} \subseteq S$ , then Cay(G, S) contains a hamiltonian cycle.

Observe that the normal Cayley graphs with vertex set a group generated by two elements have girth 4. The results are obtained via a generalization of known methods for hamiltonicity of Cayley graphs of girth 4 (see [5, 11, 13, 14]). For general concepts, we may refer the reader to [2, 16].

# 2. NOTATION AND PREVIOUS RESULTS

In order to prove the main theorems, we need some definitions and previous results.

**Theorem 3** [17]. Let G be a simple, non-abelian and finite group. G can be generated by two elements.

In all this section let  $G = \langle \delta_1, \delta_2 \rangle$  be a simple, non-abelian and finite group and Cay(G, S) be a normal Cayley graph with connection set S such that  $\{\delta_1, \delta_2\} \subseteq$ S. Let  $G_0 = \langle \delta_1 \rangle$ , and let

$$\mathcal{P} = \{a_0 G_0 \cup a_1 G_0, \dots, a_n G_0\}$$

be the partition of G in cosets induced by the subgroup  $G_0$  (with  $a_0$  the identity element of G). For each  $0 \leq i \leq n$ ,  $C(a_iG_0)$  will denote the subdigraph of Cay(G,S) induced by the set of vertices  $a_iG_0$ . Given two isomorphic vertex disjoint subgraphs H and H' of Cay(G,S), we will say that H and H' are attached if there is an isomorphism  $\Psi$  between H and H' such that for every  $x \in V(H)$ ,  $\{x, \Psi(x)\}$  is an edge of Cay(G, S). **Lemma 4.** For every  $0 \le i, j \le n$ ,  $C(a_iG_0) \cong C(a_jG_0)$ . Moreover, for every  $0 \le i \le n$  and  $\delta \in \{\delta_1, \delta_2\}$ ,  $C(a_iG_0)$  and  $C(\delta a_iG_0)$  are attached.

**Proof.** Given  $a_i, a_j$  let  $\Phi : a_i G_0 \to a_j G_0$  be defined, for each  $g \in G_0$ , as  $\Phi(a_i g) = a_j g$ . If  $\Phi(a_i g) = \Phi(a_i g_1)$  then  $a_j g = a_j g_1$ , so  $g = g_1$ . Therefore  $\Phi$  is injective and since all cosets have the same cardinality,  $\Phi$  is bijective. If  $a_i g_1$  and  $a_i g_2$  are adjacent in  $C(a_i G_0)$ , then  $g_1^{-1} a_i^{-1} a_i g_2 = g_1^{-1} g_2 \in S$ . Therefore

$$\Phi(a_ig_1)^{-1}\Phi(a_ig_2) = g_1^{-1}a_j^{-1}a_jg_2 = g_1^{-1}g_2 \in S$$

and then  $\Phi(a_ig_1)$  and  $\Phi(a_ig_2)$  are adjacent in  $C(a_jG_0)$ , and the first part of the lemma follows. For the second part, let  $a \in a_iG_0$  and  $\delta a \in \delta a_iG_0$ . Clearly the map  $a \to \delta a$  define an isomorphism between  $C(a_iG_0)$  and  $C(\delta a_iG_0)$  and since S is normal,  $a^{-1}\delta a \in S$ , therefore  $\{a, \delta a\}$  is an edge in Cay(G, S) (see Figure 1), and the lemma follows.



Figure 1

As a word on  $\{\delta_1, \delta_2\}$  we will understand a product  $s_1 s_2 \cdots s_{n-1} s_n$  of powers of  $\delta_1$  and  $\delta_2$ , where two consecutive elements in the product are not powers of the same elements, that is to say, if  $s_i \in \langle a \rangle$  then  $s_{i+1}, s_{i-1} \notin \langle a \rangle$ . The length of a word  $s_1 s_2 \cdots s_{n-1} s_n$  is n. Since  $G = \langle \delta_1, \delta_2 \rangle$ , it follows that for each  $\alpha \in G$ ,  $\alpha = s_1 s_2 \cdots s_{n-1} s_n$  for some word  $s_1 s_2 \cdots s_{n-1} s_n$  on  $\{\delta_1, \delta_2\}$ . For each  $\alpha \in G$ , let  $\ell(\alpha)$  be the minimum length of a word on  $\{\delta_1, \delta_2\}$  such that  $\alpha = s_1 s_2 \cdots s_{n-1} s_n$ .

Let  $\mathcal{H}_0 = \{G_0\}$  and, for each  $k \ge 1$ , let  $\mathcal{H}_k = \{a_i G_0 \in \mathcal{P} : \ell(a_i) = k\}.$ 

Given a coset  $a_iG_0 \in \mathcal{P}$ , let  $L[a_iG_0] = \{a_iG_0\} \cup \{\delta^r a_iG_0 \in \mathcal{P} : \delta \in \{\delta_1, \delta_2\}, r \geq 1\}$ . Observe that if  $\ell(a_i) = k$ , then for every  $aG_0 \in L[a_iG_0] \setminus \{a_iG_0\}, \ell(a) = k+1$ , and the number of cosets in the set  $\{\delta^r a_iG_0 \in \mathcal{P} : \delta \in \{\delta_1, \delta_2\}, r \geq 1\}$  depends on the commutativity of the words  $\delta a_1\delta_1^j, \delta^2 a_1\delta_1^j, \ldots, \delta^{n-1}a_1\delta_1^j$  for  $j \geq 1$ .

**Lemma 5.** If for some  $k \geq 1$ ,  $\delta^k a_i G_0 \in L[a_i G_0]$ , then  $\delta^{k-1} a_i G_0 \in L[a_i G_0]$ .

**Proof.** Let us suppose that for some  $k \ge 1$ ,  $\delta^k a_i G_0 \in L[a_i G_0]$  and let  $b \in G$  such that  $\delta^{k-1}a_i \in bG_0 \in \mathcal{P}$ . Thus,  $\delta^{k-1}a_i = b\delta_1^j$  and therefore  $\delta^k a_i = \delta b\delta_1^j$  which implies that  $\delta^k a_i \in \delta bG_0 = \delta^k a_i G_0$ . Hence  $\delta^k a_i = \delta b$  and  $\delta^{k-1}a_i = b$ , and the result follows.

Observe that for each  $k \geq 0$ , the set  $\{L[a_iG_0] : a_iG_0 \in \mathcal{H}_k\}$  is a partition of  $\mathcal{H}_k \cup \mathcal{H}_{k+1}$ . Given a coset  $a_iG_0 \in \mathcal{P}$ , the subgraph of Cay(G,S) induced by  $L[a_iG_0]$  will be called a *leaf*.

Given a leaf M induced by  $L[a_iG_0] = \{a_iG_0, \delta a_iG_0, \dots, \delta^m a_iG_0\}$  (with  $\delta \in \{\delta_1, \delta_2\}$ ), and a pair of elements  $a_i\delta_1^t, a_i\delta_1^{t+1} \in a_iG_0$ , a path of Cay(G, S) with vertex-set  $\{a_i\delta_1^t, a_i\delta_1^{t+1}\} \cup \bigcup_{j=1}^m \delta^j a_iG_0$  which starts at  $a_i\delta_1^t$ , ends at  $a_i\delta_1^{t+1}$  and such that for every  $1 \leq j \leq m$ , there is  $s_j$  such that  $\{\delta^j a_i\delta_1^{s_j}, \delta^j a_i\delta_1^{s_j+1}\}$  is an edge of the path P, will be called an  $(a_i\delta_1^t, a_i\delta_1^{t+1}, M)$ -complete path.

**Lemma 6.** Let M be a leaf of Cay(G, S) induced by

$$L[a_iG_0] = \left\{ a_iG_0, \delta a_iG_0, \dots, \delta^m a_iG_0 \right\}$$

(with  $\delta \in \{\delta_1, \delta_2\}$ ). For every pair of elements  $a_i \delta_1^t, a_i \delta_1^{t+1} \in a_i G_0$  there is an  $(a_i \delta_1^t, a_i \delta_1^{t+1}, M)$ -complete path.

**Proof.** From Lemma 4 we see that any two "consecutive" subgraphs of the leaf,  $C(\delta^t a_i G_0)$  and  $C(\delta^{t+1} a_i G_0)$ , are attached, and again, by Lemma 4, each subgraph of the leaf is isomorphic to  $C(G_0)$ , which is a cycle of the form  $(e, \delta_1, \delta_1^2, \ldots, \delta_1^n = e)$ . Since  $G = \langle \delta_1, \delta_2 \rangle$ , from here it follows that for each  $1 \leq k \leq m$ ,

$$\left(\delta^k a_i, \delta^k a_i \delta_1, \delta^k a_i \delta_1^2, \dots, \delta^k a_i \delta_1^n = \delta^k a_i\right)$$

is a hamiltonian cycle of  $C(\delta^k a_i G_0)$ .

Let  $a_i \delta_1^t, a_i \delta_1^{t+1} \in a_i G_0$  and  $\delta \in \{\delta_1, \delta_2\}$ . To simplify the notation, let  $\alpha_0 = a_i \delta_1^t, \beta_0 = a_i \delta_1^{t+1}, \epsilon_0 = a_i \delta_1^{t-1}$ , and for each  $1 \le k \le m$ , let  $\alpha_k = \delta^k a_i \delta_1^t, \beta_k = \delta^k a_i \delta_1^{t+1}$  and  $\epsilon_k = \delta^k a_i \delta_1^{t-1}$ .

Case 1. n is even (see Figure 2).



Figure 2

Let

$$P = (\alpha_0, \alpha_1, \dots, \alpha_m = \delta^m a_i \delta_1^t, \delta^m a_i \delta_1^{t-1}, \dots, (\delta^m a_i \delta_1^{t+1} = \beta_m), (\beta_{m-1} = \delta^{m-1} a_i \delta_1^{t+1}), \delta^{m-1} a_i \delta_1^{t+2}, \dots, (\delta^{m-1} a_i \delta_1^{t-1} = \epsilon_{m-1}), (\epsilon_{m-2} = \delta^{m-2} a_i \delta_1^{t-1}), \delta^{m-2} a_i \delta_1^{t-2}, \dots, (\delta^{m-2} a_i \delta_1^{t+1} = \beta_{m-2}), \dots, (\epsilon_1 = \delta a_i \delta_1^{t-1}), \delta a_i \delta_1^{t-2}, \dots, (\delta a_i \delta_1^{t+1} = \beta_1), \beta_0).$$

734

Case 2. n is odd (see Figure 3).



Figure 3

Let

$$P = (\alpha_0, \alpha_1, \dots, \alpha_m = \delta^m a_i \delta_1^t, \delta^m a_i \delta_1^{t+1}, \dots, (\delta^m a_i \delta_1^{t-1} = \epsilon_m), (\epsilon_{m-1} = \delta^{m-1} a_i \delta_1^{t-1}), \delta^{m-1} a_i \delta_1^{t-2}, \dots, (\delta^{m-1} a_i \delta_1^{t+1} = \beta_{m-1}), (\beta_{m-2} = \delta^{m-2} a_i \delta_1^{t+1}), \delta^{m-2} a_i \delta_1^{t+2}, \dots, (\delta^{m-2} a_i \delta_1^{t-1} = \epsilon_{m-2}), \dots, (\epsilon_1 = \delta a_i \delta_1^{t-1}), \delta a_i \delta_1^{t-2}, \dots, (\delta a_i \delta_1^{t+1} = \beta_1), \beta_0).$$

From here the result follows.

## 3. The Proofs of the Main Results

**Proof of Theorem 1.** Let  $G = \langle \delta_1, \delta_2 \rangle$  be a non-abelian simple group and Cay(G, S) be a normal Cayley graph with  $\{\delta_1, \delta_2\} \subseteq S$ . Let  $G_0 = \langle \delta_1 \rangle$ , and let  $\mathcal{P} = \{G_0, a_1G_0, \ldots, a_nG_0\}$  be the partition of G in cosets induced by the subgroup  $G_0$ .

Let  $\mathcal{H}_0 = \{G_0\}$  and, for each  $k \ge 1$ , let  $\mathcal{H}_k = \{a_i G_0 \in \mathcal{P} : \ell(a_i) = k\}$ . Since G is finite, it follows that for some  $p \ge 1$ ,  $G = \bigcup_{j=0}^p \left(\bigcup_{A \in \mathcal{H}_j} A\right)$ .

We will prove the result by showing, by induction on k, that for every  $k \ge 1$ the subgraph of Cay(G, S) induced by

$$\bigcup_{j=0}^k \left(\bigcup_{A \in \mathcal{H}_j} A\right)$$

contains a hamiltonian cycle C such that for each  $a_j G_0 \in \mathcal{H}_k$ , there is  $s_j$  such that  $\left\{a_i \delta_1^{s_j}, a_i \delta_1^{s_j+1}\right\}$  is an edge of C. For k = 1, observe that  $\mathcal{H}_0 = \{G_0\}$  and  $\mathcal{H}_1 = \left\{\delta_2 G_0, \delta_2^2 G_0, \dots, \delta_2^m G_0\right\}$ .

For k = 1, observe that  $\mathcal{H}_0 = \{G_0\}$  and  $\mathcal{H}_1 = \{\delta_2 G_0, \delta_2^2 G_0, \dots, \delta_2^m G_0\}$ . Thus, the subgraph M of Cay(G, S) induced by  $\bigcup_{j=0}^1 \left(\bigcup_{A \in \mathcal{H}_j} A\right)$  is the leaf of Cay(G, S) induced by  $L[G_0] = \{G_0, \delta_2 G_0, \delta_2^2 G_0, \dots, \delta_2^m G_0\}$ . Let  $e, \delta_1 \in G_0$ . By Lemma 6 there is an  $(e, \delta_1, M)$ -complete path P. Therefore  $C = P \circ (\delta_1, \delta_1^2, \ldots, \delta_1^{n-1}, e)$  is a hamiltonian cycle of M such that for every  $1 \leq j \leq m$ , there is  $s_j$  such that  $\left\{\delta_2^j \delta_1^{s_j}, \delta_2^j \delta_1^{s_j+1}\right\} \subseteq \delta_2^j G_0$  is an edge of C.

Suppose that the statement is true for  $1 \leq m \leq k$ ; let Q be the subgraph of Cay(G, S) induced by  $\bigcup_{j=0}^{k+1} \left(\bigcup_{A \in \mathcal{H}_j} A\right)$  and let Q' be the subgraph of Cay(G, S) induced by  $\bigcup_{j=0}^{k} \left(\bigcup_{A \in \mathcal{H}_j} A\right)$ . By induction hypothesis, there is a hamiltonian cycle C of Q' such that for each  $a_j G_0 \in \mathcal{H}_k$ , there is  $s_j$  such that  $\left\{a_j \delta_1^{s_j}, a_j \delta_1^{s_j+1}\right\}$  is an edge of C.

For each  $a_j G_0 \in \mathcal{H}_k$ , by Lemma 6, there is an  $\left(a_j \delta_1^{s_j}, a_j \delta_1^{s_j+1}, M\right)$ -complete path P with M the leaf induced by  $L[a_j G_0] = \left\{a_j G_0, \delta a_j G_0, \delta^2 a_j G_0, \ldots, \delta^m a_j G_0\right\}$ Therefore, by deleting from C the edge  $\left\{a_j \delta_1^{s_j}, a_j \delta_1^{s_j+1}\right\}$ , and attach to  $C \setminus \left\{a_j \delta_1^t, a_j \delta_1^{t+1}\right\}$  the path P we obtain a hamiltonian cycle C' of the subgraph of Cay(G, S) induced by  $V(Q') \cup V(M)$ , and such that for each  $1 \leq i \leq m$  there is  $s_i$  such that  $\left\{\delta^i a_j \delta_1^{s_i+1}\right\}$  is an edge of C'. Following this procedure for each coset in  $\mathcal{H}_k$ , since  $\{L[a_i G_0] : a_i G_0 \in \mathcal{H}_k\}$  is a partition of  $\mathcal{H}_k \cup \mathcal{H}_{k+1}$ , we obtain a hamiltonian cycle C of Q such that for each  $a_j G_0 \in \mathcal{H}_{k+1}$ , there is  $s_j$ such that  $\left\{a_j \delta_1^{s_j}, a_j \delta_1^{s_j+1}\right\}$  is an edge of C. From here, the result follows.

**Proof of Theorem 2.** We will prove the theorem by induction on the order of the group. For |G| = 3, we see that  $G \cong Z_3$  and the only possible normal Cayley graphs are  $Cay(Z_3, \{1\})$  and  $Cay(G, \{1, 2\})$  which are both hamiltonian graphs.

Let G be a finite group of order greater than 3,  $G = G_0 \supseteq G_1, \ldots, G_{l-1} \supseteq G_l$ be a composition series of G, and let  $\{\delta_0, \ldots, \delta_{l+1}\} \subseteq G$  such that, for each  $0 \leq i \leq l, G_i/G_{i+1} = \langle \delta_i G_{i+1}, \delta_{i+1} G_{i+1} \rangle$ . Let Cay(G, S) be a normal Cayley graph with  $\{\delta_0, \ldots, \delta_{l+1}\} \subseteq S$ .

Let  $S/G_1 = \{sG_1 : s \in S\}$  and consider the Cayley graph  $Cay(G/G_1, S/G_1)$ . If  $G/G_1$  is an abelian group, it is known that  $Cay(G/G_1, S/G_1)$  contains a hamiltonian cycle (see [4]). If  $Cay(G/G_1, S/G_1)$  is not an abelian group, consider the following.

# Claim 1. $Cay(G/G_1, S/G_1)$ is a normal Cayley graph.

**Proof.** Let  $g \in G$  and  $s \in S$ . Since  $G_1$  is a normal subgroup it follows that  $g^{-1}G_1sG_1gG_1 = g^{-1}sgG_1$  and since S is a normal connection set,  $g^{-1}sg = s_1 \in S$ . Therefore  $g^{-1}sgG_1 = s_1G_1 \in S/G_1$  and the claim follows.

Thus, by Claim 1,  $Cay(G/G_1, S/G_1)$  is a normal Cayley graph;  $G/G_1 = \langle \delta_0 G_1, \delta_1 G_1 \rangle$  is a simple non-abelian group and, by hypothesis,  $\{\delta_0, \delta_1\} \subseteq S$  which implies that  $\{\delta_0 G_1, \delta_1 G_1\} \subseteq S/G_1$ . Therefore, from Theorem 1 it follows that there is a hamiltonian cycle in  $Cay(G/G_1, S/G_1)$ .

Let  $C = (G_1, g_1G_1, \ldots, g_nG_1, G_1)$ , with  $n = |G/G_1|$ , be a hamiltonian cycle in  $Cay(G/G_1, S/G_1)$  (see Figure 4).



On the other hand, let  $S |_{G_1} = S \cap G_1$  and consider the Cayley graph  $Cay(G_1, S |_{G_1})$ .

Claim 2.  $Cay(G_1, S \mid_{G_1})$  is a normal Cayley graph.

**Proof.** Since S is a normal connection set and  $G_1$  is a normal subgroup of G we see that  $g^{-1}Sg = S$  and  $g^{-1}G_1g = G_1$ , so  $g^{-1}(S \mid_{G_1})g = g^{-1}(S \cap G_1)g = S \cap G_1 = S \mid_{G_1}$ .

Claim 3.  $\{\delta_1, \ldots, \delta_{l+1}\} \subseteq S \mid_{G_1}$ .

**Proof.** Since for each  $i \in \{0, \ldots, l+1\}$  we have  $G_i/G_{i+1} = \langle \delta_i G_{i+1}, \delta_{i+1} G_{i+1} \rangle$  it follows that  $\delta_i \in G_i \subset G_1$  and  $\delta_i \in G_1$  for all  $1 \le i \le l+1$ .

Clearly  $|G_1| < |G_0|$ , and since  $G_1 \succeq G_2, \ldots, G_{l-1} \succeq G_l$  is a composition series of  $G_1$ , by Claims 2 and 3 and by induction hypothesis we see that there is a hamiltonian cycle  $\mathcal{C}'$  in  $C(G_1, S \mid_{G_1})$ . Let  $\mathcal{C}' = (1, n_1, n_2, \ldots, n_{i-1}, 1)$  with  $i = |G_1|$ .

Let  $g_lG_1$  and  $g_{l+1}G_1$  be two consecutive vertices of the hamiltonian cycle C of  $Cay(G/G_1, S/G_1)$ . By definition  $(g_lG_1)^{-1}g_{l+1}G_1 \in S/G_1$  which implies that  $G_1g_l^{-1}g_{l+1}G_1 = s_1G_1$  with  $s_1 \in S$ . Thus  $g_l^{-1}g_{l+1} = s_1n_{l_1}$  with  $n_{l_1} \in G_1$  and then  $g_l^{-1}g_{l+1}n_{l_1}^{-1} = s_1 \in S$ . Therefore, for every  $n_j \in G_1$ , we see that

$$n_j^{-1}g_l^{-1}g_{l+1}n_{l_1}^{-1}n_j = n_j^{-1}s_1n_j \in S,$$

which implies that for every  $n_j \in G_1$ ,  $g_l n_j$  is adjacent to  $g_{l+1} n_{l_1}^{-1} n_j$  in Cay(G, S).

Observe that the map  $g_l n_j \to g_{l+1} n_{l_1}^{-1} n_j$  defines a bijection between  $g_l G_1$ and  $g_{l+1}G_1$ , and that, given  $\alpha, \beta \in G_1$ , we see that  $(g_l \alpha)^{-1} g_l \beta = \alpha^{-1} \beta \in S$  if and only if

738

$$\left(g_{l+1}n_{l_1}^{-1}\alpha\right)^{-1}\left(g_{l+1}n_{l_1}^{-1}\beta\right) = \alpha^{-1}n_{l_1}g_{l+1}^{-1}g_{l+1}n_{l_1}^{-1}\beta = \alpha^{-1}\beta \in S,$$

which implies that the subgraphs of Cay(G, S) induced by  $g_lG_1$  and  $g_{l+1}G_1$  are attached (see Figure 5).



Figure 5

From here, and by an analogous argument than in the proof for the case k = 1 in Theorem 1, we obtain a hamiltonian cycle in Cay(G, S) (see Figure 6).



Figure 6

## References

- [1] N. Alon and Y. Roichman, Random Cayley graphs and expanders, Random Structures Algorithms 5 (1994) 271–284. doi:10.1002/rsa.3240050203
- [2] J.A. Bondy and U.S.R. Murty, Graph Theory (Springer, New York, 2008).

- [3] J. Bourgain and A. Gamburd, Uniform expansion bounds for Cayley graphs of SL<sub>2</sub>(F<sub>p</sub>), Ann. of Math. 167 (2008) 625–642. doi:10.4007/annals.2008.167.625
- [4] C.C. Chen and N. Quimpo, On strongly hamiltonian abelian group graphs, Combin. Math. VIII (Geelong, 1980) Lecture Notes in Math. 884 (Springer, Berlin-New York, 1981) 23–34.
- [5] E. Durnberger, Connected Cayley graphs of semi-direct products of cyclic groups of prime order by abelian groups are hamiltonian, Discrete Math. 46 (1983) 55–68. doi:10.1016/0012-365X(83)90270-4
- [6] E. Ghaderpour and D. Witte Morris, Cayley graphs on nilpotent groups with cyclic commutator subgroup are hamiltonian, Ars Math. Contemp. 7 (2014) 55–72. doi:10.26493/1855-3974.280.8d3
- H.H. Glover, K. Kutnar, A. Malnič and D. Marušič, Hamilton cycles in (2, odd, 3)-Cayley graphs, Proc. Lond. Math. Soc. 104 (2012) 1171–1197. doi:10.1112/plms/pdr042
- [8] H.H. Glover, K. Kutnar and D. Marušič, Hamiltonian cycles in cubic Cayley graphs: the < 2, 4k, 3 > case, J. Algebraic Combin. 30 (2009) 447–475. doi:10.1007/s10801-009-0172-5
- [9] H.H. Glover and D. Marušič, Hamiltonicity of cubic Cayley graph, J. Eur. Math. Soc. 9 (2007) 775–787.
- [10] H.H. Glover and T.Y. Yang, A Hamilton cycle in the Cayley graph of the (2, p, 3)presentation of PSL2(p), Discrete Math. 160 (1996) 149–163. doi:10.1016/0012-365X(95)00155-P
- [11] K. Keating and D. Witte, On Hamilton cycles in Cayley graphs in groups with cyclic commutator subgroup, Annals of Discrete Math. 27 (1985) 89–102.
- [12] K. Kutnar, D. Marušič, D. Morris, J. Morris and P. Šparl, Hamiltonian cycles in Cayley graphs of small order, Ars Math. Contemp. 5 (2012) 27–71. doi:10.26493/1855-3974.177.341
- [13] D. Marušič, Hamilonian circuits in Cayley graphs, Discrete Math. 46 (1983) 49–54. doi:10.1016/0012-365X(83)90269-8
- I. Pak and R. Radoičić, Hamiltonian paths in Cayley graphs, Discrete Math. 309 (2009) 5501–5508.
  doi:0.1016/j.disc.2009.02.018
- [15] C. Praeger, *Finite normal edge-transitive Cayley graphs*, Bull. Aust. Math. Soc. **60** (1999) 207–220. doi:10.1017/S0004972700036340
- [16] J.J. Rotman, An Introduction to the Theory of Groups, Fourth Edition (Springer-Verlag, New York, 1995).
- [17] F. Menegazzo, The number of generator of a finite group, Irish Math. Soc. Bull. 50 (2003) 117–128.

- [18] P.E. Schupp, On the structure of hamiltonian cycles in Cayley graphs of finite quotients of the modular group, Theoret. Comput. Sci. 204 (1998) 233-248. doi:10.1016/S0304-3975(98)00041-3
- [19] C. Wang, D. Wang and M. Xu, Normal Cayley graphs of finite groups, Sci. China Ser. A 41 (1998) 242–251. doi:10.1007/BF02879042
- [20] D. Witte Morris, Odd-order Cayley graphs with commutator subgroup of order pq are hamiltonian, Ars Math. Contemp. 8 (2015) 1–28. doi:10.26493/1855-3974.330.0e6

Received 30 December 2017 Revised 12 June 2018 Accepted 19 March 2019