Discussiones Mathematicae Graph Theory 42 (2022) 5–13 https://doi.org/10.7151/dmgt.2234

DEGREE SUM CONDITION FOR THE EXISTENCE OF SPANNING *k*-TREES IN STAR-FREE GRAPHS

Michitaka Furuya

College of Liberal Arts and Sciences Kitasato University 1-15-1 Kitasato, Minami-ku, Sagamihara, Kanagawa 252-0373, Japan e-mail: michitaka.furuya@gmail.com

Shun-ichi Maezawa

Graduate School of Environment and Information Sciences Yokohama National University 79-1 Tokiwadai, Hodogaya-ku, Yokohama, Kanagawa 240-8501, Japan e-mail: maezawa-shunichi-bg@ynu.jp

Ryota Matsubara, Haruhide Matsuda

Department of Mathematics Shibaura Institute of Technology 307 Fukasaku, Minuma-ku, Saitama 337-8577, Japan

e-mail: ryota@sic.shibaura-it.ac.jp hmatsuda@sic.shibaura-it.ac.jp

Shoichi Tsuchiya

School of Network and Information Senshu University 2-1-1 Higashimita, Tama-ku, Kawasaki-shi, Kanagawa 214-8580, Japan **e-mail:** s.tsuchiya@isc.senshu-u.ac.jp

AND

TAKAMASA YASHIMA

Depatment of Mathmatics Keio University 3-14-1 Hiyoshi, Kohoku-ku, Yokohama-shi, Kanagawa 223-8522, Japan **e-mail:** takamasa.yashima@gmail.com

Abstract

For an integer $k \ge 2$, a k-tree T is defined as a tree with maximum degree at most k. If a k-tree T spans a graph G, then T is called a spanning k-tree of G. Since a spanning 2-tree is a Hamiltonian path, a spanning k-tree is an extended concept of a Hamiltonian path. The first result, implying the existence of k-trees in star-free graphs, was by Caro, Krasikov, and Roditty in 1985, and independently, Jackson and Wormald in 1990, who proved that for any integer k with $k \ge 3$, every connected $K_{1,k}$ -free graph contains a spanning k-tree. In this paper, we focus on a sharp condition that guarantees the existence of a spanning k-tree in $K_{1,k+1}$ -free graphs. In particular, we show that every connected $K_{1,k+1}$ -free graph G has a spanning k-tree if the degree sum of any 3k-3 independent vertices in G is at least |G|-2, where |G| is the order of G.

Keywords: spanning tree, k-tree, star-free, degree sum condition.

2010 Mathematics Subject Classification: 05C05, 05C50.

1. INTRODUCTION AND MAIN RESULT

In this paper, we consider only finite and simple graphs. Let G be a graph. We denote the order of G by |G|. For a vertex $x \in V(G)$, we denote the degree of x in G by $\deg_G(x)$ and the set of vertices adjacent to x in G by $N_G(x)$. The independence number of a graph G is denoted by $\alpha(G)$. For an integer $k \geq 2$ and a graph G, we define

$$\sigma_k(G) = \min_{S \subseteq V(G)} \left\{ \sum_{x \in S} \deg_G(x) \mid S \text{ is an independent set of } k \text{ vertices} \right\}$$

if $\alpha(G) \ge k$, and define $\sigma_k(G) = \infty$ if $\alpha(G) < k$.

Let $K_{1,m}$ denote the star with *m* leaves. For a graph *G* and a given graph *H*, *G* is called *H*-free if *G* contains no induced subgraph isomorphic to *H*.

The existence of a Hamiltonian path in a given graph has been much studied. In particular, if a graph satisfies any of a number of density conditions, a Hamiltonian path is guaranteed to exist. The following result is one of the best known among these density conditions.

Theorem 1 (Ore [5]). Let G be a graph. If $\sigma_2(G) \ge |G| - 1$, then G has a Hamiltonian path.

This theorem has led to many new results and conjectures concerning paths and cycles in graphs. One direction is motivated by the fact that a Hamiltonian path is a spanning tree with small maximum degree. So it is natural to ask how Theorem 1 might be generalized to guarantee the existence of a spanning tree with maximum degree at most $k \geq 3$.

For an integer $k \ge 2$, a *k*-tree *T* is defined as a tree of maximum degree at most *k*. If a *k*-tree *T* spans a graph *G*, then *T* is called a *spanning k*-tree of *G*. Note that a spanning 2-tree is a Hamiltonian path.

For general graphs, Win gave a degree sum condition for the existence of a spanning k-tree as a generalization of Theorem 1 as follows.

Theorem 2 (Win [6]). Let $k \ge 2$ be an integer, and let G be a connected graph. If $\sigma_k(G) \ge |G| - 1$, then G has a spanning k-tree.

By restricting graphs to be star-free, Caro, Krasikov, and Roditty in 1985 and independently, Jackson and Wormald in 1990, obtained the following result, which guarantees the existence of a spanning k-tree.

Theorem 3 (Caro, Krasikov, and Roditty [2], Jackson and Wormald [3]). For an integer $k \geq 3$, every connected $K_{1,k}$ -free graph contains a spanning k-tree.

Theorem 3 is best possible in the sense that there exist infinitely many connected $K_{1,k+1}$ -free graphs which have no spanning k-tree. Thus some additional conditions are needed for connected $K_{1,k+1}$ -free graphs to have a spanning ktree. In fact, for the case when k = 2, Liu and Tian in 1986 and independently, Broersma in 1988, obtained the following result.

Theorem 4 (Liu, Tian, and Wu [4], Broersma [1]). Let G be a connected $K_{1,3}$ -free graph. If

$$\sigma_3(G) \ge |G| - 2,$$

then G has a Hamiltonian path.

The purpose of this paper is to give a degree sum condition for connected $K_{1,k+1}$ -free graphs to have a spanning k-tree. Our main result is the following.

Theorem 5. Let k be an integer with $k \ge 2$. If a connected $K_{1,k+1}$ -free graph G satisfies

$$\sigma_{3k-3}(G) \ge |G| - 2,$$

then G has a spanning k-tree.

Theorem 5 gives a generalization of Theorem 4. By Theorem 5, we also obtain an upper bound on the independence number $\alpha(G)$ for $K_{1,k+1}$ -free graphs to have a spanning k-tree.

Corollary 6. Let k be an integer with $k \ge 2$. If a connected $K_{1,k+1}$ -free graph G satisfies

 $\alpha(G) \le 3k - 4,$

then G has a spanning k-tree.

The degree sum condition of Theorem 5 is sharp as shown in Section 2 and the example also shows the sharpness of the independence number in Corollary 6.

2. Sharpness of Main Theorem

Let K_n denote the complete graph of order n. For two graphs G and H, let $G \cup H$ be the union of G and H.

We show that the lower bounds of $\sigma_{3k-3}(G)$ in Theorem 5 and the independence number in Corollary 6 are best possible. In fact, we give the following example.



Figure 1. An infinite family of connected $K_{1,k+1}$ -free graphs G having no spanning k-tree and satisfying $\sigma_{3k-3}(G) = |G| - 3$.

Let $k \ge 2$ and $m \ge 1$ be integers. Let T be a triangle with $V(T) = \{x_1, x_2, x_3\}$. For each i = 1, 2, 3, define a graph H_i as k - 1 disjoint copies of K_m . The graph G is obtained by joining x_i and all the vertices in $V(H_i)$ for each i = 1, 2, 3. Then G has no induced subgraph isomorphic to $K_{1,k+1}$ and |G| = 3m(k-1) + 3. Since $\alpha(H_1 \cup H_2 \cup H_3) = 3k - 3$, we can choose 3k - 3 independent vertices one by one from each complete graph K_m . Then $\sigma_{3k-3}(G) = 3m(k-1) = |G| - 3$. For any spanning tree T of G, one of three vertices x_1, x_2 and x_3 must have degree more than k in T. Hence G has no spanning k-tree, and thus the lower bounds of $\sigma_{3k-3}(G)$ in Theorem 5 and the independence number in Corollary 6 are sharp.

Note that the graphs in Figure 1 show that $K_{1,k}$ -freeness in Theorem 3 cannot be replaced by $K_{1,k+1}$ -freeness.

3. Proof of Theorem 5

Let T be a tree and let v be a vertex of T. The *outdirected tree* with respect to (T, v) is the directed tree obtained from T in which all the edges are directed away from v. The *out-neighborhood* $N_{T,v}^+(x)$ of a vertex x of T is the set of vertices adjacent from x in the outdirected tree with respect to (T, v).

Proof of Theorem 5

Let k be an integer with $k \geq 2$, and let G be a connected $K_{1,k+1}$ -free graph satisfying $\sigma_{3k-3}(G) \geq |G| - 2$. The case k = 2 follows from Theorem 4. Thus we consider the case when $k \geq 3$. Let T be a maximal k-tree of G. Suppose that T is not a spanning tree of G. Then G has a vertex u_0 not contained in T which is adjacent to a vertex v in V(T).

Claim 7. $\deg_T(v) = k$.

Proof. Suppose that $\deg_T(v) \neq k$. Since T is a k-tree, $\deg_T(v) < k$. Then $T + vu_0$ is a k-tree of order |V(T)| + 1. This contradicts the maximality of T. Hence $\deg_T(v) = k$.

Let S_1, S_2, \ldots, S_k denote the components of T - v. For each $1 \leq i \leq k$, let s_i be the vertex of S_i which is adjacent to v in T. Note that $\deg_{S_i}(s_i) \leq k-1$ for each i.

Claim 8. For each $1 \leq i \leq k$, u_0 is nonadjacent to s_i in G.

Proof. Suppose that $u_0 s_i \in E(G)$ for some $1 \le i \le k$. Then $T + vu_0 + u_0 s_i - vs_i$ is a k-tree of order |V(T)| + 1, which contradicts the maximality of T.

Since v is a common neighbor of $u_0, s_1, s_2, \ldots, s_k$ in G, by the $K_{1,k+1}$ -freeness of G and Claim 8, s_i and s_j are adjacent in G for some $1 \le i < j \le k$. Without loss of generality, we may assume that $s_{k-1}s_k \in E(G) \setminus E(T)$.

Claim 9. $\deg_T(s_{k-1}) = \deg_T(s_k) = k$.

Proof. By symmetry, it suffices to show that $\deg_T(s_k) = k$. If $\deg_T(s_k) \neq k$, then deg_T(s_k) < k because T is a k-tree, and hence $T + s_{k-1}s_k + u_0v - vs_k$ is a k-tree of order |V(T)| + 1. This contradicts the maximality of T.

As seen in Figure 2, we redefine $T_i = S_i$ and $t_i = s_i$ for each $1 \le i \le k-2$ and let $T_{k-1}, \ldots, T_{2k-3}$ and $T_{2k-2}, \ldots, T_{3k-4}$ be the components of $S_{k-1} - s_{k-1}$ and $S_k - s_k$, respectively. Let $t_{k-1}, \ldots, t_{2k-3}$ (respectively, $t_{2k-2}, \ldots, t_{3k-4}$) denote the vertices of $T_{k-1}, \ldots, T_{2k-3}$ (respectively, $T_{2k-2}, \ldots, T_{3k-4}$) which are adjacent to s_{k-1} (respectively, s_k) in T. Since $T_1, T_2, \ldots, T_{3k-4}$ are vertex-disjoint k-trees, we can choose a leaf $u_i \in V(T_i)$ of T for each $1 \leq i \leq 3k - 4$. By the maximality of T and $\deg_T(u_i) = 1$, $N_G(u_i) \subseteq V(T)$ for each $1 \le i \le 3k - 4$.

Claim 10. The set $\{u_0, u_1, \ldots, u_{3k-4}\}$ is an independent set of G.

Proof. For $1 \leq i \leq 3k-4$, since $N_G(u_i) \subseteq V(T)$, we have $u_0u_i \notin E(G)$. Suppose that $u_i u_j \in E(G)$ for some $1 \le i < j \le 3k - 4$. Consider the following tree T_A ,

$$T_A = \begin{cases} T + u_i u_j + u_0 v - v t_i & \text{if } 1 \le i \le k - 2, \\ T + u_i u_j + u_0 v + s_{k-1} s_k - v s_k - s_{k-1} t_i & \text{if } k - 1 \le i \le 2k - 3, \\ T + u_i u_j + u_0 v + s_{k-1} s_k - v s_{k-1} - s_k t_i & \text{if } 2k - 2 \le i \le 3k - 4. \end{cases}$$



Figure 2. A maximal k-tree T.

Then T_A is a k-tree of order |V(T)| + 1, which contradicts the maximality of T. Hence the claim holds.

For each $1 \leq i \leq 3k - 4$, define

$$W_i = \left(\bigcup_{0 \le j \le 3k-4, j \ne i} N_G(u_j)\right) \cap V(T_i).$$

Claim 11. For each $1 \leq i \leq 3k - 4$, $t_i \notin W_i$.

Proof. If $t_i \in W_i$ for some $1 \le i \le 3k - 4$, then t_i is adjacent to a leaf u_j of T_j with $j \ne i$ or to the vertex u_0 . Consider the following tree T_B ,

$$T_B = \begin{cases} T + t_i u_j + u_0 v - v t_i & \text{if } 1 \le i \le k - 2, \\ T + t_i u_j + s_{k-1} s_k + u_0 v - s_{k-1} t_i - v s_k & \text{if } k - 1 \le i \le 2k - 3, \\ T + t_i u_j + s_{k-1} s_k + u_0 v - s_k t_i - v s_{k-1} & \text{if } 2k - 2 \le i \le 3k - 4. \end{cases}$$

Then T_B is a k-tree of order |V(T)| + 1, which contradicts the maximality of T. Consequently, $t_i \notin W_i$ for each $1 \le i \le 3k - 4$.

Claim 12. For each $1 \le i \le 3k - 4$, any vertex $w \in W_i$ satisfies the following three statements:

- (i) $\deg_T(w) = k;$
- (ii) no vertex u_j with $1 \le j \le 3k 4$ is adjacent to any vertex of $N^+_{T_i,u_i}(w)$ in G; and
- (iii) $|(N_G(w) \cap \{u_0, u_1, u_2, \dots, u_{3k-4}\}) \setminus \{u_i\}| \le k-1.$

Degree Sum Condition for the Existence of Spanning k-Trees ... 11

Proof. (i) Suppose that $\deg_T(w) \neq k$ for some $w \in W_i$ with $1 \leq i \leq 3k - 4$. Since T is a k-tree, $\deg_T(w) < k$. By the definition of W_i , w is adjacent to a vertex u_j with $j \neq i$ in G (possibly, j = 0). Consider the following tree T_C ,

$$T_{C} = \begin{cases} T + u_{j}w + u_{0}v - vt_{i} & \text{if } 1 \leq i \leq k-2, \\ T + u_{j}w + s_{k-1}s_{k} + u_{0}v - s_{k-1}t_{i} - vs_{k} & \text{if } k-1 \leq i \leq 2k-3, \\ T + u_{j}w + s_{k-1}s_{k} + u_{0}v - s_{k}t_{i} - vs_{k-1} & \text{if } 2k-2 \leq i \leq 3k-4. \end{cases}$$

Then T_C is a k-tree of order |V(T)| + 1, which contradicts the maximality of T. Hence $\deg_T(w) = k$ as desired.

(ii) Suppose that for some $1 \leq j \leq 3k - 4$, u_j is adjacent to a vertex $w^+ \in N^+_{T_i,u_i}(w)$ in G. By the definition of W_i , w is adjacent to a leaf u_ℓ with $\ell \neq i$ or to the vertex u_0 . Note that $w \neq t_i$ by Claim 11. Consider the following k-tree T_D ,

$$T_D = \begin{cases} T + u_\ell w + u_j w^+ + u_0 v - vt_i - ww^+ & \text{if } 1 \le i \le k-2, \\ T + u_\ell w + u_j w^+ + s_{k-1} s_k + u_0 v - vs_k - s_{k-1} t_i - ww^+ & \text{if } k-1 \le i \le 2k-3, \\ T + u_\ell w + u_j w^+ + s_{k-1} s_k + u_0 v - vs_{k-1} - s_k t_i - ww^+ & \text{if } 2k-2 \le i \le 3k-4. \end{cases}$$

Then T_D is a k-tree of order |V(T)| + 1. This contradicts the maximality of T.

(iii) To the contrary, assume that $|(N_G(w) \cap \{u_0, u_1, u_2, \dots, u_{3k-4}\}) \setminus \{u_i\}| \ge k$. Since deg_T(w) = $k \ge 3$ by Claim 12(i), a vertex $w_1 \in N_{T_i,u_i}^+(w)$ exists. Note that both w_1 is different from any u_j with $j \ne i$ because $w_1 \in V(T_i)$ and $(\{u_0, u_1, u_2, \dots, u_{3k-4}\} \setminus \{u_i\}) \cap V(T_i) = \emptyset$. Then w_1 and k vertices in $(N_G(w) \cap \{u_0, u_1, u_2, \dots, u_{3k-4}\}) \setminus \{u_i\}$ are all neighbors of w in G. Moreover, Claims 10 and 12(ii) assart that w_1 and k vertices in $(N_G(w) \cap \{u_0, u_1, u_2, \dots, u_{3k-4}\}) \setminus \{u_i\}$ are independent in G. This contradicts the assumption that G is $K_{1,k+1}$ -free. Hence $|(N_G(w) \cap \{u_0, u_1, u_2, \dots, u_{3k-4}\}) \setminus \{u_i\}| \le k-1$.

Claim 13. We have $|N_G(s_i) \cap \{u_0, u_1, \dots, u_{3k-4}\}| \le k-1$ for each i = k-1 and i = k.

Proof. We first prove that

$$N_G(s_k) \cap \{u_0, u_1, \dots, u_{3k-4}\} \subseteq \{u_{2k-2}, \dots, u_{3k-4}\}.$$

By Claim 8, $s_k u_0 \notin E(G)$. If $s_k u_i \in E(G)$ for some $1 \leq i \leq 2k-3$, then $T+s_k u_i+u_0v-vs_k$ is a k-tree of order |V(T)|+1. This contradicts the maximality of T. Hence $N_G(s_k) \cap \{u_0, u_1, \ldots, u_{3k-4}\} \subseteq \{u_{2k-2}, \ldots, u_{3k-4}\}$ as desired. This implies that $|N_G(s_k) \cap \{u_0, u_1, \ldots, u_{3k-4}\}| \leq k-1$. By symmetry, applying the preceding argument, we obtain the claim for the case when i = k-1.

Claim 14. $|N_G(v) \cap \{u_0, u_1, \dots, u_{3k-4}\}| \le k-1.$

Proof. We show that $N_G(v) \cap \{u_0, u_1, \dots, u_{3k-4}\} \subseteq \{u_0, \dots, u_{k-2}\}$. Suppose that $vu_i \in E(G)$ for some $k - 1 \leq i \leq 3k - 4$. Then $T + u_iv + s_{k-1}s_k + u_0v - s_{k-1}v - s_kv$ is a k-tree of order |V(T)| + 1. This contradicts the maximality of T. Hence $N_G(v) \cap \{u_0, u_1, \dots, u_{3k-4}\} \subseteq \{u_0, \dots, u_{k-2}\}$. Thus $|N_G(v) \cap \{u_0, u_1, \dots, u_{3k-4}\} \leq \{u_0, \dots, u_{k-2}\}$.

By Claim 12(i), $|N^+_{T_i,u_i}(w)| = k - 1$ for any $w \in W_i$ with $1 \le i \le 3k - 4$. It follows from Claim 12(ii) that

(1)
$$|N_G(u_i) \cap V(T_i)| \leq |V(T_i)| - (k-1)|W_i| - |\{u_i\}|$$
$$= |V(T_i)| - (k-1)|W_i| - 1.$$

For each $0 \le j \le 3k - 4$ with $j \ne i$, Claim 12(iii) asserts that

(2)
$$\sum_{0 \le j \le 3k-4, j \ne i} |N_G(u_j) \cap V(T_i)| \le (k-1)|W_i|.$$

By (1) and (2), we obtain

$$\sum_{0 \le j \le 3k-4} |N_G(u_j) \cap V(T_i)| \le |V(T_i)| - 1.$$

Hence we obtain

(3)

$$\sum_{1 \le i \le 3k-4} \sum_{0 \le j \le 3k-4} |N_G(u_j) \cap V(T_i)| \le \sum_{1 \le i \le 3k-4} (|V(T_i)| - 1)$$

$$\le |T| - |\{s_{k-1}, s_k, v\}| - (3k - 4)$$

$$= |T| - 3k + 1.$$

By (3), Claims (13) and (14),

$$\sum_{0 \le i \le 3k-4} \deg_G(u_i) \le |T| - 3k + 1 + (k-1)|\{s_{k-1}, s_k, v\}| + |N_{G-V(T)}(u_0)|$$
$$\le |T| - 2 + |G| - |T| - |\{u_0\}| = |G| - 3.$$

This contradicts the degree sum condition of Theorem 5 and hence the proof of Theorem 5 is completed. $\hfill\blacksquare$

Acknowledgments

The authors would like to thank the anonymous referee for valuable comments and suggestions. This work was partially supported by JSPS KAKENHI Grant numbers JP18K13449 (to M.F.), JP15K04980 (to H.M.), JP19K14584 (to S.T.), and by Grant for Basic Science Research Projects from The Sumitomo Foundation (to S.T). Degree Sum Condition for the Existence of Spanning k-Trees ... 13

References

- H.J. Broersma, Hamilton Cycles in Graphs and Related Topics (Ph.D. Thesis, University of Twente, 1988).
- [2] Y. Caro, I. Krasikov and Y. Roditty, Spanning trees and some edge reconstructible graphs, Ars Combin. 20 A (1985) 109–118.
- [3] B. Jackson and N.C. Wormald, k-walks of graphs, Australas. J. Combin. 2 (1990) 135–146.
- [4] Y. Liu, F. Tian and Z. Wu, Some results on longest paths and cycles in K_{1,3}-free graphs, J. Changsha Railway Inst. 4 (1986) 105–106.
- [5] O. Ore, Notes on Hamilton circuits, Amer. Math. Monthly 67 (1960) 55. https://doi.org/10.2307/2308928
- S. Win, Existenz von Gerüsten mit vorgeschriebenem Maximalgrad in Graphen, Abh. Math. Sem. Univ. Hamburg 43 (1975) 263–267. https://doi.org/10.1007/BF02995957

Received 12 February 2018 Revised 5 April 2019 Accepted 10 June 2019