Discussiones Mathematicae Graph Theory 42 (2022) 309–313 https://doi.org/10.7151/dmgt.2249

ON HAMILTONIAN CYCLES IN CLAW-FREE CUBIC GRAPHS

ELENA MOHR AND DIETER RAUTENBACH

Institute of Optimization and Operations Research Ulm University, Germany e-mail: {elena.mohr,dieter.rautenbach}@uni-ulm.de

Abstract

We show that every claw-free cubic graph of order n at least 8 has at most $2^{\lfloor \frac{n}{4} \rfloor}$ Hamiltonian cycles, and we also characterize all extremal graphs. **Keywords:** Hamiltonian cycle, claw-free graph, cubic graph. **2010 Mathematics Subject Classification:** 05C70.

1. INTRODUCTION

Chia and Thomassen [2] proved that every cubic multigraph of order n at least 4 has at most $2^{n/2}$ Hamiltonian cycles. They asked whether there is a cubic graph of order n at least 6 that has more than $2^{n/3}$ Hamiltonian cycles. As they observed the unique connected cubic graph of order n, where n is a multiple of 6, that arises by adding a perfect matching to the disjoint union of n/6 copies of $K_{3,3} - e$ has exactly $2^{n/3}$ Hamiltonian cycles, that is, this bound can be achieved. In the present note, we show that every claw-free cubic graph of order n at least 8 has at most $2^{\lfloor \frac{n}{4} \rfloor}$ Hamiltonian cycles, and we also characterize all extremal graphs, which are structurally similar to the above graphs based on $K_{3,3} - e$. Recall that a graph is *claw-free* if it does not contain $K_{1,3}$ as an induced subgraph.



Figure 1. The diamond D and the graph D'.

Let \mathcal{G} be the set of connected cubic graphs G of some order n such that

- if $n \equiv 0 \pmod{4}$, then G arises by adding a matching M to the disjoint union of n/4 copies of the diamond D, and
- if n ≡ 2 (mod 4), then G arises by adding a matching M to the disjoint union of one copy of D' and (n − 6)/4 copies of the diamond D.

Note that, for every even order n at least 4, the set \mathcal{G} contains exactly one graph of order n.

Our result is the following.

Theorem 1. If G is a claw-free cubic graph of order n at least 8, then G has at most $2^{\lfloor \frac{n}{4} \rfloor}$ Hamiltonian cycles with equality if and only if $G \in \mathcal{G}$.

We need two preparatory lemmas. For some non-negative integer k, let f_k denote the k-th Fibonacci number, that is, $f_0 = 0$, $f_1 = 1$, and $f_k = f_{k-1} + f_{k-2}$ for $k \ge 2$. The following fact is known [1]; in order to keep the paper self-contained, we add a simple proof.

Lemma 2. For every integer k at least 3, there are $f_{k+1} + f_{k-1}$ sets F of edges of the cycle C_k of order k such that $C_k - F$ has no isolated vertex.

Proof. Let $C_k = u_1 \cdots u_k u_1$, and let \mathcal{F} be the set of all considered sets F, that is, we need to show that $a_k = f_{k+1} + f_{k-1}$ for $a_k = |\mathcal{F}|$. Let b_k be the number of sets F' of edges of the path $P_{k+1} : v_1 \cdots v_{k+1}$ of order k+1 such that none of the vertices v_1, \ldots, v_k is isolated in $P_{k+1} - F'$; note that v_{k+1} is allowed to be isolated in $P_{k+1} - F'$. Clearly, $b_0 = 1$, $b_1 = 1$, and $b_2 = 2$. Furthermore, considering the edge $v_i v_{i+1}$ of P_{k+1} minimizing i that belongs to such a set F', it follows that $b_k = b_{k-2} + b_{k-3} + \cdots + b_0 + 1$, which implies $b_k = b_{k-1} + b_{k-2}$. The initial values and the recursion imply that $b_k = f_{k+1}$. Now, $a_k = 2b_{k-2} + b_{k-1} = 2f_{k-1} + f_k = f_{k+1} + f_{k-1}$, because there are

- b_{k-2} sets in \mathcal{F} that contain u_1u_2 (consider the path $u_2\cdots u_k$),
- b_{k-2} sets in \mathcal{F} that contain $u_k u_1$ (consider the path $u_1 \cdots u_{k-1}$), and
- b_{k-1} sets in \mathcal{F} that contain neither u_1u_2 nor u_ku_1 (consider the path $u_1\cdots u_k$).

Lemma 3. $2^{\lfloor \frac{6\ell}{4} \rfloor} > f_{2\ell+1} + f_{2\ell-1}$ for every $\ell \in \mathbb{N} \setminus \{1,3\}$.

Proof. For small values of ℓ , this is easily verified. For $\ell \geq 5$, this follows easily by induction using

$$2^{\left\lfloor\frac{6\ell}{4}\right\rfloor} \ge 2^{\frac{3\ell-1}{2}} > 0.707 \cdot 2.828^{\ell}$$

and

$$f_{2\ell+1} + f_{2\ell-1} = \frac{1}{\sqrt{5}} \left(\left(\frac{1+\sqrt{5}}{2} \right)^{2\ell+1} - \left(\frac{1-\sqrt{5}}{2} \right)^{2\ell+1} + \left(\frac{1+\sqrt{5}}{2} \right)^{2\ell-1} - \left(\frac{1-\sqrt{5}}{2} \right)^{2\ell-1} \right)$$
$$= \left(\frac{1+\sqrt{5}}{2} \right)^{2\ell} + \left(\frac{1-\sqrt{5}}{2} \right)^{2\ell} < 2.619^{\ell} + 0.382.$$

Proof of Theorem 1. If G lies in \mathcal{G} and has order n at least 8, then every Hamiltonian cycle of G contains each edge of the matching M mentioned in the definition of \mathcal{G} . Since the diamond D and the graph D' both contain exactly two paths between their two vertices of degree 2, it follows that G has exactly $2^{\lfloor n/4 \rfloor}$ Hamiltonian cycles. It remains to show that every claw-free cubic graph G of order n at least 8 has at most $2^{\lfloor n/4 \rfloor}$ Hamiltonian cycles with equality only if $G \in \mathcal{G}$. Suppose, for a contradiction, that G is a counterexample of minimum order n. Let $C: u_1 u_2 \cdots u_n u_1$ be some Hamiltonian cycle of G, where we identify indices modulo n. The edges in $E(G) \setminus E(C)$ are *chords* of C, and a chord $u_i u_j$ is *short* if u_i and u_j have distance 2 on C.

Suppose, for a contradiction, that u_2u_4 and u_3u_5 are short chords, that is, there are two crossing short chords. If u_1 and u_6 are adjacent, then the clawfreeness implies that u_1 and u_6 have a common neighbor, which implies the contradiction that u_7 equals u_n , that is, n = 7. Hence, the vertices u_1 and u_6 are not adjacent, and $G' = G - \{u_2, u_3, u_4, u_5\} + u_1 u_6$ is a Hamiltonian cubic claw-free graph of order n' = n - 4. If n' = 4, then G' is K_4 , and G arises by adding a matching to the disjoint union of two copies of D, which implies the contradiction $G \in \mathcal{G}$. If n' = 6, then the claw-freeness of G implies that G' arises by adding the perfect matching $\{u_1u_6, u_7u_{10}, u_8u_9\}$ to the disjoint union of the two triangles $u_1u_9u_{10}u_1$ and $u_6u_7u_8u_6$. It follows that G arises by adding a matching to the disjoint union of one copy of D' and one copy of D, which implies the contradiction $G \in \mathcal{G}$. Now, if $n' \geq 8$, then the choice of G implies that G' is no counterexample. Note that Hamiltonian cycles of G' that do not contain the edge u_1u_6 do not correspond to Hamiltonian cycles of G, and that Hamiltonian cycles of G' that contain the edge u_1u_6 correspond to two distinct Hamiltonian cycles of G; one using the path $u_1u_2u_3u_4u_5u_6$ and one using the path $u_1u_2u_4u_3u_5u_6$. This implies that G has at most $2 \cdot 2^{\lfloor n'/4 \rfloor} = 2^{\lfloor n/4 \rfloor}$ Hamiltonian cycles with equality only if every Hamiltonian cycle of G' contains the edge u_1u_6 and G' has $2^{\lfloor n'/4 \rfloor}$ Hamiltonian cycles, that is, $G' \in \mathcal{G}$. Since G is claw-free, the edge u_1u_6 is one of the edges of G' that belong to some 2-edge-cut of G', which implies the contradiction $G \in \mathcal{G}$. Altogether, we obtain, that two crossing short chords do not exist.

Since G is claw-free, it follows that, for every two consecutive vertices on C, either one vertex is incident with a short chord and one vertex is incident with a non-short chord, or both vertices are incident with two non-crossing short chords. This implies that n is a multiple of 3, that is, n = 3k for some positive

integer k. By symmetry, we may assume that $u_{3i-2}u_{3i}$ is a short chord for every $i \in [k]$, where [k] is the set of positive integers at most k. For every i in [k], every Hamiltonian cycle of G uses one or both of the edges $u_{3i-3}u_{3i-2}$ and $u_{3i}u_{3i+1}$. More precisely, if some Hamiltonian cycle of G uses both these edges, then it contains the subpath $u_{3i-3}u_{3i-2}u_{3i-1}u_{3i}u_{3i+1}$, if it uses $u_{3i-3}u_{3i-2}$ but not $u_{3i}u_{3i+1}$, then it contains the subpath $u_{3i-3}u_{3i-2}u_{3i}u_{3i-1}$, and if it uses $u_{3i}u_{3i+1}$ but not $u_{3i-3}u_{3i-2}$, then it contains the subpath $u_{3i-3}u_{3i-2}u_{3i}u_{3i-1}$, and if it uses $u_{3i}u_{3i+1}$ but not $u_{3i-3}u_{3i-2}$, then it contains the subpath $u_{3i-1}u_{3i-2}u_{3i}u_{3i+1}$. This implies that every Hamiltonian cycle C' of G is uniquely determined by the intersection of E(C') with the set $\{u_{3i}u_{3i+1}: i \in [k]\}$, and that $\{u_{3i}u_{3i+1}: i \in [k]\} \setminus E(C')$ does not contain two of the edges in $\{u_{3i}u_{3i+1}: i \in [k]\}$ that appear consecutively on C. By Lemma 2, it follows that the number of Hamiltonian cycles of G is at most $f_{\frac{n}{3}+1} + f_{\frac{n}{3}-1}$. Since G is cubic, the order n of G is even, which implies that k is even, that is, $k = 2\ell$ and $n = 6\ell$ for some integer $\ell \geq 2$. By Lemma 3, $2^{\lfloor n/4 \rfloor} = 2^{\lfloor 6\ell/4 \rfloor} > f_{2\ell+1} + f_{2\ell-1} = f_{\frac{n}{3}+1} + f_{\frac{n}{3}-1}$ unless $\ell = 3$, that is, n = 18. For n = 18, there are four different possibilites for the structure of G shown in Figure 2.



Figure 2. Every Hamiltonian cycle of the top left graph uses the four edges that lie in 2-edge-cuts of this graph, which easily implies that it has exactly four Hamiltonian cycles. Every Hamiltonian cycle of the top right graph uses either both the edges u_8u_{14} and $u_{11}u_{17}$ or none of these two edges, which easily implies that it has exactly four Hamiltonian cycles. The bottom left graph has one Hamiltonian cycle using no non-short chord and two using all three non-short chords, which implies that it has exactly three Hamiltonian cycles. The bottom right graph has one Hamiltonian cycle using no nonshort chord, three Hamiltonian cycles using two non-short chords, and two Hamiltonian cycles using all three non-short chord, which implies that it has exactly six Hamiltonian cycles.

Each of these graphs has at most six Hamiltonian cycles. Since this is less than $2^{\lfloor 18/4 \rfloor} = 16$, the proof is complete.

References

- [1] A.T. Benjamin and J.J. Quinn, *Recounting Fibonacci and Lucas identities*, College Math. J. **30** (1999) 359–366. https://doi.org/10.1080/07468342.1999.11974086
- [2] G.L. Chia and C. Thomassen, On the number of longest and almost longest cycles in cubic graphs, Ars Combin. 104 (2012) 307–320.

Received 16 April 2019 Revised 23 September 2019 Accepted 27 September 2019