Discussiones Mathematicae Graph Theory 42 (2022) 263–275 https://doi.org/10.7151/dmgt.2252

PACKING TREES IN COMPLETE BIPARTITE GRAPHS

JIEYAN WANG

Moscow High School Moscow, Idaho, USA 83843 e-mail: jieyan.wang17@gmail.com

Abstract

An embedding of a graph H in a graph G is an injection (i.e., a oneto-one function) σ from the vertices of H to the vertices of G such that $\sigma(x)\sigma(y)$ is an edge of G for all edges xy of H. The image of H in G under σ is denoted by $\sigma(H)$. A k-packing of a graph H in a graph G is a sequence $(\sigma_1, \sigma_2, \ldots, \sigma_k)$ of embeddings of H in G such that $\sigma_1(H), \sigma_2(H), \ldots, \sigma_k(H)$ are edge disjoint. We prove that for any tree T of order n, there is a 4-packing of T in a complete bipartite graph of order at most n + 12.

Keywords: packing, placement, edge-disjoint tree, bipartite graph. 2010 Mathematics Subject Classification: 05C70.

1. INTRODUCTION

Graphs in this article are simple and finite, which have no multiple edges nor loops. We use [4] for standard terminology and notation in graph theory. For any graph G, we use V(G) and E(G) to denote the vertex set and the edge set of G, respectively. If G is a bipartite graph with a given bipartition, then denote it by G = (X, Y; E). A tree is a connected graph without cycles. A star of order n is a tree of order n with a vertex of degree n - 1. We use B_n to represent a complete bipartite graph $K_{a,b}$ with n = a + b. Thus B_n is not uniquely defined for $n \ge 4$ because there is more than one positive integer partition for n = a + b.

An embedding of a graph H in a graph G is an injection (i.e., a one-to-one function) σ from the vertices of H to the vertices of G such that $\sigma(x)\sigma(y)$ is an edge of G for all edges xy of H. The image of H in G under σ is denoted by $\sigma(H)$. A *k*-packing of a graph H in a graph G is a sequence $(\sigma_1, \sigma_2, \ldots, \sigma_k)$ of k embeddings of H in G such that $\sigma_1(H), \sigma_2(H), \ldots, \sigma_k(H)$ are edge disjoint.

There are many research results and theorems about packings of graphs. For surveys, see [9] and [10]. In the following, we list several results and conjectures about packing of trees in complete graphs and complete bipartite graphs, which are the motivation of our current work. The following Theorem 1 was proved by Burns and Schuster [1] in 1978.

Theorem 1 [1]. If T is a tree of order n that is not a star, then there is a 2packing of T in K_n .

For a tree of order $n \ge 2$ to have a 3-packing in K_n , K_n must have at least 3(n-1) edges as a tree of order n has n-1 edges. Thus $n(n-1)/2 \ge 3(n-1)$ and so $n \ge 6$. In 1993, Wang and Sauer [6] characterized all the trees of order n that have 3-packings in K_n .

Theorem 2 [6]. If T is a tree of order $n \ge 6$, then with three exceptions, there is a 3-packing of T in K_n .

For a tree of order $n \ge 2$ to have a 4-packing in K_n , there must be $n \ge 8$ and its maximum degree must be at most n-4. In 2014, Haler and Wang [3] further characterized all the trees of order n that have 4-packings in K_n .

Theorem 3 [3]. If T is a tree of order $n \ge 8$ with maximum degree at most n-4, then with four exceptions, there is a 4-packing of T in K_n .

We refer readers to [6] and [3] for the exceptions in Theorem 2 and Theorem 3. The following conjecture proposed by Gyárfás and Lehel [2] is well-known as *Tree Packing Conjecture* in 1978. So far the results and theorems that are obtained by many mathematicians on this conjecture are still far from a complete solution to the conjecture.

Tree Packing Conjecture. For any sequence T_1, T_2, \ldots, T_n of trees such that T_i is a tree of order *i* for each $1 \le i \le n$, there is a packing of T_1, T_2, \ldots, T_n in K_n .

Since T_1, T_2, \ldots, T_n together have n(n-1)/2 edges, the above conjecture implies that K_n can be decomposed into edge disjoint trees T_1, T_2, \ldots, T_n . Wang [7] proved the following Theorem 4 in 1996.

Theorem 4 [7]. If T is a tree of order n, then there is a 2-packing of T in a complete bipartite graph B_{n+1} .

In 2003, Orchel [5] further characterized all the trees of order n that have 2-packings in a complete bipartite graph B_n . In 2009, Wang [8] proposed the following conjecture.

Conjecture 5 [8]. If T is a tree of order n, then for each positive integer k, there is a k-packing of T in a complete bipartite graph B_{n+k-1} .

264

In the above conjecture, if k = n, then n copies of T together contain exactly n(n-1) edges, and this is the number of edges of $K_{n,n-1}$. Thus, Conjecture 5 implies that the complete bipartite graph $K_{n,n-1}$ can be decomposed into n edge disjoint copies of T for any given tree T of order n. So far, the results about packing trees into complete bipartite graphs are also far from a complete solution to this conjecture.

To support this conjecture, Wang [8] proved the following theorem.

Theorem 6 [8]. If T is a tree of order n, then there is a 3-packing of T in a complete bipartite graph B_{n+2} .

Thus, Conjecture 5 is true for k = 2, 3 by Theorem 4 and Theorem 5. Although this is a hard conjecture, our goal is to make some progress on it in the case k = 4. In this case, the conjecture says that there is a 4-packing of any given tree of order n in a complete bipartite graph B_{n+3} . We have not found any counterexample to the conjecture in the case k = 4. Based on our experience working with k = 4, we find that we can prove that there is a 4-packing of any given tree of order n in a complete bipartite graph B_{n+12} . As for how to reduce B_{n+12} further to a B_{n+l} for some $3 \le l \le 11$, it remains open for future. In this article, we will prove the following theorem.

Main Theorem. If T is a tree of order n, then there is a 4-packing of T in a complete bipartite graph B_{n+12} .

In the process of proving the Main Theorem, we often construct a 4-packing (d, b, g, r) of a subgraph H of T. We then modify this 4-packing (d, b, g, r) and extend it to a 4-packing of T in a B_{n+12} . While doing so, we need keep the order of the bipartition of H to be agreeable with that of T because H may have more than one bipartitions. Therefore, for the sake of convenience, we regard a bipartite graph as a bipartite graph with a given ordered bipartition. Thus, if (X, Y) is the given bipartition of G, then for any subgraph H of G, it is already decided that $(X \cap V(H), Y \cap V(H))$ is the given bipartition of H.

Since we are dealing with embeddings of bipartite graphs in B_n , we need to keep the order of the two partites under embeddings, too. Therefore, we adopt this convention through out this article from now on. For an embedding σ of a bipartite graph G (or a tree T) in B_n , we mean that σ is an injection from V(G)into $V(B_n)$ such that $\sigma(U) \subseteq X$ and $\sigma(W) \subseteq Y$ where (U, W) and (X, Y) are the given bipartitions of G and B_n , respectively. By doing so, we say that two vertices of G have the same parity if they belong to the same partite of G and otherwise we say that they have the opposite parity.

For a 4-packing of G in B_n , we often use (d, b, g, r) to represent the four embeddings and we say that d(G) is a dark copy of G, b(G) is a blue copy of G, g(G) is a green copy of G, and r(G) is a red copy of G. For a 4-packing (d, b, g, r) of G in B_n , we say that a vertex x is 4-placed if d(x), b(x), g(x) and r(x) are distinct.

A *linear graph* is a graph such that each of its components is a path.

Given a bipartite graph G, we say that two or more vertices of G are strongly independent if no two of them are adjacent and no two of them have any common neighbor. A leaf of G is a vertex of G that has degree 1. A node of G is a vertex of G that is adjacent to a leaf of G. A supernode of G is a vertex x of G such that, with possibly one exception, every neighbor of x is a leaf of G. If G is a tree but not a star, we readily see that G has at least two distinct supernodes by observing a longest path of G, that is, the second vertex and the second to the last vertex on the longest path are two supernodes.

For a tree T, the *trunk* of T, denoted by T^* , is the subtree of T obtained from T by removing all the leaves of T. By this definition, we see that a vertex of T^* is a leaf of T^* if and only if it is a supernode of T.

2. Proof of the Main Theorem

We use induction on the order of trees to prove the Main Theorem. The theorem is obviously true for a tree of order 1. Assume that the Main Theorem holds for trees of order less than n with $n \ge 2$. Let T = (V, E) be a tree of order n. The idea of the proof is as follows. If T has four strongly independent leaves u, x, y, and z of the same parity, we apply induction on $T - \{u, x, y, z\}$ to get a 4-packing of $T - \{u, x, y, z\}$ in a $B_{n-4+12} = B_{n+8}$. This 4-packing can be easily extended to a 4-packing of T in B_{n+12} by the following Lemma 1. If T does not have four strongly independent leaves of the same parity, then T has at most six supernodes and so T^* has at most six leaves. This will allow us to classify the structure of T into several cases. We then prove that the Main Theorem holds for each of these cases.

Clearly, $K_{4,n-1}$ has four edge disjoint copies of $K_{1,n-1}$. Thus, if T is a star, there is a 4-packing of T in a B_{n+3} , and the Main Theorem holds for T. Therefore, we may assume in the following that T is not a star.

Let n^* be the order of T^* . Let ξ_i be the number of vertices of T^* with degree i in T^* . Then ξ_1 is the number of supernodes of T with

(1)
$$\xi_1 \ge 2 \text{ and } n^* = \xi_1 + \xi_2 + \dots + \xi_{n^*-1}.$$

Since the degree sum of a graph is equal to two times the number of its edges, we have

(2)
$$\xi_1 + 2\xi_2 + \dots + (n^* - 1)\xi_{n^* - 1} = 2(n^* - 1).$$

Combining (1) and (2), we obtain the number ξ_1 of supernodes of T as follows

(3)
$$\xi_1 = 2 + \xi_3 + 2\xi_4 + 3\xi_5 + \dots + (n^* - 3)\xi_{n^* - 1}.$$

266

Let $\lambda(T)$ be the largest number of k such that T has k nodes of the same parity. We now divide the proof of the Main Theorem into two parts.

Part I.
$$\lambda(T) \geq 4$$

In this case, we need the following Lemma 1 which was proven in [3].

Lemma 1 (Lemma 2.4 of [3]). Let G be a complete bipartite graph $K_{4,m}$ where $m \geq 4$. Let A and B be the two partite sets of G with |A| = 4 and |B| = m. Let B_1 , B_2 , B_3 , and B_4 be any given four subsets of B with $|B_i| = 4$ for all $i \in \{1, 2, 3, 4\}$. Then G has four disjoint matchings M_1 , M_2 , M_3 , and M_4 such that M_i matches B_i into A for all $i \in \{1, 2, 3, 4\}$.

We apply Lemma 1 as follows. Let y_1, y_2, y_3 , and y_4 be four distinct nodes of T with the same parity. Let x_1, x_2, x_3 , and x_4 be four strongly independent leaves of T such that x_i is adjacent with y_i for each $i \in \{1, 2, 3, 4\}$. Let $H = T - \{x_1, x_2, x_3, x_4\}$. Then H has order n - 4. By the induction hypothesis, there is a 4-packing $(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$ of H in a $B_{n-4+12} = B_{n+8}$. We may choose $V(B_{n+8})$ such that x_i is not in B_{n+8} for $i \in \{1, 2, 3, 4\}$. Let A = $\{x_1, x_2, x_3, x_4\}$ and $B_i = \{\sigma_i(y_1), \sigma_i(y_2), \sigma_i(y_3), \sigma_i(y_4)\}$ for each $i \in \{1, 2, 3, 4\}$. Let $B = B_1 \cup B_2 \cup B_3 \cup B_4$ and say |B| = m. Let G be the complete bipartite graph with partites A and B. By Lemma 1, there exist four disjoint matchings M_1, M_2, M_3 and M_4 such that M_i matches B_i into A for all $i \in$ $\{1, 2, 3, 4\}$. For each $i \in \{1, 2, 3, 4\}$, let f_i be the permutation of A such that $M_i = \{f_i(x_1)\sigma_i(y_1), f_i(x_2)\sigma_i(y_2), f_i(x_3)\sigma_i(y_3), f_i(x_4)\sigma_i(y_4)\}$.

We now add A to B_{n+8} to form a B_{n+12} such that A has the opposite parity with $B_i(1 \le i \le 4)$. For each $i \in \{1, 2, 3, 4\}$, we extend σ_i to an embedding of T in B_{n+12} such that $\sigma_i(x_1) = f_i(x_1)$, $\sigma_i(x_2) = f_i(x_2)$, $\sigma_i(x_3) = f_i(x_3)$ and $\sigma_i(x_4) = f_i(x_4)$. This means that we extend a copy $\sigma_i(H)$ of H to a copy $\sigma_i(T)$ of T by adding four independent edges to $\sigma_i(H)$ from M_i which matches B_i into A. As M_1, M_2, M_3 and M_4 are disjoint, we conclude that $\sigma_1(T), \sigma_2(T), \sigma_3(T)$ and $\sigma_4(T)$ are edge disjoint, i.e., $(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$ is a 4-packing of T in B_{n+12} . Thus, the Main Theorem holds for T.

Part II.
$$\lambda(T) \leq 3$$

In this case, we will apply the following lemmas to prove the Main Theorem. We need introduce a notation F' for every subgraph F of T. F' is obtained from F by adding to F all those edges of T which are not in F but join two vertices of F in T. We call F' the induced subgraph by F in T.

Lemma 2. Let H be a subgraph of T such that each vertex of T - V(H) is a leaf of T and is adjacent with some vertex of H. If there is a 4-packing of H' in a B_p

such that each vertex y of H' with $xy \in E$ for some $x \in V(T) \setminus V(H)$ is 4-placed, then there is a 4-packing of T in a B_{p+q} where q = |V(T)| - |V(H)| such that each 4-placed vertex of H remains as 4-placed.

Proof. Let $(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$ be a 4-packing of H' in a B_p such that if $xy \in E(T)$ with $x \in V(T) \setminus V(H)$, then y is 4-placed. Note that $\sigma_1(y)$, $\sigma_2(y)$, $\sigma_3(y)$ and $\sigma_4(y)$ have the same parity in B_p for all $y \in V(H)$. We obtain B_{p+q} by adding each $x \in V(T) \setminus V(H)$ to B_p such that if $xy \in E(T)$, then x and $\sigma_1(y)$ have the opposite parity. Then for each $i \in \{1, 2, 3, 4\}$, we extend σ_i to an embedding of T in B_{p+q} such that $\sigma_i(x) = x$ for each $x \in V(T) \setminus V(H)$. That is, if $xy \in E(T)$ with $x \in V(T) \setminus V(H)$, then $x\sigma_i(y)$ is an edge of $\sigma_i(T)$. Since y is 4-placed for each edge $xy \in E(T)$ with $x \in V(T) \setminus V(H)$, i.e., $x\sigma_1(y), x\sigma_2(y), x\sigma_3(y)$ and $x\sigma_4(y)$ are distinct, we conclude that $(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$ is a 4-packing of T in B_{p+q} .

Lemma 3. Let $P = x_1x_2\cdots x_t$ be a path with $4 \le t \le 6$ or $9 \le t \le 10$. Then there is a 4-packing (d, b, g, r) of P in a B_{t+4} such that every vertex of P is 4-placed.

Proof. Let $V(B_{t+4}) = \{x_1, x_2, \ldots, x_{t+4}\}$ such that $\{x_i | i \text{ is odd}\}$ and $\{x_j | j \text{ is even}\}$ form a bipartition of B_{t+4} . We exhibit (d, b, g, r) as follows. If t = 4, we define

> $d(P) = x_1 x_2 x_7 x_4 \text{ with } d(x_1) = x_1,$ $b(P) = x_3 x_4 x_1 x_6 \text{ with } b(x_1) = x_3,$ $g(P) = x_5 x_6 x_3 x_8 \text{ with } g(x_1) = x_5,$ $r(P) = x_7 x_8 x_5 x_2 \text{ with } r(x_1) = x_7.$

If t = 5, we define

 $d(P) = x_1 x_2 x_7 x_4 x_5 \text{ with } d(x_1) = x_1,$ $b(P) = x_3 x_4 x_9 x_6 x_7 \text{ with } b(x_1) = x_3,$ $g(P) = x_5 x_6 x_3 x_8 x_9 \text{ with } g(x_1) = x_5,$ $r(P) = x_7 x_8 x_5 x_2 x_3 \text{ with } r(x_1) = x_7.$

If t = 6, we define

 $d(P) = x_1 x_2 x_9 x_4 x_7 x_6 \text{ with } d(x_1) = x_1,$ $b(P) = x_3 x_4 x_1 x_6 x_9 x_8 \text{ with } b(x_1) = x_3,$ $g(P) = x_5 x_6 x_3 x_8 x_1 x_{10} \text{ with } g(x_1) = x_5,$ $r(P) = x_7 x_8 x_5 x_{10} x_3 x_2 \text{ with } r(x_1) = x_7.$ If t = 9, we define

$$\begin{aligned} d(P) &= x_1 x_2 x_{13} x_4 x_{11} x_6 x_9 x_{10} x_5 \text{ with } d(x_1) = x_1, \\ b(P) &= x_7 x_4 x_1 x_6 x_{13} x_8 x_{11} x_2 x_9 \text{ with } b(x_1) = x_7, \\ g(P) &= x_5 x_6 x_3 x_8 x_1 x_{10} x_7 x_{12} x_{13} \text{ with } g(x_1) = x_5, \\ r(P) &= x_3 x_{10} x_{11} x_{12} x_9 x_4 x_5 x_8 x_7 \text{ with } r(x_1) = x_3. \end{aligned}$$

If t = 10, we define

$$d(P) = x_1 x_2 x_{13} x_{12} x_{11} x_6 x_9 x_8 x_7 x_{14} \text{ with } d(x_1) = x_1,$$

$$b(P) = x_3 x_4 x_1 x_6 x_{13} x_8 x_{11} x_{10} x_9 x_{12} \text{ with } b(x_1) = x_3,$$

$$g(P) = x_5 x_6 x_3 x_8 x_1 x_{10} x_{13} x_{14} x_{11} x_4 \text{ with } g(x_1) = x_5,$$

$$r(P) = x_7 x_{10} x_5 x_4 x_9 x_{14} x_1 x_{12} x_3 x_2 \text{ with } r(x_1) = x_7.$$

Remark. Lemma 3 does not include the case $t \in \{7, 8\}$. Later in the proof, when Lemma 3 is needed and $t \in \{7, 8\}$, we have deleted an endvertex from P or added some new vertices to P to obtain a linear graph whose order is neither 7 nor 8 and then apply Lemma 3 to this linear graph.

Lemma 4. Let p be an integer with $p \ge 8$ and $P = x_1 x_2 \cdots x_{2p}$ be a path of order 2p. Then there is a 4-packing (d, b, g, r) of P in a B_{2p} such that each vertex of P is 4-placed.

Proof. Let B_{2p} have the bipartition $\{x_1, x_3, \ldots, x_{2p-1}\}$ and $\{x_2, x_4, \ldots, x_{2p}\}$. Draw B_{2p} in a plane as follows. Label the vertices of a regular 2p-gon as x_1, \ldots, x_{2p} in order clockwise in an xy-plane and draw edges of B_{2p} as stright line segements. Furthermore, the center of the regular 2p-gon is at the origin (0, 0). The distance from the center to the vertices is 1 and the vertex x_{2p} is at (0, 1). Thus, all the vertices are on the circle of radius 1 with its center at (0, 0). Note that the vertex x_p is at (0, -1).

Take all the edges of B_{2p} which are parallel to $x_{2p}x_1$ or x_1x_2 . These edges form a Hamiltonian cycle C of B_{2p} . If we remove the edge x_1x_{2p} from C, we get a Hamiltonian path from x_1 to x_{2p} in B_{2p} . Label this path as $L_1 = x_{i_1}x_{i_2}\cdots x_{i_{2p}}$ with $x_1 = x_{i_1}$, which can be viewed as a permutation of $\{x_1, x_2, \ldots, x_{2p}\}$ as well. Similarly, for each $x_i \in \{x_3, x_5, x_7\}$, using $x_{i-1}x_i$ and x_ix_{i+1} in place of $x_{2p}x_1$ and x_1x_2 to repeat the above action, we get a Hamiltonian path L_i from x_i to x_{i-1} . Define $d(P) = L_1$ with $d(x_1) = x_1$, $b(P) = L_3$ with $b(x_1) = x_3$, $g(P) = L_5$ with $g(x_1) = x_5$ and $r(P) = L_7$ with $r(x_1) = x_7$.

In fact, we may rotate L_1 clockwise by $(360/p)^\circ$ three times to get L_3, L_5 , and L_7 as follows

$$L_{3} = x_{i_{1}+2}x_{i_{2}+2}\cdots x_{i_{2p}+2},$$

$$L_{5} = x_{i_{1}+4}x_{i_{2}+4}\cdots x_{i_{2p}+4},$$

$$L_{7} = x_{i_{1}+6}x_{i_{2}+6}\cdots x_{i_{2p}+6}.$$

Note that in the above writing, we define $x_{2p+i} = x_i$ for each $1 \le i \le 6$. As $2p \ge 16$, this implies that each vertex of P is 4-placed. To see that L_1, L_3, L_5 , and L_7 are edge disjoint, we observe that the edges of the regular 2p-gon incident with vertices x_1, x_3, x_5 , and x_7 are the chords of the half unit circle on the right side of y-axis because $p \ge 8$. Therefore, these edges have distinct slopes and so L_1, L_3, L_5 , and L_7 are edge disjoint.

Since a linear graph is always a subgraph of a path, the following lemma is an easy observation.

Lemma 5. Let r, s, p and t be four positive integers such that $r \leq s \leq 2p \leq t$. Suppose that there is 4-packing of a path of order s in a B_{2p} such that each vertex of the path is 4-placed. Then there is a 4-packing of any linear graph of order r in a B_t such that every vertex of the linear graph is 4-placed.

We are ready to prove the Main Theorem. As $\lambda(T) \leq 3$, it follows that $\xi_1 \leq 2\lambda \leq 6$. By (3), we see that $\xi_i = 0$ for $i \geq 7$. Thus,

(4)
$$2 \le \xi_1 \le 6 \text{ and } \xi_1 = 2 + \xi_3 + 2\xi_4 + 3\xi_5 + 4\xi_6.$$

First, suppose that $\xi_1 = 2$. Then $\xi_i = 0$ for all $i \ge 3$. Thus, T^* is a path. By Lemmas 3 through 5, there is a 4-packing of T^* in a B_{n^*+6} such that each vertex of T^* is 4-placed. By Lemma 2, there is a 4-packing of T in B_{n+6} where n is the order of T. Therefore, we may assume that $\xi_1 \ge 3$, i.e., $\xi_i \ge 1$ for some $3 \le i \le 6$.

We define some terminology. For each supernode u of T, there is a unique path $u_1u_2\cdots u_kv$ in T^* with $u = u_1$ such that $deg_{T^*}(v) \ge 3$ and $deg_{T^*}(u_i) = 2$ for all $2 \le i \le k$. In this case, we say that the supernode u belongs to v. Moreover, if k is even, we say that $u_1u_2\cdots u_k$ is an arm of v. If k is odd, we choose a fixed vertex u' of T that is a leaf of T with $u'u_1 \in E$, and then we say that $u'u_1u_2\cdots u_k$ is an arm of v. Clearly, by this definition, any arm is a path of even order, and any two arms do not intersect. We now divide the remaining proof into the following five cases.

Case 1. There is exactly one vertex u of T^* with $deg_{T^*}(u) \geq 3$. Thus, all the ξ_1 supernodes of T belong to u. Let H be the union of the ξ_1 arms of u. Let (A, B) denote the bipartition of H. As each arm has an even order, |A| = |B| = h for some $h \geq 3$. By Lemmas 3 through 5, there is a 4-packing (d, b, g, r) of H in a B_{2h+6} such that each vertex of H is 4-placed. We may assume that H is a subgraph of B_{2h+6} and d is the identity embedding. Add u and three new vertices ω_1, ω_2 and ω_3 to B_{2h+6} to form a B_{2h+10} such that these four vertices have the same parity and H + u is a subgraph of B_{2h+10} . Then extend (d, b, g, r) to be a 4-packing of H + u in B_{2h+10} such that d(u) = u, $b(u) = \omega_1, g(u) = \omega_2$ and $r(u) = \omega_3$. Thus, (d, b, g, r) is a 4-packing of H + u is B_{2h+10} . By Lemma 2, there is a 4-packing of T in a $B_{2h+10+q} = B_{n+9}$ where q = |V(T)| - |V(H+u)| = n - 2h - 1. This completes Case 1.

By Case 1, we may assume that T^* has at least two vertices of degree at least 3 in T^* . As $\xi_1 \leq 6$, we see that $\xi_6 = 0$ by (4). Let L be a longest path of T^* from a vertex of degree at least 3 to another vertex of degree at least 3 in T^* . Say $L = uy_1y_2\cdots y_tv$. Say $deg_{T^*}(u) = i$ and $deg_{T^*}(v) = j$. Then there are i - 1 supernodes of T belonging to u and j - 1 supernodes of T belonging to v.

Let $L_1 = x_1 x_2 \cdots x_a$ be a path of T passing through u such that x_2 and x_{a-1} are two supernodes of T belonging to u and x_1 and x_a are two leaves of T. Similarly, let $L_2 = z_1 z_2 \cdots z_b$ be a path of T passing through v such that z_2 and z_{b-1} are two supernodes belonging to v. Since each arm has an even order and L_1 contains two arms and the vertex u, L_1 has order at least 5. Similarly, L_2 has order at least 5.

Let A and B denote the two partites of $L_1 \cup L_2$ with $|A| \ge |B|$. As $\lambda(T) \le 3$, we see that if L_1 has an odd order, then x_1 and x_a have the same parity and so one of z_1 and z_b must have the opposite parity with x_1 . This implies that $|A| - |B| \le 1$. If |A| = |B|, Let H_1 be $L_1 \cup L_2$, and if |A| - |B| = 1, let $H_1 = L_1 \cup L_2 - w$ where w is a leaf of $L_1 \cup L_2$ which belongs to A. Thus, the two partites of H_1 has the same size. Say H_1 has order 2h. Then $h \ge 5$. Let H_2 be the union of all the arms of u or v that are not on any of L_1 and L_2 . Let (C, D)denote the bipartition of H_2 . Then the two partites of H_2 have the same size. Say H_2 has order 2k.

Case 2. There are exactly two distinct vertices of T^* with degree at least 3 in T^* . If t is even, let $H = H_1 \cup H_2 \cup (y_1y_2 \cdots y_t)$. If t is odd and T has a leaf y' adjacent to y_t , let $H = H_1 \cup H_2 \cup (y_1y_2 \cdots y_ty')$, and if t is odd and T does not has a leaf y' adjacent to y_t , let $H = H_1 \cup H_2 \cup (y_1y_2 \cdots y_{t-1})$. Then, H is a linear subgraph of T and its two partites have the same size. Say |V(H)| = m. As $m \ge 10$ and by Lemmas 3 through 5, there is a 4-packing (d, b, g, r) of H in a B_{m+4} such that all vertices of H are 4-placed. We may assume that H is a subgraph of B_{m+4} and d is the identity embedding. This 4-packing of H does not consider the edges of T^* that join a component of H to another component of H. If these edges are added back to H to form H', this 4-packing of H cannot be guaranteed to be a 4-packing of H'. These edges are incident with at least one u, v, and y_t. Therefore, we need to modify (d, b, g, r) in the following. First, if y_t is not in H, add y_t to H and define $d(y_t) = b(y_t) = g(y_t) = r(y_t) = y_t$.

Let m' be the order of $H + y_t$. Note that if y_t is already in H, then m' = m. Then add six new vertices $\omega_1, \ldots, \omega_6$ to $B_{m'+4}$ to form a $B_{m'+10}$ such that ω_1 , ω_2 and ω_3 have the same parity with u and ω_4 , ω_5 and ω_6 has the same parity with v. Change the images of u and v under b, g and r as follows. Let

$$b(u) = \omega_1, g(u) = \omega_2$$
, and $r(u) = \omega_3$,

$$b(v) = \omega_4, g(v) = \omega_5$$
, and $r(v) = \omega_6$.

The images of all the other vertices of $(H + y_t)'$ are not changed under (d, b, g, r). Clearly, if y_t is not in H, then y_t has only two neighbors y_{t-1} and v in T, and

$$\{d(y_{t-1}), b(y_{t-1}), g(y_{t-1}), r(y_{t-1})\} \cap \{d(v), b(v), g(v), r(v)\} = \emptyset.$$

Thus, (d, b, g, r) becomes a 4-packing of $(H + y_t)'$ in $B_{m'+10}$ such that all the vertices of H are 4-placed. By Lemma 2, T has a 4-packing in a B_{n+10} . This completes Case 2.

By Case 2, we may assume that T^* has at least three vertices of degree at least 3 in T^* . As T^* is connected, any vertex of degree 3 in T^* is connected by a path to a vertex on $y_1 \cdots y_t$ in T^* . Thus, y_r has degree at least 3 in T^* for some $1 \leq r \leq t$. As $\xi_1 \leq 6$ and by (4), we see that $\xi_5 = 0$ and if $deg_{T^*}(y_r) = 4$, then $deg_{T^*}(x) < 3$ for all $x \in V(T^*) \setminus \{u, v, y_r\}$. In the following cases, the arguments are very much similar to the argument in Case 2.

Case 3. Either $deg_{T^*}(y_r) = 4$ or $deg_{T^*}(y_s) = 3$ for some $1 \leq s \leq t$ and $s \neq r$. We define a new number a such that a = r if $deg_{T^*}(y_r) = 4$ and a = s if $deg_{T^*}(y_r) = 3$ and $deg_{T^*}(y_s) = 3$ for some $1 \leq s \leq t$ and $s \neq r$. Thus, T has two paths $Q_1 = u_1u_2\cdots u_ly_r$ and $Q_2 = v_1v_2\cdots v_cy_a$ such that $u_1u_2\cdots u_ly_r$ and $v_1v_2\cdots v_c$ are vertex disjoint and u_2 and v_2 are two supernodes of T belonging y_r and y_a , respectively. We may assume that $r \leq a$. Let

$$L_3 = u_1 \cdots u_l y_r y_{r+1} \cdots y_a v_c v_{c-1} \cdots v_1.$$

As $\lambda(T) \leq 3$, it is easy to see that if two of L_1 , L_2 and L_3 have odd orders, then the leaves of one of them have the opposite parity with the leaves of the other and so the third path must have an even order. This implies that the two partites of $L_1 \cup L_2 \cup L_3$ have the same size or their sizes differ by 1. If the two partites of $L_1 \cup L_2 \cup L_3$ have the same size, we define $J_1 = L_1 \cup L_2 \cup L_3$. If the two partites of $L_1 \cup L_2 \cup L_3$ differ by 1 in size, we delete one leaf of $L_1 \cup L_2 \cup L_3$ from the larger partite and denote the resulting linear subgraph by J_1 . Thus, J_1 has an even order 2q for some $q \geq 8$.

For each of $y_{r-1}y_{r-2}\cdots y_1$ and $y_{a+1}y_{a+2}\cdots y_t$, we delete its last vertex if it has an odd order and otherwise, we keep it. Thus, we obtain two paths of even orders and let J_2 be the union of these two paths. Let $H = J_1 \cup J_2$. By Lemma 4 and Lemma 5, there is a 4-packing (d, b, g, r) of H in a B_m such that all vertices of H are 4-placed, where m is the order of H. We may assume that H is a subgraph of B_m and d is the identity embedding.

Add y_1 , y_t and 12 new vertices $\omega_1, \ldots, \omega_{12}$ to B_m to obtain a B_{p+12} such that ω_1, ω_2 and ω_3 have the same parity with $u; \omega_4, \omega_5$ and ω_6 has the same parity with $y_{r-1}; \omega_7, \omega_8$ and ω_9 have the same parity with y_{a+1} ; and ω_{10}, ω_{11} and ω_{12}

272

have the same parity with v, where p is the order of $H + y_1 + y_t$. Note that for $i \in \{1, t\}$, if y_i is already in H, adding y_i to H does not increase the order of the resulting graph.

For each $y_i \in \{y_1, y_t\}$, if $y_i \notin V(H)$, we define $d(y_i) = b(y_i) = g(y_i) = r(y_i)$. Change the images of u, y_{r-1}, y_{a+1} and v under b, g, and r as follows. Let

$$b(u) = \omega_1, g(u) = \omega_2, r(u) = \omega_3,$$

$$b(y_{r-1}) = \omega_4, g(y_{r-1}) = \omega_5, r(y_{r-1}) = \omega_6,$$

$$b(y_{a+1}) = \omega_7, g(y_{a+1}) = \omega_8, r(y_{a+1}) = \omega_9,$$

$$b(v) = \omega_{10}, g(v) = \omega_{11}, r(v) = \omega_{12}.$$

The images of all the other vertices of $H+y_1+y_t$ are not changed. Thus, (d, b, g, r) is a 4-packing of $(H + y_1 + y_t)'$ in a B_{p+12} such that all the vertices of H are 4-placed. As $\lambda(T) \leq 3$, every leaf of T is adjacent to one of the six supernodes of T. By Lemma 2, there is a 4-packing of T in a B_{n+12} . This completes Case 3.

Case 4. There are exactly three distinct vertices of T^* with degree at least 3 in T^* . By Case 3, we may assume that $deg_{T^*}(y_r) = 3$. Let $Q_1 = u_1 \cdots u_l y_r$ be a path of T such that u_2 is supernode of T belonging to y_r . Consider three paths $y_{r-1}y_{r-2}\cdots y_1, y_{r+1}y_{r+2}\cdots y_t$ and $u_1\cdots u_l y_r$. Let $I_1 = u_1u_2\cdots u_l y_r y_{r-1}\cdots y_1$. Then I_1 has order at least 3. Let $I_2 = y_{r+1}y_{r+2}\cdots y_t$ if this path has an even order. If $y_{r+1}y_{r+2}\cdots y_t$ has an odd order and there is a leaf of T adjacent to y_t , let $I_2 = y_{r+1}\cdots y_t y'$. Otherwise, let $I_2 = y_{r+1}y_{r+2}\cdots y_{t-1}$. Let $H = H_1 \cup H_2 \cup I_1 \cup I_2$. Then H has an order at least 13. By Lemma 4 and Lemma 5, there is a 4-packing (d, b, g, r) of H in a B_{m+3} such that all vertices of H are 4-placed, where m is the order of H. We may assume that H is a subgraph of B_{m+3} and d is the identity embedding. Add y_t and nine new vertices ω_1, \ldots, w_9 to B_m and obtain a B_{p+12} , where p is the order of $H + y_t$, such that $H + y_t$ is a subgraph of $B_{p+12}, \omega_1, \omega_2$ and ω_3 have the same parity with $u; \omega_4, \omega_5$ and ω_6 have the same parity with v; ω_7, ω_8 and ω_9 have the same parity with y_r .

If y_t is not in H, we define $d(y_t) = b(y_t) = g(y_t) = r(y_t) = y_t$. Change the images of u, v and y_r under b, g, and r as follows.

$$b(u) = \omega_1, g(u) = \omega_2, r(u) = \omega_3,$$

$$b(v) = \omega_4, g(v) = \omega_5, r(v) = \omega_6,$$

$$b(y_r) = \omega_7, g(y_r) = \omega_8, r(y_r) = \omega_9.$$

The images of all the other vertices of $H + y_t$ are not changed. Thus, (d, b, g, r) is a 4-packing of $(H + y_t)'$ in a B_{p+12} such that all the vertices of H are 4-placed. By Lemma 2, there is a 4-packing of T in a B_{n+12} . This completes Case 4.

Case 5. There are at least four distinct vertices of T^* with degree at least 3 in T^* . By (4), we see that $\xi_3 = 4$ and $\xi_4 = \xi_5 = 0$. Let w be the fourth vertex

of degree 3. By Case 3, we may assume that w is not on $y_1y_2\cdots y_t$. As T^* is connected, there is a path $Q = y_r u_1 \cdots u_l w$ from y_r to w such that y_i is not on this path for all $i \in \{1, 2, \ldots, t\} \setminus \{r\}$.

Let $L_3 = v_1 v_2 \cdots v_c$ be a path of T passing through w such that v_2 and v_{c-1} are two supsernodes of T. Clearly, $c \geq 5$. Let $J_1 = L_1 \cup L_2 \cup L_3$ if the two partites have the same size. If their sizes differ by 1, we delete a leaf of $L_1 \cup L_2 \cup L_3$ to obtain a subgraph with its two partites having the same size and denote this resulting graph by J_1 . As in Case 3, we see that J_1 has order 2q for some $q \geq 8$.

Consider three paths $y_{r-1}y_{r-2}\cdots y_1$, $y_{r+1}y_{r+2}\cdots y_t$ and $y_ru_1u_2\cdots u_l$. For each of these three paths, we delete its last vertex if it has an odd order and otherwise, we keep it. Thus, we obtain three paths of even orders and let J_2 be the union of these three paths of even order. Let $H = J_1 \cup J_2$. By Lemma 4 and Lemma 5, there is a 4-packing (d, b, g, r) of H in a B_m such that all vertices of Hare 4-placed, where m is the order of H. We may assume that H is a subgraph of B_m and d is the identity embedding.

Let the order of $H + y_1 + y_t + u_l$ be p. Add y_1, y_t, u_l and 12 new vertices $\omega_1, \ldots, \omega_{12}$ to B_m to obtain a B_{p+12} such that ω_1, ω_2 and ω_3 have the same parity with u; ω_4, ω_5 and ω_6 have the same parity with v; ω_7, ω_8 and ω_9 have the same parity with w; and ω_{10}, ω_{11} and ω_{12} have the same parity with y_r . Furthermore, $H + y_1 + y_t + u_l$ is a subgraph of B_{p+12} .

For each $x \in \{y_1, y_t, u_l\}$, if x is not in H, we define d(x) = b(x) = g(x) = r(x). Change or define the images of u, v, w, and y_r under d, b, g, and r as follows. Let $d(y_r) = y_r$ and

$$b(u) = \omega_1, g(u) = \omega_2, r(u) = \omega_3,$$

$$b(v) = \omega_4, g(v) = \omega_5, r(v) = \omega_6,$$

$$b(w) = \omega_7, g(w) = \omega_8, r(w) = \omega_9,$$

$$b(y_r) = \omega_{10}, g(y_r) = \omega_{11}, r(y_r) = \omega_{12}$$

The images of all the other vertices of H are not changed. Thus, (d, b, g, r) is a 4-packing of $(H + y_1 + y_t + u_l)'$ in B_{p+12} such that all the vertices of $H + y_r$ are 4-placed. As $\lambda(T) \leq 3$ and $\xi_1 = 6$, each leaf of T is adjacent to one of the six supernodes of T. By Lemma 2, there is a 4-packing of T in a B_{n+12} .

This proves the Main Theorem.

Acknowledgment

This work is advised and mentored by Teacher P. Blount and Dr H. Wang. The author is grateful for their advice and encouragement.

References

- [1] D. Burns and S. Schuster, *Embedding* (p, p 1)-graphs in their complements, Israel J. Math. **30** (1978) 313–320. https://doi.org/10.1007/BF02761996
- [2] A. Gyárfás and J. Lehel, Packing trees of different order into K_n , Colloq. Math. Soc. János Bolyai **18** (1978) 463–469.
- [3] S.P. Haler and H. Wang, Packing four copies of a tree into a complete graph, Australas. J. Combin. 59 (2014) 323–332.
- J.M. Harris, J. Hirst and M.J. Mossinghoff, Combinatorics and Graph Theory (Springer, New York, 2010). https://doi.org/10.1007/978-0-387-79711-3
- [5] B. Orchel, 2-placement of (p,q)-trees, Discuss. Math. Graph Theory 23 (2003) 23–26. https://doi.org/10.7151/dmgt.1183
- [6] H. Wang and N. Sauer, Packing three copies of a tree into a complete graph, European J. Combin. 14 (1993) 137–142. https://doi.org/10.1006/eujc.1993.1018
- H. Wang, Packing two forests into a bipartite graph, J. Graph Theory 23 (1996) 209–213. https://doi.org/10.1002/(SICI)1097-0118(199610)23:2;209::AID-JGT12;3.0.CO;2-B
- [8] H. Wang, Packing three copies of a tree into a complete bipartite graph, Ann. Comb. 13 (2009) 261–269. https://doi.org/10.1007/s00026-009-0022-0
- [9] M. Woźniak, Packing of graphs and permutations—a survey, Discrete Math. 276 (2004) 379–391. https://doi.org/10.1016/S0012-365X(03)00296-6
- [10] H.P. Yap, Packing of graphs—a survey, Discrete Math. 72 (1988) 395–404. https://doi.org/10.1016/0012-365X(88)90232-4

Received 19 March 2019 Revised 24 September 2019 Accepted 26 September 2019