

TOTAL COLORING OF CLAW-FREE PLANAR GRAPHS

ZUOSONG LIANG

*School of Management
Qufu Normal University
Rizhao 276826, China*

e-mail: liangzuosong@126.com

Abstract

A total coloring of a graph is an assignment of colors to both its vertices and edges so that adjacent or incident elements acquire distinct colors. Let $\Delta(G)$ be the maximum degree of G . Vizing conjectured that every graph has a total $(\Delta + 2)$ -coloring. This Total Coloring Conjecture remains open even for planar graphs, for which the only open case is $\Delta = 6$. Claw-free planar graphs have $\Delta \leq 6$. In this paper, we prove that the Total Coloring Conjecture holds for claw-free planar graphs.

Keywords: total coloring, total coloring conjecture, planar graph, claw.

2010 Mathematics Subject Classification: 05C15.

1. INTRODUCTION

All graphs considered here are finite, simple and nonempty. Let $G = (V, E)$ be a graph with *vertex set* V and *edge set* E . The number of vertices of G is called the *order* of G . For a vertex $v \in V$, the *open neighborhood* $N(v)$ of v is defined as the set of vertices adjacent to v , i.e., $N(v) = \{u \mid uv \in E\}$. The *closed neighborhood* of v is $N[v] = N(v) \cup \{v\}$. Every vertex in $N(v)$ is also called a *neighbor* of v . The *degree* of v is equal to $|N(v)|$, denoted by $d_G(v)$ or simply $d(v)$. By $\delta(G)$ and $\Delta(G)$, we denote the *minimum degree* and the *maximum degree* of the graph G , respectively. For a subset $S \subseteq V$, the *closed neighborhood* of S is $N[S] = \bigcup_{v \in S} N[v]$ and the *closed 2-neighborhood* of S is $N_2[S] = N[N[S]]$. For a subset $X \subseteq V$, the subgraph induced by X is denoted by $G[X]$. The set of edges between X and Y in E is denoted by $E(X, Y)$ for $X, Y \subseteq V$. As usual, $K_{m,n}$ denotes a complete bipartite graph with classes of cardinality m and n , and K_n and C_n denote the complete graph and cycle of order n . The graph $K_{1,3}$ is

also called a *claw*, and K_3 a *triangle*. Given a graph F , a graph G is F -free if it does not contain F as an induced subgraph. In particular, a $K_{1,3}$ -free graph is *claw-free*. For a family $\{F_1, \dots, F_k\}$ of graphs, we say that G is (F_1, \dots, F_k) -free if it is F_i -free for all i . By starting with a disjoint union of two graphs G and H and adding edges joining every vertex of G to every vertex of H , one obtains the *join* of G and H , denoted by $G \vee H$. The n -wheel is the graph $C_n \vee K_1$ and the *double n -wheel* is the graph $C_n \vee \overline{K_2}$ ($n \geq 4$).

Given a graph G , an *element* of G is a member of $V(G) \cup E(G)$. Let two elements of a graph G be *adjacent* if they are either adjacent or incident in the traditional sense. Given a graph G , a *total k -coloring* of G is a function that takes each element to $\{1, 2, \dots, k\}$ such that adjacent distinct elements receive distinct colors. In 1968, Vizing [17] (see also [2]) made the following conjecture, known as the Total Coloring Conjecture.

Conjecture. *Every graph has a total $(\Delta + 2)$ -coloring.*

This conjecture is trivial for $\Delta \leq 2$. Rosenfeld [13] and Vijayaditya [16] proved it for $\Delta = 3$. Kostochka proved the $\Delta = 4$ [9] and $\Delta = 5$ [11] cases. The conjecture remains open even for planar graphs, but more is known. Borodin [4] proved it for planar graphs with $\Delta \geq 9$. The $\Delta = 8$ case was proved for planar graphs by Yap [21] and Andersen [1]. The $\Delta = 7$ case was proved for planar graphs by Sanders and Zhao [14]. Thus the only open case for planar graphs is the $\Delta = 6$ case. An extensive study on total coloring is done in [3, 5–8, 10, 18–20] and elsewhere. Claw-free planar graphs have maximum degree at most 6 [12]. In this paper, we prove that the Total Coloring Conjecture holds for claw-free planar graphs.

2. TOTAL COLORING ON THE CLAW-FREE PLANAR GRAPHS

First we introduce some notation and lemmas which are useful for the total coloring of claw-free planar graphs. Let a k -vertex be a vertex of degree k . Let an *at most k -vertex*, or simply a k^- -vertex, be a vertex of degree at most k . Given a graph and integers j_1, j_2, \dots, j_i , let a $(j_1^-, j_2^-, \dots, j_i^-)$ -vertex be an i -vertex v of G such that, for each $1 \leq m \leq i$, there is a j_m^- -vertex y_m of G and the vertices y_1, y_2, \dots, y_i are distinct neighbors of v .

Lemma 1 [15]. *If G is a (claw, K_4)-free planar graph, then $\Delta(G) \leq 5$ and for every vertex v of degree 5 in G , $G[N[v]]$ is a 5-wheel.*

Lemma 2. *Let v be a $(6^-, 6^-, 5^-)$ -vertex, or a $(6, 6, 6)$ -vertex such that $G[N[v]]$ is not a claw, in a graph G . If $G - v$ has a total 8-coloring, then G is total 8-colorable.*

Proof. Let v be a $(6^-, 6^-, 6^-)$ -vertex of a graph G . Given a total 8-coloring ϕ of $G - v$, we will attempt to extend ϕ to a total 8-coloring of G . Let $N(v) = \{y_1, y_2, y_3\}$, and for $i \in \{1, 2, 3\}$ let L_{vy_i} be the set of colors that are not used on y_i or its incident edges and so are available for use on the edge vy_i . Then $|L_{vy_i}| \geq 2$ for each i , since $d_{G-v}(y_i) \leq 5$. Clearly, these edges can be properly colored unless L_{vy_i} is the same set of two colors, say $L_{vy_i} = \{1, 2\}$, for each i . This is impossible if v is a $(6^-, 6^-, 5^-)$ -vertex, so assume that v is a $(6, 6, 6)$ -vertex such that $G[N[v]]$ is not a claw. Without loss of generality, y_2y_3 is an edge of G and $\phi(y_2y_3) = 3$. Recolor y_2y_3 with color 1, and then color vy_1 , vy_2 and vy_3 with colors 1, 2 and 3, respectively. Once the edges incident with v are colored, there are at most six colors that cannot be used on v , and so v can be colored. This gives a total 8-coloring of G , as required. ■

Lemma 3. *Let v be a $(7^-, 6^-, 5^-, 4^-)$ -vertex in a graph G such that $G[N[v]]$ contains neither a claw nor a K_4 . If $G - v$ has a total 8-coloring, then G is total 8-colorable.*

Proof. Let $N(v) = \{y_1, y_2, y_3, y_4\}$ where $d(y_i) \leq 8 - i$ for each i . Given a total coloring ϕ of $G - v$ using a set C of eight colors, let F be the set of colors that are used on the vertices in $N(v)$, and for $i \in \{1, 2, 3, 4\}$ let L_{vy_i} be the set of colors that are available for use on the edge vy_i , as in the previous proof. Then $|L_{vy_i}| \geq i$ for each i , and so the edges vy_1, vy_2, vy_3, vy_4 can be properly colored in this order. If possible, do this so that at least one of these edges has a color in F . Call the new coloring ϕ' .

It is now possible to color v unless every color is used on an element adjacent to v . For this to happen, it must be that $|F| = 4$ and all the lists L_{vy_i} are subsets of $C \setminus F$. In particular, $L_{vy_4} = C \setminus F$. Since $G[N[v]]$ contains neither a claw nor a K_4 , G must contain an edge $y_r y_4$ for some $r \in \{1, 2, 3\}$. Clearly $\phi(y_r y_4) \notin L_{vy_4}$, and so $\phi(y_r y_4) \in F$, while $\phi'(vy_r) \in L_{vy_r} \subseteq L_{vy_4}$. Interchange the colors of vy_r and $y_r y_4$. Since vy_r now has a color in F , there are at most seven different colors that are unavailable for v , and so v can now be colored. This gives a total 8-coloring of G , as required. ■

Theorem 4. *Every claw-free planar graph is total 8-colorable.*

Proof. Let G be a claw-free planar graph. We prove the theorem by induction on the size $m = |E(G)|$. Suppose that the theorem holds when G has fewer than m edges. In the following, we will prove the theorem when G has m edges.

If G has a cut vertex v , we can easily see that the theorem holds. So we may assume that G is 2-connected. In addition, if G has no K_4 , then $\Delta(G) \leq 5$ by Lemma 1, and so the theorem holds since the Total Coloring Conjecture holds when $\Delta(G) \leq 5$ [11]. So let $K = [x_1 x_2 x_3 x_4]$ be a K_4 of G , where x_1 is inside the cycle $C = [x_2 x_3 x_4]$ in the embedding of G in the plane. Let G' be the plane

graph induced by the vertices inside and on C , and choose K so that G' has as fewer vertices as possible. Then K is the only K_4 in G' .

Since every claw-free planar graph has maximum degree at most 6, the result follows from Lemma 2 if G contains a 3^- -vertex. Thus we may assume that $\delta(G) \geq 4$. Without loss of generality, we may assume that x_1 has a neighbor u inside the triangle $T = [x_1x_2x_3]$ in the embedding of G in the plane. Let $V_{\text{in}}(T)$ denote the set of vertices inside the triangle T , and let $G_T = G[V_{\text{in}}(T) \cup V(T)]$, the plane graph induced by the vertices inside and on T . We will make frequent use of the following facts.

(F1) G_T is K_4 -free. This is because K is the only K_4 in G' .

(F2) Every vertex of $V_{\text{in}}(T)$ is adjacent to at most two vertices of T , by (F1).

(F3) Every vertex of $V_{\text{in}}(T)$ has degree 4 or 5, since $\delta(G) \geq 4$ and $\Delta(G_T) \leq 5$ by Lemma 1.

(F4) We may assume that no two 4-vertices in $V_{\text{in}}(T)$ are adjacent. Indeed, if they are, then each of them is a $(6^-, 6^-, 5^-, 4^-)$ -vertex by (F2) and (F3), and so the result follows by Lemma 3.

(F5) For $i = 1, 2, 3$, the neighbors of x_i in $V_{\text{in}}(T)$ induce a complete graph. Otherwise, there would be a claw centered on x_i (including the edge x_ix_4).

(F6) Every vertex of T is adjacent to 0, 1 or 2 adjacent vertices in $V_{\text{in}}(T)$. This follows from (F1) and (F5).

(F7) Every 5-vertex in $V_{\text{in}}(T)$ is adjacent to 0 or 2 vertices of T . Indeed, let v be a 5-vertex in $V_{\text{in}}(T)$ that is adjacent to x_1 (say). By Lemma 1, $G[N[v]]$ is a 5-wheel. Thus $G[N[v]]$ contains two vertices v_1, v_2 that are adjacent to both x_1 and v but not to each other. At least one of v_1, v_2 must be in T , by (F5). But the neighbors x_1, v_1, v_2 of v cannot all be in T , by (F2). Thus v is adjacent to exactly two vertices of T .

Recall that x_1 has a neighbor $u \in V_{\text{in}}(T)$. We consider two cases.

Case 1. $d(u) = 5$. By (F7), u is adjacent to exactly one of x_2 and x_3 , say x_2 . By Lemma 2, $G[N[u]]$ is a 5-wheel. Let $x_1x_2u_1u_2u_3x_1$ be the 5-cycle of $G[N[u]]$ (see Figure 1). Then u and u_3 are the two vertices of $V_{\text{in}}(T)$ that are adjacent to x_1 , and u and u_1 are the two vertices of $V_{\text{in}}(T)$ that are adjacent to x_2 .

Case 1.1. $d(u_1) = 5$. Note that u_1 is adjacent to x_2 but cannot be adjacent to x_1 , and so u_1 is adjacent to x_3 by (F7). So let $x_2x_3wu_2ux_2$ be the 5-cycle of $G[N[u_1]]$ (see Figure 1, left). Each vertex of T is now adjacent to two vertices of $V_{\text{in}}(T)$, and so cannot be adjacent to any further vertex of $V_{\text{in}}(T)$, by (F6). Thus any further neighbor of u_3 cannot be adjacent to x_1 and so must be adjacent to u_2 , to avoid a claw centered on u_3 . If $d(u_2) = 4$, then the only possible further neighbor of u_3 is w , and so u_2, u_3 are adjacent 4-vertices, contrary to (F4). If however $d(u_2) = 5$, then the additional neighbor q of u_2 can be adjacent only to u_2, u_3, w of the vertices so far named (since it cannot be adjacent to x_1 or x_3), and so q is a 3-vertex by the claw-freeness, which is not permissible, by (F3).




Figure 1. Cases 1.1 and 1.2 of Theorem 5.

Case 1.2. $d(u_1) = 4$. Then $d(u_2) = 5$, by (F4). Let $uu_1w_1w_2u_3u$ be the 5-cycle of $G[N[u_2]]$ (see Figure 1, right). Note that x_1 and x_2 each already have two neighbors in $V_{\text{in}}(T)$, and so u_2 has no neighbors in T , by (F6) and (F7). Thus $w_1 \in V_{\text{in}}(T)$ and so $d(w_1) = 5$, by (F4), since $d(u_1) = 4$. But it is impossible for $G[N[w_1]]$ to be a 5-wheel since $d(u_1) = 4$ and there is no edge x_2w_1 .

Case 2. $d(u) = 4$. Let the neighbors of u be x_1, u_1, u_2 and u_3 . We may assume that $u_2, u_3 \in V_{\text{in}}(T)$ and possibly $u_1 = x_2$ or $u_1 = x_3$, by (F2). Then $d(u_2) = 5$ and $d(u_3) = 5$, by (F4). If x_1 is adjacent to the 5-vertex u_2 or u_3 , by choosing u_2 or u_3 instead of u , we are back in Case 1. In addition, if x_2 or x_3 is adjacent to the 5-vertex u_2 or u_3 , by choosing x_2 or x_3 instead of x_1 , we are also back in Case 1. Thus we assume that $N[\{u_2, u_3\}] \subseteq V_{\text{in}}(T)$. But since $G[N[u_2]]$ and $G[N[u_3]]$ are 5-wheels, u must be adjacent to at least four vertices of $N[\{u_2, u_3\}]$ as well as to x_1 , and this is impossible since $d(u) = 4$.

This completes the proof of Theorem 4. ■

Note that the only open case for the Total Coloring Conjecture in planar graphs is $\Delta = 6$. By Theorem 4, immediately, we have the following theorem.

Theorem 5. *The Total Coloring Conjecture holds in claw-free planar graphs.*

Acknowledgements

The author would like to thank the referees for valuable comments and suggestions. Especially, the author has followed closely the version suggested by one referee. This research was supported by the National Nature Science Foundation of China (No. 11601262).

REFERENCES

- [1] L. Andersen, Total Colouring of Simple Graphs, Master's Thesis (University of Aalborg, 1993).
- [2] M. Behzad, Graphs and Their Chromatic Numbers, PhD Thesis (Michigan State University, 1965).

- [3] V.A. Bojarshinov, *Edge and total coloring of interval graphs*, Discrete Appl. Math. **114** (2001) 23–28.
[https://doi.org/10.1016/S0166-218X\(00\)00358-9](https://doi.org/10.1016/S0166-218X(00)00358-9)
- [4] O.V. Borodin, *On the total coloring of planar graphs*, J. Reine Angew. Math. **394** (1989) 180–185.
<https://doi.org/10.1515/crll.1989.394.180>
- [5] O.V. Borodin, A.V. Kostochka and D.R. Woodall, *List edge and list total colourings of multigraphs*, J. Combin. Theory Ser. B **71** (1997) 184–204.
<https://doi.org/10.1006/jctb.1997.1780>
- [6] B.-L. Chen, H.-L. Fu and M.T. Ko, *Total chromatic number and chromatic index of split graphs*, J. Combin. Math. Combin. Comput. **17** (1995) 137–146.
- [7] C.M.H. de Figueiredo, J. Meidanis and C.P. de Mello, *Total-chromatic number and chromatic index of dually chordal graphs*, Inform. Process. Lett. **70** (1999) 147–152.
[https://doi.org/10.1016/S0020-0190\(99\)00050-2](https://doi.org/10.1016/S0020-0190(99)00050-2)
- [8] Ł. Kowalik, J.S. Sereni and R. Škrekovski, *Total-coloring of plane graphs with maximum degree nine*, SIAM J. Discrete Math. **22** (2008) 1462–1479.
<https://doi.org/10.1137/070688389>
- [9] A.V. Kostochka, *The total coloring of a multigraph with maximal degree 4*, Discrete Math. **17** (1977) 161–163.
[https://doi.org/10.1016/0012-365X\(77\)90146-7](https://doi.org/10.1016/0012-365X(77)90146-7)
- [10] A.V. Kostochka, *Exact upper bound for the total chromatic number of a graph*, Proc. 24th Int. Wiss. Koll. Tech. Hochsch. Ilmenau (1979) 33–36, in Russian.
- [11] A.V. Kostochka, *The total chromatic number of any multigraph with maximum degree five is at most seven*, Discrete Math. **162** (1996) 199–214.
[https://doi.org/10.1016/0012-365X\(95\)00286-6](https://doi.org/10.1016/0012-365X(95)00286-6)
- [12] M.D. Plummer, *Extending matchings in claw-free graphs*, Discrete Math. **125** (1994) 301–307.
[https://doi.org/10.1016/0012-365X\(94\)90171-6](https://doi.org/10.1016/0012-365X(94)90171-6)
- [13] M. Rosenfeld, *On the total coloring of certain graphs*, Israel J. Math. **9** (1971) 396–402.
<https://doi.org/10.1007/BF02771690>
- [14] D.P. Sanders and Y. Zhao, *On total 9-coloring planar graphs of maximum degree seven*, J. Graph Theory **31** (1999) 67–73.
[https://doi.org/10.1002/\(SICI\)1097-0118\(199905\)31:1;1-7::AID-JGT6;3.0.CO;2-C](https://doi.org/10.1002/(SICI)1097-0118(199905)31:1;1-7::AID-JGT6;3.0.CO;2-C)
- [15] E.F. Shan, Z.S. Liang and L.Y. Kang, *Clique-transversal sets and clique-coloring in planar graphs*, European J. Combin. **36** (2014) 367–376.
<https://doi.org/10.1016/j.ejc.2013.08.003>
- [16] N. Vijayaditya, *On total chromatic number of a graph*, J. London Math. Soc. (2) **3** (1971) 405–408.
<https://doi.org/10.1112/jlms/s2-3.3.405>

- [17] V.G. Vizing, *Some unsolved problems in graph theory*, Uspekhi Mat. Nauk **23** (1968) 117–134, English translation in Russian Math. Surveys **23** (1968) 125–141, in Russian.
<https://doi.org/10.1070/RM1968v023n06ABEH001252>
- [18] W.F. Wang, *Total chromatic number of planar graphs with maximum degree ten*, J. Graph Theory **54** (2007) 91–102.
<https://doi.org/10.1002/jgt.20195>
- [19] B. Wang and J.-L. Wu, *Total colorings of planar graphs with maximum degree seven and without intersecting 3-cycles*, Discrete Math. **311** (2011) 2025–2030.
<https://doi.org/10.1016/j.disc.2011.05.038>
- [20] P. Wang and J.-L. Wu, *A note on total colorings of planar graphs without 4-cycles*, Discuss. Math. Graph Theory **24** (2004) 125–135.
<https://doi.org/10.7151/dmgt.1219>
- [21] H.-P. Yap, *Total Colourings of Graphs*, in: Lecture Notes in Math. **1623** (Springer, Berlin, Heidelberg, 1996).
<https://doi.org/10.1007/BFb0092895>

Received 14 May 2019

Revised 19 January 2020

Accepted 19 January 2020