Discussiones Mathematicae Graph Theory 43 (2023) 95–114 https://doi.org/10.7151/dmgt.2352

DOMINATION PARAMETERS OF THE UNITARY CAYLEY GRAPH OF $\mathbb{Z}/n\mathbb{Z}$

Amanda Burcroff

University of Michigan 500 S. State St. Ann Arbor, MI 48109 United States

e-mail: burcroff@umich.edu

Abstract

The unitary Cayley graph of $\mathbb{Z}/n\mathbb{Z}$, denoted X_n , is the graph with vertex set $\{0, \ldots, n-1\}$ where vertices a and b are adjacent if and only if gcd(a-b,n) = 1. We answer a question of Defant and Iyer by constructing a family of infinitely many integers n such that $\gamma_t(X_n) \leq g(n) - 2$, where γ_t denotes the total domination number and g denotes the Jacobsthal function. We determine the irredundance number, domination number, and lower independence number of certain direct products of complete graphs and give bounds for these parameters for any direct product of complete graphs. We provide upper bounds on the size of irredundant sets in direct products of balanced, complete multipartite graphs which are asymptotically correct for the unitary Cayley graphs of integers with a bounded smallest prime factor.

Keywords: unitary Cayley graph, domination chain, direct product, complete balanced multipartite graph.

2010 Mathematics Subject Classification: 05C69, 05C76.

1. INTRODUCTION

For a group Γ and a set $S = S^{-1} \subseteq \Gamma$ not containing the identity element, the Cayley graph $\operatorname{Cay}(\Gamma; S)$ is the undirected graph with vertices labeled by Γ and edge set $\{\{a, b\} : a - b \in S\}$. If Γ is a commutative ring with unity, the unitary Cayley graph X_{Γ} is the Cayley graph $\operatorname{Cay}(\Gamma, U_{\Gamma})$, where U_{Γ} is the set of units in Γ . More information on Cayley graphs and unitary Cayley graphs can be found in the algebraic graph theory texts by Biggs [5] and by Godsil and Royle [17]. We are interested in $X_n := X_{\mathbb{Z}/n\mathbb{Z}}$, that is, the graph on $\{0, \ldots, n-1\}$ where vertices

a and b are connected by an edge if and only if gcd(a - b, n) = 1. Observe that X_n is vertex-transitive and regular of degree $\phi(n)$, where ϕ denotes the Euler totient function.

These graphs were cast into the limelight in 1989, when Erdős and Evans showed that every finite simple graph G is isomorphic to an induced subgraph of X_n for some positive integer n, in which case they say G is representable modulo n. The representation number rep(G) of a graph G is the minimum positive integer such that G is representable modulo n. The representation numbers of many classes of graphs have been determined, including progress related to complete multipartite graphs and $\mathbb{Z}/n\mathbb{Z}$ [3, 13, 14]. See Section 7.6 of [15] for a survey of representation numbers and additional references.

In this paper, we consider domination parameters of the unitary Cayley graph of $\mathbb{Z}/n\mathbb{Z}$, including variants of the domination, irredundance, and independence numbers. Recall that a set $S \subseteq V(G)$ is called *independent* if it contains no pair of adjacent vertices. A set $S \subseteq V(G)$ is called *dominating* if every vertex of G is either contained in S or adjacent to a vertex of S. A set $S \subseteq V(G)$ is called *irredundant* if for each $v \in S$, either v is isolated in S or v has a neighbor $u \notin S$ such that u is not adjacent to any vertex of $S \setminus \{v\}$. These notions yield the following graph parameters for a graph G.

- The *irredundance number* ir(G) is the minimum size of a maximal irredundant set.
- The domination number $\gamma(G)$ is the minimum size of a dominating set.
- The lower independence number i(G), also known as the independent domination number, is the minimum size of a maximal independent set.
- The independence number $\alpha(G)$ is the maximum size of an independent set.
- The upper domination number $\Gamma(G)$ is the maximum size of a minimal dominating set.
- The upper irredundance number IR(G) is the maximum size of an irredundant set.

For any graph G, we have the following chain of inequalities, known as the *domination chain*

$$\operatorname{ir}(G) \le \gamma(G) \le i(G) \le \alpha(G) \le \Gamma(G) \le \operatorname{IR}(G).$$

Hundreds of papers have been written showing that some inequalities in this chain are equalities for certain classes of graphs, see Section 3.5 of [18] for more details. Many of these results were unified in 1994, when Cheston and Fricke showed that $\alpha(G) = \text{IR}(G)$ for any strongly perfect graph G [9].

It has been shown in [2] that the unitary Cayley graph of any finite commutative ring is a direct product of balanced, complete multipartite graphs, so we will often work in this more general setting. The direct product (sometimes called the tensor product or Kronecker product) of graphs G_1, \ldots, G_t , denoted by $\prod_{i=1}^t G_i$, is the graph with vertex set given by the Cartesian product of the vertex sets of G_i , i.e., $\prod_{i=1}^t V(G_i)$, where (g_1, \ldots, g_t) and (g'_1, \ldots, g'_t) in $\prod_{i=1}^t V(G_i)$ are adjacent if and only g_i and g'_i are adjacent in G_i for all $i \in \{1, \ldots, t\}$. The direct product of two graphs G and H is often denoted by $G \times H$ or $G \otimes H$. The balanced, complete b-partite graph with partite set size a, denoted K[a, b], is the graph on ab vertices partitioned into b partite sets of size a such that two vertices are adjacent if and only if they lie in different partite sets. If p is a prime and α is a positive integer, note that $X_{p^{\alpha}} \cong K[p^{\alpha-1}, p]$. An application of the Chinese remainder theorem shows that for an integer n with prime factorization $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_t^{\alpha_t}$, we have $X_n \cong \prod_{i=1}^t X_{p_i}^{\alpha_i} \cong \prod_{i=1}^t K[p_i^{\alpha_i-1}, p_i]$.

In this paper, we build upon the work of Defant and Iyer [11] to determine domination parameters of the unitary Cayley graphs of $\mathbb{Z}/n\mathbb{Z}$. Let g(n) denote the minimum positive integer m such that every set of m consecutive integers contains an integer which is coprime to n; this arithmetic function is known as the Jacobsthal function. The total domination number of a graph G, denoted by $\gamma_t(G)$, is the minimum size of a set S in G such that every vertex is adjacent to a member of S. Defant observed in [10] that there exist integers n such that $\gamma_t(X_n) \leq g(n) - 1$. In Section 3, we answer two questions of Defant and Iyer in the positive. The first asks whether there exists a single integer n such that $\gamma_t(X_n) \leq g(n) - 2$, and the second asks whether there exist integers n with arbitrarily many distinct prime factors such that $\gamma(X_n) \leq g(n) - 2$. We construct integers n with arbitrarily many distinct prime factors such that X_n contains a dominating cycle of size g(n) - 2; this answers both questions of Defant and Iyer since a dominating cycle is necessarily a total dominating set.

In Section 4, we provide bounds on the irredundance, domination, and lower independence numbers of direct products of complete graphs and determine these parameters in certain cases. One application of this work is the construction of some infinite families of integers n where $ir(X_n) = \gamma(X_n) = i(X_n)$. We provide an upper bound on the lower independence number of X_n in Corollary 4.12, thus demonstrating an error in a result of Uma Maheswari and Maheswari [20, Theorem 4.2]. Defant and Iyer [11] recently determined the value of $\gamma(\prod_{i=1}^{4} K_{n_i})$ in several cases; we compute this parameter in all cases. In Section 5, we provide upper bounds on the sizes of irredundant sets in direct products of balanced, complete multipartite graphs. In the case of unitary Cayley graphs of $\mathbb{Z}/n\mathbb{Z}$, we prove the following bound.

Corollary 5.6. Let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_t^{\alpha_t}$, where $p_1 < \cdots < p_t$. Then

$$\operatorname{IR}(X_n) \le \left(1 + 2 \cdot \frac{p_1}{p_t} \cdot \frac{1}{p_1^{\alpha_1 - 1} p_2^{\alpha_2 - 1} \cdots p_t^{\alpha_t - 1}}\right) \alpha(X_n).$$

In a preprint of the current article, we conjectured that the upper irredundance number was equal to the independence number for all direct products of balanced, complete multipartite graphs. This was a strengthening of an earlier conjecture of Defant and Iyer [11, Conjecture 4.2]. Since the online release of that preprint, Alon and Defant [4, Theorem 1.2] have confirmed the stronger conjecture for all but 37 such products, thus improving upon Corollary 5.6.

We conclude with a discussion of possible directions for further study and pose several open questions in Section 6.

2. Preliminaries

A graph G is a set of vertices V(G) along with a set of undirected edges E(G), excluding loops. For any $U \subseteq V(G)$, the subgraph of G induced by U, denoted G[U], is the graph with vertex set U and whose edge set is precisely the edge set E(G) restricted to $U \times U$.

Let N(v) denote the neighborhood of a vertex v, the set of all vertices adjacent to v. Let N[v] denote the *closed neighborhood* of a vertex v, that is, the set N(v) along with the vertex v itself. For $S \subseteq V(G)$, let $N[S] = \bigcup_{v \in S} N[v]$. A vertex $u \in V(G)$ is a *private neighbor* of a vertex $v \in S \subseteq V(G)$ if $u \in$ $(N[v] \setminus N[S \setminus \{v\}])$. Note u can equal v. Let pn[v] = pn[v; S] denote the set of private neighbors of $v \in S$. Let $pn[S] = \bigcup_{v \in S} pn[v; S]$. Note that a set $S \subseteq V(G)$ is irredundant if and only if every $v \in S$ has a private neighbor, and $S \subseteq V(G)$ is dominating if and only if N[S] = V(G).

Let $\mathbb{N} = \{1, 2, 3, ...\}$. For $n \in \mathbb{N}$, let $\omega(n)$ denote the number of distinct prime factors of n. For $S \subseteq \mathbb{N}$, let $\omega(S) = \{\omega(n) : n \in S\}$.

3. Dominating Cycles in the Unitary Cayley Graphs of $\mathbb{Z}/n\mathbb{Z}$

It was shown by Maheswari and Manjuri [21] that the value of the Jacobsthal function g(n) is an upper bound for the domination number of X_n , the unitary Cayley graph of $\mathbb{Z}/n\mathbb{Z}$. Defant and Iyer [11] note that the stronger inequality $\gamma_t(X_n) \leq g(n)$ holds and that these quantities can differ by 1 for n with arbitrarily many distinct prime factors.

We consider a variation of the domination number, introduced by Veldman [25] in 1983. The cycle domination number of a graph G, denoted $\gamma_c(G)$, is the minimum size of a dominating cycle in G, provided that such a cycle exists. Note that $\gamma_c(G) \geq \gamma_t(G) \geq \gamma(G)$ for any graph G. Since $(0, 1, \ldots, n-1)$ is a cycle in X_n , the cycle domination number of the unitary Cayley graph of $\mathbb{Z}/n\mathbb{Z}$ exists for all $n \in \mathbb{N}$.

In the following theorem, we exhibit an infinite family of integers n such that

 $g(n) - \gamma_t(X_n) \ge g(n) - \gamma_c(X_n) \ge 2$. Let $M_{c,j}$ be the set of positive integers n such that $g(n) - \gamma_c(X_n) \ge j$. Similarly, let $M_{t,j}$ be the set of positive integers n such that $g(n) - \gamma_t(X_n) \ge j$.

Theorem 3.1. The set $\omega(M_{c,2})$ is unbounded.

Proof. Let q be a prime such that $q \equiv 1 \mod 3$. Let $k = \frac{2q-2}{3}$, and let p_1, \ldots, p_k be primes such that $2q + 10 < p_1 < \cdots < p_k$. Let $n = 6q \prod_{j=1}^k p_j$.

We begin by showing $g(n) \ge 2q + 8$. For $i \in \{0, \dots, 2q + 6\}$, let

$$a_{i} = \begin{cases} 2 & \text{if } i \equiv 0 \mod 2, \\ 3 & \text{if } i \equiv 1 \mod 6, \\ q & \text{if } i = 3 \text{ or } i = 2q+3, \\ \min(\{p_{1}, \dots, p_{k}\} \setminus \{a_{0}, \dots, a_{i-1}\}) & \text{otherwise.} \end{cases}$$

Let $L(x) = \{i \in \{0, \dots, 2q+6\} : a_i = x\}$. Note that $|L(2)| = q+4, |L(3)| = \frac{q+5}{3}$, and |L(q)| = 2, and $|L(p_j)| = 1$ for each $j \in \{1, \ldots, k\}$. By the Chinese remainder theorem, there exists a $z \in \mathbb{Z}$ such that $z \equiv -i \mod a_i$ for all *i*. We note that the Chinese remainder theorem can be applied because the a_i 's are all either coprime or equal, and in the case that $a_i = a_j$ for $i \neq j$, we have $-i \equiv -j$ mod a_i . Hence it is equivalent to consider the set of conditions that $z \equiv -i$ mod a_i , where *i* ranges over a subset $I \subset \{0, \ldots, 2q+6\}$ is chosen such that the a_i 's are distinct and $\{a_i\}_{i \in I} = \{a_i\}_{0 \le i \le 2q+6}$. Having chosen such a $z \in \mathbb{Z}$, the set $\{z+i: 0 \le i \le 2q+6\}$ is a set of 2q+7 integers, none of which are relatively prime to n. Hence $g(n) \ge 2q + 8$.

Let y be the unique vertex of X_n such that $y \equiv 0 \mod 2$, $y \equiv 2 \mod 3$, $y \equiv -1 \mod q$, and $y \equiv -1 \mod p_i$ for all $1 \leq i \leq k$. Let z be the unique vertex of X_n such that $z \equiv 1 \mod 2$, $z \equiv 0 \mod 3$, $z \equiv -2 \mod q$, and $z \equiv -2$ mod p_i for all $1 \le i \le k$. Let $D = \{0, 1, \dots, 2q + 3, y, z\}$. We will show that the vertices of D form a cycle dominating set of X_n . Since $|D| = 2q + 6 \le g(n) - 2$, this will prove $n \in M_{c,2}$.

Suppose a vertex x is not adjacent to any element of $D \setminus \{y, z\}$. We will show that x is adjacent to either y or z. The set $S = \{x, x - 1, x - 2, \dots, x - (2q + 3)\}$ consists of 2q + 4 consecutive integers, none of which are coprime to n. For $r \in \mathbb{N}$, let $B(r) = \{s \in S : s \equiv 0 \mod r\}$. Let

$$B(2,3) = \{s \in S : s \equiv 0 \mod 2 \text{ or } s \equiv 0 \mod 3\}.$$

Observe that $|B(2,3)| = \frac{2}{3}(2q+4) = \frac{4q+8}{3}$, $|B(q)| \le 3$, and $|B(p_i)| \le 1$ for all p_i . Since

$$S = B(2,3) \cup B(q) \cup \bigcup_{i=1}^{\kappa} B(p_i),$$

we have

$$2q + 4 = \left| B(2,3) \cup B(q) \cup \bigcup_{i=1}^{k} B(p_i) \right| \le |B(2,3)| + |B(q)| + \sum_{i=1}^{k} |B(p_i)| \le \frac{4q + 8}{3} + 3 + \frac{2q - 2}{3} = 2q + 5.$$

This calculation implies that $|B(q) \setminus B(2,3)| \ge 2$. Let ℓ denote the minimum nonnegative integer such that $x - \ell \in B(q)$. If $\ell > 3$, then one element of $\{x - \ell, x - \ell - q\} = B(q)$ is even, hence $|B(q) \setminus B(2,3)| < 2$. Therefore, we must have $\ell \le 3$ with $B(q) \cap B(2,3) = \{x - \ell - q\}$. Hence the sets B(2,3), $B(q) \setminus \{x - \ell - q\}$, and $\bigcup_{i=1}^{k} B(p_i)$ are disjoint.

Thus, exactly one of x, x - 1, x - 2, x - 3 is contained in B(q). In particular, $x \not\equiv -1, -2 \mod q$. For each $1 \leq i \leq k$, from the fact that $|B(p_i)| = 1$ and the assumption that $p_i \geq 2q + 10$, we can conclude that $x \not\equiv -1, -2 \mod p_i$.

If x is not adjacent to y, then either $x \equiv 2 \mod 3$ or $x \equiv 0 \mod 2$. We will show that, under these conditions, $x \equiv 2 \mod 6$ or $x \equiv 4 \mod 6$. Since $z \equiv 3 \mod 6$, this is enough to show that x is adjacent to z.

Suppose $x \equiv 2 \mod 3$.

- If $x \equiv 0 \mod 2$, then $x \equiv 2 \mod 6$.
- If $x \equiv 1 \mod 2$, then $x \equiv 5 \mod 6$. Thus $x 1, x 2, x 3 \in B(2, 3)$, so $x \in B(q)$. Hence $x - 2q \in B(q)$. However, $x - 2q \equiv 2 - 2 \equiv 0 \mod 3$, contradicting the disjointedness of B(2, 3) and $B(q) \setminus \{x - \ell - q\}$.

Now suppose $x \not\equiv 2 \mod 3$ and $x \equiv 0 \mod 2$.

- If $x \equiv 1 \mod 3$, then $x \equiv 4 \mod 6$.
- If $x \equiv 0 \mod 3$, then $x \equiv 0 \mod 6$. Thus $x, x 2, x 3 \in B(2,3)$. Hence $x 1 \in B(q)$. However, $(x 1) 2q \equiv 2 2 \equiv 0 \mod 3$, contradicting the disjointedness of B(2,3) and $B(q) \setminus \{x \ell q\}$.

Thus, if x is not adjacent to y, then $x \equiv 4 \mod 6$ or $x \equiv 2 \mod 6$. We can conclude that D is indeed a total dominating set, and we have

$$\gamma_t(X_n) \le 2q + 6 = (2q + 8) - 2 \le g(n) - 2.$$

Lastly, note that y is adjacent to 1, z is adjacent to 2, and y is adjacent to z. Therefore, $(0, 1, y, z, 2, 3, \dots, 2q + 3)$ is a dominating cycle.

We will briefly expand upon the motivation behind the construction in Theorem 3.1. Fix an integer d and a prime q. Let $[n]_d$ denote the smallest nonnegative integer equivalent to n modulo d. Let $R_{q,d}$ be the set of integers $x \in \{0, \ldots, d-1\}$ such that x and $[x - 2q]_d$ are relatively prime to d. Let $R_{q,d,k} = \{x + \ell : x \in R_{q,d} \text{ and } 0 \le \ell \le k-1\}$. A key property used in both the construction of integers n for which $\gamma_t(X_n) \le g(n) - 1$ by Defant and Iyer [11] (using d = 2 and k = 2) and the construction in Theorem 3.1 (using d = 6 and k = 4) is that $R_{q,d,k}$ can be covered by relatively few vertices in X_d , namely 1 vertex in [11] and 2 in Theorem 3.1. Note that k is the minimum integer such that, for x not adjacent the consecutive vertices of the constructed dominating set, one of $x, x - 1, \ldots, x - (k-1)$ is divisible by q. If other such triples of integers (q, d, k) can be found, similar constructions could yield other families of integers in $M_{c,2}$ and perhaps even $M_{c,j}$ for $j \ge 3$.

Budadoddi and Mallikarjuna Reddy claim in [8] that the cycle dominating number (see Section 3) of X_n is given by the Jacobsthal function g(n), provided n is neither a prime power nor twice a prime power. Theorem 3.1 shows that this is not the case; in fact, there are integers n with $\omega(n)$ arbitrarily large such that $g(n) - \gamma_c(X_n) \ge 2$.

In the family constructed in Theorem 3.1, we see that $\gamma_c(X_n) \leq g(n)$. However, this inequality does not hold for all integers. For example, it is easily seen that $\gamma_c(X_6) = 6$ while g(6) = 4. We do not know if there exist infinitely many integers for which $\gamma_c(X_n) > g(n)$. The construction of a dominating set in X_n of size g(n) by Manjuri and Maheswari [21] shows that $\gamma_c(X_n) \leq g(n)$ whenever gcd(n, g(n)) = 1.

Theorem 3.1 also answers two questions of Defant and Iyer from [11], the first asking whether $M_{t,2}$ is nonempty and the second asking whether there exist integers n with $\omega(n)$ arbitrarily large such that $\gamma(X_n) \leq g(n) - 2$.

Corollary 3.2. There exist integers n with arbitrarily many distinct prime factors such that $\gamma(X_n) \leq \gamma_t(X_n) \leq g(n) - 2$.

This leads to the natural next question: does there exist a single integer such that $\gamma_t(X_n) \leq g(n) - 3$?

4. Lower Domination Parameters in Products of Complete Graphs

In this section, we consider the quantities in the lower portion of the domination chain for products of complete graphs. It is often useful to think of vertices in $\prod_{i=1}^{t} K_{n_i}$ as *t*-tuples of integers where the *i*th-entry is in the range $\{0, \ldots, n_i - 1\}$, where two vertices are adjacent if and only if their corresponding vectors differ in every coordinate. For squarefree positive integers *n*, we refer to vertices in X_n as integers and tuples interchangeably.

4.1. Irredundant sets in products of complete graphs

We will make use of two previous results; the first from Defant and Iyer in [11],

and the second from Bollobás and Cockayne, as well as Allan and Laskar, independently, in [6, 1].

Theorem 4.1 [11]. Let $G = \prod_{i=1}^{t} K_{n_i}$, where $2 \le n_1 \le n_2 \le \cdots \le n_t$, $t \ge 4$, and $n_2 \ge 3$. We have

$$\gamma(G) \ge t + 1 + \left\lfloor \frac{t-1}{n_1 - 1} \right\rfloor.$$

Theorem 4.2 [1, 6]. For any graph G, $ir(G) \ge \frac{1}{2}(\gamma(G) + 1)$.

As a direct consequence of these two theorems, we have the following lower bound on the irredundance numbers of direct products of at least 4 complete graphs.

Corollary 4.3 [1, 6, 11]. Let $G = \prod_{i=1}^{t} K_{n_i}$, where $2 \le n_1 \le n_2 \le \cdots \le n_t$, $t \ge 4$, and $n_2 \ge 3$. We have $\operatorname{ir}(G) \ge \frac{1}{2} \left(t + \left\lfloor \frac{t-1}{n_1-1} \right\rfloor \right) + 1$.

For direct products of at most 3 complete graphs, we compute the irredundance number exactly.

Theorem 4.4. Let $G = \prod_{i=1}^{t} K_{n_i}$, where $2 \le n_1 \le n_2 \le \cdots \le n_t$. If t = 1, then ir(G) = 1. If t = 2, then

$$\operatorname{ir}(G) = \begin{cases} 2, & \text{if } n_1 = 2, \\ 3, & \text{if } n_1 \ge 3. \end{cases}$$

If t = 3, then ir(G) = 4.

Proof. As $ir(G) \leq \gamma(G)$, the calculation of the domination number for $G = \prod_{i=1}^{t} K_{n_i}$ for $t \leq 3$ by Mekiš [22] proves that these irredundance numbers are at most the stated values. The case t = 1 is trivial, since any single vertex is a maximal irredundant set.

Let t = 2 and $n_1 = 2$. Fix an irredundant set $\{(x_1, y_1)\}$. We claim this set is not a maximal irredundant set. This follows from the fact that $\{(x_1, y_1), (1 - x_1, y_1)\}$ is also irredundant, as each vertex is its own private neighbor. Therefore, ir $(G) \ge 2$.

Let t = 2, and suppose $n_1 \ge 3$. Clearly no vertex of G is dominating, so ir(G) > 1. Fix an irredundant set $S = \{(x_1, y_1), (x_2, y_2)\}$. Suppose these two vertices are equal in some coordinate, without loss of generality $x_1 = x_2$. Fix $y_3 \in \{1, 2, 3\} \setminus \{y_1, y_2\}$. The set $\{(x_1, y_1), (x_2, y_2), (x_1, y_3)\}$ is an independent hence irredundant set. Thus, we can assume $x_1 \ne x_2$ and $y_1 \ne y_2$. Let S' be the set $\{(x_1, y_1), (x_2, y_2), (x_1, x_2)\}$ and $z_2 \in \{1, 2, 3\} \setminus \{y_1, y_2\}$. Then (z_1, y_2) is a private neighbor of (x_1, y_1) in S', (x_1, z_2) is a private neighbor.

of (x_2, y_2) in S', and (x_1, y_2) is its own private neighbor in S'. Therefore, the minimum size of a maximal irredundant set in G is at least 3.

Let t = 3. Suppose, seeking a contradiction, that

$$S = \{(x_1, y_1, z_1), (x_2, y_2, z_2), (x_3, y_3, z_3)\}$$

is a maximal irredundant set in G. We can assume these three vertices are not all equal in any coordinate, otherwise S can be extended to an independent set of size 4 by taking a fourth vector which is also equal in that coordinate.

Suppose that S is independent; without loss of generality assume $x_1 = x_2 \neq x_3$, $y_1 = y_3 \neq y_2$, and $z_2 = z_3 \neq z_1$. Note that if $n_3 = 2$, every irredundant set is independent. The point (x_3, y_2, z_1) is not in S, and we have that $S \cup \{(x_3, y_2, z_1)\}$ is independent. This contradicts that S is a maximal irredundant set.

Thus, we can assume S is not independent; without loss of generality assume

$$S = \{(0,0,0), (1,1,1), (x_3, y_3, z_3)\}.$$

If each of x_3 , y_3 , and z_3 is greater than 1, then $S \cup \{(0,1,z_3)\}$ is irredundant, contradicting the maximality of S. Thus, we can assume $x_3 \in \{0,1\}$. If $y_3 \ge 2$ and $z_3 \ge 2$, then $S \cup \{(x_3,0,1)\}$ is irredundant, which contradicts the maximality of S. Therefore, at least one of y_3, z_3 is in the set $\{0,1\}$, so without loss of generality we shall assume that $y_3 \in \{0,1\}$. Moreover, we may assume that $x_3 = 0$. However, either $S \cup \{(0,0,1)\}$ or $S \cup \{(0,1,0)\}$ is irredundant as well. Therefore, we can conclude $ir(G) \ge 4$.

4.2. Dominating sets in products of complete graphs

We provide an upper bound on the domination number of any product of t complete graphs which is exponential in t. This is an improvement on the upper bound yielded by a theorem of Brešar, Klavžar, and Rall [7], stating that $\gamma(G \times H) \leq 3\gamma(G)\gamma(H)$ for any graphs G and H. This implies $\gamma\left(\prod_{i=1}^{t} K_{n_i}\right) \leq 3^{t-1}$. We show that $\gamma\left(\prod_{i=1}^{t} K_{n_i}\right) \leq 3 \cdot 2^{t-2}$.

Theorem 4.5. Let $G = \prod_{i=1}^{t} K_{n_i}$. Let M be a family of vertices in $\{0,1\}^t \subseteq V(G)$ such that no two vertices in M are equal in t-1 coordinates or different in all t coordinates. Then $\{0,1\}^t \setminus M$ is a dominating set for G.

Proof. Let $D = \{0,1\}^t \setminus M$. Suppose $v \in \{0,1\}^t$. By the requirement that no two vertices in M differ in all coordinates, at least one of v and its Boolean complement $(1, \ldots, 1) - v$ is in D. Hence v is dominated by D.

Suppose $u \in V(G) \setminus \{0, 1\}^t$, that is, u has at least one coordinate which is not in the set $\{0, 1\}$. Fix $\ell \in \{1, \ldots, t\}$ to be some coordinate in which u is neither 0 nor 1. Observe that there exist two vertices w_1 and w_2 in $\{0, 1\}^t$, differing

in only the ℓ^{th} coordinate, which differ from u in every coordinate. Namely, for $i \neq \ell$ we can choose some element $a_i \in \{0, 1\}$ such that a_i is not equal to the i^{th} entry of u. Then we let $w_1 = (a_1, \ldots, a_{\ell-1}, 0, a_{\ell+1}, \ldots, a_t)$ and $w_2 = (a_1, \ldots, a_{\ell-1}, 1, a_{\ell+1}, \ldots, a_t)$. Since w_1 and w_2 differ from u in every coordinate, w_1 and w_2 are both adjacent to u in G. Since w_1 and w_2 are equal in t-1 coordinates, at least one of them is in D. Therefore, u is dominated by D. We conclude that D is a dominating set of G.

Let A(t, d, t - 1) denote the maximum number of binary vectors of length t such that any two distinct vectors have Hamming distance between d and t - 1, inclusive. Let A(t, d) denote the maximum number of binary vectors such that no two vectors have Hamming distance less than d. By taking a set witnessing A(t, d) and throwing out one of the vectors in any pair of Boolean complements, we obtain the following bound.

A classic result in coding theory is the Gilbert-Varshamov lower bound [16, 24] on A(t, d), originally stated for alphabets of prime power size.

Theorem 4.6 [16, 24]. If k satisfies

$$2^k < \frac{2^t}{\sum_{j=0}^{d-2} {t \choose j}}$$

then $A(t,d) \geq 2^k$.

Corollary 4.7. Let $G = \prod_{i=1}^{t} K_{n_i}$. We have

$$\gamma(G) \le 2^t - A(t, 2, t-1) \le 3 \cdot 2^{t-2}.$$

Proof. By Theorem 4.6, we have $A(t,2) \geq 2^{t-1}$. Let M be a set of length t binary vectors such that no two vectors are equal in t-1 coordinates and $|M| = 2^{t-1}$. We can delete one vector in each pair of Boolean complements in M to obtain a set $M' \subseteq M$ of size at least 2^{t-2} such that no two vectors are equal in t-1 coordinates nor differ in all coordinates. Theorem 4.5 implies $\{0,1\}^t \setminus M'$ is a dominating set, hence

$$\gamma(G) \le 2^t - 2^{t-2} = 3 \cdot 2^{t-2}.$$

We now determine the domination number of a product of four complete graphs, extending the results of Defant and Iyer in [11] and Mekiš in [22]. Defant and Iyer determined $\gamma_t \left(\prod_{i=1}^4 K_{n_i}\right)$ in the cases when $n_1 = 2$, $n_3 > 4 = n_1$, or $n_2 - 2 > 3 = n_1$. Mekiš determined $\gamma \left(\prod_{i=1}^4 K_{n_i}\right)$ in the case that $n_1 \ge 5$. **Theorem 4.8.** If $2 \le n_1 \le n_2 \le n_3 \le n_4$ and $n_2 \ge 3$, then

$$\gamma\left(\prod_{i=1}^{4} K_{n_{i}}\right) = \begin{cases} 8 & \text{if } n_{1} = 2, \text{ or if } n_{1} = n_{2} = 3, \\ 7 & \text{if } n_{1} = 3, n_{2} \in \{4, 5\}, \text{ or if } n_{1} = n_{3} = 4, n_{4} \in \{4, 5\}, \\ 6 & \text{if } n_{1} = 3, n_{2} > 5, \text{ or if } n_{1} = 4, (n_{3}, n_{4}) \notin \{(4, 4), (4, 5)\}, \\ 5 & \text{if } n_{1} \ge 5. \end{cases}$$

Moreover, if $n_1 \ge 3$, then $\gamma\left(\prod_{i=1}^4 K_{n_i}\right) = \gamma_t\left(\prod_{i=1}^4 K_{n_i}\right)$.

Proof. Let $G = \prod_{i=1}^{4} K_{n_i}$. We will proceed through the cases in increasing order of domination number. The case $n_1 \geq 5$ is determined by Corollary 2.2 of [22]. Furthermore, the same result yields that if $n_1 < 5$, then $\gamma(G) \geq 6$.

Consider those cases when the theorem claims $\gamma(G) = \gamma_t(G) = 6$. The cases $n_1 = 4, n_3 \ge 5$ or $n_1 = 3, n_2 > 5$ follow from Theorem 2.8 of [11], using the fact that the dominating sets constructed by Defant and Iyer are in fact total dominating sets. To show that $\gamma(G) = \gamma_t(G) = 6$ when $n_1 = 4$ and $n_4 \ge 6$, we construct a total dominating set D. Let

$$D = \{(0, 0, 0, 0), (0, 1, 1, 1), (1, 0, 1, 2), (1, 1, 0, 3), (2, 2, 2, 4), (3, 3, 3, 5)\}.$$

Observe that there are no two pairs of vertices in D such that each pair is equal in a different coordinate. Moreover, no three vertices of D are equal in any coordinate. Therefore any vertex of G is adjacent to a vertex of D, hence D is a total dominating set of G.

Next, we handle the cases in which $\gamma(G) = \gamma_t(G)$ are claimed to equal 7. We begin by showing that $\gamma(G) \geq 7$ if $n_1 = 3, n_2 \leq 5$ or $n_1 = n_2 = n_3 = 4, n_4 \in \{4, 5\}$. The lower bound in the case $n_1 = 3, n_2 \leq 5, n_3 \geq 5$ follows from Theorem 2.8 of [11]. Thus we need only handle the case $n_3 \leq 4$. Fix $\tilde{D} \subseteq V(G)$ such that $|\tilde{D}| = 6$. It is straightforward to show that if three vertices of \tilde{D} are equal in some coordinate, then there are at least 6 vertices which are not adjacent to a vertex of \tilde{D} , hence \tilde{D} is not dominating. Observe that there must be at least seven tuples $(\{v, w\}, i) \in {\tilde{D} \choose 2} \times \{1, 2, 3, 4\}$ such that the i^{th} coordinates of v and w are equal. It is straightforward to show that there exist two such tuples $(\{v, w\}, i)$ and $(\{\hat{v}, \hat{w}\}, \hat{i})$ such that $\{v, w\} \cap \{\hat{v}, \hat{w}\} = \emptyset$ and $i \neq \hat{i}$. Thus the vertex which is equal to v in the i^{th} coordinate, \hat{v} in the \hat{i}^{th} coordinate, and each vertex of $\tilde{D} \setminus \{v, w, \hat{v}, \hat{w}\}$ in the remaining two coordinates is not dominated by \tilde{D} . We can conclude that $\gamma(G) \geq 7$ in the claimed cases. For the upper bound, consider the set

$$D = \{(0,0,0,0), (0,1,1,1), (0,2,2,2), (1,0,1,1), (1,1,2,2), (1,2,0,0), (2,3,3,3)\}.$$

Observe that if we take any subset of D determined by omitting those vertices with a fixed first entry, the projection of the remaining vertices to the last three coordinates forms a total dominating set of $K_{n_2} \times K_{n_3} \times K_{n_4}$. Thus D is a total dominating set of G.

Lastly, we consider those cases in which $\gamma(G) = \gamma_t(G)$ are claimed to equal 8. When $n_1 = 2$, the result follows from Theorem 2.9 of [11]. When $n_1 = 3$, we obtain the upper bound by constructing an explicit dominating set. Let

$$D = \{(0,0,0,0), (0,0,1,1), (1,1,0,1), (1,1,1,0), \\(1,2,2,2), (2,2,0,1), (2,2,1,0), (2,1,2,2)\}.$$

A similar technique to that used earlier of removing vertices with a fixed entry in one coordinate and verifying that the projection of the remaining vertices to the remaining coordinates forms a total dominating set shows that D is indeed a total dominating set of G. For the lower bound when $n_1 = n_2 = 3$, suppose there exists a dominating set of size 7. By the pigeonhole principal, at least 3 vertices are equal in the first coordinate, and it is easy to see that at most 3 vertices are equal in any coordinate. The projection of the remaining 4 vertices to $K_3 \times K_{n_3} \times K_{n_4}$ must be adjacent to 3 vertices of $K_3 \times K_{n_3} \times K_{n_4}$, which is not possible (as the only dominating sets of $K_3 \times K_{n_3} \times K_{n_4}$ are independent).

4.3. Maximal independent sets in products of complete graphs

We begin by calculating the lower independence numbers of products of two or three complete graphs.

Theorem 4.9. Let $G = \prod_{i=1}^{t} K_{n_i}$ for $2 \le n_1 \le \cdots \le n_t$. We have

$$i(G) = \begin{cases} n_1 & \text{if } t = 2, \\ 4 & \text{if } t = 3. \end{cases}$$

Proof. Suppose t = 2. If $n_1 = 2$, observe that $\{(0,0), (1,0)\}$ is a maximal independent set and $i(G) \ge \gamma(G) \ge 2$, hence $i(G) = n_1 = 2$. For $n_1 \ge 3$, we will show that the maximal independent sets are precisely the fibers under projection onto some coordinate. Since the minimum size of such fibers is n_1 , occurring when the projection is onto the second coordinate, this is sufficient.

Fix three independent vertices $(x_1, x_2), (y_1, y_2), (z_1, z_2) \in V(G)$. It is impossible to have both $x_1 = y_1 \neq z_1$ and $y_2 = z_2 \neq x_2$, as in this case $x_1 \neq z_1$ and $x_2 \neq z_2$. Thus, every set of three independent vertices must be equal in some coordinate. We can conclude that the maximal independent sets must all be equal in some coordinate, hence can be extended to a fiber under the projection onto that coordinate. Therefore $i(G) = n_1$.

Now we handle the case t = 3. Observe that the set

$$\{(0,0,0),(1,0,1),(1,1,0),(0,1,1)\}$$

forms a maximal independent set. Theorem 4.4 provides the lower bound $4 = \gamma(G) \leq i(G)$, so we can conclude i(G) = 4.

Corollary 4.10. For all integers $n_1, n_2, n_3 \ge 3$, we have

ir
$$(K_2 \times K_{n_1}) = \gamma (K_2 \times K_{n_1}) = i (K_2 \times K_{n_1}),$$

ir $(K_3 \times K_{n_1}) = \gamma (K_3 \times K_{n_1}) = i (K_3 \times K_{n_1}),$
ir $(K_{n_1} \times K_{n_2} \times K_{n_3}) = \gamma (K_{n_1} \times K_{n_2} \times K_{n_3}) = i (K_{n_1} \times K_{n_2} \times K_{n_3}).$

We also briefly note an error in a paper by Uma Maheswari and Maheswari [20, Theorem 4.2]. Their paper correctly shows that $i(X_n) \leq \frac{n}{p_t}$, where p_t is the largest prime divisor of n, but they incorrectly claim that equality holds for all n. We show that the gap between this upper bound and the true value of $i(X_n)$ can be arbitrarily large.

Theorem 4.11. Let p_1, \ldots, p_t be distinct primes, where $t \ge 3$. For $n = p_1 \cdots p_t$, we have $i(X_n) \le 4p_1p_2 \cdots p_{t-3}$.

Proof. Recall from the proof of Theorem 4.9 that

$$E = \{(0,0,0), (1,0,1), (1,1,0), (0,1,1)\}$$

is an independent dominating set of $K_{p_{t-2}} \times K_{p_{t-1}} \times K_{p_t}$. We claim that the set

$$D = \left(\prod_{i=1}^{t-3} \{0, \dots, p_i - 1\}\right) \times \{(0, 0, 0), (1, 0, 1), (1, 1, 0), (0, 1, 1)\}$$

is an independent dominating set of $\prod_{i=1}^{t} K_{p_i}$. Since we require our primes p_1, \ldots, p_t to be distinct, we have $X_n = \prod_{i=1}^{t} K[1, p_i] = \prod_{i=1}^{t} K_{p_i}$.

For any vertex x equal to an element of D in the last three coordinates, we have $x \in D$. For any vertex x not equal to an element of E in the last three coordinates, we can form a vertex $y \in D$ adjacent to x by taking $y_i = x_i + 1$ in the first t - 3 coordinates and choosing a dominating element of E in the last three. Lastly, we note that D is independent as any two vertices of D will be equal in at least one of the last three coordinates.

Therefore, the upper bound given by Uma Maheswari and Maheswari [20, Theorem 4.2] is tight only when n has at most 2 prime divisors.

Corollary 4.12. For any $t \geq 3$ and $\epsilon > 0$, there exists a positive integer n such that $\omega(n) = t$ and $\frac{i(X_n)}{n/p_t} < \epsilon$, where p_t is the largest prime divisor of n.

5. Upper Domination Parameters of Products of Complete Multipartite Graphs

We now shift our focus to the upper portion of the domination chain and broaden our scope to products of balanced, complete multipartite graphs. In [11], Defant and Iyer initiated the investigation of the upper domination parameters of products of balanced, complete multipartite graphs, proving the following result.

Theorem 5.1 [11]. Let $G = \prod_{i=1}^{t} K[a_i, b_i]$ with $2 \le b_1 \le \cdots \le b_t$. If $b_1 = 2$ or $t \le 3$, then $\alpha(G) = IR(G)$.

In fact, the original result stated only that $\alpha(G) = \Gamma(G)$, but their methods never use the hypothesis that D is dominating, only that D is irredundant. In this section, we provide upper bounds for IR(G) in products of balanced, complete multipartite graphs not covered by Theorem 5.1. Since the release of the preprint of the current article, Alon and Defant [4] have improved upon these bounds in all but finitely many cases using recent progress in the study of vertex isoperimetry. In particular, they proved the following theorem.

Theorem 5.2 [4, Theorem 1.2]. Let G be a product of complete, balanced multipartite graphs. For all but at most 37 exceptional cases (listed in [4]), we have $\alpha(G) = IR(G)$.

Our approach, though ultimately yielding obtains a weaker result, is shorter and fairly elementary.

Fix an irredundant set $S \subseteq V(G)$. If $v \in S$ is isolated in G[S], we say v is *lonely*; otherwise, we say v is *social*. Let Lon(S) denote the set of lonely vertices in S, and let Soc(S) denote the set of social vertices in S. Observe that if $v \in S$ is social, then $\text{pn}[v; S] \subseteq V(G) \setminus S$.

As we did in the case of products of complete graphs, it is often useful to associate a *t*-tuple of integers to each vertex in $\prod_{i=1}^{t} K[a_i, b_i]$. Label the vertices of $K[a_i, b_i]$ by elements of $\mathbb{Z}/a_i b_i \mathbb{Z}$, where two vertices are adjacent if they are not congruent modulo b_i . Associate to $v \in \prod_{i=1}^{t} K[a_i, b_i]$ a vector of integers $(v(1), \ldots, v(t))$, where the v(i) is the element of $\{0, \ldots, a_i b_i - 1\}$ associated to the projection of v onto the i^{th} coordinate. Let p(v) be the vector $(v(1) \mod b_1, v(2) \mod b_2, \ldots, v(t) \mod b_t)$. Note that vertices u and v are adjacent in $\prod_{i=1}^{t} K[a_i, b_i]$ if and only if p(u) differs from p(v) in every coordinate.

Lemma 5.3. Let $G = \prod_{i=1}^{t} K[a_i, b_i]$, where $2 \leq b_1 \leq \cdots \leq b_t$. Let S be an irredundant set in G, and let $T \subseteq pn[S]$ be chosen so that each vertex of S has exactly one private neighbor in T and $S \cap T$ consists only of lonely vertices. If there exist three vertices $v_1, v_2, v_3 \in T$ such that the vectors $p(v_1), p(v_2), p(v_3)$ vary in exactly one coordinate, then v_1, v_2 , and v_3 are lonely vertices of S.

Proof. Suppose at least one of v_1, v_2, v_3 , say v_1 , is in $pn[S] \setminus Lon[S]$. Then v_1 is adjacent to $p(v_1)$. Since $p(v_1)$, $p(v_2)$, and $p(v_3)$ differ in exactly one coordinate, any vertex which is adjacent to at least one of $p(v_1), p(v_2), p(v_3)$ is adjacent to at least two of $p(v_1), p(v_2), p(v_3)$. Thus v_1 is adjacent to at least one of $p(v_2), p(v_3) \in S$, contradicting that v_1 is a private neighbor of $p(v_1)$.

Lemma 5.4. Let $G = \prod_{i=1}^{t} K[a_i, b_i]$, where $2 \leq b_1 \leq \cdots \leq b_t$. Let S be an irredundant set in G, and let $T \subseteq pn[S]$ be chosen so that each vertex of S has exactly one private neighbor in T. If there are two vertices $v_1, v_2 \in T$ such that $p(v_1) = p(v_2)$, then v_1 and v_2 are lonely vertices of S.

Proof. Note that $N(v_1) = N(v_2)$. Suppose, seeking a contradiction, that $v_1 \in T \setminus \text{Lon}[S]$. Then there exists some $w \in S$ such that w is adjacent to v_1 and v_2 , so v_2 cannot be lonely. Hence v_2 must also be a private neighbor of some vertex in S, but as is it already a neighbor of $w \in S$, this contradicts our assumption that each vertex in S has a unique private neighbor in T.

Theorem 5.5. If $G = \prod_{i=1}^{t} K[a_i, b_i]$, where $2 \leq b_1 \leq \cdots \leq b_t$, then

$$\operatorname{IR}(G) \le \frac{1}{b_1} \prod_{i=1}^t a_i b_i + \frac{2}{b_t} \prod_{i=1}^t b_i.$$

Proof. Suppose $S \subseteq V(G)$ is irredundant. Observe that $|\operatorname{Lon}(S)| \leq \alpha(G) = \frac{1}{b_1} \prod_{i=1}^t a_i b_i$. Let $T \subseteq \operatorname{pn}[S]$ be chosen so that each vertex of S has exactly one private neighbor in T and $S \cap T$ consists of lonely vertices. Note that such a set T always exists, since the latter condition is equivalent to, for each $s \in S$, choosing a vertex in the open neighborhood of s rather than s itself. By Lemma 5.4, no two vertices of $T \setminus \operatorname{Lon}[S]$ have the same associated vector. Moreover, by Lemma 5.3, no three vertices of $T \setminus \operatorname{Lon}[S]$ have associated vectors which agree in all but the last coordinate. Hence

$$|\operatorname{Soc}(S)| = |T \setminus \operatorname{Lon}[S]| \le \frac{2}{b_t} \prod_{i=1}^t b_i.$$

We can conclude

$$|S| = |\operatorname{Lon}(S)| + |\operatorname{Soc}(S)| \le \frac{1}{b_1} \prod_{i=1}^t a_i b_i + \frac{2}{b_t} \prod_{i=1}^t b_i.$$

In particular, for the case of unitary Cayley graphs of $\mathbb{Z}/n\mathbb{Z}$, we have the following bound.

Corollary 5.6. If $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_t^{\alpha_t}$, where $p_1 < \cdots < p_t$ are primes, then

$$\operatorname{IR}(X_n) \le \left(1 + 2 \cdot \frac{p_1}{p_t} \cdot \frac{1}{p_1^{\alpha_1 - 1} p_2^{\alpha_2 - 1} \cdots p_t^{\alpha_t - 1}}\right) \alpha(X_n).$$

If we look at sets of integers with a bounded smallest prime factor, we can show that $\operatorname{IR}(X_n)$ and $\alpha(X_n)$ are asymptotically the same. An integer x is called r-rough (or r-jagged) for a positive integer r if every prime factor of x is at least r. Let $D_r = \{x \in \mathbb{N} : \exists d \in \{2, 3, \ldots, r\}$ such that $d \mid x\}$; that is, D_r is the set of positive integers that are not (r + 1)-rough.

Corollary 5.7. For each $r \in \mathbb{N}$, we have

$$\lim_{\substack{n \in D_r \\ n \to \infty}} \frac{\mathrm{IR}(X_n)}{\alpha(X_n)} = 1$$

Proof. Let $n = p_1^{\alpha_1} \cdots p_n^{\alpha_n}$, where $p_1 < \cdots < p_n$ are primes. Hence $X_n \cong \prod_{i=1}^t K[a_i, b_i]$, where $a_i = p_i^{\alpha_i - 1}$ and $b_i = p_i$. By the condition $n \in D_r$, we know $b_1 \leq r$. Hence

$$\frac{\text{IR}(X_n)}{\alpha(X_n)} \le 1 + \frac{2b_1}{b_t \prod_{i=1}^t a_i} \le 1 + \frac{2r}{b_t \prod_{i=1}^t a_i}$$

Note that $b_t \prod_{i=1}^t a_i$ tends to infinity as n tends to infinity. Since $IR(X_n) \ge \alpha(X_n)$ for all n, the limit goes to 1.

We note that the bound in Theorem 5.5 is trivial if $a_1 = \cdots = a_t = 1$ and $b_1 = \cdots = b_t = 3$, that is, when G is a direct product of triangles. This case motivates the following theorem, which provides a nontrivial (though, usually worse) upper bound for the upper irredundance number of any direct product of balanced, complete multipartite graphs.

Theorem 5.8. If $G = \prod_{i=1}^{t} K[a_i, b_i]$, where $2 \leq b_1 \leq \cdots \leq b_t$, then

$$\operatorname{IR}(G) \le \frac{b_1}{2b_1 - 1} \prod_{i=1}^t a_i b_i.$$

Proof. Let $S \subseteq V(G)$ have size greater than $\frac{1}{2-\frac{1}{b_1}}\prod_{i=1}^t a_i b_i$. Suppose, seeking a contradiction, that S is irredundant.

Let P_j be the set of vertices $v \in S$ such that v(1) = j. By the Pigeonhole Principle, $|P_j| \geq \frac{1}{b_1}|S|$ for some $j \in \{0, \ldots, b_1 - 1\}$; without loss of generality, suppose $|P_0| \geq \frac{1}{b_1}|S|$.

For each $k \in \{1, \ldots, b_1 - 1\}$, let Q_k denote the set of private neighbors $u \in \bigcup_{v \in (S \setminus P_0)} \operatorname{pn}[v, S]$ such that u(1) = k. Observe that of the $|S| - |P_0|$ vertices



Figure 1. An example of how the sets P_0 , Q_1 , and R are determined by a fixed irredundant set $S = \{(0,0), (2,0), (2,2)\}$ in $K_3 \times K_3$. For a vertex v in the figure, v(0) is given by its row and v(1) by its column, hence two vertices are adjacent if and only if they lie in different rows and columns.

in $S \setminus P_0$, at most $\frac{1}{b_1} \prod_{i=1}^t a_i b_i - |P_0|$ have private neighbors u such that u(1) = 0. By the Pigeonhole Principle,

$$|Q_k| \ge \frac{1}{b_1 - 1} \left(|S| - |P_0| - |\{u \in \operatorname{pn}[S \setminus P_0] : u(1) = 0\}| \right)$$
$$\ge \frac{1}{b_1 - 1} \left(|S| - \frac{1}{b_1} \prod_{i=1}^t a_i b_i \right)$$

for some $k \in \{1, \ldots, b_1 - 1\}$; without loss of generality, assume

$$|Q_1| \ge \frac{1}{b_1 - 1} \left(|S| - \frac{1}{b_1} \prod_{i=1}^t a_i b_i \right)$$

Let R be the set of vertices u such that p(u) = (0, v(2) + 1, ..., v(t) + 1) for some $v \in Q_1$. That is,

$$R = \bigcup_{v \in Q_1} p^{-1}((0, v(2) + 1, \dots, v(t) + 1)).$$

Since the associated vectors are disjoint, each member of R is adjacent to a member of Q_1 . Using the fact that P_0 is disjoint from Q_1 and that Q_1 consists of private neighbors of $S \setminus P_0$, we conclude that P_0 and R are disjoint; lest a member of P_0 is adjacent to a private neighbor of $S \setminus P_0$. Since P_0 and R both consist of vectors u satisfying $u_1 = 0$, we have $|P_0| + |R| \leq \frac{1}{b_1} \prod_{i=1}^t a_i b_i$. However, we reach a contradiction by noting that

$$\begin{aligned} |P_0| + |R| &= |P_0| + |Q_1| \ge \frac{1}{b_1} |S| + \frac{1}{b_1 - 1} \left(|S| - \frac{1}{b_1} \prod_{i=1}^t a_i b_i \right) \\ &> \left(\frac{1}{2b_1 - 1} + \frac{1}{b_1 - 1} \left(\frac{b_1}{2b_1 - 1} - \frac{1}{b_1} \right) \right) \prod_{i=1}^t a_i b_i = \frac{1}{b_1} \prod_{i=1}^t a_i b_i. \end{aligned}$$

We conclude that S is not irredundant.

6. Further Directions

In Section 3, we raise the question of whether there exist infinitely many integers n such that $\gamma_c(X_n) > g(n)$ and, if so, whether such integers can have arbitrarily many distinct prime factors. At the end of this same section, we ask whether there exists a single integer n such that $\gamma_t(X_n) \leq g(n) - 3$.

We calculate the irredundance and lower independence numbers of direct products of at most three complete graphs in Section 4. It remains open to determine these parameters for larger products of complete graphs. From these calculations, it follows that $ir(X_n) = i(X_n)$ when n is prime, n = 2p, or n = 3pfor some prime p, or n is squarefree with exactly three prime divisors. We pose the problem of finding other squarefree integers n for which equality is achieved in the lower portion of the domination chain.

We note that the irredundance, domination, and lower independence numbers of $K_a \times K_b$ depend on min(a, b), whereas in the case of $K_a \times K_b \times K_c$ these parameters are independent of a, b, and c. We show that the domination number of a direct product of four complete graphs is dependent on the size of all four graphs in the product. We pose the question of determining for which integers t do the irredundance, domination, or lower independence numbers of $\prod_{i=1}^{t} K_{n_i}$ depend on all of n_1, n_2, \ldots, n_t , where $n_1 \leq n_2 \leq \cdots \leq n_t$.

In 2007, Klotz and Sander [19] introduced the notion of the gcd-graph $X_n(D)$, the graph on $\{0, \ldots, n-1\}$ where vertices x and y are adjacent if and only if $gcd(|x-y|, n) \in D \subseteq \mathbb{N}$. In particular, Klotz and Sander show that all eigenvalues of $X_n(D)$ are integral. It may be interesting to investigate domination parameters in the more general case of gcd-graphs.

Acknowledgements

The author would like to thank Joe Gallian for his tireless efforts to foster a productive and engaging mathematical community. She extends her gratitude to Colin Defant for introducing her to this problem and providing incredible support throughout her research. The author would also like to thank the anonymous referees for their helpful comments and simplifications to several proofs, as well as Harish Vemuri for a useful discussion about Theorem 4.8.

This research was conducted at the University of Minnesota, Duluth REU and was supported by NSF/DMS grant 1650947 and NSA grant H98230-18-1-0010.

References

 R.B. Allan and R. Laskar, On domination and some related topics in graph theory, in: Proceedings of the Ninth Southeast Conference on Combinatorics, Graph Theory, and Computing, Util. Math. (1978) 43–56.

- R. Akhtar, M. Boggess, T. Jackson-Henderson, I. Jiménez, R. Karpman, A. Kinzel and D. Pritikin, On the unitary Cayley graph of a finite ring, Electron. J. Combin. 16 (2009) #R117. https://doi.org/10.37236/206
- R. Akhtar, A.B. Evans and D. Pritikin, Representation numbers of complete multipartite graphs, Discrete Math. **312** (2012) 1158–1165. https://doi.org/10.1016/j.disc.2011.12.003
- N. Alon and C. Defant, Isoperimetry, stability, and irredundance in direct products, Discrete Math. 343 (2020) 111902. https://doi.org/10.1016/j.disc.2020.111902
- N. Biggs, Algebraic Graph Theory, Second Edition (Cambridge University Press, Cambridge, 1993). https://doi.org/10.1017/CBO9780511608704
- B. Bollobás and E.J. Cockayne, Graph-theoretic parameters concerning domination, independence, and irredundance, J. Graph Theory 3 (1979) 241-249. https://doi.org/10.1002/jgt.3190030306
- B. Brešar, S. Klavžar and D.F. Rall, Dominating direct products of graphs, Discrete Math. 307 (2007) 1636–1642. https://doi.org/10.1016/j.disc.2006.09.013
- [8] K. Budadoddi and A. Mallikarjuna Reddy, Cycle dominating sets of Euler totient Cayley graphs, Math. Sci. Int. Res. J. 3 (2014) 727–730.
- G.A. Cheston and G.H. Fricke, Classes of graphs for which the upper fractional domination equals independence, upper domination, and upper irredundance, Discrete Appl. Math. 55 (1994) 241–258. https://doi.org/10.1016/0166-218X(94)90011-6
- C. Defant, Unitary Cayley graphs of Dedekind domain quotients, AKCE Int. J. Graphs Comb. 13 (2016) 65–75. https://doi.org/10.1016/j.akcej.2016.03.001
- C. Defant and S. Iyer, Domination and upper domination of direct product graphs, Discrete Math. 341 (2018) 2742–2752. https://doi.org/10.1016/j.disc.2018.06.031
- [12] P. Erdős and A.B. Evans, Representations of graphs and orthogonal Latin square graphs, J. Graph Theory 13 (1989) 593–595. https://doi.org/10.1002/jgt.3190130509
- [13] A.B. Evans, Representations of disjoint unions of complete graphs, Discrete Math. 307 (2007) 1191–1198. https://doi.org/10.1016/j.disc.2006.07.029
- [14] A.B. Evans, G. Isaak and D.A. Narayan, *Representations of graphs modulo n*, Discrete Math. 223 (2000) 109–123. https://doi.org/10.1016/S0012-365X(00)00062-5

- [15] J.A. Gallian, Graph labeling, A dynamic survey, Electron. J. Combin. (2018) #DS6. https://doi.org/10.37236/27
- E.N. Gilbert, A comparison of signalling alphabets, The Bell System Technical Journal **31** (1952) 504–522. https://doi.org/10.1002/j.1538-7305.1952.tb01393.x
- C. Godsil and G. Royle, Algebraic Graph Theory, in: Grad. Texts in Math. 207 (Springer-Verlag, New York, 2001). https://doi.org/10.1007/978-1-4613-0163-9
- [18] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, Fundamentals of Domination in Graphs (Marcel Dekker, Inc, New York, 1998). https://doi.org/10.1201/9781482246582
- W. Klotz and T. Sander, Some properties of unitary Cayley graphs, Electron. J. Combin. 14 (2007) #R45. https://doi.org/10.37236/963
- [20] S.U. Maheswari and B. Maheswari, Independent domination number of Euler totient Cayley graphs and arithmetic graphs, Int. J. Adv. Res. Eng. Technol. 7(3) (2016) 56-65.
- M. Manjuri and B. Maheswari, Strong dominating sets of some arithmetic graphs, Int. J. Comput. Appl. 83(3) (2013) 36–40. https://doi.org/10.5120/14431-2577
- [22] G. Mekiš, Lower bounds for the domination number and the total domination number of direct product graphs, Discrete Math. **310** (2010) 3310–3317. https://doi.org/10.1016/j.disc.2010.07.015
- [23] M. Valencia-Pabon, Idiomatic partitions of direct products of complete graphs, Discrete Math. **310** (2010) 1118–1122. https://doi.org/10.1016/j.disc.2009.10.012
- [24] R.R. Varshamov, Estimate of the number of signals in error correcting codes, Dokl. Akad. Nauk 117 (1957) 739–741.
- [25] H.J. Veldman, Existence of dominating cycles and paths, Discrete Math. 43 (1983) 281–296. https://doi.org/10.1016/0012-365X(83)90165-6
- H. Vemuri, Domination in direct products of complete graphs, Discrete Appl. Math. 285 (2020) 473–482. https://doi.org/10.1016/j.dam.2020.06.015

Received 22 June 2019 Revised 21 July 2020 Accepted 25 July 2020