

## ON THE SIZES OF $(k, l)$ -EDGE-MAXIMAL $r$ -UNIFORM HYPERGRAPHS

YINGZHI TIAN<sup>1</sup>, HONG-JIAN LAI<sup>2</sup>, JIXIANG MENG<sup>1</sup>

AND

MURONG XU<sup>3</sup>

<sup>1</sup>*College of Mathematics and System Sciences  
Xinjiang University, Urumqi, Xinjiang 830046, PR China*

<sup>2</sup>*Department of Mathematics  
West Virginia University, Morgantown, WV 26506, USA*

<sup>3</sup>*Department of Mathematics  
The Ohio State University, Columbus, OH 43210, USA*

**e-mail:** tianyzhxj@163.com  
Hong-jian.Lai@mail.wvu.edu  
mjx@xju.edu.cn  
xu.3646@osu.edu

### Abstract

Let  $H = (V, E)$  be a hypergraph, where  $V$  is a set of vertices and  $E$  is a set of non-empty subsets of  $V$  called edges. If all edges of  $H$  have the same cardinality  $r$ , then  $H$  is an  $r$ -uniform hypergraph; if  $E$  consists of all  $r$ -subsets of  $V$ , then  $H$  is a complete  $r$ -uniform hypergraph, denoted by  $K_n^r$ , where  $n = |V|$ . An  $r$ -uniform hypergraph  $H = (V, E)$  is  $(k, l)$ -edge-maximal if every subhypergraph  $H'$  of  $H$  with  $|V(H')| \geq l$  has edge-connectivity at most  $k$ , but for any edge  $e \in E(K_n^r) \setminus E(H)$ ,  $H + e$  contains at least one subhypergraph  $H''$  with  $|V(H'')| \geq l$  and edge-connectivity at least  $k + 1$ . In this paper, we obtain the lower bounds and the upper bounds of the sizes of  $(k, l)$ -edge-maximal hypergraphs. Furthermore, we show that these bounds are best possible.

**Keywords:** edge-connectivity,  $(k, l)$ -edge-maximal hypergraphs,  $r$ -uniform hypergraphs.

**2010 Mathematics Subject Classification:** 05C65, 05C35.

## 1. INTRODUCTION

In this paper, we consider finite simple graphs. For graph-theoretical terminologies and notation not defined here, we follow [3]. For a graph  $G$ , we use  $\kappa'(G)$  to denote the *edge-connectivity* of  $G$ . The *complement* of a graph  $G$  is denoted by  $G^c$ . For  $X \subseteq E(G^c)$ ,  $G + X$  is the graph with vertex set  $V(G)$  and edge set  $E(G) \cup X$ . We will use  $G + e$  for  $G + \{e\}$ . The *floor* of a real number  $x$ , denoted by  $\lfloor x \rfloor$ , is the greatest integer not larger than  $x$ ; the *ceiling* of a real number  $x$ , denoted by  $\lceil x \rceil$ , is the least integer greater than or equal to  $x$ . For two integers  $n$  and  $k$ , we define  $\binom{n}{k} = \frac{n!}{k!(n-k)!}$  when  $k \leq n$  and  $\binom{n}{k} = 0$  when  $k > n$ .

For generalizing a prior result of Mader [7], Boesch and McHugh [2] introduced the following definitions. For integers  $k$  and  $l$  with  $l > k \geq 2$ , a graph  $G$  with  $n = |V(G)| \geq l$  is a  $(k, l)$ -graph if  $\kappa'(G') \leq k$  for any  $G' \subseteq G$  with  $|V(G')| \geq l$ . A  $(k, l)$ -graph  $G$  is  $(k, l)$ -edge-maximal if, for any  $e \in E(G^c)$ ,  $G + e$  has a subgraph  $G'$  with  $|V(G')| \geq l$  and  $\kappa'(G') \geq k + 1$ .  $(k, k + 1)$ -edge-maximal graphs have been studied in [5, 7, 8], among others.

**Theorem 1.1.** *Let  $k \geq 2$  be an integer, and  $G$  be a  $(k, k + 1)$ -edge-maximal graph on  $n > k + 1$  vertices. Each of the following holds.*

- (i) (Mader [7])  $|E(G)| \leq (n - k)k + \binom{k}{2}$ . Furthermore, this bound is best possible.
- (ii) (Lai [5])  $|E(G)| \geq (n - 1)k - \binom{k}{2} \lfloor \frac{n}{k+2} \rfloor$ . Furthermore, this bound is best possible.

In [2], Boesch and McHugh extended Theorem 1(i) to  $(k, l)$ -edge-maximal graphs.

**Theorem 1.2** (Boesch and McHugh [2]). *Let  $G$  be a graph of order  $n$  and let  $n \geq l \geq k + 1$ . Let  $p, q \geq 0$  be integers such that  $n = p(l - 1) + q$  with  $0 \leq q < l - 1$ . If  $G$  is a  $(k, l)$ -edge-maximal graph, then*

$$|E(G)| \leq \begin{cases} \frac{p(l-1)(l-2)}{2} + (p-1+q)k, & l-1 > 2k \text{ and } q \leq 2k, \\ \frac{p(l-1)(l-2)}{2} + pk + \frac{q(q-1)}{2}, & l-1 > 2k \text{ and } q > 2k, \\ \frac{(l-1)(l-2)}{2} + (n-l+1)k, & l-1 \leq 2k. \end{cases}$$

Furthermore, these bounds are best possible.

In [6], Lai and Zhang extended Theorem 1(ii) to  $(k, l)$ -edge-maximal graphs.

**Theorem 1.3** (Lai and Zhang [6]). *Let  $G$  be a graph of order  $n$  and let  $n \geq l \geq k + 3 \geq 5$ . Let  $p, q \geq 0$  be integers such that  $n = p(l - 1) + q$  with  $0 \leq q < l - 1$ .*

If  $G$  is a  $(k, l)$ -edge-maximal graph, then

$$|E(G)| \geq \begin{cases} \frac{(l-1)(l-2)}{2} + (n-l+1)k, & l \leq n < 2k+4, \\ (n-1)k - \lfloor \frac{n}{k+2} \rfloor \frac{(k+1)^2-3(k+1)}{2}, & l \leq 2k+4 \leq n, \\ (n-2a+1)k + a(a-1) - \lfloor \frac{n-2a}{k+2} \rfloor \frac{(k+1)^2-3(k+1)}{2}, & n \geq l = 2a \geq 2k+5, \\ (n-2b)k + b^2 - \lfloor \frac{n-2b-1}{k+2} \rfloor \frac{(k+1)^2-3(k+1)}{2}, & n \geq l = 2b+1 \geq 2k+5. \end{cases}$$

Furthermore, these bounds are best possible.

Let  $H = (V, E)$  be a *hypergraph*, where  $V$  is a finite set and  $E$  is a set of non-empty subsets of  $V$ , called *edges*. An edge of cardinality 2 is just a graph edge. For a vertex  $u \in V$  and an edge  $e \in E$ , we say  $u$  is *incident with*  $e$  or  $e$  is *incident with*  $u$  if  $u \in e$ . If all edges of  $H$  have the same cardinality  $r$ , then  $H$  is an  *$r$ -uniform hypergraph*; if  $E$  consists of all  $r$ -subsets of  $V$ , then  $H$  is a *complete  $r$ -uniform hypergraph*, denoted by  $K_n^r$ , where  $n = |V|$ . For  $n < r$ , the complete  $r$ -uniform hypergraph  $K_n^r$  is just the hypergraph with  $n$  vertices and no edges. The *complement* of an  $r$ -uniform hypergraph  $H = (V, E)$ , denoted by  $H^c$ , is the  $r$ -uniform hypergraph with vertex set  $V$  and edge set consisting of all  $r$ -subsets of  $V$  not in  $E$ . A hypergraph  $H' = (V', E')$  is called a *subhypergraph* of  $H = (V, E)$ , denoted by  $H' \subseteq H$ , if  $V' \subseteq V$  and  $E' \subseteq E$ . For  $X \subseteq E(H^c)$ ,  $H + X$  is the hypergraph with vertex set  $V(H)$  and edge set  $E(H) \cup X$ ; for  $X' \subseteq E(H)$ ,  $H - X'$  is the hypergraph with vertex set  $V(H)$  and edge set  $E(H) \setminus X'$ . We use  $H + e$  for  $H + \{e\}$  and  $H - e'$  for  $H - \{e'\}$  when  $e \in E(H^c)$  and  $e' \in E(H)$ . For  $Y \subseteq V(H)$ , we use  $H[Y]$  to denote the hypergraph *induced* by  $Y$ , where  $V(H[Y]) = Y$  and  $E(H[Y]) = \{e \in E(H) : e \subseteq Y\}$ .  $H - Y$  is the hypergraph induced by  $V(H) \setminus Y$ .

For a hypergraph  $H = (V, E)$  and two disjoint vertex subsets  $X, Y \subseteq V$ , let  $E_H[X, Y]$  be the set of edges intersecting both  $X$  and  $Y$  and  $d_H(X, Y) = |E_H[X, Y]|$ . We use  $E_H(X)$  and  $d_H(X)$  for  $E_H[X, V \setminus X]$  and  $d_H(X, V \setminus X)$ , respectively. If  $X = \{u\}$ , we use  $E_H(u)$  and  $d_H(u)$  for  $E_H(\{u\})$  and  $d_H(\{u\})$ , respectively. We call  $d_H(u)$  the *degree* of  $u$  in  $H$ . The *minimum degree*  $\delta(H)$  of  $H$  is defined as  $\min\{d_H(u) : u \in V\}$ ; the *maximum degree*  $\Delta(H)$  of  $H$  is defined as  $\max\{d_H(u) : u \in V\}$ . When  $\delta(H) = \Delta(H) = k$ , we call  $H$   *$k$ -regular*.

For a nonempty proper vertex subset  $X$  of a hypergraph  $H$ , we call  $E_H(X)$  an *edge-cut* of  $H$ . The *edge-connectivity*  $\kappa'(H)$  of a hypergraph  $H$  is  $\min\{d_H(X) : \emptyset \neq X \subsetneq V(H)\}$ . By definition,  $\kappa'(H) \leq \delta(H)$ . We call a hypergraph  $H$   *$k$ -edge-connected* if  $\kappa'(H) \geq k$ . A hypergraph is connected if it is 1-edge-connected. A maximal connected subhypergraph of  $H$  is called a *component* of  $H$ . An  $r$ -uniform hypergraph  $H = (V, E)$  is  *$(k, l)$ -edge-maximal* if every subhypergraph  $H'$  of  $H$  with  $|V(H')| \geq l$  has edge-connectivity at most  $k$ , but for any edge  $e \in E(H^c)$ ,  $H + e$  contains at least one subhypergraph  $H''$  with  $|V(H'')| \geq l$

and edge-connectivity at least  $k + 1$ . If  $H$  is a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph with  $n = |V(H)| < l$ , then  $H \cong K_n^r$ . For results on the connectivity of hypergraphs, see [1, 4] for references.

In order to construct the complete  $r$ -uniform hypergraph with the maximum number of vertices and degree at most  $k$ , we introduce the parameter  $t = t(k, r)$ , which is determined by  $k$  and  $r$ .

**Definition 1.4.** For two integers  $k$  and  $r$  with  $k, r \geq 2$ , define  $t = t(k, r)$  to be the largest integer such that  $\binom{t-1}{r-1} \leq k$ . That is,  $t$  is the integer satisfying  $\binom{t-1}{r-1} \leq k < \binom{t}{r-1}$ .

In [9], the authors determined, for given integers  $n$ ,  $k$  and  $r$ , the extremal sizes of  $(k, t)$ -edge-maximal  $r$ -uniform hypergraphs on  $n$  vertices.

**Theorem 1.5** (Tian, Xu, Lai and Meng [9]). *Let  $H$  be a  $(k, t)$ -edge-maximal  $r$ -uniform hypergraph such that  $n \geq t$  and  $k, r \geq 2$ , where  $n = |V(H)|$  and  $t = t(k, r)$ . Then each of the following holds.*

- (i)  $|E(H)| \leq \binom{t}{r} + (n - t)k$ . Furthermore, this bound is best possible.
- (ii)  $|E(H)| \geq (n - 1)k - ((t - 1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ . Furthermore, this bound is best possible.

The main goal of this research is to extend these results in [9]. For given integers  $n$ ,  $k$  and  $r$ , the extremal sizes of a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph on  $n$  vertices are determined, where  $l \geq t + 1$ . Section 2 below is devoted to the study of some properties of  $(k, l)$ -edge-maximal  $r$ -uniform hypergraphs. In section 3, we give the upper bounds of the sizes of  $(k, l)$ -edge-maximal  $r$ -uniform hypergraphs and illustrate that these bounds are best possible. We obtain the lower bounds of the sizes of  $(k, l)$ -edge-maximal  $r$ -uniform hypergraphs and show that these bounds are best possible in section 4.

## 2. PROPERTIES OF $(k, l)$ -EDGE-MAXIMAL $r$ -UNIFORM HYPERGRAPHS

Because of Theorem 1.5, we assume  $l \geq t + 1$  in this paper.

**Lemma 2.1.** *Let  $H = (V, E)$  be a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph such that  $n \geq l \geq t + 1$  and  $k, r \geq 2$ , where  $n = |V(H)|$  and  $t = t(k, r)$ . Assume  $X$  is a proper nonempty subset of  $V(H)$  such that  $\kappa'(H) = |E_H(X)|$ . Then each of the following holds.*

- (i)  $E_{H^c}(X) \neq \emptyset$ .
- (ii)  $\kappa'(H) = |E_H(X)| = k$ .

**Proof.** Let  $n_1 = |X|$  and  $n_2 = |V(H) \setminus X|$ . Then  $n = n_1 + n_2$ . Since  $H$  is  $(k, l)$ -edge-maximal, we have  $\kappa'(H) \leq k$ .

(i) Assume  $E_{H^c}(X) = \emptyset$ . Then  $E_H(X)$  consists of all  $r$ -subsets of  $V(H)$  intersecting both  $X$  and  $V(H) \setminus X$ . Thus

$$|E_H(X)| = \sum_{s=1}^{r-1} \binom{n_1}{s} \binom{n_2}{r-s} = \binom{n}{r} - \binom{n_1}{r} - \binom{n_2}{r}.$$

Let  $g(x) = \binom{x}{r} + \binom{n-x}{r}$ . It is routine to verify that  $g(x)$  is a decreasing function when  $1 \leq x \leq n/2$ . If  $\min\{n_1, n_2\} \geq 2$ , then by  $\min\{n_1, n_2\} \leq n/2$ , we have

$$(1) \quad \begin{aligned} \kappa'(H) = |E_H(X)| &= \binom{n}{r} - \binom{n_1}{r} - \binom{n_2}{r} \geq \binom{n}{r} - \binom{2}{r} - \binom{n-2}{r} > \binom{n-1}{r-1} \\ &\geq \delta(H), \end{aligned}$$

which contradicts to  $\kappa'(H) \leq \delta(H)$ . Now we assume  $\min\{n_1, n_2\} = 1$ .

Then

$$\begin{aligned} \kappa'(H) = |E_H(X)| &= \binom{n}{r} - \binom{n_1}{r} - \binom{n_2}{r} = \binom{n}{r} - \binom{1}{r} - \binom{n-1}{r} = \binom{n-1}{r-1} \\ &\geq \delta(H), \end{aligned}$$

which implies  $\kappa'(H) = \delta(H) = \binom{n-1}{r-1}$  and so  $H$  is a complete  $r$ -uniform hypergraph. Thus  $\kappa'(H) = \binom{n-1}{r-1} \geq \binom{l-1}{r-1} \geq \binom{t}{r-1} > k$ , contrary to  $\kappa'(H) \leq k$ . Therefore  $E_{H^c}(X) \neq \emptyset$  holds.

(ii) By (i), we have  $E_{H^c}(X) \neq \emptyset$ . Pick an edge  $e \in E_{H^c}(X)$ . Since  $H$  is  $(k, l)$ -edge-maximal, there is a subhypergraph  $H' \subseteq H + e$  such that  $|V(H')| \geq l$  and  $\kappa'(H') \geq k + 1$ . We have  $e \in H'$  by  $H$  is  $(k, l)$ -edge-maximal. It follows that  $(E_H(X) \cup \{e\}) \cap E(H')$  is an edge-cut of  $H'$ . Thus  $|E_H(X)| + 1 \geq |E_H(X) \cup \{e\}| \geq \kappa'(H') \geq k + 1$ , implying  $\kappa'(H) = |E_H(X)| \geq k$ . By  $\kappa'(H) \leq k$ , we obtain  $\kappa'(H) = |E_H(X)| = k$ . ■

**Lemma 2.2.** Let  $H = (V, E)$  be a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph such that  $n \geq l \geq t + 1$  and  $k, r \geq 2$ , where  $n = |V(H)|$  and  $t = t(k, r)$ . Assume  $X$  is a proper nonempty subset of  $V(H)$  such that  $|E_H(X)| = k$ . Then each of the following holds.

- (i) If  $|X| \leq r - 1$ , then  $H[X]$  contains no edges in  $E(H)$ , and each edge of  $E_H(X)$  contains  $X$  as a subset.
- (ii) If  $r \leq |X| \leq l - 1$ , then  $H[X]$  is a complete  $r$ -uniform hypergraph and  $|X| \geq t$ .
- (iii) If  $|X| \geq l$ , then  $H[X]$  is also a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph.

**Proof.** (i) Since  $H$  is an  $r$ -uniform hypergraph,  $H[X]$  contains no edges in  $E[H]$  if  $|X| \leq r - 1$ . By Lemma 2.1 and  $|E_H(X)| = k$ , we obtain  $k = |E_H(X)| \geq \delta(H) \geq \kappa'(H) = k$ , implying each edge of  $E_H(X)$  contains  $X$  as a subset.

(ii) Assume  $r \leq |X| \leq l - 1$ . If  $H[X]$  is not complete, then there is an edge  $e \in E(H[X]^c) \subseteq E(H^c)$  and so  $H + e$  has no subhypergraph  $H'$  with  $|V(H')| \geq l$  and  $\kappa'(H') \geq k + 1$ , contrary to the assumption that  $H$  is  $(k, l)$ -edge-maximal. Hence  $H[X]$  must be complete.

On the contrary, assume  $|X| < t$ . Since  $\delta(H) \geq \kappa'(H) = |E_H(X)| = k$  and  $\binom{t-1}{r-1} \leq k < \binom{t}{r-1}$ , in order to ensure each vertex in  $X$  has degree at least  $k$  in  $H$ , we must have  $|X| = t - 1$  and  $k = \binom{t-1}{r-1}$ . Moreover, each vertex in  $X$  is incident with exact  $\binom{t-2}{r-2}$  edges in  $E_H(X)$ , and thus  $d_H(u) = k$  for each  $u \in X$ . By Lemma 2.1(i), there is an  $e$  intersecting both  $X$  and  $V(H) \setminus X$  but  $e \notin E_H(X)$ . Since  $|X| \geq r$ , there is a vertex  $w \in X$  such that  $w$  is not incident with  $e$ . Then  $d_{H+e}(w) = k$ . This implies  $w$  is not contained in a  $(k + 1)$ -edge-connected subhypergraph of  $H + e$ . But then each vertex in  $X \setminus \{w\}$  has at most degree  $k$  in  $(H + e) - w$ , and thus each vertex in  $X \setminus \{w\}$  is not contained in a  $(k + 1)$ -edge-connected subhypergraph of  $H + e$ . This implies that there is no  $(k + 1)$ -edge-connected subhypergraph with at least  $l$  vertices in  $H + e$ , a contradiction. Thus we have  $|X| \geq t$ .

(iii) Assume  $|X| \geq l$ . If  $H[X]$  is complete, then  $\kappa'(H[X]) = \delta(H[X]) = \binom{|X|-1}{r-1} \geq \binom{l-1}{r-1} \geq \binom{t}{r-1} > k$ , contrary to the definition of  $(k, l)$ -edge-maximal hypergraph. Thus  $H[X]$  is not complete. For any edge  $e \in E(H[X]^c) \subseteq E(H^c)$ ,  $H + e$  has a subhypergraph  $H'$  with  $|V(H')| \geq l$  and  $\kappa'(H') \geq k + 1$ . Since  $|E_H(X)| = k$ , we have  $E_H(X) \cap E(H') = \emptyset$ . As  $e \in E(H') \cap E(H[X]^c)$ , we conclude that  $H'$  is a subhypergraph of  $H[X] + e$ , and so  $H[X]$  is a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph. ■

### 3. THE UPPER BOUNDS OF THE SIZES OF $(k, l)$ -EDGE-MAXIMAL $r$ -UNIFORM HYPERGRAPHS

We first extend the definition of star-like- $(k, l)$  graphs in [2] to hypergraphs.

**Definition 3.1.** Let  $k, l, r$  be integers such that  $k, r \geq 2$  and  $l \geq t + 1$ , where  $t = t(k, r)$ . Star-like- $(k, l)$   $r$ -uniform hypergraphs are defined constructively as follows. Start with a complete  $r$ -uniform hypergraph  $K_{l-1}^r$ . Call it the *nucleus*. Attach a single vertex  $K_1$  or a complete  $r$ -uniform hypergraph  $K_i^r$  ( $r \leq i \leq l - 1$ ) to this nucleus using  $k$  edges joining  $K_1$  or  $K_i^r$  to the nucleus. Call this attached hypergraph a *satellite*. Attach an arbitrary number of such satellites to the nucleus in the same manner. We call  $r$ -uniform hypergraphs constructed in this manner *star-like- $(k, l)$   $r$ -uniform hypergraphs*. We use  $SH(k, l, r)$  to denote the collection of all star-like- $(k, l)$   $r$ -uniform hypergraphs. See Figure 1 for example.

**Definition 3.2.** For integers  $k, r \geq 2$ , let  $s = s(k, r)$  be the largest integer such that  $k + \binom{s}{r} \leq ks$ .

**Remark 3.3.** Since  $k + \binom{t}{r} = k + \frac{t}{r} \binom{t-1}{r-1} \leq k + \frac{t}{r} k \leq kt$ , we have  $t \leq s$ , where  $t = t(k, r)$  and  $s = s(k, r)$ .




Figure 1. An example of a star-like- $(k, l)$   $r$ -uniform hypergraph.

**Definition 3.4.** Let  $n, k, l, r$  be integers such that  $k, r \geq 2$  and  $n \geq l \geq t + 1$ , where  $t = t(k, r)$ . Let  $p, q \geq 0$  be integers such that  $n = p(l - 1) + q$  with  $0 \leq q < l - 1$ . We construct a class of star-like- $(k, l)$   $r$ -uniform hypergraphs on  $n$  vertices as follows.

(i) If  $l - 1 > s$  (where  $s = s(k, r)$ ), then a star-like- $(k, l)$   $r$ -uniform hypergraph consists of  $p$  copies of  $K_{l-1}^r$ , one serving as the nucleus, the rest as satellites, together with addition satellites determined as follows: (i-a) if  $q > s$ , the single additional satellite is  $K_q^r$ ; (i-b) if  $q \leq s$ , the additional satellites are  $q$  copies of  $K_1$ , each attached to the nucleus by  $k$  edges.

(ii) If  $l - 1 \leq s$  (where  $s = s(k, r)$ ), then the nucleus is  $K_{l-1}^r$ . The satellites are  $n - (l - 1)$  copies of  $K_1$ , each attached to the nucleus by  $k$  edges.

We denote the collection of all star-like- $(k, l)$   $r$ -uniform hypergraphs on  $n$  vertices constructed in Definition 3.4 by  $MSH(n; k, l, r)$ . Note that all hypergraphs in  $MSH(n; k, l, r)$  have the same number of edges, denoted this number by  $|E(MSH(n; k, l, r))|$  for brevity. By definition, we have

$$|E(MSH(n; k, l, r))| = \begin{cases} p \binom{l-1}{r} + pk + \binom{q}{r}, & l - 1 > s \text{ and } q > s, \\ p \binom{l-1}{r} + (p - 1 + q)k, & l - 1 > s \text{ and } q \leq s, \\ \binom{l-1}{r} + (n - l + 1)k, & l - 1 \leq s. \end{cases}$$

The following theorem shows that  $MSH(n; k, l, r)$  is a class of star-like- $(k, l)$   $r$ -uniform hypergraphs with the maximum number of edges among all star-like- $(k, l)$   $r$ -uniform hypergraphs on  $n$  vertices.

**Theorem 3.5.** *Let  $n, k, l, r$  be integers such that  $k, r \geq 2$  and  $n \geq l \geq t + 1$ , where  $t = t(k, r)$ . Let  $p, q \geq 0$  be integers such that  $n = p(l - 1) + q$  with  $0 \leq q < l - 1$ . For each star-like- $(k, l)$   $r$ -uniform hypergraph  $H$  on  $n$  vertices, we have  $|E(H)| \leq |E(MSH(n; k, l, r))|$ .*

**Proof.** The idea of the proof is that we transform  $H$  into one of the hypergraph in  $MSH(n; k, l, r)$  by appropriate addition and deletion of edges. The edge transformations applied always yield an increase (not necessary net increase) in the number of edges. The proof uses two basic techniques: *splitting* and *grouping*. The splitting operation replaces a single  $K_i^r$  ( $i \geq r$ ) by  $i$   $K_1$ -satellites. The grouping operations move vertices from smaller to larger satellites (with corresponding edge additions and deletions), or cluster a set of  $K_1$ -satellites into a single large satellite. We define the *satellite spectrum* of a star-like- $(k, l)$  hypergraph  $H$  as  $(S_1, S_r, \dots, S_{l-1})$ , where  $S_i$ ,  $i \in \{1, r, \dots, l - 1\}$ , is the number of satellites of  $H$  with  $i$  vertices. We consider two cases.

*Case 1.*  $l - 1 > s$ , where  $s = s(k, r)$ . If  $H$  contains a  $K_i^r$ -satellite,  $r \leq i \leq s$ , then perform the *splitting operation*. That is, replace  $K_i^r$ -satellite (and the  $k$  edges connecting it to the nucleus) by  $i$   $K_1$ -satellites (together with the  $k$  edges that join each of them to the nucleus). The new hypergraph contains at least as many edges as  $H$ . The argument is as follows. The number of edges associated with the original  $K_i^r$ -satellite is  $k + \binom{i}{r}$ . On the other hand, the satellites introduced by the splitting operation contribute  $ik$  edges. Since  $i \leq s$ , we have  $k + \binom{i}{r} \leq ik$  by Definition 3.2. Thus the hypergraph produced by the splitting operation contains at least as many edges as  $H$ . If we repeat this process on any remaining  $K_i^r$ -satellites,  $r \leq i \leq s$ , we eventually obtain a transformed hypergraph with satellite spectrum satisfying  $S_i = 0$  for  $r \leq i \leq s$ .

If there are two satellites  $K_i^r$  and  $K_j^r$  satisfying  $s < i \leq j < l - 1$ , we perform the following grouping operation (call *Grouping operation 1*). Since  $i > s \geq t$  and  $\binom{i-1}{r-1} \geq \binom{t}{r-1} > k$ , we can assume that there is a vertex  $u$  in  $K_i^r$  not adjacent to the nucleus. Delete the edges connecting  $u$  to  $K_i^r$ . Add edges from  $u$  to  $K_j^r$  such that  $V(K_j^r) \cup \{u\}$  induces a complete  $r$ -uniform hypergraph. The operation is edge-increasing because while it removes  $\binom{i-1}{r-1}$  edges, it adds  $\binom{j}{r-1}$  edges. The  $(K_i^r, K_j^r)$  pair of satellites become a  $(K_{i-1}^r, K_{j+1}^r)$  pair. If  $i - 1 \leq s$ , then we apply the splitting operation to  $K_{i-1}^r$ . We repeat the grouping operation until at most one satellite remains in the range  $s < i < l - 1$ . Now the satellite spectrum of the resulting hypergraph is simple. Except  $S_1 \geq 0$ ,  $S_{l-1} \geq 0$ , and some  $S_{i_0}$  ( $s < i_0 < l - 1$ ) may be 1 or 0, all other entries must be zero.

If  $S_1$  and  $S_{i_0}$  are positive, then we apply a second grouping operation (call *Grouping operation 2*). Select one  $K_1$ -satellite. Delete the  $k$  edges connecting the  $K_1$ -satellite to the nucleus. Add  $\binom{i_0}{r-1}$  edges connecting the  $K_1$ -satellite to  $K_{i_0}^r$ . This transformation is edge-increasing since  $\binom{i_0}{r-1} > \binom{s}{r-1} \geq \binom{t}{r-1} > k$ . We



repeat this operation until the supply of  $K_1$ -satellites has been exhausted or the original  $K_{i_0}^r$  has been augmented to  $K_{l-1}^r$ -satellite.

If  $S_1 > s$ , we apply one further grouping operation (call *Grouping operation 3*). Let  $S_1 = p_1(l-1) + q_1$ ,  $0 \leq q_1 < l-1$ . Replace the  $S_1$   $K_1$ -satellites by  $p_1$   $K_{l-1}^r$ -satellites, and by either  $q_1$   $K_1$ -satellites or 1  $K_{q_1}$ -satellite, as  $q_1 \leq s$  or not. By Definition 3.2, this operation is edge increasing.

This completes the edge transformation of the original hypergraph in Case 1. For the resultant hypergraph, either  $S_{i_0} = 1$  for some  $s < i_0 < l-1$  and  $S_1 = 0$ , or all  $S_i$  for  $r \leq i \leq l-2$  are zero, which correspond to the case (i-a) or the case (i-b) in Definition 3.4.

*Case 2.*  $l-1 \leq s$ , where  $s = s(k, r)$ . We apply splitting operation to all satellites. By Definition 3.2, the resultant hypergraph, which correspond to the case (ii) in Definition 3.4, is edge increasing.

This completes the proof of the theorem.  $\blacksquare$

**Theorem 3.6.** *Let  $n, k, l, r$  be integers such that  $k, r \geq 2$  and  $n \geq l \geq t+1$ , where  $t = t(k, r)$ . Let  $p, q \geq 0$  be integers such that  $n = p(l-1) + q$  with  $0 \leq q < l-1$ . If  $H$  is a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph on  $n$  vertices, then*

$$|E(H)| \leq |E(MSH(n; k, l, r))| = \begin{cases} p \binom{l-1}{r} + pk + \binom{q}{r}, & l-1 > s \text{ and } q > s, \\ p \binom{l-1}{r} + (p-1+q)k, & l-1 > s \text{ and } q \leq s, \\ \binom{l-1}{r} + (n-l+1)k, & l-1 \leq s, \end{cases}$$

where  $s = s(k, r)$ .

**Proof.** By Theorem 3.5, we only need to prove that there is a star-like- $(k, l)$   $r$ -uniform hypergraph  $H'$  on  $n$  vertices such that  $|E(H)| \leq |E(H')|$ . The proof is by induction on  $n$ .

If  $n = l$ , let  $H'$  be a star-like- $(k, l)$   $r$ -uniform hypergraph with the nucleus  $K_{l-1}^r$  and a single  $K_1$ -satellite. Since  $\kappa'(K_l^r) = \binom{l-1}{r-1}$ , we need to delete at least  $\binom{l-1}{r-1} - k$  edges such that the remaining hypergraph have edge-connectivity at most  $k$ . Since  $\kappa'(H) = k$  (by Lemma 2.1), we have  $|E(K_l^r)| - |E(H)| \geq \binom{l-1}{r-1} - k$ , implying  $|E(H)| \leq |E(H')|$ .

Now we assume  $n > l$ , and assume that for any  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph with less than  $n$  vertices, there is a star-like- $(k, l)$   $r$ -uniform hypergraph having the same number of vertices and at least as many edges as the given hypergraph.

Let  $F$  be a minimum edge-cut  $H$ . By Lemma 2.1, we have  $|F| = k$ . We consider two cases in the following.

*Case 1.* There is a component, say  $H_1$ , of  $H - F$  such that  $|V(H_1)| = 1$ . Let  $H_2 = H - V(H_1)$ . Then  $|V(H_2)| = n-1 \geq l$ . By Lemma 2.2(iii),  $H_2$  is

$(k, l)$ -edge-maximal. By induction assumption, there is a star-like- $(k, l)$   $r$ -uniform hypergraph, say  $H'_2$ , such that  $|V(H'_2)| = |V(H_2)|$  and  $|E(H'_2)| \geq |E(H_2)|$ . Let  $H'$  be the star-like- $(k, l)$   $r$ -uniform hypergraph obtained from  $H'_2$  by adding a  $K_1$ -satellite. Since  $|E(H)| = k + |E(H_2)|$ , we have  $|E(H)| \leq |E(H')|$ .

*Case 2.* Each component of  $H - F$  has at least two vertices. Then, by Lemma 2.2, each component of  $H - F$  is either a complete  $r$ -uniform hypergraph with at least  $t$  vertices, or a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph with at least  $l$  vertices.

Let  $H_1$  be a component of  $H - F$  and  $H_2 = H - V(H_1)$ . Assume  $n_1 = |V(H_1)|$  and  $n_2 = |V(H_2)|$ . Then  $n_1 + n_2 = n$ .

*Subcase 2.1.*  $n_1 \geq l$  and  $n_2 \geq l$ . By Lemma 2.2, both  $H_1$  and  $H_2$  are  $(k, l)$ -edge-maximal  $r$ -uniform hypergraphs. By induction assumption, there are two star-like- $(k, l)$   $r$ -uniform hypergraphs, say  $H'_1$  and  $H'_2$ , such that  $|V(H'_i)| = |V(H_i)|$  and  $|E(H'_i)| \geq |E(H_i)|$  for  $i = 1, 2$ . Let  $H'$  be a star-like- $(k, l)$   $r$ -uniform hypergraph obtained from  $H'_1$  and  $H'_2$  by moving the satellites of  $H'_1$  to  $H'_2$  and changing the nucleus of  $H'_1$  to be a satellite of  $H'_2$ . Then  $|E(H')| = |E(H'_1)| + |E(H'_2)| + k$ , and thus  $|E(H)| \leq |E(H')|$  holds.

*Subcase 2.2.*  $n_1 \geq l$  and  $n_2 < l$ . By Lemma 2.2,  $H_1$  is a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph and  $H_2$  is a complete  $r$ -uniform hypergraph with at least  $t$  vertices. By induction assumption, there is star-like- $(k, l)$   $r$ -uniform hypergraphs, say  $H'_1$ , such that  $|V(H'_1)| = |V(H_1)|$  and  $|E(H'_1)| \geq |E(H_1)|$ . Let  $H'$  be a star-like- $(k, l)$   $r$ -uniform hypergraph obtained from  $H'_1$  by adding  $H_2$  to be a satellite of  $H_1$ . Then  $|E(H')| = |E(H'_1)| + |E(H_2)| + k$ , and thus  $|E(H)| \leq |E(H')|$  holds.

*Subcase 2.3.*  $n_1 < l$  and  $n_2 < l$ . By Lemma 2.2, both  $H_1$  and  $H_2$  are complete  $r$ -uniform hypergraphs with at least  $t$  vertices. Since  $|E(H)| = \binom{n_1}{r} + \binom{n_2}{r} + k \leq \binom{l-1}{r} + \binom{n-l+1}{r} + k$ , we obtain that each star-like- $(k, l)$   $r$ -uniform hypergraph on  $n$  vertices having at least as many edges as  $H$  in this case. Therefore, the proof of this case follows.

This completes the proof of Theorem 3.6. ■

In the following theorem, we will show that each hypergraph in  $MSH(n; k, l, r)$  is  $(k, l)$ -edge-maximal. So the upper bounds given in Theorem 3.6 are best possible.

**Theorem 3.7.** *Let  $n, k, l, r$  be integers such that  $k, r \geq 2$  and  $n \geq l \geq t + 1$ , where  $t = t(k, r)$ . If  $H \in MSH(n; k, l, r)$ , then  $H$  is  $(k, l)$ -edge-maximal.*

**Proof.** If  $l - 1 = t$ , then all satellites of  $H$  are  $K_1$ -satellites by  $s \geq t$ , where  $s = s(k, r)$ . By Lemma 3.1 in [9],  $H$  is  $(k, l)$ -edge-maximal. Thus, in the following, we assume  $l - 1 > t$ .

By definition, there is no subhypergraph  $H'$  of  $H$  such that  $|V(H')| \geq l$  and  $\kappa'(H') > k$ . We will prove the theorem by induction on  $n$ . If  $n = l$ , then  $H$  is a star-like- $(k, l)$   $r$ -uniform hypergraph with the nucleus  $K_{l-1}^r$  and a  $K_1$ -satellite. Since  $\kappa'(K_{l-1}^r) = \binom{l-2}{r-1} \geq \binom{t}{r-1} > k$ , for any  $e \in E(H^c)$ , we have  $\kappa'(H + e) > k$ . Thus  $H$  is  $(k, l)$ -edge-maximal.

Now suppose  $n > l$ . We assume that each hypergraph in  $MSH(n'; k, l, r)$ , where  $n' < n$ , is  $(k, l)$ -edge-maximal. In the following, we will show that each  $H$  in  $MSH(n; k, l, r)$  is also  $(k, l)$ -edge-maximal.

By contradiction, assume that there is an edge  $e \in E(H^c)$  such that  $H + e$  contains no subhypergraph  $H'$  satisfying  $|V(H')| \geq l$  and  $\kappa'(H') > k$ . Let  $F$  be an edge-cut in  $H + e$  with cardinality at most  $k$ . Since  $\kappa'(K_{l-1}^r) = \binom{l-2}{r-1} \geq \binom{t}{r-1} > k$  and  $\kappa'(K_q^r) = \binom{q-1}{r-1} \geq \binom{s}{r-1} \geq \binom{t}{r-1} > k$  when  $q > s$ , we obtain that  $F$  is exact the edge-cut joining some satellite and the nucleus. Thus there is a component, say  $H_1$ , of  $H - F$ , such that  $H_1$  is the hypergraph obtained from  $H$  by deleting one satellite and  $e \in H_1^c$ . By induction assumption,  $H_1 + e$  contains a subhypergraph  $H'_1$  such that  $|V(H'_1)| \geq l$  and  $\kappa'(H'_1) > k$ . But  $H'_1$  is also a subhypergraph of  $H + e$ , a contradiction. ■

#### 4. THE LOWER BOUNDS OF THE SIZES OF $(k, l)$ -EDGE-MAXIMAL $r$ -UNIFORM HYPERGRAPHS

The following lemma will be needed in proving the main result in this section.

**Lemma 4.1.** *Let  $n, a, k, r$  be integers such that  $k, r \geq 2$  and  $n \geq a \geq t$ , where  $t = t(k, r)$ . We have the following two inequalities.*

- (i)  $\binom{n}{r} \geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ .
- (ii)  $\binom{n}{r} \geq (n-a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-a}{t} \rfloor$ .

**Proof.** Since  $|E(K_n^r)| = \binom{n}{r}$ , the lemma will hold if we can construct two  $r$ -uniform hypergraphs  $H$  and  $H'$  on  $n$  vertices such that  $|E(H)| = (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$  and  $|E(H')| = (n-a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-a}{t} \rfloor$ .

Let  $H$  be a  $r$ -uniform star-like hypergraph with the nucleus  $K_t^r$ ,  $\lfloor \frac{n}{t} \rfloor - 1$   $K_t^r$ -satellites and  $n - t \lfloor \frac{n}{t} \rfloor$   $K_1$ -satellites, adding  $k$  edges joining each satellite to the nucleus. It is routine to count that  $|E(H)| = (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ .

Let  $H'$  be a  $r$ -uniform star-like hypergraph with the nucleus  $K_a^r$ ,  $\lfloor \frac{n-a}{t} \rfloor$   $K_t^r$ -satellites and  $n - a - t \lfloor \frac{n-a}{t} \rfloor$   $K_1$ -satellites, adding  $k$  edges joining each satellite to the nucleus. Then  $|E(H')| = (n-a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-a}{t} \rfloor$ . ■

It is routine to verify the following lemma.

**Lemma 4.2.** *For given integers  $n$  and  $r$  with  $n \geq r \geq 2$ , the function  $g(x) = \binom{x}{r} + \binom{n-x}{r}$  is decreasing in the range  $1 \leq x \leq n/2$ .*

**Theorem 4.3.** *Let  $n, k, l, r$  be integers such that  $k, r \geq 2$  and  $n \geq l \geq t+1$ , where  $t = t(k, r)$ . If  $H$  is a  $(k, l)$ -edge-maximal  $r$ -uniform hypergraph on  $n$  vertices, then*

$$|E(H)| \geq \begin{cases} \binom{l-1}{r} + (n-l+1)k, & l \leq n < 2t, \\ (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor, & l \leq 2t \leq n, \\ (n-2a+1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor, & n \geq l = 2a \geq 2t+1, \\ (n-2b)k + \binom{b}{r} + \binom{b+1}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2b-1}{t} \rfloor, & n \geq l = 2b+1 \geq 2t+1. \end{cases}$$

**Proof.** Let  $F$  be a minimum edge-cut of  $H$ . By Lemma 2.1,  $|F| = k$ . Assume  $H_1$  is a minimum component of  $H - F$  and  $H_2 = H - V(H_1)$ . Let  $n_1 = |V(H_1)|$  and  $n_2 = |V(H_2)|$ . Then  $n = n_1 + n_2$  and  $n_1 \leq n_2$ .

(i) For  $l \leq n < 2t$ , we have  $n_1 < t$ , and then  $n_1 = 1$  by Lemma 2.2. If  $n = l$ , then, by Lemma 2.2,  $H_2$  is a complete  $r$ -uniform hypergraph on  $l-1$  vertices. Thus  $|E(H)| = k + \binom{l-1}{r} = \binom{l-1}{r} + (n-l+1)k$ . If  $n > l$ , by Lemma 2.2,  $H_2$  is  $(k, l)$ -edge-maximal. By induction on  $n$ , assume  $|E(H_2)| \geq \binom{l-1}{r} + (n_2-l+1)k$ . Therefore,  $|E(H)| = |F| + |E(H_2)| \geq k + \binom{l-1}{r} + (n_2-l+1)k = \binom{l-1}{r} + (n-l+1)k$ .

(ii) We now assume  $l \leq 2t \leq n$ . We shall prove this case by induction on  $n$ .

If  $n = 2t$ , then either  $n_1 = n_2 = t$ , or  $n_1 = 1$  and  $n_2 = n-1$  by Lemma 2.2. When  $n_1 = n_2 = t$ , then  $H_1$  and  $H_2$  are complete, and thus  $|E(H)| = \binom{t}{r} + \binom{t}{r} + k = (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ . Assume  $n_1 = 1$  and  $n_2 = n-1$ . If  $n = l$ , then, by Lemma 2.2,  $H_2$  is a complete  $r$ -uniform hypergraph. By Lemma 4.2,  $\binom{n-1}{r} = \binom{1}{r} + \binom{n-1}{r} \geq \binom{t}{r} + \binom{t}{r}$ . Thus  $|E(H)| = k + \binom{n-1}{r} \geq k + 2\binom{t}{r} = (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ . If  $n > l$ , by Lemma 2.2,  $H_2$  is  $(k, l)$ -edge-maximal. Assume, by induction hypothesis,  $|E(H_2)| \geq (n_2-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2}{t} \rfloor$ . Therefore,  $|E(H)| = |E(H_2)| + k \geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-1}{t} \rfloor \geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ , the last inequality holds because  $(t-1)k - \binom{t}{r} \geq (t-1)\binom{t-1}{r-1} - \frac{t}{r}\binom{t-1}{r-1} \geq 0$ .

Assume that  $n > 2t$ . If  $n_1 = 1$ , then  $n_2 = n-1 \geq 2t$ . By induction assumption,  $|E(H_2)| \geq (n_2-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2}{t} \rfloor$ . Thus  $|E(H)| = |E(H_2)| + k \geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-1}{t} \rfloor \geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor$ . So we assume  $n_1 \geq t$ .

**Claim.**  $|E(H_i)| \geq (n_i-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_i}{t} \rfloor$  for  $i \in \{1, 2\}$ .

If  $|V(H_i)| \geq l$ , then by induction assumption, we have  $|E(H_i)| \geq (n_i-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_i}{t} \rfloor$ . If  $t \leq |V(H_i)| \leq l-1$ , then by Lemma 2.2,  $H_i$  is a complete  $r$ -uniform hypergraph. Thus, by Lemma 4.1(i),  $|E(H_i)| = \binom{n_i}{r} \geq (n_i-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_i}{t} \rfloor$ .

By this claim, we have

$$\begin{aligned} |E(H)| &= |E(H_1)| + |E(H_2)| + k \\ &\geq (n_1-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1}{t} \rfloor + (n_2-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2}{t} \rfloor + k \\ &= (n-1)k - ((t-1)k - \binom{t}{r}) (\lfloor \frac{n_1}{t} \rfloor + \lfloor \frac{n_2}{t} \rfloor) \end{aligned}$$

$$\begin{aligned} &\geq (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1+n_2}{t} \rfloor \quad (\text{By } (t-1)k - \binom{t}{r} \geq 0) \\ &= (n-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n}{t} \rfloor. \end{aligned}$$

(iii) We then assume  $n \geq l = 2a \geq 2t + 1$ . By induction on  $n$ , we will prove this case.

If  $n = l$ , then either  $n_1 = 1, n_2 = n - 1$  or  $t \leq n_1 \leq n_2 \leq n - t$ . When  $n_1 = 1$  and  $n_2 = n - 1 = l - 1$ , then  $H_2$  is complete by Lemma 2.2. Since  $\binom{l-1}{r} = \binom{1}{r} + \binom{l-1}{r} = \binom{1}{r} + \binom{2a-1}{r} \geq \binom{a}{r} + \binom{a}{r}$  (by Lemma 4.2), we have  $|E(H)| = |E(H_2)| + k = k + \binom{n-1}{r} \geq k + 2\binom{a}{r} = (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor$ . When  $t \leq n_1 \leq n_2 \leq l - t$ , by Lemma 2.2, both  $H_1$  and  $H_2$  are complete. Since  $n_1 \leq n/2 = a$ , we have  $\binom{n_1}{r} + \binom{n_2}{r} \geq \binom{a}{r} + \binom{a}{r}$  by Lemma 4.2. Thus  $|E(H)| = |E(H_1)| + |E(H_2)| + k = \binom{n_1}{r} + \binom{n_2}{r} + k \geq k + 2\binom{a}{r} = (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor$ .

Thus we assume  $n > l$ . If  $n_1 = 1$ , then  $n_2 = n - 1 \geq l$ . By induction assumption,  $|E(H_2)| \geq (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor$ . Thus  $|E(H)| = |E(H_2)| + k \geq (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor + k \geq (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor$ . So we assume  $n_1 \geq t$  and consider three cases in the following.

*Case 1.*  $n_1 \geq l$ . By induction assumption, we have  $|E(H_i)| \geq (n_i - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_i-2a}{t} \rfloor$  for  $i = 1, 2$ . By setting  $n$  to be  $a$  in Lemma 4.1(i), we have  $\binom{a}{r} \geq (a-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{a}{t} \rfloor$ . Thus

$$\begin{aligned} |E(H)| &= |E(H_1)| + |E(H_2)| + k \\ &\geq (n_1 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1-2a}{t} \rfloor \\ &\quad + (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor + k \\ &\geq (n_1 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1-2a}{t} \rfloor + (n_2 - 2a + 1)k \\ &\quad + 2((a-1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{a}{t} \rfloor) - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor + k \\ &\geq (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor. \end{aligned}$$

*Case 2.*  $n_1 < l$  and  $n_2 \geq l$ . By  $n_1 < l$  and  $n_2 \geq l$ , we have  $H_1$  is complete and  $H_1$  is  $k$ -edge-maximal by Lemma 2.2. By induction assumption,  $|E(H_2)| \geq (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor$ . Setting  $n$  to be  $n_1$  in Lemma 4.1(i), we have  $\binom{n_1}{r} \geq (n_1 - 1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1}{t} \rfloor$  by  $n_1 \geq t$ . Thus

$$\begin{aligned} |E(H)| &= |E(H_1)| + |E(H_2)| + k \\ &\geq \binom{n_1}{r} + (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor + k \\ &\geq (n_1 - 1)k - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1}{t} \rfloor \\ &\quad + (n_2 - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2-2a}{t} \rfloor + k \\ &\geq (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n-2a}{t} \rfloor. \end{aligned}$$

*Case 3.*  $n_1 \leq n_2 < l$ . By  $n_1 \leq n_2 < l$ , we obtain that both  $H_1$  and  $H_2$  are complete by Lemma 2.2. If  $n_1 \geq a$ , then by setting  $n$  to be  $n_i$  in Lemma 4.1(ii), we have  $\binom{n_i}{r} \geq (n_i - a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_i - a}{t} \rfloor$  for  $i = 1, 2$ . Thus

$$\begin{aligned} |E(H)| &= |E(H_1)| + |E(H_2)| + k = \binom{n_1}{r} + \binom{n_2}{r} + k \\ &\geq (n_1 - a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1 - a}{t} \rfloor \\ &\quad + (n_2 - a)k + \binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_2 - a}{t} \rfloor + k \\ &\geq (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n - 2a}{t} \rfloor. \end{aligned}$$

If  $n_1 \leq a$ , then  $n_2 = n - n_1 \geq 2a - n_1 \geq t$ . By setting  $n$  to be  $n_2$  and  $a$  to be  $2a - n_1$  in Lemma 4.1(ii), we have  $\binom{n_2}{r} \geq (n_1 + n_2 - 2a)k + \binom{2a - n_1}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1 + n_2 - 2a}{t} \rfloor$ . Together with  $\binom{n_1}{r} + \binom{2a - n_1}{r} \geq 2\binom{a}{r}$  by Lemma 4.2, we have

$$\begin{aligned} |E(H)| &= |E(H_1)| + |E(H_2)| + k = \binom{n_1}{r} + \binom{n_2}{r} + k \\ &\geq \binom{n_1}{r} + (n_1 + n_2 - 2a)k + \binom{2a - n_1}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n_1 + n_2 - 2a}{t} \rfloor + k \\ &\geq (n - 2a + 1)k + 2\binom{a}{r} - ((t-1)k - \binom{t}{r}) \lfloor \frac{n - 2a}{t} \rfloor. \end{aligned}$$

Therefore, the proof for (iii) of Theorem 4.3 is complete.

(iv) The proof for (iv) of Theorem 4.3 is similar to that (iii) of Theorem 4.3, thus we omit the proof here.  $\blacksquare$

**Remark 4.4.** For  $l \leq n < 2t$ , the star-like- $(k, l)$  hypergraphs in  $MSH(n; k, l, r)$  show that the bound given in Theorem 4.3(i) is best possible. Hypergraphs constructed in Definition 3 [9] illustrate that the bound given in Theorem 4.3(ii) is best possible. The following example will show that bounds given in Theorem 4.3(iii) and (iv) are also best possible.

**Definition 4.5.** Let  $k, t, r$  be integers such that  $t > r > 2$ ,  $k = \binom{t-1}{r-1}$  and  $kr \geq 2t$ . Assume  $n$  and  $l$  are integers satisfying  $n = l + pt$  and  $l \geq 2t + 2$ , where  $p \geq 0$ . Let  $a = \lceil \frac{l}{2} \rceil$  and  $b = \lfloor \frac{l}{2} \rfloor$ . Then  $a, b \geq t + 1$ .

Let  $H_0$  be an  $r$ -uniform hypergraph obtained from the disjoint union of  $K_a^r$  and  $K_b^r$  by adding  $k$  edges joining  $K_a^r$  and  $K_b^r$ . Let  $H$  be a star-like  $r$ -uniform hypergraph with the nucleus  $H_0$  and  $p$   $K_t^r$ -satellites, adding  $k$  edges from each satellite to the nucleus such that (i) each vertex in the satellite adjacent to some added edge (we can do this by  $kr \geq 2t$ ); (ii) not all of the added  $k$  edges are incident with the same complete subhypergraph  $K_a^r$  or  $K_b^r$ .

**Theorem 4.6.** *Let  $H$  be an  $r$ -uniform hypergraph constructed in Definition 4.5. Then  $H$  is  $(k, l)$ -edge-maximal.*

**Proof.** By definition, there is no subhypergraph  $H'$  in  $H$  such that  $|V(H')| \geq l$  and  $\kappa'(H') \geq k + 1$ . We will prove the theorem by induction on  $p$ . If  $p = 0$ , then  $|V(H)| = l$ . Since  $\kappa'(K_a^r) = \binom{a-1}{r-1} \geq \binom{t}{r-1} > k$  and  $\kappa'(K_b^r) = \binom{b-1}{r-1} \geq \binom{t}{r-1} > k$ ,

the only edge-cut with  $k$  edges of  $H$  is these edges connecting  $K_a^r$  and  $K_b^r$ . For any  $e \in E(H^c)$ , we have  $e \in E_{H^c}[V(K_a^r), V(K_b^r)]$ . Thus every edge-cut of  $H + e$  has cardinality at least  $k+1$ , that is,  $\kappa'(H+e) \geq k+1$ . Thus  $H$  is  $(k, l)$ -edge-maximal.

Now suppose  $p \geq 1$ . We assume that each hypergraph constructed in Definition 4.5 with less than  $l + pt$  vertices is  $(k, l)$ -edge-maximal. In the following, we will show that each  $H$  in Definition 4.5 with  $l + pt$  vertices is also  $(k, l)$ -edge-maximal.

On the contrary, assume that there is an edge  $e \in E(H^c)$  such that  $H + e$  contains no subhypergraph  $H'$  such that  $|V(H')| \geq l$  and  $\kappa'(H') \geq k+1$ . Let  $F$  be an edge-cut in  $H + e$  with cardinality at most  $k$ . By Definition 4.5(i), we have  $\delta(H) \geq k+1$ . By (1) in the proof of Lemma 2.1 and Definition 4.5(ii), we obtain that edge-cuts in  $H + e$  with cardinality at most  $k$  are these  $k$  edges joining one satellite to the nucleus. Thus there is a component, say  $H_1$ , of  $H - F$ , such that  $H_1$  is the hypergraph obtained from  $H$  by deleting one satellite and  $e \in H_1^c$ . By induction assumption,  $H_1 + e$  contains a subhypergraph  $H'_1$  such that  $|V(H'_1)| \geq l$  and  $\kappa'(H'_1) > k$ . But  $H'_1$  is also a subhypergraph of  $H + e$ , a contradiction. ■

### Acknowledgments

The research of Yingzhi Tian is partially supported by National Natural Science Foundation of China (No.11861066) and Tianshan Youth Project of Xinjiang (2018Q066). The research of Hong-Jian Lai is partially supported by National Natural Science Foundation of China (Nos. 11771039 and 11771443). The research of Jixiang Meng is partially supported by National Natural Science Foundation of China (No. 11531011).

### REFERENCES

- [1] M.A. Bahmanian and M. Šajna, *Connection and separation in hypergraphs*, Theory Appl. Graphs **2** (2015) Article 5.  
<https://doi.org/10.20429/tag.2015.020205>
- [2] F.T. Boesch and J.A.M. McHuge, *An edge extremal result for subcohesion*, J. Combin. Theory Ser. B **38** (1985) 1–7.  
[https://doi.org/10.1016/0095-8956\(85\)90087-5](https://doi.org/10.1016/0095-8956(85)90087-5)
- [3] J.A. Bondy and U.S.R. Murty, *Graph Theory*, Grad. Texts in Math. **244** (Springer, Berlin, 2008).
- [4] M. Dewar, D. Pike and J. Proos, *Connectivity in hypergraphs*, Canad. Math. Bull. **61** (2018) 252–271.  
<https://doi.org/10.4153/CMB-2018-005-9>
- [5] H.-J. Lai, *The size of strength-maximal graphs*, J. Graph Theory **14** (1990) 187–197.  
<https://doi.org/10.1002/jgt.3190140207>

- [6] H.-J. Lai and C.-Z. Zhang, *Edge-maximal  $(k, i)$ -graphs*, J. Graph Theory **18** (1994) 227–240.  
<https://doi.org/10.1002/jgt.3190180303>
- [7] W. Mader, *Minimale  $n$ -fach kantenzusammenhängende Graphen*, Math. Ann. **191** (1971) 21–28.  
<https://doi.org/10.1007/BF01433466>
- [8] D.W. Matula,  *$k$ -components, clusters, and slicings in graphs*, SIAM J. Appl. Math. **22** (1972) 459–480.  
<https://doi.org/10.1137/0122040>
- [9] Y.Z. Tian, L.Q. Xu, H.-J. Lai and J.X. Meng, *On the sizes of  $k$ -edge-maximal  $r$ -uniform hypergraphs* (2018).  
arXiv:1802.08843v3

Received 27 December 2019

Revised 19 August 2020

Accepted 19 August 2020