

THE GRAPH GRABBING GAME ON BLOW-UPS OF TREES AND CYCLES

SOPON BORIBOON AND TEERADEJ KITTIPASSORN

*Department of Mathematics and Computer Science
Faculty of Science, Chulalongkorn University
Bangkok 10330, Thailand*

e-mail: soponboriboon@gmail.com
teeradej.k@chula.ac.th

Abstract

The graph grabbing game is played on a non-negatively weighted connected graph by Alice and Bob who alternately claim a non-cut vertex from the remaining graph, where Alice plays first, to maximize the weights on their respective claimed vertices at the end of the game when all vertices have been claimed. Seacrest and Seacrest conjectured that Alice can secure at least half of the total weight of every weighted connected bipartite even graph. Later, Egawa, Enomoto and Matsumoto partially confirmed this conjecture by showing that Alice wins the game on a class of weighted connected bipartite even graphs called $K_{m,n}$ -trees. We extend the result on this class to include a number of graphs, e.g. even blow-ups of trees and cycles.

Keywords: games on graphs, two-player games, graph grabbing games, blow-ups of graphs.

2020 Mathematics Subject Classification: 05C57.

1. INTRODUCTION

A vertex v of a connected graph G is a *cut vertex* if $G - v$ is disconnected. A graph G is *even* (respectively, *odd*) if the number of vertices of G is even (respectively, odd). A *weighted graph* G is a graph G with a weighted function $w : V(G) \rightarrow \mathbb{R}^+ \cup \{0\}$.

The *graph grabbing game* is played on a non-negatively weighted connected graph by two players: Alice and Bob alternately claim a non-cut vertex from the remaining graph and collect the weight on the vertex, where Alice plays first. The

aim of each player is to maximize the weights on their respective claimed vertices at the end of the game when all vertices have been claimed. Alice *wins* the game if she gains at least half of the total weight of the graph.

The first version of the graph grabbing game appeared in the first problem in Winkler's puzzle book (2003) [12], where he gave a winning strategy for Alice on every weighted even path and he observed that there is a weighted odd path on which Alice cannot win. In 2009, Rosenfeld [10] proposed the game for trees and call it the *gold grabbing game*. In 2011, Micek and Walczak [8] generalized the game to general graphs and call it the graph grabbing game. They showed that Alice can secure at least a quarter of the total weight of every weighted even tree and they conjectured that Alice can in fact secure at least half of the total weight of every weighted even tree. Later in 2012, Seacrest and Seacrest [11] solved this conjecture by considering a vertex-rooted version of the game and they posed the following conjecture.

Conjecture 1 [11]. *Alice wins the game on every weighted connected bipartite even graph.*

In 2018, Egawa, Enomoto and Matsumoto [3] gave a supporting evidence for this conjecture. They generalized the proof of Seacrest and Seacrest by considering a set-rooted version of the game to prove that Alice wins the game on every weighted even $K_{m,n}$ -tree, namely a bipartite graph obtained from a complete bipartite graph $K_{m,n}$ on $[m+n]$ and trees T_1, \dots, T_{m+n} by identifying vertex i of $K_{m,n}$ with exactly one vertex of T_i for each $i \in [m+n]$, where $[k]$ means the set of the natural numbers from one to k .

For a graph G with vertices v_1, \dots, v_k and non-empty sets V_1, \dots, V_k , a *blow-up* $B(G)$ of G is a graph obtained from G by replacing v_1, \dots, v_k with V_1, \dots, V_k , respectively, where, for each $i, j \in [k]$, vertices $x \in V_i$ and $y \in V_j$ are adjacent in $B(G)$ if and only if v_i and v_j are adjacent in G . For a graph G on $[k]$ and trees T_1, \dots, T_k , a G -tree is a graph obtained from G by identifying vertex i of G with exactly one vertex of T_i for each $i \in [k]$. For a tree T , we note that a $B(T)$ -tree and $B(C_{2n})$ are connected bipartite graphs, and a $B(T)$ -tree is a $K_{m,n}$ -tree when T is the path on two vertices, (see Figure 1).

In this paper, we partially confirm Conjecture 1 as follows.

Theorem 2. *Alice wins the game on every weighted even $B(T)$ -tree, where T is a tree.*

Corollary 3. *Alice wins the game on every weighted even $B(C_n)$.*

For a graph G and a set $S \subseteq V(G)$, let $N_G(S)$ denote the neighborhood of S , i.e., the set of vertices having a neighbor in S . The proof is based on the method of Egawa, Enomoto and Matsumoto, where their main lemmas dealt with the

score of the game on a $K_{m,n}$ -tree rooted at a partite class. We generalize their method by considering instead the scores of the game on an H -tree rooted at V_i and the game on the H -tree rooted at $N_H(V_i)$, where H is a blow-up of a tree.




Figure 1. Examples of a tree T , a blow-up $B(T)$ and a $B(T)$ -tree.

The rest of this paper is organized as follows. In Section 2, we recall some observations and a lemma on $K_{m,n}$ -trees given by Egawa, Enomoto and Matsumoto. Section 3 is devoted to proving Theorem 2 and then applying it to prove Corollary 3. In Section 4, we give some concluding remarks.

2. PRELIMINARIES

In this section, we prepare some observations and a lemma on $K_{m,n}$ -trees which will be useful for the proof of Theorem 2.

We first give definitions of a rooted version of the graph grabbing game and some related terms introduced by Egawa, Enomoto and Matsumoto. For a weighted graph G , a *root set* S of G is a set of vertices intersecting every component of G and the *game on G rooted at S* is a graph grabbing game, where each player does not have to claim a non-cut vertex, but instead they claim a vertex v such that every component of $G - v$ contains at least one vertex in S . Therefore, a move v in the game on G is *feasible* if $G - v$ is connected, and a move v in the game on G rooted at S is *feasible* if every component of $G - v$ contains at least one vertex in S . A move v in the game on G (rooted at S) is *optimal* if there is an optimal strategy in the game on G (rooted at S) having v as the first move. The first (respectively, second) player is called *Player 1* (respectively, *Player 2*). The last (respectively, second from last) player is called *Player -1* (respectively, *Player -2*). For $k \in \{1, 2, -1, -2\}$, assuming that both players play optimally, let $N(G, k)$ denote the score of Player k in the game on G and let $R(G, S, k)$ denote the score of Player k in the game on G rooted at S and we write $R(G, v, k)$ for $R(G, \{v\}, k)$. For a set S and an element x , we write $S - x$ for $S \setminus \{x\}$.

Egawa, Enomoto and Matsumoto observed some relationships between the scores of both players in the normal version and the rooted version of the game. Note that the equation/inequality in the brackets in each observation is an equivalent form of the first one because of the fact that, assuming that both players play optimally, the sum of their scores equals the total weight of the graph.

Observation 4 [3]. *If x is a feasible move in the game on G , then*

$$N(G, 2) \leq N(G - x, 1) \quad (\Leftrightarrow N(G, 1) \geq N(G - x, 2) + w(x)).$$

If x is an optimal move in the game on G , then

$$N(G, 2) = N(G - x, 1) \quad (\Leftrightarrow N(G, 1) = N(G - x, 2) + w(x)).$$

Observation 5 [3]. *Let S be a root set of G . If x is a feasible move in the game on G rooted at S , then*

$$R(G, S, 2) \leq R(G - x, S - x, 1) \quad (\Leftrightarrow R(G, S, 1) \geq R(G - x, S - x, 2) + w(x)).$$

If x is an optimal move in the game on G rooted at S , then

$$R(G, S, 2) = R(G - x, S - x, 1) \quad (\Leftrightarrow R(G, S, 1) = R(G - x, S - x, 2) + w(x)).$$

Observation 6 [3]. *If v is a root of G , then*

$$\begin{aligned} R(G, v, -2) &= R(G - v, N_G(v), -1) \\ (\Leftrightarrow R(G, v, -1) &= R(G - v, N_G(v), -2) + w(v)). \end{aligned}$$

The next lemma is a part of their main results which will help us in the proof.

Lemma 7 [3]. *Let G be a $K_{m,n}$ -tree with partite classes X, Y of size $m, n \geq 1$, respectively. Then*

$$R(G, Y, -2) \leq N(G, -2) \quad (\Leftrightarrow R(G, Y, -1) \geq N(G, -1)).$$

3. PROOFS

In this section, we start by proving Lemma 8 which will be used repeatedly in the proof of our main lemmas, namely, Lemmas 9 and 10. We then prove Theorem 2 by applying the main lemmas and deduce Corollary 3 from Theorem 2.

The following lemma shows the relationship between the scores of both players in the game on an even graph rooted at two different sets of some structure.

Lemma 8. *Let G_1 and G_2 be subgraphs of an even graph G such that $V(G_1)$ and $V(G_2)$ partition $V(G)$. If $U_1 = V(G_1) \cap N_G(V(G_2))$ and $U_2 = V(G_2) \cap N_G(V(G_1))$ are root sets of G_1 and G_2 , respectively, and every vertex in U_1 is joined to every vertex in U_2 , (see Figure 2), then*

$$8.1. R(G, U_1, 1) \geq R(G_1, U_1, -2) + R(G_2, U_2, -1),$$

$$8.2. R(G, U_1, 1) \geq R(G, U_2, 2).$$




Figure 2. The graph G in Lemma 8.

Proof. First, we shall prove Lemma 8.1 by considering a strategy for Alice who plays first in the game on G rooted at U_1 . She plays optimally as Player -2 in the game on G_1 rooted at U_1 and plays optimally as Player -1 in the game on G_2 rooted at U_2 . Since $|V(G_1)| + |V(G_2)|$ is even, she plays as Player 1 in one game and as Player 2 in the other. Now, we check that Alice's moves are feasible in the game on G rooted at U_1 , and Bob's moves are feasible in the game on G_1 rooted at U_1 and the game on G_2 rooted at U_2 . Indeed, after each move of Alice, every remaining component of G_1 and G_2 contains a vertex in U_1 and U_2 , respectively. Together with the fact that every vertex in U_2 is joined to the remaining subset of U_1 , we can conclude that every remaining component of G contains a vertex in U_1 . That is, her moves are feasible in the game on G rooted at U_1 . On the other hand, after each move of Bob, every remaining component of G contains a vertex of U_1 . Since the edges between G_1 and G_2 have endpoints only in U_1 and U_2 , every remaining component of G_1 or G_2 contains a vertex in U_1 or U_2 , respectively. That is, his moves are feasible in the game on G_1 rooted at U_1 and the game on G_2 rooted at U_2 . Hence

$$R(G, U_1, 1) \geq R(G_1, U_1, -2) + R(G_2, U_2, -1),$$

which completes the proof of Lemma 8.1. By symmetry, we have

$$R(G, U_2, 1) \geq R(G_1, U_1, -1) + R(G_2, U_2, -2),$$

which is equivalent to

$$R(G, U_2, 2) \leq R(G_1, U_1, -2) + R(G_2, U_2, -1),$$

by considering the total weight of G, G_1 and G_2 . Together with Lemma 8.1, we have

$$R(G, U_2, 2) \leq R(G_1, U_1, -2) + R(G_2, U_2, -1) \leq R(G, U_1, 1),$$

which completes the proof of Lemma 8.2. ■

We are now ready to prove the main lemmas which generalize the results on $K_{m,n}$ -trees to $B(T)$ -trees relating the scores of both players in the normal version and the rooted version of the game.

Lemma 9. *Let H be a blow-up graph of a tree with sets of vertices V_1, \dots, V_k and let G be an H -tree.*

$$\begin{aligned} 9.1. \text{ For a vertex } v \in V(G), \quad R(G, v, -2) &\leq N(G, -2) \\ &(\Leftrightarrow R(G, v, -1) \geq N(G, -1)). \end{aligned}$$

$$\begin{aligned} 9.2. \text{ For each } i \in [k], \quad R(G, V_i, -2) &\leq N(G, -2) \\ &(\Leftrightarrow R(G, V_i, -1) \geq N(G, -1)). \end{aligned}$$

$$\begin{aligned} 9.3. \text{ For each } i \in [k], \quad R(G, N_H(V_i), -2) &\leq N(G, -2) \\ &(\Leftrightarrow R(G, N_H(V_i), -1) \geq N(G, -1)). \end{aligned}$$

Lemma 10. *Let H be a blow-up graph of a tree with sets of vertices V_1, \dots, V_k and let G be an even H -tree.*

$$\begin{aligned} 10.1. \text{ For a vertex } v \in V(G), \quad R(G, v, 1) &\geq N(G, 2) \\ &(\Leftrightarrow R(G, v, 2) \leq N(G, 1)). \end{aligned}$$

$$\begin{aligned} 10.2. \text{ For each } i \in [k], \quad R(G, V_i, 1) &\geq N(G, 2) \\ &(\Leftrightarrow R(G, V_i, 2) \leq N(G, 1)). \end{aligned}$$

$$\begin{aligned} 10.3. \text{ For each } i \in [k], \quad R(G, N_H(V_i), 1) &\geq N(G, 2) \\ &(\Leftrightarrow R(G, N_H(V_i), 2) \leq N(G, 1)). \end{aligned}$$

We prove Lemmas 9 and 10 simultaneously by induction on $n = |V(G)|$. It is easy to check that Lemmas 9 and 10 hold for $n \leq 2$. Now, we let $n \geq 3$ and suppose that Lemmas 9 and 10 hold for $|V(G)| < n$. We remark that the following fact will be used throughout the proofs. Let G be an H -tree, where H is a blow-up of a tree and let v be a vertex in G . Then $G - v$ is an H' -tree, where H' is a blow-up of some tree if and only if $G - v$ is connected.

Proof of Lemma 9.1. Let $v \in V(G)$.

Case 1. G is even. Let a be an optimal move in the game on G rooted at v . Therefore, $a \neq v$ and a is feasible in the game on G . So $G - a$ is connected. Then

$$\begin{aligned} R(G, v, -1 = 2) &= R(G - a, v, 1 = -1) && \text{(Observation 5)} \\ &\geq N(G - a, -1 = 1) && \text{(Lemma 9.1 by induction)} \\ &\geq N(G, 2 = -1) && \text{(Observation 4).} \end{aligned}$$

Case 2. G is odd. Let b be an optimal move in the game on G . So $G - b$ is connected.

Case 2.1. $b \neq v$. Now, b is a feasible move in the game on G rooted at v . Then

$$\begin{aligned} R(G, v, -2 = 2) &\leq R(G - b, v, 1 = -2) && \text{(Observation 5)} \\ &\leq N(G - b, -2 = 1) && \text{(Lemma 9.1 by induction)} \\ &= N(G, 2 = -2) && \text{(Observation 4).} \end{aligned}$$

Case 2.2. $b = v$ and v is a leaf. Let u be the unique neighbor of v . Then

$$\begin{aligned} R(G, v, -2) &= R(G - v, u, -1 = 2) && \text{(Observation 6)} \\ &\leq N(G - v, 1) && \text{(Lemma 10.1 by induction)} \\ &= N(G, 2 = -2) && \text{(Observation 4 and } b = v\text{).} \end{aligned}$$

Case 2.3. $b = v$ and v is not a leaf. Therefore, $v \in V_i$ for some $i \in [k]$ and $N_G(v) = N_H(V_i)$. Then

$$\begin{aligned} R(G, v, -2) &= R(G - v, N_G(v) = N_H(V_i), -1 = 2) && \text{(Observation 6)} \\ &\leq N(G - v, 1) && \text{(Lemma 10.3 by induction)} \\ &= N(G, 2 = -2) && \text{(Observation 4 and } b = v\text{).} \end{aligned}$$

■

Proof of Lemma 9.2. Let $i \in [k]$. If $|V_i| = 1$, then we are done by Lemma 9.1. Now, suppose that $|V_i| \geq 2$.

Case 1. G is odd. Let b be an optimal move in the game on G . So $G - b$ is connected. Since $|V_i| \geq 2$, we have $V_i - b \neq \emptyset$. Therefore, b is a feasible move in the game on G rooted at V_i . Then


$$\begin{aligned} N(G, -2 = 2) &= N(G - b, 1 = -2) && \text{(Observation 4)} \\ &\geq R(G - b, V_i - b, -2 = 1) && \text{(Lemma 9.2 by induction)} \\ &\geq R(G, V_i, 2 = -2) && \text{(Observation 5).} \end{aligned}$$

Case 2. G is even. Let a be an optimal move in the game on G rooted at V_i .

Case 2.1. a is a feasible move in the game on G . So $G - a$ is connected. Then

$$\begin{aligned} R(G, V_i, -1 = 2) &= R(G - a, V_i - a, 1 = -1) && \text{(Observation 5)} \\ &\geq N(G - a, -1 = 1) && \text{(Lemma 9.2 by induction)} \\ &\geq N(G, 2 = -1) && \text{(Observation 4).} \end{aligned}$$

Case 2.2. a is not a feasible move in the game on G .

Figure 3. The graph G in Case 2.2 of Lemma 9.2.

So $G - a$ is disconnected. Since a is a feasible move in the game on G rooted at V_i , we have $a \in V_j$ for some $j \in [k]$ and $N_G(V_j) = N_H(V_j)$. Since $G - a$ is disconnected, $V_j = \{a\}$ and a is not a leaf. Suppose that $i = j$. Then every component of $G - a$ does not contain a vertex in V_i , a contradiction. Hence $i \neq j$. Suppose that there is a vertex set V_ℓ , where $\ell \notin \{i, j\}$. Then either $G - a$ is connected or there is a component of $G - a$ which does not contain a vertex in V_i , a contradiction. Hence $V_j = \{a\}$ for some $j \neq i$, $N_H(V_j) = V_i$ and $N_H(V_i) = V_j$, (see Figure 3). Therefore, G is a $K_{m,n}$ -tree with partite classes V_i and V_j . Then, by Lemma 7,

$$N(G, -1) \leq R(G, V_i, -1). \quad \blacksquare$$

Proof of Lemma 9.3. We remark that the proofs of Lemmas 9.1 and 9.2 do not use Lemma 9.3. Let $i \in [k]$. If $|N_H(V_i)| = 1$ or $N_H(V_i) = V_j$ for some $j \in [k]$, then we are done by Lemmas 9.1 or 9.2, respectively. Now, suppose that $|N_H(V_i)| \geq 2$ and V_i is joined to at least two sets in V_1, \dots, V_k .

Case 1. G is odd. Let b be an optimal move in the game on G . So $G - b$ is connected. Since $|N_H(V_i)| \geq 2$, we have $N_H(V_i) - b \neq \emptyset$. Then b is a feasible move in the game on G rooted at $N_H(V_i)$. Then

$$\begin{aligned} N(G, -2 = 2) &= N(G - b, 1 = -2) && \text{(Observation 4)} \\ &\geq R(G - b, N_H(V_i) - b, -2 = 1) && \text{(Lemma 9.3 by induction)} \\ &\geq R(G, N_H(V_i), 2 = -2) && \text{(Observation 5).} \end{aligned}$$

Case 2. G is even. Let a be an optimal move in the game on G rooted at $N_H(V_i)$.

Case 2.1. a is a feasible move in the game on G . So $G - a$ is connected. Then

$$\begin{aligned} R(G, N_H(V_i), -1 = 2) &= R(G - a, N_H(V_i) - a, 1 = -1) && \text{(Observation 5)} \\ &\geq N(G - a, -1 = 1) && \text{(Lemma 9.3 by induction)} \\ &\geq N(G, 2 = -1) && \text{(Observation 4).} \end{aligned}$$

Case 2.2. a is not a feasible move in the game on G .




Figure 4. The graph G in Case 2.2 of Lemma 9.3.

So $G - a$ is disconnected. Since a is a feasible move in the game on G rooted at $N_H(V_i)$, we have $a \in V_\ell$ for some $\ell \in [k]$ and $N_G(V_\ell) = N_H(V_\ell)$. Since $G - a$ is disconnected, $V_\ell = \{a\}$ and a is not a leaf. Suppose that $i \neq \ell$. Since V_i is joined to at least two sets, V_i and $N_H(V_i)$ lie in the same component of $G - a$, but other components of $G - a$ do not contain a vertex in $N_H(V_i)$, which is a contradiction. Hence $V_i = \{a\}$. Let $V_j \subseteq N_H(V_i)$ and let G_1 be the union of components in $G - a$ containing some vertices of V_j and let $G_2 = G - a - G_1$. By assumption, G_2 is not empty.

First, we shall show that

$$R(G, N_H(V_i), -1) \geq R(G_1, V_j, -1) + R(G_2, N_H(V_i) \setminus V_j, -1),$$

by considering a strategy for Bob who plays second in the game on G rooted at $N_H(V_i)$ after Alice grabs a . He plays optimally as Player -1 in the game on G_1 rooted at V_j and plays optimally as Player -1 in the game on G_2 rooted at $N_H(V_i) \setminus V_j$. Since $|V(G_1)| + |V(G_2)|$ is odd, he plays as Player 1 in one game and as Player 2 in the other. Now, we check that Bob's moves are feasible in the game on G rooted at $N_H(V_i)$ and Alice's moves are feasible in the game on G_1 rooted at V_j and the game on G_2 rooted at $N_H(V_i) \setminus V_j$. Indeed, after each move of Bob, every remaining component in G_1 or G_2 contains a vertex in V_j or $N_H(V_i) \setminus V_j$, respectively. Then every remaining component of G contains a vertex in $N_H(V_i)$. That is, his moves are feasible in the game on G rooted at $N_H(V_i)$. On the other hand, after each move of Alice, every remaining component of G contains a vertex in $N_H(V_i)$. Then every remaining component of G_1 or G_2 contains a vertex in V_j or $N_H(V_i) \setminus V_j$, respectively. That is, her moves are feasible in the game on G_1 rooted at V_j and the game on G_2 rooted at $N_H(V_i) \setminus V_j$. Hence

$$(1) \quad R(G, N_H(V_i), -1) \geq R(G_1, V_j, -1) + R(G_2, N_H(V_i) \setminus V_j, -1).$$

Next, we let $H_1 = G_1$ and $H_2 = G - G_1$. We observe that $V_j = V(H_1) \cap N_G(V(H_2))$ and $\{a\} = V(H_2) \cap N_G(V(H_1))$ are root sets of H_1 and H_2 , respec-

tively, and a is adjacent to all vertices in V_j , (see Figure 4). Hence

$$\begin{aligned} R(G, V_j, -2 = 1) &\geq R(G_1, V_j, -2) + R(G - G_1, a, -1) && \text{(Lemma 8.1)} \\ &= R(G_1, V_j, -2) + R(G_2, N_H(V_i) \setminus V_j, -2) + w(a) && \text{(Observation 6),} \end{aligned}$$

which is equivalent to

$$(2) \quad R(G, V_j, -1) \leq R(G_1, V_j, -1) + R(G_2, N_H(V_i) \setminus V_j, -1),$$

by considering the total weight of G, G_1 and G_2 . Then

$$\begin{aligned} N(G, -1) &\leq R(G, V_j, -1) && \text{(Lemma 9.2)} \\ &\leq R(G_1, V_j, -1) + R(G_2, N_H(V_i) \setminus V_j, -1) && \text{(Inequality (2))} \\ &\leq R(G, N_H(V_i), -1) && \text{(Inequality (1)).} \quad \blacksquare \end{aligned}$$




Figure 5. The graph G in Lemma 10.3.

Proof of Lemma 10.3. For $i \in [k]$, let G_1 be the union of components of $G - V_i$ containing some vertices of $N_H(V_i)$ and let $G_2 = G - G_1$. We observe that $N_H(V_i) = V(G_1) \cap N_G(V(G_2))$ and $V_i = V(G_2) \cap N_G(V(G_1))$ are root sets of G_1 and G_2 , respectively, and every vertex in $N_H(V_i)$ is joined to every vertex in V_i , (see Figure 5). Then

$$\begin{aligned} N(G, 2 = -1) &\leq R(G, V_i, -1 = 2) && \text{(Lemma 9.2)} \\ &\leq R(G, N_H(V_i), 1) && \text{(Lemma 8.2).} \quad \blacksquare \end{aligned}$$

Proof of Lemma 10.2. For $i \in [k]$, let G_1 be the union of components of $G - N_H(V_i)$ containing some vertices of V_i and let $G_2 = G - G_1$. We observe that $V_i = V(G_1) \cap N_G(V(G_2))$ and $N_H(V_i) = V(G_2) \cap N_G(V(G_1))$ are root sets of G_1 and G_2 , respectively, and every vertex in V_i is joined to every vertex in $N_H(V_i)$. Then

$$\begin{aligned} N(G, 2 = -1) &\leq R(G, N_H(V_i), -1 = 2) && \text{(Lemma 9.3)} \\ &\leq R(G, V_i, 1) && \text{(Lemma 8.2).} \quad \blacksquare \end{aligned}$$

Proof of Lemma 10.1. Let $v \in V(G)$.

Case 1. There is a cut edge uv incident to v .




Figure 6. The graph G in Case 1 of Lemma 10.1.

Let G_1 be the component of $G - uv$ containing v and let $G_2 = G - G_1$. We observe that $\{v\} = V(G_1) \cap N_G(V(G_2))$ and $\{u\} = V(G_2) \cap N_G(V(G_1))$ are root sets of G_1 and G_2 , respectively, and v is adjacent to u , (see Figure 6). Then

$$\begin{aligned} R(G, v, 1) &\geq R(G, u, 2 = -1) && \text{(Lemma 8.2)} \\ &\geq N(G, -1 = 2) && \text{(Lemma 9.1).} \end{aligned}$$

Case 2. There is no cut edge incident to v . Then $v \in V_j$ for some $j \in [k]$ and $N_G(v) = N_H(V_j)$.

Case 2.1. $|V_j| \geq 2$. Therefore, v is a feasible move in the game on G . So $G - v$ is connected. Then

$$\begin{aligned} R(G, v, 1 = -2) &= R(G - v, N_G(v) = N_H(V_j), -1) && \text{(Observation 6)} \\ &\geq N(G - v, -1 = 1) && \text{(Lemma 9.3 by induction)} \\ &\geq N(G, 2) && \text{(Observation 4).} \end{aligned}$$

Case 2.2. $|V_j| = 1$. Then, by Lemma 10.2,

$$R(G, v, 1) = R(G, V_j, 1) \geq N(G, 2). \quad \blacksquare$$

We proceed to prove our main theorem.

Proof of Theorem 2. Let G be an even $B(T)$ -tree, where T is a tree and let $v \in V(G)$. Then, by Lemmas 9.1 and 10.1, it follows that

$$N(G, 2 = -1) \leq R(G, v, -1 = 2) \leq N(G, 1).$$

Therefore, Alice wins the game on G . \blacksquare

We now deduce Corollary 3 from Theorem 2.

Proof of Corollary 3. We give a proof by induction on the number of vertices. Let G be an even blow-up of a cycle. We note that every vertex of G is a non-cut

vertex. Alice claims a maximum weighted vertex of G in her first move, say a vertex a . Let b be the vertex claimed by Bob in his first move. Then $G - \{a, b\}$ is an even blow-up of either a path or a cycle. If $G - \{a, b\}$ is an even blow-up of a path, then Alice wins the game on $G - \{a, b\}$ by Theorem 2. Otherwise, Alice wins the game on $G - \{a, b\}$ by the induction hypothesis. In both cases, since $w(a) \geq w(b)$, Alice wins the game on G . ■

4. CONCLUDING REMARKS

We provide two new classes, namely $B(T)$ -trees and $B(C_{2n})$, of bipartite even graphs which satisfy Conjecture 1. However, this conjecture is still open. It was shown in [3] that Lemmas 9.1 and 10.1 are not true for general bipartite graphs, therefore this method cannot be directly used to solve the full conjecture. There are several variants of the graph grabbing game, for example, the graph sharing game (see [1, 2, 5, 6, 9]), the graph grabbing game on $\{0, 1\}$ -weighted graphs (see [4]), and the convex grabbing game (see [7]), where a few problems are left open.

Acknowledgment

The first author is grateful for financial support from the Science Achievement Scholarship of Thailand.

REFERENCES

- [1] S. Chaplick, P. Micek, T. Ueckerdt and V. Wiechert, *A note on concurrent graph sharing games*, *Integers* **16** (2016) #G1.
- [2] J. Cibulka, J. Kynčl, V. Mészáros, R. Stolař, and P. Valtr, *Graph sharing games: complexity and connectivity*, *Theoret. Comput. Sci.* **494** (2013) 49–62.
<https://doi.org/10.1016/j.tcs.2012.12.029>
- [3] Y. Egawa, H. Enomoto and N. Matsumoto, *The graph grabbing game on $K_{m,n}$ -trees*, *Discrete Math.* **341** (2018) 1555–1560.
<https://doi.org/10.1016/j.disc.2018.02.023>
- [4] S. Eoh and J. Choi, *The graph grabbing game on $\{0, 1\}$ -weighted graphs*, *Results Appl. Math.* **3** (2019) # 100028.
<https://doi.org/10.1016/j.rinam.2019.100028>
- [5] A. Gałol, P. Micek and B. Walczak, *Graph sharing game and the structure of weighted graphs with a forbidden subdivision*, *J. Graph Theory* **85** (2017) 22–50.
<https://doi.org/10.1002/jgt.22045>
- [6] K. Knauer, P. Micek and T. Ueckerdt, *How to eat 4/9 of a pizza*, *Discrete Math.* **311** (2011) 1635–1645.
<https://doi.org/10.1016/j.disc.2011.03.015>

- [7] N. Matsumoto, T. Nakamigawa and T. Sakuma, *Convex grabbing game of the point set on the plane*, Graphs Combin. **36** (2020) 51–62.
<https://doi.org/10.1007/s00373-019-02117-z>
- [8] P. Micek and B. Walczak, *A graph-grabbing game*, Combin. Probab. Comput. **20** (2011) 623–629.
<https://doi.org/10.1017/S0963548311000071>
- [9] P. Micek and B. Walczak, *Parity in graph sharing games*, Discrete Math. **312** (2012) 1788–1795.
<https://doi.org/10.1016/j.disc.2012.01.037>
- [10] M. Rosenfeld, *A gold-grabbing game*, *Open Problem Garden* (2009).
http://www.openproblemgarden.org/op/a_gold_grabbing_game
- [11] D.E. Seacrest and T. Seacrest, *Grabbing the gold*, Discrete Math. **312** (2012) 1804–1806.
<https://doi.org/10.1016/j.disc.2012.01.010>
- [12] P. Winkler, *Mathematical Puzzles: A Connoisseur's Collection* (A K Peters, Ltd., Natick, MA, 2004).

Received 7 December 2020

Revised 25 April 2021

Accepted 5 May 2021

Available online 24 June 2021