Discussiones Mathematicae Graph Theory 42 (2022) 1383–1388 https://doi.org/10.7151/dmgt.2437

# A NOTE ON PACKING OF UNIFORM HYPERGRAPHS

JERZY KONARSKI, MARIUSZ WOŹNIAK

AND

Andrzej Żak

AGH University of Science and Technology

e-mail: konarski@agh.edu.pl mwozniak@agh.edu.pl zakandrz@agh.edu.pl

## Abstract

We say that two *n*-vertex hypergraphs  $H_1$  and  $H_2$  pack if they can be found as edge-disjoint subhypergraphs of the complete hypergraph  $K_n$ . Whilst the problem of packing of graphs (i.e., 2-uniform hypergraphs) has been studied extensively since seventies, much less is known about packing of k-uniform hypergraphs for  $k \geq 3$ . Naroski [Packing of nonuniform hypergraphs - product and sum of sizes conditions, Discuss. Math. Graph Theory 29 (2009) 651–656] defined the parameter  $m_k(n)$  to be the smallest number m such that there exist two n-vertex k-uniform hypergraphs with total number of edges equal to m which do not pack, and conjectured that  $m_k(n) = \Theta(n^{k-1})$ . In this note we show that this conjecture is far from being truth. Namely, we prove that the growth rate of  $m_k(n)$  is of order  $n^{k/2}$  exactly for even k's and asymptotically for odd k's.

Keywords: packing, hypergraphs.

2010 Mathematics Subject Classification: 05C35.

#### 1. INTRODUCTION

By a hypergraph H we mean a pair (V(H), E(H)) where V(H) is a finite set (elements of V(H) are called *vertices*) and E(H) is a family of subsets of V(H)(members of E(H) are called *edges*). A hypergraph is *complete* if its set of edges consists of all subsets of V(H). The complete hypergraph on n vertices is denoted  $K_n$ . Furthermore, we use the term *k*-uniform hypergraph to refer to hypergraphs with all edges consisting of exactly k vertices. Let  $H_1$  and  $H_2$  be two hypergraphs such that  $|V(H_1)| = |V(H_2)| = n$ .

We say that  $H_1$  and  $H_2$  pack into  $K_n$  (in short pack) if  $H_1$  and  $H_2$  can be found as edge-disjoint subhypergraphs in  $K_n$ . Clearly, the more edges  $H_1$  and  $H_2$  have, the harder is to pack them. Hence a natural question is to determine the bound on the common number of edges that guarantees the packing property of any two hypergraphs satisfying the bound. To this end let us define  $m_k(n)$  to be the least integer m such that there exist two n-vertex k-uniform hypergraphs  $H_1$  and  $H_2$  with  $|E(H_1)| + |E(H_2)| = m$  which do not pack. It is trivial that

 $m_1(n) = n + 1$ 

and it was shown by Sauer and Spencer [9] that

$$m_2(n) = \left\lceil \frac{3}{2}n \right\rceil - 1,$$

(see also, among others, [1, 2, 4, 8] for generalizations). For  $k \ge 3$ , Naroski [5] proved that

(1) 
$$m_k(n) \ge 2\sqrt{\binom{n}{k}},$$

and conjectured that  $m_k(n) = \Theta(n^{k-1})$  (earlier Pilśniak and Woźniak [7] conjectured the same for k = 3). It turns out that this is far from being truth. In this note we establish the growth order of  $m_k(n)$  exactly for even k's, and asymptotically for odd k's by proving the following theorem.

**Theorem 1.** If k is even then

$$m_k(n) \le \binom{n-k/2}{k/2} + \frac{\binom{n+k^{k/2}-1}{k/2}}{\binom{k}{k/2}}.$$

If k is odd then, as n tends to infinity,

$$m_k(n) \le \left(\frac{k^{k-1}+1}{(k-1)!} + o(1)\right) n^{(k^2-k-1)/(2k-3)}.$$

Note that for k = 2, the bound agrees with the result of Sauer and Spencer.

## 2. Proof of Theorem 1

For any positive integer n let  $[n] = \{1, ..., n\}$ . In the proof we will use the following famous results, one very old and one relatively new.

**Theorem 2** (Chinese Remainder Theorem [6]). Let  $a_i, b_i, i = 1, ..., r$ , be any integers. If  $a_i \equiv a_j \mod \text{gcd}(b_i, b_j)$ , then there exists exactly one  $0 \leq x < \text{lcm}(b_1, ..., b_r)$  satisfying

$$x \equiv a_i \mod b_i \quad i = 1, \ldots, r_i$$

A t- $(n, k, \lambda)$ -design on a set X of size n is a collection T of k-element subsets of X such that every t elements of X are contained in exactly  $\lambda$  sets of T. Recently Keevash [3] proved the following deep existence theorem.

**Theorem 3** ([3]). A t- $(n, k, \lambda)$ -design on a set X exists if and only if for every  $0 \le i \le t-1$ 

(2) 
$$\binom{k-i}{t-i}$$
 divides  $\lambda \binom{n-i}{t-i}$ ,

apart from a finite number of exceptional n given fixed  $k, t, \lambda$ .

In the sequel we will identify the design with the hypergraph (X,T). Note that if (X,T) is a t- $(n,k,\lambda)$ -design, then

(3) 
$$|T| = \frac{\lambda \binom{n}{t}}{\binom{k}{t}}$$

since each t-subset of X is contained in exactly  $\lambda$  edges (i.e., elements of T), and each such edge is counted  $\binom{k}{t}$  many times.

Now we are ready to give a proof of Theorem 1.

**Proof of Theorem 1.** We are going to construct two *n*-vertex hypergraphs  $H_1$  and  $H_2$  which do not pack and have few edges. Let  $V(H_1) = V(H_2) = [n]$ .

Consider first the case when k is even, i.e., k = 2t. Then let

$$E(H_1) = \left\{ e \in \binom{[n]}{k} : [t] \subset e \right\}.$$

If n satisfies the divisibility conditions (2), then let

$$H_2$$
 be a  $t$ - $(n, k, 1)$ -design.

Since [t] forms an edge with every t-subset of  $V(H_1) \setminus [t]$  and every t-subset of  $V(H_2)$  is contained in exactly one edge of  $H_2$ , it is not possible to find edge disjoint copies of  $H_1$  and  $H_2$  in  $K_n$ . Hence  $H_1$  and  $H_2$  do not pack. In the case when (2) is not satisfied, we have to modify a bit the construction of  $H_2$ . First

we will find the smallest possible number x such that n + x satisfies (2) with n replaced by n + x. Note that

$$\binom{k-i}{t-i} \left| \binom{n+x-i}{t-i} \Leftrightarrow \frac{(n+x-i)!}{(t-i)!(n+x-t)!} = p \frac{(k-i)!}{(t-i)!(k-t)!} \right|$$
$$\Leftrightarrow (n+x-i)\cdots(n+x-t+1) = p(k-i)\cdots(k-t+1)$$

for some integer p. Hence, in order to assure that n + x satisfies (2) it suffices that

(4) 
$$k - i|n + x - i \text{ for } i = 0, \dots, t - 1.$$

In other words

(5) 
$$x \equiv i - n \mod (k - i) \text{ for } i = 0, \dots, t - 1.$$

Note that if g|(k-i) and g|(k-j), then g|(i-j). Thus,  $i-n \equiv j-n \mod g$ . Hence, by the Chinese Remainder Theorem, there exist

$$(6) x \le k^t - 1$$

that satisfies (4). Thus, by Theorem 3, as  $H'_2$  we can take a t-(n + x, k, 1)-design on the set [n+x]. In order to construct  $H_2$  we replace each k-subset e' of  $H'_2$  such that  $e' \not\subset [n]$  by a k-set  $e = (e' \cap [n]) \cup f$  where f is an arbitrary  $|e' \setminus [n]|$ -subset of  $[n] \setminus e'$ . Thus

(7) 
$$|E(H_2)| = |E(H'_2)|$$
 and

(8) each t-subset of  $V(H_2)$  is contained in at least one edge of  $H_2$ .

Hence,  $H_1$  and  $H_2$  do not pack. Furthermore, by (7), (6) and (3), we have

$$|E(H_1)| + |E(H_2)| \le {\binom{n-k/2}{k/2}} + \frac{{\binom{n+k^{k/2}-1}{k/2}}}{{\binom{k}{k/2}}},$$

which proves the statement in the case when k is even.

In the case when k is odd one can obtain the bound  $m_k(n) = O(n^{(k+1)/2})$  in a similar way. However, we can do better via a different construction. Let

(9) 
$$s = |n^{(k-2)/(2k-3)}|$$

and a, b be non-negative integers satisfying

(10) 
$$n = a\lfloor n/s \rfloor + b\lceil n/s \rceil, \quad a+b = s.$$

1386

Define

$$E(H_1) = \left\{ e \in \binom{[n]}{k} : |e \cap [s(k-2)+1]| \ge k-1 \right\}.$$

By a similar reasoning as in the construction of  $H_2$ , we can deduce the existence of a (k-1)- $(\lceil n/s \rceil + x, k, 1)$ -design with  $x < k^{k-1}$ . We identify the design with a hypergraph H' on  $\lceil n/s \rceil + x$  vertices and then modify it (again in the same way as before) to get a  $\lfloor n/s \rfloor$ -vertex hypergraph  $H_{\lfloor n/s \rfloor}$  and a  $\lceil n/s \rceil$ -vertex hypergraph  $H_{\lceil n/s \rceil}$  satisfying

(11) |E(H)| = |E(H')| and

(12) each (k-1)-subset of V(H) is contained in at least one edge of H,

where  $H \in \{H_{\lfloor n/s \rfloor}, H_{\lceil n/s \rceil}\}$ . Now let

$$H_2 = aH_{\lfloor n/s \rfloor} + bH_{\lceil n/s \rceil},$$

that means  $H_2$  is a disjoint union of a hypergraphs  $H_{\lfloor n/s \rfloor}$  and b hypergraphs  $H_{\lceil n/s \rceil}$ . By a Pigeonhole Principle, the subset [s(k-2)+1] of the vertex set of every copy of  $H_1$  in  $K_n$  intersects the vertex set of some  $H_{\lfloor n/s \rfloor}$  or  $H_{\lceil n/s \rceil}$  in every copy of  $H_2$  in  $K_n$  in at least k-1 vertices. Let I be such an intersection for fixed copies of  $H_1$  and  $H_2$  in  $K_n$ . By the construction of  $H_1$ , every k-subset containing I is an edge of  $H_1$ , while by (11), at least one k-subset containing I is an edge of  $H_2$ . Hence, the copies have at least one common edge. Thus,  $H_1$  and  $H_2$  do not pack. Furthermore, by (9), (3) and (6),

$$\begin{aligned} |E(H_1)| + |E(H_2)| &\leq \binom{s(k-2)+1}{k-1} (n-s(k-2)-1) + s\binom{\lceil n/s \rceil + x - 1}{k-1} \\ &\leq \frac{(ks)^{k-1}}{(k-1)!} n + s \frac{(n/s+x)^{k-1}}{(k-1)!} \\ &= \frac{k^{k-1}+1}{(k-1)!} n^{(k^2-k-1)/(2k-3)} + o\left(n^{(k^2-k-1)/(2k-3)}\right). \end{aligned}$$

This proves the statement in the case when k is odd.

# References

- [1] B. Bollobás, Extremal Graph Theory (Academic Press, London-New York, 1978).
- B. Bollobás and S.E. Eldridge, Packing of graphs and applications to computational complexity, J. Combin. Theory Ser. B 25 (1978) 105–124. https://doi.org/10.1016/0095-8956(78)90030-8

- [3] P. Keevash, The existence of designs (2014). arXiv:1401.3665
- [4] A. Kostochka, C. Stocker and P. Hamburger, A hypergraph version of a graph packing theorem by Bollobás and Eldridge, J. Graph Theory 74 (2013) 222–235. https://doi.org/10.1002/jgt.21706
- P. Naroski, Packing of nonuniform hypergraphs—product and sum of sizes conditions, Discuss. Math. Graph Theory 29 (2009) 651–656. https://doi.org/10.7151/dmgt.1471
- [6] O. Ore, The general Chinese reminder theorem, Amer. Math. Monthly 59 (1952) 365–370. https://doi.org/10.1080/00029890.1952.11988142
- M. Pilśniak and M. Woźniak, A note on packing of two copies of a hypergraph, Discuss. Math. Graph Theory 27 (2007) 45–49. https://doi.org/10.7151/dmgt.1343
- [8] M. Pilśniak and M. Woźniak, On packing of two copies of a hypergraph, Discrete Math. Theor. Comput. Sci. 13 (2011) 67–74. https://doi.org/10.46298/dmtcs.537
- [9] N. Sauer and J. Spencer, *Edge disjoint placement of graphs*, J. Combin. Theory Ser. B 25 (1978) 295–302. https://doi.org/10.1016/0095-8956(78)90005-9

Received 24 June 2021 Revised 2 November 2021 Accepted 2 November 2021