

DEGREE SUM CONDITION FOR VERTEX-DISJOINT 5-CYCLES

MAOQUN WANG AND JIANGUO QIAN¹

*School of Mathematical Sciences
Xiamen University
Xiamen, Fujian 361005, P.R. China
e-mail: jgqian@xmu.edu.cn*

Abstract

Let n and k be two integers and G a graph with $n = 5k$ vertices. Wang proved that if $\delta(G) \geq 3k$, then G contains k vertex disjoint cycles of length 5. In 2018, Chiba and Yamashita asked whether the degree condition can be replaced by degree sum condition. In this paper, we give a positive answer to this question.

Keywords: degree sum conditions; vertex disjoint 5-cycles.

2020 Mathematics Subject Classification: 05C38, 05C70, 05C75.

1. INTRODUCTION

Let G be a graph. We denote by $V(G)$ and $E(G)$ the vertex set and edge set of G , respectively, and call $|V(G)|$ the *order* of G . An edge joining two vertices x and y is denoted by xy . A class of subgraphs of G is *vertex disjoint*, or simply *disjoint*, if no two subgraphs in the class have a common vertex. The *length* of a cycle C (or a path P) is the number of edges on C (or P). For $X \subseteq V(G)$, the *neighbour set* of X , denoted by $N_G(X)$ or $N(X)$ with no confusion, is the set of the vertices not in X but adjacent to at least one vertex in X . In particular, if X consists of a single vertex x , then we call $|N_G(\{x\})|$ the *degree* of x and denote it by $d_G(x)$. We use $\delta(G)$ to denote the minimum degree of vertices in G . Define

$$\sigma_2(G) = \min\{d(x) + d(y) : x, y \in V(G), x \neq y, xy \notin E(G)\}.$$

The degree condition for the existence of cycle(s) with specified length(s) is one of the most elementary concerns in graph theory. A classic result should be

¹Corresponding author.

the one given by Dirac [4] in 1952, which says that every graph with n vertices and minimum degree at least $n/2$ has a Hamilton cycle. Since then, this result has been generalized to various forms in terms of degree condition or degree sum condition. Corrádi and Hajnal [2] considered the maximum number of disjoint cycles in a graph and proved that if a graph G has at least $3k$ vertices and minimum degree at least $2k$, then G contains k disjoint cycles. In particular, when G has exactly $3k$ vertices, then G contains k disjoint triangles. Erdős and Faudree [6] conjectured that if G has $4k$ vertices and minimum degree $\delta(G) \geq 2k$, then G contains k vertex disjoint cycles of length 4. This conjecture was later confirmed by Wang [7]. In general, El-Zahar posed the following conjecture.

Conjecture 1 (El-Zahar, [5]). *Let G be a graph of order $n = n_1 + n_2 + \cdots + n_k$ with $n_i \geq 3$ for each $i \in \{1, 2, \dots, k\}$. If $\delta(G) \geq \lceil \frac{n_1}{2} \rceil + \lceil \frac{n_2}{2} \rceil + \cdots + \lceil \frac{n_k}{2} \rceil$, then G has k disjoint cycles of length n_1, n_2, \dots, n_k , where, for a real number r , $\lceil r \rceil$ is the least integer not less than r .*

In the same paper, El-Zahar also proved that the above conjecture is true for $k = 2$. In [1], Abbasi confirmed the conjecture for sufficiently large graphs by using the regularity lemma. Wang [8] proved the conjecture for the case when $n = 5k$ and $n_i = 5$ ($1 \leq i \leq k$) as follows.

Theorem 2 (Wang, [8]). *Let k be a positive integer and G a graph of order $n = 5k$. If $\delta(G) \geq (n+k)/2$, then G contains k vertex disjoint cycles of length 5.*

In 2018, Chiba and Yamashita [3] posed the following question.

Question (Chiba and Yamashita, [3]). *Let k be a positive integer and G a graph of order $n = 5k$. Is it true that if $\sigma_2(G) \geq n+k$, then G contains k vertex disjoint cycles of length 5?*

In this paper, we give a positive answer to this question.

Theorem 3. *Let k be a positive integer and G a graph of order $n = 5k$. If $\sigma_2(G) \geq n+k$, then G contains k vertex disjoint cycles of length 5.*

2. PRELIMINARIES

For $X, Y \subseteq V(G)$, we denote by $N(X)|Y$ the neighbour set of X restricted on Y , i.e., $N(X)|Y = N(X) \cap Y$. For simplicity, if X consists of a single vertex x , we simply write $N(\{x\})|Y$ as $N(x)|Y$, and if H is a subgraph of G , then we simply write $N(X)|V(H)$ as $N(X)|H$. A *chord* of a cycle C is an edge not on C that joins two vertices of C , and the number of chords of C is denoted by $\tau(C)$. Further, for $x \in V(C)$, we use $\tau(x, C)$ to denote the number of chords of C that are incident with x . For a graph H , we say that

G contains H if G has a subgraph isomorphic to H and denote by $G \supseteq H$. For two graphs G and G' , we denote by $G \uplus G'$ the vertex disjoint union of G and G' . For simplicity, we write the vertex disjoint union of k copies of a graph H as kH . For two disjoint vertex subsets, or subgraphs, A and B of G , we define $E(A, B)$ to be the set of all the edges of G between A and B and denote $e(A, B) = |E(A, B)|$. For a vertex subset X of G , we denote by $[X]$ the subgraph of G induced by X . Further, for subgraphs G_1, G_2, \dots, G_t of G , we write $[V(G_1) \cup V(G_2) \cup \dots \cup V(G_t)] = [G_1, G_2, \dots, G_t]$. For $t + 1$ disjoint subgraphs H, L_1, \dots, L_t of G , we call $\{H, L_1, \dots, L_t\}$ a *family* of G if $L_i \cong C_5$ for all $i \in \{1, 2, \dots, t\}$. In particular, we call a family $\{H, L_1, \dots, L_t\}$ *optimal* if $\sum_{i=1}^t \tau(L'_i) \leq \sum_{i=1}^t \tau(L_i)$ for any family $\{H', L'_1, \dots, L'_t\}$ of $[H, L_1, \dots, L_t]$ with $H' \cong H$ and $L'_i \cong C_5$ for all $i \in \{1, 2, \dots, t\}$.

As usual, for $i \geq 3$, we denote by K_i, C_i and P_i the complete graph, cycle and path of order i , respectively. Following the notations in [8], let $B, F, F_1, F_2, F_3, F_4, F_5, K_4^+$ be the graphs as illustrated in Figure 1. Let L be a 5-cycle, $u \in V(L)$ and $x \in V(G) \setminus V(L)$. We write $x \rightarrow (L, u)$ if $[L - u + x] \supseteq C_5$. In particular, if $x \rightarrow (L, u)$ for all $u \in V(L)$, then we write $x \rightarrow L$. Further, for $\{v_1, v_2\} \subseteq V(G)$, we write $x \rightarrow (L, u; \{v_1, v_2\})$ if $x \rightarrow (L, u)$ and u is adjacent to both v_1 and v_2 .




Figure 1. The subgraphs $B, F, F_1, F_2, F_3, F_4, F_5, K_4^+$.

To prove the main theorem, we introduce the following lemmas.

Lemma 4 [8]. *For a graph G , the following statements hold.*

- Let B_1 and B_2 be two disjoint subgraphs of G with $B_1 \cong B_2 \cong B$ and R the set of the four vertices of degree 2 in B_1 . If $e(R, B_2) \geq 13$, then $[B_1, B_2] \supseteq 2C_5$ or $[B_1, B_2] \supseteq B \uplus C_5$.
- Let D and L be two disjoint subgraphs of G with $D \cong B$ and $L \cong C_5$ and x the unique vertex of degree 4 in D . If $e(D - x, L) \geq 13$, then $[D, L] \supseteq 2C_5$.
- Let D and L be two disjoint subgraphs of G with $D \cong F_2$ and $L \cong C_5$ and R the set of the three vertices of degree 2 in D . If $e(R, L) \geq 10$, then

$$[D, L] \supseteq F_1 \uplus C_5.$$

- (d) Let P and L be two disjoint subgraphs of G with $P \cong P_5$ and $L \cong C_5$. If $e(P, L) \geq 16$, $[P, L] \not\supseteq 2C_5$ and $\{P, L\}$ is optimal, then $[P, L] \supseteq F \uplus C_5$.
- (e) Let $R \subseteq V(G)$ and L be a 5-cycle in $G - R$. If $|R| = 4$ and $e(R, L) \geq 13$, then $x \rightarrow (L, y; \{x_1, x_2\})$ for some $y \in V(L)$, $x \in R$ and $\{x_1, x_2\} \subseteq R \setminus \{x\}$; or there are vertex labellings $R = \{x_1, x_2, x_3, x_4\}$ and $L = y_1y_2y_3y_4y_5$ such that $N(x_1) \cap L = N(x_2) \cap L = \{y_1, y_2, y_3, y_4\}$, $N(x_3) \cap L = \{y_1, y_4, y_5\}$ and $N(x_4) \cap L = \{y_1, y_4\}$.

Throughout the following, when we speak of a subgraph isomorphic to one of that in Figure 1, we always assume that its vertices are labelled as indicated in the figure. Further, for two vertex subsets $\{x_1, x_2, \dots, x_k\}$ and $\{y_1, y_2, \dots, y_h\}$, we write $e(\{x_1, x_2, \dots, x_k\}, \{y_1, y_2, \dots, y_h\})$ by $e(x_1x_2 \cdots x_k, y_1y_2 \cdots y_h)$ for simplicity.

Lemma 5 [8]. *For a graph G , the following statements hold.*

- (a) Let D and L be two disjoint subgraphs of G with $D \cong F$ and $L \cong C_5$. If $\{D, L\}$ is optimal and $e(D, L) \geq 16$, then $[D, L]$ contains one of $F_1 \uplus C_5$, $F_2 \uplus C_5$, $B \uplus C_5$ and $2C_5$; or the cycle L has a labelling $L = y_1y_2y_3y_4y_5$ satisfying the following property
 \mathbf{P}_1 : $e(x_1, L) = 0$, $e(x_2x_4, L) = 10$, $N(x_3) \cap L = N(x_5) \cap L = \{y_1, y_2, y_4\}$, $\tau(L) = 4$ and $y_3y_5 \notin E(G)$.
- (b) Let D , L and L' be three disjoint subgraphs of G with $D \cong F$, $L \cong L' \cong C_5$ and $L = y_1y_2y_3y_4y_5$. If D and L satisfy the property \mathbf{P}_1 and $e(x_1x_3y_3y_5, L') \geq 13$, then $[D, L, L']$ contains either $F_1 \uplus 2C_5$ or $3C_5$.

Lemma 6 [8]. *For a graph G , the following statements hold.*

- (a) Let D and L be two disjoint subgraphs of G with $D \cong F_1$ and $L \cong C_5$. If $\{D, L\}$ is optimal and $e(D, L) \geq 16$, then $[D, L]$ contains one of $K_4^+ \uplus C_5$, $K_4^+ \uplus B$, $B \uplus C_5$ and $2C_5$; or L has a labelling $L = y_1y_2y_3y_4y_5$ satisfying the following property
 \mathbf{P}_2 : $e(x_1, L) = 0$, $e(y_1y_2y_4, D - x_1) = 12$, $N(y_3) \cap D = N(y_5) \cap D = \{x_3, x_5\}$, $\tau(L) = 4$ and $y_3y_5 \notin E$.
- (b) Let D , L and L' be three disjoint subgraphs of G with $D \cong F_1$, $L \cong L' \cong C_5$ and $L = y_1y_2y_3y_4y_5$. If D and L satisfy \mathbf{P}_2 , $\{D, L, L'\}$ is optimal and $e(x_1x_4y_3y_5, L') \geq 13$, then $[D, L, L']$ contains either $K_4^+ \uplus 2C_5$ or $3C_5$.

Lemma 7. *Let D and L be two disjoint subgraphs of G with $D \cong F_3$ and $L \cong C_5$. If $e(D - x_3, L) \geq 13$, then $[D, L]$ contains either $2C_5$ or $F \uplus C_5$ or $B \uplus C_5$ or $F_4 \uplus C_5$.*

Proof. Let $L = y_1y_2y_3y_4y_5y_1$ and $R = V(D) \setminus \{x_3\}$. Suppose to the contrary that $[D, L]$ does not contain any of $2C_5$, $F \uplus C_5$, $B \uplus C_5$ and $F_4 \uplus C_5$. Suppose $e(x_4x_5, L) \geq 8$. Without loss of generality, assume that $e(x_4, L) \geq e(x_5, L)$. If $e(x_4, L) = 5$, then since $e(x_1x_2x_5, L) \geq 8$, L has a vertex y_j such that $x_ty_j, x_5y_j \in E(G)$ for some $t \in \{1, 2\}$ and so $x_4 \rightarrow (L, y_j; \{x_5, x_t\})$. Thus $[D, L]$ contains $2C_5$ or $F \uplus C_5$, a contradiction. Then $e(x_4x_5, L) = 8$ and $e(x_4, L) = e(x_5, L) = 4$, say $N(x_4) \mid L = \{y_1, y_2, y_3, y_4\}$. If $N(x_5) \mid L = \{y_1, y_2, y_3, y_5\}$ or $N(x_5) \mid L = \{y_1, y_2, y_4, y_5\}$ or $N(x_5) \mid L = \{y_1, y_3, y_4, y_5\}$ or $N(x_5) \mid L = \{y_2, y_3, y_4, y_5\}$, then since $e(x_1x_2, L) \geq 5$, L has a vertex y_j such that $x_r \rightarrow (L, y_j; \{x_s, x_t\})$ for some $t \in \{1, 2\}$ and $\{r, s\} = \{4, 5\}$. Hence $[D, L]$ contains $2C_5$ or $F \uplus C_5$, a contradiction. If $N(x_5) \mid L = \{y_1, y_2, y_3, y_4\}$, then $e(x_1x_2, y_2y_3) = 0$ for otherwise $[D, L]$ contains $2C_5$ or $F \uplus C_5$. Hence $e(x_1x_2, y_1y_4y_5) \geq 5$ and so $[x_1, x_2, y_1, y_4, y_5] \supseteq C_5$ and $[x_3, x_4, x_5, y_2, y_3] \supseteq C_5$, again a contradiction. Therefore $e(x_4x_5, L) \leq 7$ and hence $e(x_1x_2, L) \geq 6$. If $e(x_1, L) = 1$, say $x_1y_1 \in E(G)$, then $e(y_2y_3, x_4x_5) \leq 1$ (otherwise $[y_1, y_4, y_5, x_1, x_2] \supseteq C_5$ and $[x_3, x_4, x_5, y_2, y_3]$ contains C_5 or F or B). Similarly, $e(y_4y_5, x_4x_5) \leq 1$. Hence $e(R, L) \leq 10$, a contradiction. Now suppose $e(x_1, L) \geq 4$, say $N(x_1) \mid L \supseteq \{y_1, y_2, y_3, y_4\}$. If $x_1 \rightarrow (L, y_i)$ for some $i \in \{1, 2, 3, 4, 5\}$, then $e(y_i, x_4x_5) = 0$ (otherwise $[L - y_i + x_1] \supseteq C_5$ and $[D + y_i - x_1] \supseteq F_4$). This implies that $N(x_1) \mid L = \{y_1, y_2, y_3, y_4\}$, $e(x_4x_5, y_1y_4) = 4$, $e(x_2, L) = 5$. Therefore, $[y_2, y_3, x_1, x_2, x_3] \supseteq F$ and $[y_1, y_4, y_5, x_4, x_5] \supseteq C_5$, again a contradiction.

Case 1. $e(x_1, L) = 2$. In this case, $e(x_2, L) \geq 4$ as $e(x_1x_2, L) \geq 6$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2\}$, where and herein after, the subscript of a vertex y_i on C_5 means in i modulo 5 if $i > 5$. If $x_2y_4 \in E(G)$, then $y_1y_5y_4x_2x_1y_1 \cong y_2y_3y_4x_2x_1y_2 \cong C_5$. Hence, $e(y_1y_5, x_4x_5) \leq 1$ and $e(y_2y_3, x_4x_5) \leq 1$, for otherwise, $[y_2, y_3, x_3, x_4, x_5]$ and $[y_1, y_5, x_3, x_4, x_5]$ would contain C_5 or F or B . This means that $e(R, L) \leq 11$, a contradiction. We now assume $x_2y_4 \notin E(G)$ and, hence $N(x_2) \mid L = \{y_1, y_2, y_3, y_5\}$. We claim that $e(y_3y_4, x_4x_5) \leq 1$. If not, then $[y_1, y_2, y_5, x_1, x_2] \supseteq C_5$, $[y_3, y_4, x_3, x_4, x_5]$ contains C_5 or F or B , a contradiction. Similarly, $e(y_4y_5, x_4x_5) \leq 1$. Hence $e(R, L) \leq 12$, a contradiction. Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_3\}$. As $x_1 \rightarrow (L, y_2)$, $e(y_2, x_4x_5) = 0$. Recall that $e(x_2, L) \geq 4$. We have $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$. Therefore, $e(y_4y_5, x_4x_5) \leq 1$, for otherwise $[y_4, y_5, x_3, x_4, x_5]$ would contain C_5 or F or B . It follows that $e(R, L) \leq 12$, again a contradiction.

Case 2. $e(x_1, L) = 3$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L, y_2)$, $e(y_2, x_4x_5) = 0$. Note that $[x_2, x_1, y_2, y_3, y_4] \supseteq F_4$ and $[x_2, x_1, y_1, y_2, y_5] \supseteq F_4$. Then $e(y_3y_4, x_4x_5) \leq 2$ and $e(y_5y_1, x_4x_5) \leq 2$ for otherwise $[x_3, x_4, x_5, y_3, y_4] \supseteq C_5$ and $[x_3, x_4, x_5, y_1, y_5] \supseteq C_5$. Hence $e(R, L) \leq 12$, a contradiction. Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_4\}$. As $x_1 \rightarrow$

(L, y_i) for all $i \in \{3, 5\}$, $e(y_i, x_4x_5) = 0$. It follows that $e(x_2, y_3y_4y_5) \geq 2$ and so $[x_1, x_2, y_3, y_4, y_5] \supseteq F$. Then $e(y_1y_2, x_4x_5) \leq 2$ as $[x_3, x_4, x_5, y_1, y_2] \not\supseteq C_5$. Consequently, $e(R, L) \leq 12$, a contradiction. ■

Lemma 8. *Let D and L be two disjoint subgraphs of G such that $D \cong F_4$ and $L \cong C_5$. If $e(D - x_3, L) \geq 13$, then $[D, L]$ contains either $2C_5$ or $F \uplus C_5$ or $B \uplus C_5$.*

Proof. Let $L = y_1y_2y_3y_4y_5y_1$ and $R = V(D) \setminus \{x_3\}$. Suppose to the contrary that $[D, L]$ does not contain any of $2C_5$, $F \uplus C_5$ and $B \uplus C_5$. Without loss of generality assume that $e(x_1x_2, L) \geq e(x_4x_5, L)$. It is clear that $7 \leq e(x_1x_2, L) \leq 10$ and, hence $2 \leq e(x_1, L) \leq 5$. If $e(x_1, L) \geq 4$, say $N(x_1) \mid L \supseteq \{y_1, y_2, y_3, y_4\}$, then $e(y_i, x_2x_4x_5) \leq 1$ as $x_1 \rightarrow (L, y_i)$ for all $i \in \{2, 3, 5\}$. Further, we have $e(y_j, x_2x_4x_5) \leq 2$ for all $j \in \{1, 4\}$, for otherwise, $[D - x_1 + y_j]$ would contain C_5 and $[L - y_j + x_1]$ contain F . Hence $e(R, L) \leq 12$, a contradiction. Therefore, $e(x_1, L) \leq 3$.

Case 1. $e(x_1, L) = 2$. In this case, $e(x_2, L) = 5$ as $e(x_1x_2, L) \geq 7$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2\}$. We claim that $e(y_iy_{i+1}, x_4x_5) \leq 2$ for all $i \in \{2, 3, 4, 5\}$, for otherwise, $[(V(L) \cup \{x_1, x_2\}) \setminus \{y_i, y_{i+1}\}] \supseteq C_5$ and $[x_3, x_4, x_5, y_i, y_{i+1}] \supseteq F$. Then $e(y_1, x_4x_5) = e(y_4, x_4x_5) = 2$ as $e(x_4x_5, L) \geq 6$. It follows that $[y_2, y_3, x_1, x_2, x_3] \supseteq F$ and $[x_4, x_5, y_1, y_4, y_5] \supseteq C_5$, a contradiction. Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_3\}$. We say that $e(y_1y_2, x_4x_5) \leq 2$ for otherwise $[x_1, x_2, y_5, y_4, y_3] \supseteq C_5$ and $[x_3, x_4, x_5, y_1, y_2] \supseteq F$. Similarly, $e(y_2y_3, x_4x_5) \leq 2$ and $e(y_4y_5, x_4x_5) \leq 2$. Then $e(y_1, x_4x_5) = e(y_3, x_4x_5) = 2$ as $e(x_4x_5, L) \geq 6$. Further, we have $e(y_2, x_4x_5) = 0$ as $x_1 \rightarrow (L, y_2)$. Then $e(y_4y_5, x_4x_5) = 2$ and so $[D, L] \supseteq F \uplus C_5$, a contradiction.

Case 2. $e(x_1, L) = 3$. In this case, $e(x_2, L) \geq 4$ as $e(x_1x_2, L) \geq 7$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L, y_2)$, $e(y_2, x_4x_5) \leq 1$. Further, if $y_2x_2 \in E(G)$, then $e(y_2, x_4x_5) = 0$. If $N(x_2) \mid L \supseteq \{y_4, y_5\}$, then $[y_1, y_2, y_5, x_1, x_2] \supseteq C_5$ and $[y_2, y_3, y_4, x_1, x_2] \supseteq C_5$. Further, $e(y_1y_5, x_4x_5) \leq 2$ and $e(y_3y_4, x_4x_5) \leq 2$ for otherwise $[y_1, y_5, x_3, x_4, x_5] \supseteq F$ and $[y_3, y_4, x_3, x_4, x_5] \supseteq F$. Hence $e(R, L) = e(x_1, L) + e(x_2, L) + e(y_2, x_4x_5) + e(y_1y_5, x_4x_5) + e(y_3y_4, x_4x_5) \leq 12$, no matter whether y_2x_2 is an edge in $E(G)$ or not, a contradiction. Then $N(x_2) \mid L \not\supseteq \{y_4, y_5\}$. Without loss of generality, we assume $N(x_2) \mid L = \{y_1, y_2, y_3, y_4\}$. Then $[y_4, y_5, y_1, x_1, x_2] \supseteq C_5$ and $[y_2, y_3, y_4, x_2, x_1] \supseteq C_5$. Hence $e(y_1y_5, x_4x_5) \leq 2$ and $e(y_2y_3, x_4x_5) \leq 2$. Then $e(y_4, x_4x_5) = 2$ as $e(x_4x_5, L) \geq 6$. This implies that $[L - y_4 + x_1] \supseteq F$ and $[D - x_1 + y_4] \supseteq C_5$, a contradiction. Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_4\}$. As $x_1 \rightarrow (L, y_i)$ for all $i \in \{3, 5\}$, $e(y_i, x_2x_4x_5) \leq 1$. Then $e(y_j, x_2x_4x_5) = 3$ for some $j \in \{1, 2\}$. It follows that $[L - y_j + x_1] \supseteq F$ and $[D + y_j - x_1] \supseteq C_5$, a contradiction. ■

Lemma 9. *Let D and L be two disjoint subgraphs of G such that $D \cong F_5$ and $L \cong C_5$. If $e(D - x_2, L) \geq 13$, then $[D, L]$ contains either $2C_5$ or $F_1 \uplus C_5$ or $K_{2,3} \uplus C_5$.*

Proof. Let $L = y_1y_2y_3y_4y_5y_1$ and $R = V(D) \setminus \{x_2\}$. Suppose to the contrary that $[D, L]$ does not contain any of $2C_5$, $F_1 \uplus C_5$ and $K_{2,3} \uplus C_5$. If $e(x_1, L) \geq 4$, say $N(x_1) \mid L \supseteq \{y_1, y_2, y_3, y_4\}$, then $e(y_i, x_3x_4x_5) \leq 1$ for all $i \in \{2, 3, 5\}$ as $x_1 \rightarrow (L, y_i)$. Further, $e(y_4, x_3x_4x_5) \leq 1$ as $[x_1, y_1, y_2, y_3, y_5] \supseteq F_1$. Thus, $e(R, L) \leq 12$, a contradiction. Now suppose $e(x_1, L) = 1$, say $x_1y_1 \in E(G)$. Suppose that $x_3y_j \in E(G)$ for some $j \in \{2, 5\}$, then $[x_1, y_1, y_j, x_3, x_2] \supseteq C_5$. It follows that $e(x_4x_5, L - \{y_1, y_j\}) \leq 3$ for otherwise $[(L - \{y_1, y_j\}) \cup \{x_4, x_5\}]$ would contain C_5 or F_1 or $K_{2,3}$, a contradiction. Then $e(x_3, L) \geq e(R, L) - e(x_4x_5, L - \{y_1, y_j\}) - 4 - e(x_1, L) \geq 5$. Therefore, either $e(x_3, L) = 5$ or $e(x_3, L) \leq 3$. Similarly, either $e(x_5, L) = 5$ or $e(x_5, L) \leq 3$. If $e(x_3, L) \leq 3$ and $e(x_5, L) \leq 3$, then $e(R, L) \leq 12$, a contradiction. Without loss of generality assume that $e(x_3, L) = 5$. If $e(x_i, y_1) = 1$, $e(x_j, y_2y_5) = 2$ for some $i \in \{3, 4, 5\}$, $j \in \{3, 5\}$, $i \neq j$, then $[L - y_1 + x_j] \supseteq C_5$ and $[D + y_1 - x_j] \supseteq C_5$. This implies that $N(x_4) \mid L = \{y_2, y_3, y_4, y_5\}$ and $N(x_5) \mid L \supseteq \{y_3, y_4\}$ and so $[x_1, x_2, x_3, y_1, y_2] \supseteq C_5$ and $[x_4, x_5, y_3, y_4, y_5] \supseteq C_5$, a contradiction.

Case 1. $e(x_1, L) = 2$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2\}$. If $e(x_4, L) = 5$, then $e(x_3x_5, y_i) \leq 1$ for all $i \in \{1, 2\}$, for otherwise, $[L - y_i + x_4] \supseteq C_5$ and $[D + y_i - x_4] \supseteq K_{2,3}$. If $e(x_3x_5, y_3y_4y_5) \geq 5$, then $[x_1, x_2, x_4, y_1, y_2] \supseteq C_5$ and $[x_3, x_5, y_3, y_4, y_5] \supseteq C_5$, a contradiction. Thus, $e(x_3x_5, y_3y_4y_5) = 4$, $e(x_3x_5, y_i) = 1$ for all $i \in \{1, 2\}$. Further, either $[x_1, x_2, x_3, y_1, y_2] \supseteq C_5$ and $[x_4, x_5, y_3, y_4, y_5] \supseteq C_5$, or $[x_1, x_2, x_5, y_1, y_2] \supseteq C_5$ and $[x_3, x_4, y_3, y_4, y_5] \supseteq C_5$, a contradiction. Therefore $e(x_4, L) \leq 4$. If $e(x_4, y_1y_2) = 2$, then $e(y_{i+1}y_{i-1}, x_j) \leq 1$ for all $i \in \{1, 2\}$, $j \in \{3, 5\}$, for otherwise, $[L - y_i + x_j] \supseteq C_5$, $[D + y_i - x_j] \supseteq C_5$. Hence $e(R, L) \leq 12$, a contradiction. Therefore, $e(x_4, y_1y_2) \leq 1$. Without loss of generality assume that $e(x_3, L) \geq e(x_5, L)$. If $e(x_4, L) = 4$, say $N(x_4) \mid L = \{y_2, y_3, y_4, y_5\}$, then $[D + y_2 - x_3]$ would contain C_5 and $[L - y_2 + x_3]$ contain F_1 or C_5 , a contradiction. Therefore $e(x_4, L) \leq 3$ and hence $e(x_3x_5, L) \geq 8$. If $e(x_3, L) = 5$, then $e(x_i, y_j) = 0$ for all $i \in \{4, 5\}$, $j \in \{1, 2\}$, for otherwise, $[L - y_j + x_3] \supseteq C_5$, $[D + y_j - x_3] \supseteq C_5$. Thus $e(x_4x_5, y_3y_4y_5) = 6$. Then $[x_1, x_2, x_3, y_1, y_2] \supseteq C_5$ and $[x_4, x_5, y_3, y_4, y_5] \supseteq C_5$, a contradiction. Hence $e(x_3, L) = e(x_5, L) = 4$ and $e(x_4, L) = 3$. If $x_4y_1 \in E(G)$, then $e(x_3x_5, y_2y_5) \leq 2$ and so $e(x_3x_5, y_1y_3y_4) = 6$. This implies that $[x_1, x_2, x_3, y_2, y_3] \supseteq C_5$ and $[x_4, x_5, y_1, y_4, y_5] \supseteq C_5$, a contradiction. Therefore, $x_4y_1 \notin E(G)$. Similarly, $x_4y_2 \notin E(G)$. Then $e(x_4, y_3y_4y_5) = 3$. We say that $e(x_3x_5, y_2y_3) \leq 3$ for otherwise $[x_1, y_1, y_2, x_3, x_2] \supseteq C_5$ and $[x_4, x_5, y_3, y_4, y_5] \supseteq C_5$. Then $e(x_3x_5, y_1y_4y_5) \geq 5$ and so $[x_1, x_2, x_4, y_2, y_3] \supseteq C_5$ and $[x_3, x_5, y_1, y_4, y_5] \supseteq C_5$, a contradiction.

Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_3\}$. As $x_1 \rightarrow (L, y_2)$, $e(y_2, x_3x_5) = 0$. If $e(x_4, L) \leq 3$, then $e(x_3x_5, L - y_2) = 8$ and so $[L - y_1 + x_5] \supseteq F_1$ and $[D - x_5 + y_1] \supseteq C_5$, a contradiction. Without loss of generality assume that $e(x_3, L) \geq e(x_5, L)$. If $e(x_4, L) = 4$, then $N(x_3) \mid L = \{y_1, y_3, y_4, y_5\}$. We claim that $e(y_i, x_5) = 0$ for all $i \in \{1, 3\}$, for otherwise, $[D - x_3 + y_i] \supseteq C_5$, $[L - y_i + x_3] \supseteq F_1$. Hence $e(R, L) \leq 12$, a contradiction. If $e(x_4, L) = 5$, then $e(x_3x_5, y_1y_5y_4) \leq 4$ for otherwise $[x_1, y_2, y_3, x_4, x_2] \supseteq C_5$ and $[x_3, x_5, y_1, y_5, y_4] \supseteq C_5$. It follows that $e(x_3x_5, y_3) = 2$ and so $[L - y_3 + x_4] \supseteq C_5$ and $[D + y_3 - x_4] \supseteq K_{2,3}$, again a contradiction.

Case 2. $e(x_1, L) = 3$. First suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L, y_2)$, $e(y_2, x_3x_4x_5) \leq 1$. Further, $e(y_4, x_3x_4x_5) \leq 1$ as $[x_1, y_1, y_2, y_3, y_5] \supseteq F_1$. Similarly, $e(y_5, x_3x_4x_5) \leq 1$. Thus, $e(R, L) \leq 12$, a contradiction. Next, suppose that $N(x_1) \mid L = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L = \{y_1, y_2, y_4\}$. As $x_1 \rightarrow (L, y_i)$ for all $i \in \{3, 5\}$, $e(y_i, x_3x_5) = 0$. Then $e(x_3x_5, L) \leq 6$ and so $e(x_4, L) \geq 4$. If $e(x_4, L) = 5$, then $e(y_i, x_3x_5) \leq 1$ for all $i \in \{1, 2, 4\}$, for otherwise, $[D - x_4 + y_i] \supseteq K_{2,3}$ and $[L - y_i + x_4] \supseteq C_5$. Hence $e(R, L) \leq 11$, a contradiction. If $e(x_4, L) = 4$, then $e(x_3x_5, y_1y_2y_4) = 6$. Further, if $e(x_4, y_3y_5) = 2$, then $[x_1, x_2, x_4, y_2, y_3] \supseteq C_5$ and $[x_3, x_5, y_1, y_4, y_5] \supseteq K_{2,3}$, a contradiction. Thus $e(x_4, y_3y_5) \leq 1$. Without loss of generality assume that $N(x_4) \mid L = \{y_1, y_2, y_3, y_4\}$. Then $[y_2, y_3, x_1, x_2, x_4] \supseteq C_5$ and $[y_1, y_4, y_5, x_3, x_5] \supseteq K_{2,3}$, again a contradiction. ■

3. PROOF OF THEOREM 3

Let G be a graph of order $n = 5k$ with $\sigma_2(G) \geq n + k$. It is easy to see that G is Hamiltonian if $k = 1$. In the following, we always assume that $k \geq 2$. Suppose, for a contradiction, that $G \not\supseteq kC_5$. We may assume that G is maximal, i.e., $G + xy \supseteq kC_5$ for each pair of non-adjacent vertices x and y of G . Thus $G \supseteq P_5 \uplus (k-1)C_5$. Our proof will follow from the following lemmas.

Lemma 10. *For each $s \in \{1, 2, \dots, k\}$, $G \not\supseteq sB \uplus (k-s)C_5$.*

Proof. To the contrary, suppose that $G \supseteq sB \uplus (k-s)C_5$ for some $s \in \{1, 2, \dots, k\}$. Let s be the minimum number in $\{1, 2, \dots, k\}$ for which $G \supseteq sB \uplus (k-s)C_5$ and let $B_1, \dots, B_s, L_1, \dots, L_{k-s}$ be k disjoint subgraphs of G with $B_i \cong B$ for $i \in \{1, 2, \dots, s\}$ and $L_i \cong C_5$ for $i \in \{1, 2, \dots, k-s\}$. Let R be the set of the four vertices of degree 2 in B_1 . By Lemmas 4(a) and (b) and the minimality of s , we see that $e(R, B_i) \leq 12$ and $e(R, L_j) \leq 12$ for all $i \in \{2, 3, \dots, s\}$ and $j \in \{1, 2, \dots, k-s\}$. Therefore $\sum_{x \in R} d_G(x) \leq 12(k-1) + 8 = 12k - 4$. However, by

the degree sum condition, we have $\sum_{x \in R} d_G(x) = (d_G(x_2) + d_G(x_4)) + (d_G(x_3) + d_G(x_5)) \geq 12k$, a contradiction. ■

Lemma 11. $G \supseteq F_1 \uplus (k-1)C_5$.

Proof. First, we claim that $G \supseteq F \uplus (k-1)C_5$. Let $\{H, L_1, L_2, \dots, L_{k-1}\}$ be an optimal family of G with $H \cong P_5$. If $[H] \cong P_5 = x_1x_2x_3x_4x_5$, then $d_G(x_1) + d_G(x_5) \geq 6k$. Without loss of generality, we assume that $d_G(x_1) \geq 3k$. Since $\sigma_2(G) \geq 6k$, $\sum_{x \in V(H)} d_G(x) = d_G(x_1) + (d_G(x_2) + d_G(x_4)) + (d_G(x_3) + d_G(x_5)) \geq 15k$. Then $e(H, G - V(H)) \geq 15k - 8 = 15(k-1) + 7$. Thus $e(H, L_i) \geq 16$ for some $i \in \{1, 2, \dots, k-1\}$. By Lemma 4(d), $[H, L_i] \supseteq F \uplus C_5$ and, hence $G \supseteq F \uplus (k-1)C_5$. If $[H] \cong F_4$, then $e(x_1x_2x_4x_5, G - V(H)) \geq 12k - 8 = 12(k-1) + 4$. This implies that $e(x_1x_2x_4x_5, L_i) \geq 13$ for some $i \in \{1, 2, \dots, k-1\}$. By Lemma 8 and Lemma 10, $[H, L_i] \supseteq F \uplus C_5$ and so $G \supseteq F \uplus (k-1)C_5$. If $[H] \cong F_3$, then $e(x_1x_2x_4x_5, G - V(H)) \geq 12k - 7 = 12(k-1) + 5$. Hence $e(x_1x_2x_4x_5, L_i) \geq 13$ for some $i \in \{1, 2, \dots, k-1\}$. By Lemmas 7, 8 and 10, $[H, L_i] \supseteq F \uplus C_5$ and, therefore, $G \supseteq F \uplus (k-1)C_5$.

We first assume that $G \supseteq F_2 \uplus (k-1)C_5$ and let $\{H, L_1, L_2, \dots, L_{k-1}\}$ be a family of G with $H \cong F_2$. Note that $d_G(x_1) + d_G(x_5) \geq 6k$, say $d_G(x_1) \geq 3k$. Then $e(x_1x_3x_5, G - V(H)) = d_G(x_1) + (d_G(x_3) + d_G(x_5)) - 6 \geq 9k - 6 = 9(k-1) + 3$ and, hence $e(x_1x_3x_5, L_i) \geq 10$ for some $i \in \{1, 2, \dots, k-1\}$. So by Lemma 4(c), $[H, L_i] \supseteq F_1 \uplus C_5$ and, therefore, $G \supseteq F_1 \uplus (k-1)C_5$. We now assume that $G \not\supseteq F_2 \uplus (k-1)C_5$. Recalling that $G \supseteq F \uplus (k-1)C_5$, let $\{H, L_1, L_2, \dots, L_{k-1}\}$ be an optimal family of G with $H \cong F$. If $[H] \cong F$ or $[H] \cong K_{2,3}$, say $\{x_1, x_2, x_3, x_4, x_5\} = V(H)$ with $x_1x_3, x_1x_5, x_3x_5 \notin E(G)$ and $x_2x_4 \notin E(G)$, then $d_G(x_1) + d_G(x_5) \geq 6k$, say $d_G(x_1) \geq 3k$. Further, $e(H, G - V(H)) \geq d_G(x_1) + (d_G(x_3) + d_G(x_5)) + (d_G(x_2) + d_G(x_4)) - 12 \geq 15k - 12 = 15(k-1) + 3$. Then $e(H, L_i) \geq 16$ for some $i \in \{1, 2, \dots, k-1\}$. By Lemma 5(a) and Lemma 10, the cycle L_i has a labelling $L_i = y_1y_2y_3y_4y_5y_1$ satisfying property **P**₁. Therefore, $e(x_1x_3y_3y_5, G - V(H \cup L_i)) \geq 12k - 17 = 12(k-2) + 7$ and, hence, $e(x_1x_3y_3y_5, L_j) \geq 13$ for some $j \in \{1, \dots, k-1\} \setminus \{i\}$. So by Lemma 5(b), we have $[H, L_i, L_j] \supseteq F_1 \uplus 2C_5$ and, hence, $G \supseteq F_1 \uplus (k-1)C_5$. We next assume that $G \not\supseteq K_{2,3} \uplus (k-1)C_5$. If $[H] \cong F_5$, then $e(x_1x_3x_4x_5, G - V(H)) = (d_G(x_1) + d_G(x_4)) + (d_G(x_3) + d_G(x_5)) - 8 \geq 12k - 8 = 12(k-1) + 4$. Thus $e(x_1x_3x_4x_5, L_i) \geq 13$ for some $i \in \{1, 2, \dots, k-1\}$. So by Lemma 9, $[H, L_i] \supseteq F_1 \uplus C_5$, i.e., $G \supseteq F_1 \uplus (k-1)C_5$. ■

Lemma 12. Let $\psi = \{H, L_1, L_2, \dots, L_{k-1}\}$ be an optimal family of G with $H \cong F_1$ and let $T = \{x_2, x_4, x_5\}$. If $G \not\supseteq K_4^+ \uplus (k-1)C_5$, then for each $t \in \{1, 2, \dots, k-1\}$, the following statements hold.

- (a) If $e(x_1, L_t) = 5$, then $e(T, L_t) \leq 5$.
- (b) If $e(x_1, L_t) = 4$, then $e(T, L_t) \leq 7$.

- (c) If $e(x_1, L_t) = 3$, then $e(T, L_t) \leq 9$.
- (d) If $e(x_1, L_t) = 2$, then $e(T, L_t) \leq 11$.
- (e) If $e(x_1, L_t) = 1$, then $e(T, L_t) \leq 12$.
- (f) If $e(x_1, L_t) = 0$, then $e(T, L_t) \leq 15$.

Proof. Let $L_t = y_1y_2y_3y_4y_5y_1$ and $G_t = [H, L_t]$. If $e(x_1, L_t) = 5$, then $e(y_i, T) \leq 1$ for all $i \in \{1, 2, 3, 4, 5\}$ since $G_t \not\supseteq 2C_5$. Hence, (a) follows directly.

To prove (b), without loss of generality assume that $N(x_1) \mid L_t = \{y_1, y_2, y_3, y_4\}$. Suppose to the contrary that $e(T, L_t) \geq 8$. It is clear that $\tau(y_5, L_t) = 0$ for otherwise $x_1 \rightarrow L_t$ and so $G_t \supseteq 2C_5$. As $x_1 \rightarrow (L_t, y_i)$ for all $i \in \{2, 3, 5\}$, $e(y_i, T) \leq 1$. Hence, $e(y_1y_4, T) \geq 5$, say $e(y_4, T) = 3$ and $e(y_1, T) \geq 2$. If $e(y_5, x_2x_4) \geq 1$, then $[H - x_1 + y_5] \supseteq F_1$ and $\tau(L_t - y_5 + x_1) > \tau(L_t)$, contradicting the optimality of ψ . Thus, $e(y_5, x_2x_4) = 0$. If $y_5x_5 \in E(G)$, then $y_1x_2 \notin E(G)$ for otherwise $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$ and $[y_4, y_5, x_3, x_4, x_5] \supseteq C_5$. This means that $e(y_1, x_4x_5) = 2$ and so $[y_2, y_3, y_4, x_1, x_2] \supseteq C_5$ and $[y_1, y_5, x_3, x_4, x_5] \supseteq C_5$, a contradiction. Therefore, $y_5x_5 \notin E(G)$ and hence $e(y_1, T) = 3$, $e(y_2y_3, T) = 2$. If $e(y_2y_3, x_5) = 2$, then $[y_1, y_5, y_4, x_1, x_2] \supseteq C_5$ and $[y_2, y_3, x_3, x_4, x_5] \supseteq B$, contradicting Lemma 10. Hence, $e(y_2y_3, x_2x_4) \geq 1$, say $y_3x_2 \in E(G)$. We claim that $y_1y_3 \in E(G)$ for otherwise $[y_3, x_2, x_3, x_4, x_5] \supseteq F_1$ and $\tau(y_1y_2x_1y_4y_5y_1) > \tau(L_t)$. This implies that $[y_5, y_1, y_2, y_3, x_1] \supseteq K_4^+$ and $[y_4, x_2, x_3, x_4, x_5] \supseteq C_5$, a contradiction.

To prove (c), suppose to the contrary that $e(T, L_t) \geq 10$. Assume first $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L_t, y_2)$, $e(y_2, T) \leq 1$. For any $i \in \{1, 2, 3, 4, 5\}$, if $[(V(L_t) \cup \{x_1, x_2\}) \setminus \{y_i, y_{i+1}\}] \supseteq C_5$, then $e(y_iy_{i+1}, x_4x_5) \leq 2$ for otherwise $[x_3, x_4, x_5, y_i, y_{i+1}] \supseteq C_5$. If $e(x_2, y_4y_5) = 2$, then $e(T, L_t) = e(x_2, L_t) + e(x_4x_5, y_2) + e(x_4x_5, y_3y_4) + e(x_4x_5, y_1y_5) \leq 9$, no matter whether y_2x_2 is an edge of G or not, a contradiction. Hence $e(x_2, y_4y_5) \leq 1$, say $x_2y_4 \notin E(G)$. Further, if $e(x_2, y_3y_5) = 2$, then $e(x_4x_5, y_1y_2) \leq 2$ and $e(x_4x_5, y_4y_5) \leq 2$. Since $e(T, L_t) \geq 10$ and $x_2y_4 \notin E(G)$, $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_5\}$ and $e(y_3, x_4x_5) = 2$. Therefore, $[x_3, x_2, x_1, y_1, y_2] \supseteq K_4^+$. Then $e(y_5, x_4x_5) = 0$ as $G_t \not\supseteq K_4^+ \uplus C_5$. Therefore, $e(y_4, x_4x_5) = 2$. Then $[x_1, x_2, y_1, y_2, y_5] \supseteq C_5$ and $[x_3, x_4, x_5, y_3, y_4] \supseteq C_5$, a contradiction. Therefore, $e(x_2, y_3y_5) \leq 1$. If $x_2y_5 \in E(G)$, then $e(x_4x_5, y_1y_2) \leq 2$, $e(x_4x_5, y_3y_4) \leq 2$ and so $e(T, L_t) \leq 9$, a contradiction. Thus $x_2y_5 \notin E(G)$. If $x_2y_1 \in E(G)$, then $e(x_4x_5, y_4y_5) \leq 2$ and so $e(T, L_t) = e(x_2, L_t) + e(x_4x_5, y_2) + e(x_4x_5, y_1y_3) + e(x_4x_5, y_4y_5) \leq 9$, no matter whether y_2x_2 is an edge of G or not, a contradiction. Hence $x_2y_1 \notin E(G)$. Similarly, $x_2y_3 \notin E(G)$. Then $e(T, L_t) \leq 9$, no matter whether y_2x_2 is an edge in $E(G)$ or not, again a contradiction.

Next, assume that $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_4\}$. As $x_1 \rightarrow (L_t, y_i)$ for all $i \in \{3, 5\}$, $e(y_i, T) \leq 1$. Further, if $e(x_2, L_t) \leq 1$ or $e(x_2, L_t) = 2$, $N(x_2) \mid L_t \cap \{y_3, y_5\} \neq \emptyset$ or $e(x_2, L_t) = 3$,

$N(x_2) \mid L_t \supseteq \{y_3, y_5\}$, then $e(T, L_t) \leq 9$, a contradiction. Suppose $e(x_2, L_t) \leq 3$. If $x_2y_4 \in E(G)$, then $e(y_1y_5, x_4x_5) \leq 2$, $e(y_2y_3, x_4x_5) \leq 2$ and so $e(T, L_t) \leq 9$, a contradiction. Hence $x_2y_4 \notin E(G)$. Then $N(x_2) \mid L_t \supseteq \{y_1, y_2\}$. It follows that $e(y_1y_5, x_4x_5) \leq 2$, $e(y_2y_3, x_4x_5) \leq 2$ and so $e(T, L_t) \leq 9$, again a contradiction.

Now suppose $e(x_2, L_t) \geq 4$. If $y_3y_5 \in E(G)$, then $x_1 \rightarrow L_t$. Since $e(T, L_t) \geq 10$, there is a vertex y_i for some $i \in \{1, 2, 3, 4, 5\}$ such that $e(y_i, T) \geq 2$. Then $[H - x_1 + y_i] \supseteq C_5$ and $[L_t - y_i + x_1] \supseteq C_5$, a contradiction. Thus $y_3y_5 \notin E(G)$. If $x_2y_3 \in E(G)$ and $y_1y_3 \notin E(G)$, then $[y_3, x_2, x_3, x_4, x_5] \cong F_1$ and $\tau(L_t) < \tau(L_t - y_3 + x_1)$, a contradiction. Therefore, if $x_2y_3 \in E(G)$, then $y_1y_3 \in E(G)$. Similarly, if $x_2y_5 \in E(G)$, then $y_2y_5 \in E(G)$. We claim that $e(y_1y_2y_4, x_4x_5) = 6$ if $|N(x_2) \mid L_t| = 4$. If $x_2y_3 \in E(G)$ and $x_2y_5 \in E(G)$, then $e(y_1y_2y_4, x_4x_5) = 6$ since $e(y_i, T) \leq 1$ for all $i \in \{3, 5\}$. Without loss of generality, assume that $x_2y_3 \in E(G)$ and $x_2y_5 \notin E(G)$. If $e(y_5, x_4x_5) = 0$, then $e(y_1y_2y_4, x_4x_5) = 6$ as $e(y_3, T) \leq 1$. If $e(y_5, x_4x_5) = 2$, then $[x_1, y_1, y_2, y_3, y_4] \supseteq C_5$ and $[y_5, x_2, x_3, x_4, x_5] \supseteq C_5$, a contradiction. If $e(y_5, x_4x_5) = 1$, then $e(y_4, x_4x_5) = 0$ for otherwise $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$ and $[y_4, y_5, x_3, x_4, x_5]$ would contain C_5 or B . Therefore, $e(T, L_t) \leq 9$, a contradiction. If $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4\}$, then $[y_4, y_5, x_3, x_4, x_5] \supseteq F_1$. Further, $\tau(L_t) = 4$ as $\tau(L_t) \geq \tau(y_1y_2y_3x_2x_1y_1)$. Similarly, if $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_5\}$ or $N(x_2) \mid L_t = \{y_1, y_2, y_4, y_5\}$, then we also have $\tau(L_t) = 4$. If $N(x_2) \mid L_t = \{y_2, y_3, y_4, y_5\}$, then $[x_1, y_1, x_3, x_4, x_5] \supseteq F_1$. Further, $\{y_1y_3, y_1y_4, y_2y_5\} \subseteq E(G)$ as $\tau(L_t) \geq \tau(y_2y_3y_4y_5x_2y_2)$. If $N(x_2) \mid L_t = \{y_1, y_3, y_4, y_5\}$, then $[x_1, y_2, x_3, x_4, x_5] \supseteq F_1$. Further, $\{y_1y_3, y_2y_4, y_2y_5\} \subseteq E(G)$ as $\tau(L_t) \geq \tau(y_1x_2y_3y_4y_5y_1)$. If $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4, y_5\}$, then $e(y_1y_2y_4, x_4x_5) \geq 5$ since $e(y_i, T) \leq 1$ for all $i \in \{3, 5\}$. We claim that $\tau(L_t) = 4$ if $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4, y_5\}$. If $e(y_4, x_4x_5) = 2$, then $[x_1, y_4, x_3, x_4, x_5] \supseteq F_1$. Further, $\tau(L_t) = 4$ as $\tau(L_t) \geq \tau(y_1y_2y_3x_2y_5y_1)$. If $e(y_4, x_4x_5) \leq 1$, then $e(y_1y_2, x_4x_5) = 4$ and so $[x_1, y_1, x_3, x_4, x_5] \supseteq F_1$ and $[x_1, y_2, x_3, x_4, x_5] \supseteq F_1$. Further, $\tau(L_t) = 4$ as $\tau(L_t) \geq \tau(x_2y_2y_3y_4y_5x_2)$ and $\tau(L_t) \geq \tau(y_1x_2y_3y_4y_5y_1)$.

Let $R = \{x_1, x_3, y_3, y_5\}$. If $x_3y_3 \in E(G)$, then $x_2y_3 \notin E(G)$ and so $N(x_2) \mid L_t = \{y_1, y_2, y_4, y_5\}$. Then $[y_1, y_4, y_5, x_1, x_2] \supseteq C_5$ and $[y_2, y_3, x_3, x_4, x_5] \supseteq C_5$, a contradiction. Thus $x_3y_3 \notin E(G)$. Similarly, $x_3y_5 \notin E(G)$. Hence R is an independent set. As $e(R, G - V(G_t)) \geq 12k - 18 = 12(k - 2) + 6$, $e(R, L_i) \geq 13$ for some $i \in \{1, 2, \dots, k - 1\} \setminus \{t\}$.

Claim 1. *If $u \rightarrow (L_i, z; \{v, w\})$ for some $z \in V(L_i)$, $u \in R$ and $\{v, w\} \subseteq R \setminus \{u\}$, then $[G_t, L_i] \supseteq 3C_5$.*

Proof. We separate the proof into two cases.

Case 1. $e(x_2, y_3y_5) = 2$. In this case, $e(x_4x_5, y_3y_5) = 0$. Further, $e(x_4x_5, y_1y_2y_4) = 6$ if $|N(x_2) \mid L_t| = 4$ and $e(x_4x_5, y_1y_2y_4) \geq 5$ if $|N(x_2) \mid L_t| = 5$. Recall that $\tau(L_t) = 4$ if $|N(x_2) \mid L_t| = 5$. Hence, y_1, y_2 and y_4 are symmetric in $[L_t]$ if $|N(x_2) \mid L_t| = 5$. Without loss of generality, assume that $e(x_4x_5, y_1y_4) = 4$.

Further, by the symmetry of x_1 , y_3 and y_5 in $[L_t + x_1]$, we need only to consider the following cases. If $x_1 \rightarrow (L_i, z; \{y_3, y_5\})$ for some $z \in V(L_i)$, then $[L_i - z + x_1] \supseteq C_5$, $[y_1, y_2, y_3, y_5, z] \supseteq C_5$ and $[x_2, x_3, x_4, x_5, y_4] \supseteq C_5$. If $x_1 \rightarrow (L_i, z; \{y_3, x_3\})$ for some $z \in V(L_i)$, then $[L_i - z + x_1] \supseteq C_5$, $[y_3, x_2, x_5, x_3, z] \supseteq C_5$ and $[y_1, y_2, y_4, y_5, x_4] \supseteq C_5$. If $x_3 \rightarrow (L_i, z; \{y_3, y_5\})$ for some $z \in V(L_i)$, then $[L_i - z + x_3] \supseteq C_5$, $[y_1, y_2, y_3, y_5, z] \supseteq C_5$ and $[x_1, x_2, x_4, x_5, y_4] \supseteq C_5$.

Case 2. $e(x_2, y_3 y_5) = 1$, say $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4\}$. In this case, $e(x_4 x_5, y_1 y_2 y_4) = 6$ and $e(x_4 x_5, y_3) = 0$. Further, $e(x_4 x_5, y_5) = 0$ for otherwise $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$ and $[y_4, y_5, x_3, x_4, x_5]$ would contain C_5 or B . By the symmetry of x_1 and y_3 in $[L_t + x_1]$, we need only to consider the following cases. If $x_1 \rightarrow (L_i, z; \{y_3, y_5\})$ for some $z \in V(L_i)$, then $[L_i - z + x_1] \supseteq C_5$, $[y_1, y_2, y_3, y_5, z] \supseteq C_5$ and $[x_2, x_3, x_4, x_5, y_4] \supseteq C_5$. If $x_1 \rightarrow (L_i, z; \{y_3, x_3\})$ for some $z \in V(L_i)$, then $[L_i - z + x_1] \supseteq C_5$, $[y_3, x_2, x_5, x_3, z] \supseteq C_5$ and $[y_1, y_2, y_4, y_5, x_4] \supseteq C_5$. If $x_1 \rightarrow (L_i, z; \{y_5, x_3\})$ for some $z \in V(L_i)$, then $[L_i - z + x_1] \supseteq C_5$, $[y_4, y_5, z, x_2, x_3] \supseteq C_5$ and $[y_1, y_2, y_3, x_4, x_5] \supseteq C_5$.

If $x_3 \rightarrow (L_i, z; \{y_3, y_5\})$ for some $z \in V(L_i)$, then $[L_i - z + x_3] \supseteq C_5$, $[y_1, y_2, y_3, z, y_5] \supseteq C_5$ and $[x_1, x_2, x_4, x_5, y_4] \supseteq C_5$. If $x_3 \rightarrow (L_i, z; \{y_3, x_1\})$ for some $z \in V(L_i)$, then $[L_i - z + x_3] \supseteq C_5$, $[x_1, y_2, x_2, y_3, z] \supseteq C_5$ and $[y_1, y_5, y_4, x_4, x_5] \supseteq C_5$. If $y_5 \rightarrow (L_i, z; \{x_3, x_1\})$ for some $z \in V(L_i)$, then $[L_i - z + y_5] \supseteq C_5$, $[x_1, x_2, x_3, x_5, z] \supseteq C_5$ and $[y_1, y_2, y_3, y_4, x_4] \supseteq C_5$. If $y_5 \rightarrow (L_i, z; \{y_3, x_1\})$ for some $z \in V(L_i)$, then $[L_i - z + y_5] \supseteq C_5$, $[y_1, y_2, y_3, x_1, z] \supseteq C_5$ and $[x_2, x_3, x_4, x_5, y_4] \supseteq C_5$. \square

By Claim 1 and Lemma 4(e), there are vertex labellings $R = \{a_1, a_2, a_3, a_4\}$ and $L_i = b_1 b_2 b_3 b_4 b_5 b_1$ such that $e(a_1 a_2, b_1 b_2 b_3 b_4) = 8$, $e(a_3, b_1 b_5 b_4) = 3$ and $e(a_4, b_1 b_4) = 2$. Recall that $e(x_4 x_5, y_1 y_2 y_4) = 6$ if $|N(x_2) \mid L_t| = 4$, and $e(x_4 x_5, y_1 y_2 y_4) \geq 5$ and $\tau(L_t) = 4$ if $|N(x_2) \mid L_t| = 5$. Assume first that $|N(x_2) \mid L_t| = 5$. Then x_1, y_3 and y_5 are symmetric in $[L_t + x_1]$. Further, y_1, y_2 and y_4 are symmetric in $[L_t]$. Without loss of generality, assume that $e(x_4 x_5, y_1 y_4) = 4$. If $x_3 \in \{a_1, a_2\}$, say $\{x_3, y_3\} = \{a_1, a_2\}$, then $[x_1, b_1, y_5, b_4, b_5] \supseteq C_5$, $[x_2, x_3, x_4, x_5, y_1] \supseteq C_5$ and $[b_2, b_3, y_2, y_3, y_4] \supseteq B$, a contradiction. If $x_3 \notin \{a_1, a_2\}$, say $\{x_1, y_3\} = \{a_1, a_2\}$, then $[x_3, y_5, b_1, b_5, b_4] \supseteq C_5$, $[x_1, x_2, x_4, x_5, y_1] \supseteq C_5$ and $[b_2, b_3, y_2, y_3, y_4] \supseteq B$, again a contradiction.

Next, assume that $|N(x_2) \mid L_t| = 4$. If $e(x_2, y_3 y_5) = 2$, then x_1, y_3 and y_5 are symmetric in $[L_t + x_1]$. If $x_3 \in \{a_1, a_2\}$, say $\{x_3, y_3\} = \{a_1, a_2\}$, then $[x_1, b_1, y_5, b_4, b_5] \supseteq C_5$, $[x_2, x_3, x_4, x_5, y_1] \supseteq C_5$ and $[b_2, b_3, y_2, y_3, y_4] \supseteq B$, or $[x_1, b_1, y_5, b_4, b_5] \supseteq C_5$, $[x_2, x_3, x_4, x_5, y_2] \supseteq C_5$ and $[b_2, b_3, y_1, y_3, y_4] \supseteq B$, a contradiction. If $x_3 \notin \{a_1, a_2\}$, say $\{x_1, y_3\} = \{a_1, a_2\}$, then $[x_3, y_5, b_1, b_5, b_4] \supseteq C_5$, $[y_3, x_2, x_5, x_4, y_4] \supseteq C_5$ and $[b_2, b_3, x_1, y_1, y_2] \supseteq B$, again a contradiction. If $e(x_2, y_3 y_5) = 1$, then $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4\}$ or $N(x_2) \mid L_t = \{y_1, y_2, y_4, y_5\}$. Without loss of generality, assume that $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4\}$. By the symmetry of x_1 and y_3 in $[L_t + x_1]$, we need only to consider the following

cases. If $\{x_3, y_3\} = \{a_1, a_2\}$, then $[x_1, b_1, y_5, b_4, b_5] \supseteq C_5$, $[x_2, x_3, x_4, x_5, y_1] \supseteq C_5$ and $[b_2, b_3, y_2, y_3, y_4] \supseteq B$, a contradiction. If $\{x_3, y_5\} = \{a_1, a_2\}$, then $[x_1, b_1, y_3, b_4, b_5] \supseteq C_5$, $[x_2, x_3, x_4, x_5, y_1] \supseteq C_5$ and $[b_2, b_3, y_2, y_4, y_5] \supseteq B$, a contradiction. If $\{x_1, y_5\} = \{a_1, a_2\}$, then $[x_3, b_1, y_3, b_4, b_5] \supseteq C_5$, $[x_1, x_2, x_4, x_5, y_2] \supseteq C_5$ and $[b_2, b_3, y_1, y_4, y_5] \supseteq B$, a contradiction. If $\{x_1, y_3\} = \{a_1, a_2\}$, then $[x_3, b_1, y_5, b_4, b_5] \supseteq C_5$, $[x_1, x_2, x_4, x_5, y_4] \supseteq C_5$ and $[b_2, b_3, y_1, y_2, y_3] \supseteq B$, again a contradiction.

To prove (d), suppose to the contrary that $e(T, L_t) \geq 12$. Assume first $N(x_1) \mid L_t = \{y_i, y_{i+1}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2\}$. For any $i \in \{1, 2, 3, 4, 5\}$, if $[(V(L_t) \cup \{x_1, x_2\}) \setminus \{y_i, y_{i+1}\}] \supseteq C_5$, then $e(y_i y_{i+1}, x_4 x_5) \leq 2$ for otherwise $[x_3, x_4, x_5, y_i, y_{i+1}] \supseteq C_5$. It follows that if $x_2 y_4 \in E(G)$, then $e(y_1 y_5, x_4 x_5) \leq 2$, $e(y_2 y_3, x_4 x_5) \leq 2$ and so $e(T, L_t) \leq 11$, a contradiction. Hence $x_2 y_4 \notin E(G)$. If $e(x_2, y_3 y_5) = 2$, then $e(y_3 y_4, x_4 x_5) \leq 2$, $e(y_4 y_5, x_4 x_5) \leq 2$ and so $e(y_1 y_2, x_4 x_5) = 4$ and $e(x_2, y_1 y_2 y_3 y_5) = 4$. Further, $y_3 x_5 \notin E(G)$ and $y_5 x_4 \notin E(G)$ as $G_t \not\supseteq 2C_5$. Moreover, $x_4 y_4, x_5 y_5 \in E(G)$ as $e(T, L_t) \geq 12$. Clearly, $G_t \supseteq 2C_5$, a contradiction. Therefore, $e(x_2, y_3 y_5) \leq 1$ and hence $e(x_2, y_1 y_2 y_3 y_5) \leq 3$. If $e(x_2, y_1 y_2 y_3 y_5) = 3$, say $x_2 y_3 \in E(G)$, then $e(x_4 x_5, y_4 y_5) \leq 2$. It follows that $e(T, L_t) \leq 11$, a contradiction. Hence $e(x_2, y_1 y_2 y_3 y_5) \leq 2$. Then $e(x_4 x_5, L_t) = 10$ and so $[x_1, y_1, x_4, y_3, y_2] \supseteq C_5$ and $[y_4, y_5, x_2, x_3, x_5] \supseteq B$, a contradiction. Next, assume that $N(x_1) \mid L_t = \{y_i, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_3\}$. As $x_1 \rightarrow (L_t, y_2)$, $e(y_2, T) \leq 1$. If $e(x_2, y_1 y_3) \geq 1$, then $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and so $e(x_4 x_5, y_4 y_5) \leq 2$. Then $e(T, L_t) = e(x_2, L_t) + e(x_4 x_5, y_4 y_5) + e(x_4 x_5, y_1 y_3) + e(x_4 x_5, y_2) \leq 11$, no matter whether $x_2 y_2 \in E(G)$ or $x_2 y_2 \notin E(G)$, a contradiction. Therefore, $e(x_2, y_1 y_3) = 0$ and hence $e(T, L_t) \leq 11$, again a contradiction.

To prove (e), suppose to the contrary that $e(T, L_t) \geq 13$. Without loss of generality assume that $x_1 y_1 \in E(G)$. If $e(x_2, y_3 y_4) \geq 1$, say $x_2 y_3 \in E(G)$, then $e(y_4 y_5, x_4 x_5) \leq 2$ for otherwise $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$ and $[x_3, x_4, x_5, y_4, y_5] \supseteq C_5$. Therefore, $6 \geq e(x_4 x_5, y_1 y_2 y_3) = e(T, L_t) - e(x_2, L_t) - e(x_4 x_5, y_4 y_5) \geq 6$. Then $e(x_4 x_5, y_1 y_2 y_3) = 6$ and $e(x_2, L_t) = 5$. Further, $[x_1, y_1, y_5, y_4, x_2] \supseteq C_5$ and $[y_2, y_3, x_3, x_4, x_5] \supseteq C_5$, a contradiction. Therefore, $e(x_2, y_3 y_4) = 0$ and hence $e(x_2, L_t) = 3$, $e(x_4 x_5, L_t) = 10$. It follows that $[x_1, y_1, x_5, x_3, x_2] \supseteq C_5$ and $[y_2, y_3, y_4, y_5, x_4] \supseteq C_5$, again a contradiction. ■

Lemma 13. $G \supseteq K_4^+ \uplus (k-1)C_5$.

Proof. Suppose to the contrary that $G \not\supseteq K_4^+ \uplus (k-1)C_5$. By Lemma 11, let $\psi = \{H, L_1, L_2, \dots, L_{k-1}\}$ be an optimal family of G with $H \cong F_1$. Suppose that $e(H, L_i) \geq 16$ for some $i \in \{1, 2, \dots, k-1\}$, say $e(H, L_1) \geq 16$. By Lemma 6(a) and Lemma 10, we may first assume that there exists a labelling $L_1 = y_1 y_2 y_3 y_4 y_5 y_1$ with property **P**₂. Let $R = \{x_1, x_4, y_3, y_5\}$ and $G_1 = [H, L_1]$. Then $e(R, V(G_1) - R) \leq 16$ and so $e(R, G - V(G_1)) \geq (d_{G_1}(x_1) + d_{G_1}(x_4)) + (d_{G_1}(y_3) +$

$d_{G_1}(y_5)) - 16 \geq 12k - 16 = 12(k - 2) + 8$. Hence, $e(R, L_i) \geq 13$ for some $i \in \{2, 3, \dots, k - 1\}$. So by Lemma 6(b), $[G_1, L_i] \supseteq K_4^+ \uplus 2C_5$ and, therefore $G \supseteq K_4^+ \uplus (k - 1)C_5$, a contradiction. Next, we assume that $G \supseteq K_4^+ \uplus B \uplus (k - 2)C_5$ and let $K_4^+, B, L_1, L_2, \dots, L_{k-2}$ be k disjoint subgraphs of G with $L_i \cong C_5$ for $i \in \{1, 2, \dots, k - 2\}$. Let $B = y_1y_2y_3y_4y_5y_1$ and $R' = \{y_2, y_3, y_4, y_5\}$. We claim that $e(R', K_4^+) \leq 15$. Suppose to the contrary that $e(R', K_4^+) \geq 16$. If $e(x_1, R') \leq 1$, then $e(R', x_2x_3x_4x_5) \geq 15$. It is easy to see that $[K_4^+, B] \supseteq K_4^+ \uplus C_5$, a contradiction. If $e(x_1, R') \geq 3$, say $x_1y_2, x_1y_3, x_1y_4 \in E(G)$, then $[B - y_i + x_1] \supseteq C_5$ for all $i \in \{2, 3, 5\}$. However, $e(y_j, x_2x_3x_4x_5) \geq 2$ for some $j \in \{2, 3, 5\}$ and so $[K_4^+ - x_1 + y_j] \supseteq C_5$, a contradiction. If $e(x_1, R') = 2$, then we just need to consider $x_1y_2, x_1y_4 \in E(G)$ and $x_1y_2, x_1y_3 \in E(G)$. If $x_1y_2, x_1y_4 \in E(G)$, then $[B - y_i + x_1] \supseteq C_5$ for all $i \in \{3, 5\}$. However, $e(y_j, x_2x_3x_4x_5) \geq 2$ for some $j \in \{3, 5\}$ and so $[K_4^+ - x_1 + y_j] \supseteq C_5$, a contradiction. If $x_1y_2, x_1y_3 \in E(G)$, then $e(x_2, y_2y_3) = 0$ for otherwise $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and $[x_3, x_4, x_5, y_4, y_5] \supseteq C_5$, a contradiction. Then $e(y_2y_3, x_3x_4x_5) = 6$ and $e(y_4y_5, x_2x_3x_4x_5) = 8$ and so $[K_4^+, B] \supseteq 2C_5$, again a contradiction. Further, $e(R', G - V(K_4^+ \cup B)) \geq 12k - 15 - 8 = 12(k - 2) + 1$. Hence $e(R', L_j) \geq 13$ for some $j \in \{1, 2, \dots, k - 2\}$. So by Lemma 4(b), $[B, L_j] \supseteq 2C_5$, i.e., $G \supseteq K_4^+ \uplus (k - 1)C_5$, a contradiction. Therefore, $e(H, L_i) \leq 15$ for each $i \in \{1, 2, \dots, k - 1\}$. It follows that $d_G(x_1) > 3k$, for otherwise, we obtain $e(H, G - V(H)) = (d_G(x_1) + d_G(x_3)) + (d_G(x_2) + d_G(x_4)) + (d_G(x_1) + d_G(x_5)) - d_G(x_1) - 12 \geq 18k - d_G(x_1) - 12 \geq 15k - 12 > 15(k - 1)$, then there exists $i \in \{1, 2, \dots, k - 1\}$ such that $e(H, L_i) \geq 16$, a contradiction.

For r with $0 \leq r \leq 5$, let $\mathcal{A}_r = \{L_t \mid e(x_1, L_t) = r, 1 \leq t \leq k - 1\}$ and $a_r = |\mathcal{A}_r|$. It is clear that $a_0 + a_1 + a_2 + a_3 + a_4 + a_5 = k - 1$. Further, it can be seen that

$$(1) \quad d_G(x_1) = d_H(x_1) + \sum_{r=0}^5 \sum_{L_t \in \mathcal{A}_r} e(x_1, L_t) = 1 + a_1 + 2a_2 + 3a_3 + 4a_4 + 5a_5.$$

Let $R_1 = \{x_1, x_2, x_4, x_5\}$. By Lemma 12, we obtain

$$(2) \quad \begin{aligned} \sum_{x \in R_1} d_G(x) &= \sum_{x \in R_1} d_H(x) + \sum_{r=0}^5 \sum_{L_t \in \mathcal{A}_r} e(R_1, L_t) \\ &\leq 9 + 15a_0 + 13a_1 + 13a_2 + 12a_3 + 11a_4 + 10a_5. \end{aligned}$$

By (1) and (2), we obtain $d_G(x_1) + \sum_{x \in R_1} d_G(x) \leq 10 + 15a_0 + 14a_1 + 15a_2 + 15a_3 + 15a_4 + 15a_5 = 15k - 5 - a_1$. But by the degree sum condition, we have $d_G(x_1) + \sum_{x \in R_1} d_G(x) \geq 15k$, which is a contradiction. ■

Lemma 14. *For any family $\{H, L_1, \dots, L_{k-1}\}$ of G with $H \cong K_4^+$, $d_G(x_2) < 3k$.*

Proof. Suppose to the contrary that $d_G(x_2) \geq 3k$ for some family $\psi = \{H, L_1, \dots, L_{k-1}\}$ with $H \cong K_4^+$. Further, we assume that $\sum_{i=1}^{k-1} \tau(L'_i) \leq \sum_{i=1}^{k-1} \tau(L_i)$ for any family $\{H', L'_1, \dots, L'_{k-1}\}$ with $H' \cong K_4^+$ and $d_G(x_2) \geq 3k$. Let $Q = [x_2, x_3, x_4, x_5]$ and $T = [x_3, x_4, x_5]$. Then $Q \cong K_4$ and $T \cong K_3$. For r with $0 \leq r \leq 5$, let $\mathcal{B}_r = \{L_t \mid e(x_1, L_t) = r, 1 \leq t \leq k-1\}$ and $b_r = |\mathcal{B}_r|$. It is clear that $b_0 + b_1 + b_2 + b_3 + b_4 + b_5 = k-1$.

Claim 2. For each $t \in \{1, 2, \dots, k-1\}$, the following statements hold.

- (a) If $e(x_1, L_t) = 5$, then $e(Q, L_t) \leq 5$.
- (b) If $e(x_1, L_t) = 4$, then $e(Q, L_t) \leq 9$ except possible one L_t with $e(Q, L_t) = 10$.
- (c) If $e(x_1, L_t) = 3$, then $e(Q, L_t) \leq 12$.
- (d) If $e(x_1, L_t) = 2$, then $e(Q, L_t) \leq 15$.
- (e) If $e(x_1, L_t) = 1$, then $e(Q, L_t) \leq 16$.
- (f) If $e(x_1, L_t) = 0$, then $e(Q, L_t) \leq 20$.

Proof. Let $L_t = y_1 y_2 y_3 y_4 y_5 y_1$ and $G_t = [H, L_t]$. If $e(x_1, L_t) = 5$, then $e(y_i, Q) \leq 1$ for all $i \in \{1, 2, 3, 4, 5\}$ since $G_t \not\supseteq 2C_5$. Hence, (a) follows directly.

To prove (b), say $N(x_1) \mid L_t = \{y_1, y_2, y_3, y_4\}$. First, we claim that $\{y_1 y_3, y_2 y_4, y_1 y_4\} \subseteq E(L_t)$, $N(x_2) \mid L_t = \{y_1, y_2, y_3, y_4\}$ and $N(x_3) \mid L_t = N(x_4) \mid L_t = N(x_5) \mid L_t = \{y_1, y_4\}$ if $e(Q, L_t) \geq 10$. It is clear that $\tau(y_5, L_t) = 0$ for otherwise $x_1 \rightarrow L_t$ and so $G_t \supseteq 2C_5$. As $x_1 \rightarrow (L_t, y_i)$ for $i \in \{2, 3, 5\}$, $e(y_i, Q) \leq 1$. Hence, $e(y_1 y_4, Q) \geq 7$, say $e(y_1, Q) \geq 3$ and $e(y_4, Q) = 4$. If $x_2 y_5 \in E(G)$, then $[Q + y_5] \supseteq K_4^+$ and $\tau(x_1 y_1 y_2 y_3 y_4 x_1) > \tau(L_t)$, contradicting the definition of ψ . Hence $x_2 y_5 \notin E(G)$. We say that $e(x_i, y_j) = 0$ for all $i \in \{3, 4, 5\}$ and $j \in \{2, 3\}$. If not, then $[x_1, x_2, x_i, y_2, y_3] \supseteq C_5$ and $[(V(T) \cup \{y_1, y_4, y_5\}) \setminus \{x_i\}] \supseteq C_5$. If $e(y_5, T) \geq 1$, then $x_2 y_1 \notin E(G)$ for otherwise $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and $[x_3, x_4, x_5, y_4, y_5] \supseteq C_5$. Hence $e(y_1, T) = 3$. Then $[x_1, x_2, y_2, y_3, y_4] \supseteq C_5$ and $[x_3, x_4, x_5, y_1, y_5] \supseteq C_5$, a contradiction. Therefore, $e(y_5, T) = 0$. Then $e(y_1, Q) = e(y_4, Q) = 4$ and $e(y_2 y_3, x_2) = 2$. If $y_1 y_3 \notin E(G)$ or $y_2 y_4 \notin E(G)$, then $[x_3, x_2, x_1, y_3, y_2] \supseteq K_4^+$ and $\tau(y_1 x_4 x_5 y_4 y_5 y_1) > \tau(L_t)$, a contradiction. Hence $y_1 y_3, y_2 y_4 \in E(G)$. If $y_1 y_4 \notin E(G)$, then either $d_G(y_1) \geq 3k$ or $d_G(y_4) \geq 3k$, say $d_G(y_1) \geq 3k$. Then $[y_5, y_1, y_2, y_3, x_1] \supseteq K_4^+$ and $\tau(y_4 x_2 x_3 x_4 x_5 y_4) > \tau(L_t)$, contradicting the definition of ψ . Hence, $y_1 y_4 \in E(G)$. Next, we claim that at most one L_t with $e(x_1, L_t) = 4$ and $e(Q, L_t) = 10$. If not, then there is another one $L'_t = z_1 z_2 z_3 z_4 z_5 z_1$ with $e(x_1, L'_t) = 4$ and $e(Q, L'_t) = 10$. Without loss of generality, assume that $N(x_1) \mid L'_t = \{z_1, z_2, z_3, z_4\}$. Then $\{z_1 z_3, z_2 z_4, z_1 z_4\} \subseteq E(L'_t)$, $N(x_2) \mid L'_t = \{z_1, z_2, z_3, z_4\}$ and $N(x_3) \mid L'_t = N(x_4) \mid L'_t = N(x_5) \mid L'_t = \{z_1, z_4\}$. Therefore, $[y_1, y_5, y_4, x_4, x_5] \supseteq C_5$, $[x_2, z_4, z_5, z_1, x_3] \supseteq C_5$ and $[y_2, y_3, x_1, z_2, z_3] \supseteq B$, contradicting Lemma 10.

To prove (c), suppose to the contrary that $e(Q, L_t) \geq 13$. Assume first $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_4\}$.

As $x_1 \rightarrow (L_t, y_i)$ for all $i \in \{3, 5\}$, $e(y_i, Q) \leq 1$. Further, $y_3y_5 \notin E(G)$ as $x_1 \not\rightarrow L_t$. It follows that $e(y_1y_2y_4, Q) \geq 11$ and so $e(x_2, y_1y_4) \geq 1$, $e(x_2, y_2y_4) \geq 1$. Then $[x_1, x_2, y_1, y_5, y_4] \supseteq C_5$ and $[x_1, x_2, y_2, y_3, y_4] \supseteq C_5$. Further, as $e(y_i, T) \geq 2$ for all $i \in \{1, 2\}$, $e(y_3y_5, T) = 0$ for otherwise $[x_3, x_4, x_5, y_2, y_3] \supseteq C_5$ or $[x_3, x_4, x_5, y_1, y_5] \supseteq C_5$, a contradiction. Then $N(y_3)|Q \cup N(y_5)|Q \subseteq \{x_2\}$.

Let $R = \{x_1, x_4, y_3, y_5\}$. Then $e(R, V(G_t) - R) \leq 18$ and so $e(R, G - V(G_t)) \geq 12k - 18 = 12(k - 2) + 6$ and hence $e(R, L_i) \geq 13$ for some $i \in \{1, 2, 3, \dots, k-1\} \setminus \{t\}$. Note that $e(Q, L_t) \geq 13$ and $N(y_3)|Q \cup N(y_5)|Q \subseteq \{x_2\}$. If $u \rightarrow (L_i, z; \{v, w\})$ for some $z \in V(L_i)$, $u \in R$ and $\{v, w\} \subseteq R \setminus \{u\}$, then $[G_t, L_i] \supseteq 3C_5$, a contradiction. By Lemma 4(e), there are vertex labellings $L_i = z_1z_2z_3z_4z_5z_1$ and $R = \{a_1, a_2, a_3, a_4\}$ such that $e(a_1a_2, z_1z_2z_3z_4) = 8$, $e(a_3, z_1z_5z_4) = 3$ and $e(a_4, z_1z_4) = 2$. If $\{a_1, a_2\} = \{x_1, x_4\}$, then set $\{r, s\} = \{1, 2\}$ with $y_r \in N(x_1) \mid L_t \cap N(x_4) \mid L_t$. We can see that $[x_1, y_r, x_4, z_2, z_3] \supseteq C_5$, $[y_3, y_5, z_1, z_5, z_4] \supseteq C_5$ and $[x_2, x_3, x_5, y_s, y_4] \supseteq C_5$, a contradiction. If $\{a_1, a_2\} = \{x_1, y_i\}$ for some $i \in \{3, 5\}$, say $\{a_1, a_2\} = \{x_1, y_5\}$, then $[x_1, y_1, y_5, z_2, z_3] \supseteq C_5$, $[y_3, x_4, z_1, z_5, z_4] \supseteq C_5$ and $[y_2, y_4, x_2, x_3, x_5] \supseteq C_5$, a contradiction. If $\{a_1, a_2\} = \{x_4, y_i\}$ for some $i \in \{3, 5\}$, say $\{a_1, a_2\} = \{x_4, y_5\}$, then we set $\{r, s\} = \{1, 4\}$ with $x_4y_r \in E(G)$. It is clear that $[x_4, y_r, y_5, z_2, z_3] \supseteq C_5$, $[x_1, y_3, z_1, z_5, z_4] \supseteq C_5$ and $[x_2, x_3, x_5, y_2, y_s] \supseteq C_5$, a contradiction. Hence $\{a_1, a_2\} = \{y_3, y_5\}$. Therefore, $[y_3, y_4, y_5, z_2, z_3] \supseteq C_5$, $[x_1, x_4, z_1, z_5, z_4] \supseteq C_5$ and $[x_2, x_3, x_5, y_1, y_2] \supseteq C_5$, again a contradiction.

Next, assume that $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L_t, y_2)$, $e(y_2, Q) \leq 1$. Suppose $e(x_2, y_4y_5) \geq 1$, say $x_2y_5 \in E(G)$. Then $e(y_3y_4, T) \leq 3$ for otherwise $[x_1, x_2, y_5, y_1, y_2] \supseteq C_5$ and $[x_3, x_4, x_5, y_3, y_4] \supseteq C_5$. Further, if $x_2y_4 \in E(G)$, then similarly, $e(y_1y_5, T) \leq 3$ and so $e(Q, L_t) \leq 12$, a contradiction. Hence $x_2y_4 \notin E(G)$. Since $e(Q, L_t) \geq 13$, then $e(y_1y_5, Q) = 8$, $e(y_3y_4, T) = 3$, $e(y_2, Q) = 1$ and $x_2y_3 \in E(G)$. If $e(y_4, T) \geq 1$, then $[x_3, x_4, x_5, y_4, y_5] \supseteq C_5$ and $[y_1, y_2, y_3, x_1, x_2] \supseteq C_5$, a contradiction. Thus $e(y_4, T) = 0$ and so $e(y_3, T) = 3$. Then $[x_1, y_2, y_1, x_3, x_2] \supseteq C_5$ and $[y_3, y_4, y_5, x_4, x_5] \supseteq C_5$, a contradiction. Therefore, $e(x_2, y_4y_5) = 0$. If $e(x_2, y_1y_3) \geq 1$, then $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and so $e(y_4y_5, T) \leq 3$. It follows that $e(Q, L_t) = e(y_4y_5, T) + e(y_1y_3, T) + e(y_2, T) + e(x_2, y_1y_2y_3) \leq 12$, no matter whether y_2x_2 is an edge of $E(G)$ or not, a contradiction. Therefore, $e(x_2, y_1y_3y_4y_5) = 0$ and hence $e(T, y_1y_3y_4y_5) = 12$. Then $[x_1, y_3, x_3, y_1, y_2] \supseteq C_5$ and $[x_2, x_4, y_4, y_5, x_5] \supseteq C_5$, again a contradiction.

To prove (d), suppose to the contrary that $e(Q, L_t) \geq 16$. Assume first $N(x_1) \mid L_t = \{y_i, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_3\}$. As $x_1 \rightarrow (L_t, y_2)$, $e(y_2, Q) \leq 1$. Therefore, $e(Q, y_1y_3y_4y_5) \geq 15$ and hence $e(x_2, y_1y_3) \geq 1$. Then $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and so $e(y_4y_5, T) \leq 3$. This implies that $e(Q, y_1y_3y_4y_5) \leq 13$, a contradiction. Next, assume that $N(x_1) \mid L_t = \{y_i, y_{i+1}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2\}$. If $x_2y_4 \in E(G)$, then

$e(y_2y_3, T) \leq 3$ for otherwise $[x_1, x_2, y_4, y_5, y_1] \supseteq C_5$ and $[x_3, x_4, x_5, y_2, y_3] \supseteq C_5$, a contradiction. Similarly, if $x_2y_4 \in E(G)$, then $e(y_1y_5, T) \leq 3$. This means that $e(Q, L_t) \leq 14$, a contradiction. Hence $x_2y_4 \notin E(G)$. If $e(x_2, y_3y_5) \geq 1$, say $x_2y_5 \in E(G)$, then $[x_1, x_2, y_5, y_1, y_2] \supseteq C_5$ and so $e(y_3y_4, T) \leq 3$. Further, $e(y_1y_2y_5, Q) = 12$ as $e(Q, L_t) \geq 16$. This implies that $[x_1, y_1, x_5, x_4, x_2] \supseteq C_5$ and $[y_2, y_3, y_4, y_5, x_3] \supseteq C_5$, a contradiction. Hence $e(x_2, y_3y_5) = 0$. If $y_3x_3 \in E(G)$, then $e(y_1y_4y_5, x_4x_5) \leq 4$ for otherwise $[x_1, y_2, y_3, x_3, x_2] \supseteq C_5$ and $[x_4, x_5, y_1, y_4, y_5] \supseteq C_5$. Thus $e(Q, L_t) = e(x_2, y_1y_2) + e(x_3, L_t) + e(x_4x_5, y_1y_4y_5) + e(x_4x_5, y_2y_3) \leq 15$, a contradiction. Then $y_3x_3 \notin E(G)$. Similarly, $y_3x_4, y_3x_5 \notin E(G)$. Then $e(Q, L_t) = e(x_2, y_1y_2) + e(x_3x_4x_5, y_1y_2y_4y_5) \leq 14$, again a contradiction.

To prove (e), suppose to the contrary that $e(Q, L_t) \geq 17$. Without loss of generality assume that $x_1y_1 \in E(G)$. Suppose $e(x_2, y_3y_4) \geq 1$, say $x_2y_3 \in E(G)$. Then $[x_1, x_2, y_1, y_2, y_3] \supseteq C_5$ and so $e(y_4y_5, T) \leq 3$. Since $e(Q, L_t) \geq 17$, $e(y_1y_2y_3, Q) = 12$, $e(y_4y_5, T) = 3$ and $e(x_2, y_4y_5) = 2$. Then $[x_1, x_2, y_4, y_5, y_1] \supseteq C_5$ and $[x_3, x_4, x_5, y_2, y_3] \supseteq C_5$, a contradiction. Hence $e(x_2, y_3y_4) = 0$ and so $e(T, L_t) \geq 14$. This implies that $e(x_i, y_2y_5) = 2$ and $y_1x_j \in E(G)$ for some $\{i, j\} \subseteq \{3, 4, 5\}$ with $i \neq j$. Then $[L_t - y_1 + x_i] \supseteq C_5$ and $[H + y_1 - x_i] \supseteq C_5$, a contradiction. \square

Recall that $b_0 + b_1 + b_2 + b_3 + b_4 + b_5 = k - 1$. Further, we can see that

$$(3) \quad d_G(x_1) = d_H(x_1) + \sum_{r=0}^5 \sum_{L_t \in \mathcal{B}_r} e(x_1, L_t) = 1 + b_1 + 2b_2 + 3b_3 + 4b_4 + 5b_5.$$

By Claim 2, there may exist one $L'_t \in \mathcal{B}_4$ with $e(H, L'_t) = 14$. Further, we obtain

$$(4) \quad \begin{aligned} \sum_{x \in V(H)} d_G(x) &= \sum_{x \in V(H)} d_H(x) + \sum_{r=0}^3 \sum_{L_t \in \mathcal{B}_r} e(H, L_t) + \sum_{L_t \in \mathcal{B}_5} e(H, L_t) \\ &\quad + \sum_{L_t \in \mathcal{B}_4 \setminus \{L'_t\}} e(H, L_t) + e(H, L'_t) \\ &\leq 14 + 20b_0 + 17b_1 + 17b_2 + 15b_3 + 10b_5 + (13b_4 + 1) \\ &= 15 + 20b_0 + 17b_1 + 17b_2 + 15b_3 + 13b_4 + 10b_5. \end{aligned}$$

Combining (4) with (3), we obtain $2d_G(x_1) + \sum_{x \in V(H)} d_G(x) \leq 17 + 20b_0 + 19b_1 + 21b_2 + 21b_3 + 21b_4 + 20b_5 = 21k - b_0 - 2b_1 - b_5 - 4$. Since $d_G(x_2) \geq 3k$, we have $2d_G(x_1) + \sum_{x \in V(H)} d_G(x) = d_G(x_2) + (d_G(x_1) + d_G(x_3)) + (d_G(x_1) + d_G(x_4)) + (d_G(x_1) + d_G(x_5)) \geq 21k$, which is a contradiction. \blacksquare

Let $\psi = \{H, L_1, L_2, \dots, L_{k-1}\}$ be an optimal family of G with $H \cong K_4^+$, and let \mathcal{B}_r and b_r be defined as in the proof of Lemma 14. Let $Q = [x_2, x_3, x_4, x_5]$ and $T = [x_3, x_4, x_5]$.

Lemma 15. *For each $t \in \{1, 2, \dots, k-1\}$, the following statements hold.*

- (a) *If $e(x_1, L_t) = 5$, then $e(T, L_t) \leq 3$.*
- (b) *If $e(x_1, L_t) = 4$, then $e(T, L_t) \leq 6$.*
- (c) *If $e(x_1, L_t) = 3$, then $e(T, L_t) \leq 9$.*
- (d) *If $1 \leq e(x_1, L_t) \leq 2$, then $e(T, L_t) \leq 12$.*
- (e) *If $e(x_1, L_t) = 0$, then $e(T, L_t) \leq 15$.*

Proof. Let $L_t = y_1y_2y_3y_4y_5y_1$ and $G_t = [H, L_t]$. To prove (a), suppose to the contrary that $e(T, L_t) \geq 4$. As $G_t \not\supseteq 2C_5$, $e(y_i, Q) \leq 1$ for all $i \in \{1, 2, 3, 4, 5\}$. Further, by the optimality of ψ , we obtain $[L_t] \cong K_5$. Without loss of generality assume that $y_1x_3 \in E(G)$. Then $y_1x_2 \notin E(G)$ and so $d_G(y_1) \geq 3k$, $[Q+y_1] \supseteq K_4^+$, $[L_t - y_1 + x_1] \cong K_5$. Therefore, we may assume that $d_G(x_1) \geq 3k$. Since $e(T, L_t) \geq 4$, $e(x_l, L_t) \geq 2$ for some $l \in \{3, 4, 5\}$, say $x_3y_1, x_3y_2 \in E(G)$. Then $[x_2, x_1, y_3, y_4, y_5] \supseteq K_4^+$, $[x_3, x_4, x_5, y_1, y_2] \supseteq B$. Let $L' = [x_3, x_4, x_5, y_1, y_2]$ and $R' = \{y_1, y_2, x_4, x_5\}$. Note that $e(R', x_2x_1y_3y_4y_5) \leq 11$ and $\sum_{x \in R'} d_{L'}(x) = 8$. Then $e(R', G - V(G_t)) \geq 12k - 19 = 12(k-2) + 5$ and so $e(R', L_i) \geq 13$ for some $i \in \{1, \dots, k-1\} \setminus \{t\}$. By Lemma 4(b), $[L', L_i] \supseteq 2C_5$. This contradicts Lemma 14 as $[x_2, x_1, y_3, y_4, y_5] \supseteq K_4^+$ and $d_G(x_1) \geq 3k$.

To prove (b), suppose to the contrary that $e(T, L_t) \geq 7$. Without loss of generality assume that $N(x_1) \mid L_t = \{y_1, y_2, y_3, y_4\}$. It is clear that $\tau(y_5, L_t) = 0$ for otherwise $x_1 \rightarrow L_t$ and so $G_t \supseteq 2C_5$. As $x_1 \rightarrow (L_t, y_i)$ for $i \in \{2, 3, 5\}$, $e(y_i, T) \leq 1$. If $e(y_5, T) = 1$, then $[Q + y_5] \supseteq K_4^+$ and $\tau(L_t - y_5 + x_1) > \tau(L_t)$, contradicting the optimality of ψ . Therefore, $e(y_5, T) = 0$ and hence $e(y_1y_4, T) \geq 5$. Without loss of generality, assume that $e(y_1, T) = 3$ and $y_4x_3, y_4x_4 \in E(G)$. Further, since $e(T, L_t) \geq 7$, $e(y_2y_3, T) \geq 1$. If $x_3 \in N(y_2) \mid T \cup N(y_3) \mid T$, then $[x_1, y_2, y_3, x_3, x_2] \supseteq C_5$ and $[y_1, y_4, y_5, x_4, x_5] \supseteq C_5$, a contradiction. If $x_4 \in N(y_2) \mid T \cup N(y_3) \mid T$, then $[x_1, y_2, y_3, x_4, x_2] \supseteq C_5$ and $[y_1, y_4, y_5, x_3, x_5] \supseteq C_5$, a contradiction. If $x_5 \in N(y_2) \mid T \cup N(y_3) \mid T$, then $[x_1, y_2, y_3, x_5, x_2] \supseteq C_5$ and $[y_1, y_4, y_5, x_3, x_4] \supseteq C_5$, again a contradiction.

To prove (c), suppose to the contrary that $e(T, L_t) \geq 10$. Assume first $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+2}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_3\}$. As $x_1 \rightarrow (L_t, y_2)$, $e(y_2, T) \leq 1$. Without loss of generality assume that $e(y_1y_4, T) \geq e(y_3y_5, T)$. Then $[x_1, y_2, y_3, x_i, x_2] \supseteq C_5$ and $[(V(T) \cup \{y_1, y_4, y_5\}) \setminus \{x_i\}] \supseteq C_5$, where $x_i \in N(y_2) \mid T \cup N(y_3) \mid T$, a contradiction. Next, assume that $N(x_1) \mid L_t = \{y_i, y_{i+1}, y_{i+3}\}$ for some $i \in \{1, 2, 3, 4, 5\}$, say $N(x_1) \mid L_t = \{y_1, y_2, y_4\}$. As $x_1 \rightarrow (L_t, y_i)$ for $i \in \{3, 5\}$, $e(y_i, T) \leq 1$. Since $e(T, L_t) \geq 10$, $e(y_1y_4, x_4x_5) \geq 3$ and $e(y_3y_5, T) \geq 1$, say $x_3y_3 \in E(G)$. Then $[x_1, y_2, y_3, x_3, x_2] \supseteq C_5$ and $[y_1, y_5, y_4, x_4, x_5] \supseteq C_5$, a contradiction.

To prove (d), suppose to the contrary that $e(T, L_t) \geq 13$. Without loss of generality assume that $x_1y_1 \in E(G)$. Since $e(T, L_t) \geq 13$, $e(y_2y_5, T) \geq 4$. Assume

first that $e(y_2y_5, T) = 4$, say $x_3 \in N(y_2)|T \cap N(y_5)|T$. Further, since $e(T, L_t) \geq 13$, $e(y_1, T) = 3$. Then $[x_1, y_1, x_2, x_4, x_5] \supseteq C_5$ and $[x_3, y_2, y_3, y_4, y_5] \supseteq C_5$, a contradiction. Next, assume that $e(y_2y_5, T) \geq 5$, say $\{x_3, x_4\} \subseteq N(y_2)|T \cap N(y_5)|T$. Since $e(T, L_t) \geq 13$, $e(y_1, T) \geq 1$. If $y_1x_3 \in E(G)$, then $[x_1, y_1, x_2, x_3, x_5] \supseteq C_5$ and $[x_4, y_2, y_3, y_4, y_5] \supseteq C_5$, a contradiction. If $y_1x_4 \in E(G)$, then $[x_1, y_1, x_2, x_4, x_5] \supseteq C_5$ and $[x_3, y_2, y_3, y_4, y_5] \supseteq C_5$, a contradiction. If $y_1x_5 \in E(G)$, then $[x_1, y_1, x_2, x_4, x_5] \supseteq C_5$ and $[x_3, y_2, y_3, y_4, y_5] \supseteq C_5$, again a contradiction. ■

Let $R = \{x_1, x_3, x_4, x_5\}$. By Lemma 15, we obtain

$$(5) \quad \sum_{x \in R} d_G(x) = \sum_{x \in R} d_H(x) + \sum_{r=0}^5 \sum_{L_t \in \mathcal{B}_r} e(R, L_t) \\ \leq 10 + 15b_0 + 13b_1 + 14b_2 + 12b_3 + 10b_4 + 8b_5.$$

Combining (5) with (3), we have $2d_G(x_1) + \sum_{x \in R} d_G(x) \leq 12 + 15b_0 + 15b_1 + 18b_2 + 18b_3 + 18b_4 + 18b_5 = 18k - 3b_0 - 3b_1 - 6$. But by the degree sum condition, we have $2d_G(x_1) + \sum_{x \in R} d_G(x) \geq 18k$, a contradiction. This proves Theorem 3.

Acknowledgements

The authors would like to thank the anonymous reviewers for their careful reading of our original manuscript and constructive suggestions. This work was supported by the National Natural Science Foundation of China [Grant numbers, 11971406, 12171402]

REFERENCES

- [1] S. Abbasi, The Solution of the El-Zahar Problem (PhD Thesis, Rutgers University, 1998).
- [2] K. Corrádi and A. Hajnal, *On the maximal number of independent circuits in a graph*, Acta Math. Acad. Sci. Hungar. **14** (1963) 423–439.
<https://doi.org/10.1007/BF01895727>
- [3] S. Chiba and T. Yamashita, *Degree conditions for the existence of vertex-disjoint cycles and paths: A survey*, Graphs Combin. **34** (2018) 1–83.
<https://doi.org/10.1007/s00373-017-1873-5>
- [4] G.A. Dirac, *Some theorems on abstract graphs*, Proc. Lond. Math. Soc. (3) **s3-2** (1952) 69–81.
<https://doi.org/10.1112/plms/s3-2.1.69>
- [5] M.H. El-Zahar, *On circuits in graphs*, Discrete Math. **50** (1984) 227–230.
[https://doi.org/10.1016/0012-365X\(84\)90050-5](https://doi.org/10.1016/0012-365X(84)90050-5)
- [6] P. Erdős, Some Recent Combinatorial Problems (Technical Report, University of Bielefeld, 1990).

- [7] H. Wang, *Proof of the Erdős-Faudree conjecture on quadrilaterals*, Graphs Combin. **26** (2010) 833–877.
<https://doi.org/10.1007/s00373-010-0948-3>
- [8] H. Wang, *Disjoint 5-cycles in a graph*, Discuss. Math. Graph Theory **32** (2012) 221–242.
<https://doi.org/10.7151/dmgt.1605>

Received 1 August 2021

Revised 1 May 2022

Accepted 2 May 2022

Available online 16 May 2022