Discussiones Mathematicae Graph Theory 44 (2024) 717–726 https://doi.org/10.7151/dmgt.2464

A NOTE ON MINIMUM DEGREE, BIPARTITE HOLES, AND HAMILTONIAN PROPERTIES*

QIANNAN ZHOU^{*a*}, HAJO BROERSMA^{*b*,1}, LIGONG WANG^{*c*}, YONG LU^{*a*}

 ^a School of Mathematics and Statistics Jiangsu Normal University
Xuzhou, Jiangsu 221116, People's Republic of China
^b Faculty of EEMCS, University of Twente
P.O. Box 217, 7500 AE Enschede, The Netherlands

^cSchool of Mathematics and Statistics Northwestern Polytechnical University Xi'an, Shaanxi 710072, People's Republic of China

> e-mail: qnzhoumath@163.com h.j.broersma@utwente.nl lgwangmath@163.com luyong@jsnu.edu.cn

Abstract

We adopt the recently introduced concept of the bipartite-hole-number due to McDiarmid and Yolov, and extend their result on Hamiltonicity to other Hamiltonian properties of graphs with a large minimum degree in terms of this concept. An (s,t)-bipartite-hole in a graph G consists of two disjoint sets of vertices S and T with |S| = s and |T| = t such that E(S,T) = \emptyset . The bipartite-hole-number $\tilde{\alpha}(G)$ is the maximum integer r such that G contains an (s,t)-bipartite-hole for every pair of nonnegative integers sand t with s + t = r. Our main results are that a graph G is traceable if $\delta(G) \geq \tilde{\alpha}(G) - 1$, and Hamilton-connected if $\delta(G) \geq \tilde{\alpha}(G) + 1$, both improving the analogues of Dirac's Theorem for traceable and Hamiltonconnected graphs.

Keywords: Hamilton-connected graph, traceable graph, degree condition, bipartite-hole-number, minimum degree.

2020 Mathematics Subject Classification: 05C45, 05C07.

¹Corresponding author.

^{*}Research supported by the National Natural Science Foundation of China (No. 11871398, 11901253), the Natural Science Basic Research Plan in Shaanxi Province of China (Program No. 2018JM1032), the Fundamental Research Funds for the Central Universities (No. 3102019ghjd003), the China Scholarship Council (No. 201806290178) and the Natural Science Foundation for Colleges and Universities in Jiangsu Province of China (No. 21KJB110025).

1. INTRODUCTION

Our motivation for the presented results is a recent generalization of a classic result of Dirac [3] on Hamiltonicity (Theorem 1 below) due to McDiarmid and Yolov [8] (Theorem 2 below). We answer the natural question whether similar extensions can be established for analogues of Dirac's Theorem for traceability and Hamilton-connectivity. Throughout this note, we use Bondy and Murty [1] for terminology and notation not defined here and only consider finite simple graphs.

For a graph G, we use V(G) and E(G) to denote the vertex set and the edge set of G, respectively. For $v \in V(G)$, we use $N_G(v)$ to denote the set of neighbors of v in G, and we let $d(v) = d_G(v) = |N_G(v)|$ denote the degree of v in G. Moreover, we use $N_G[v] = N_G(v) \cup \{v\}$. If the graph G is clear from the context, we will usually drop the subscript G. Let $\delta(G)$ denote the minimum degree of (the vertices of) G. An independent set of G is a set of vertices no two of which are adjacent. The cardinality of a maximum independent set in G is called the independence number of G, and denoted by $\alpha(G)$. A spanning subgraph of a graph G is a subgraph obtained by edge deletions only. If H is a spanning subgraph of G, we use G - H to denote the graph with vertex set V(G)and edge set $E(G) \setminus E(H)$. For two disjoint nonempty subsets S and T of V(G), E[S,T] denotes the set of edges with one end in S and one end in T. The disjoint union of G and H, denoted by G + H, is the graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. The join of G and H, denoted by $G \vee H$, is the graph obtained from the disjoint union of G and H by adding edges joining every vertex of G to every vertex of H. The complement \overline{G} of G is the graph with vertex set V(G) and the property that $uv \in E(\overline{G})$ if and only if $uv \notin E(G)$.

A connected graph G is said to be k-connected if it has more than k vertices and remains connected whenever fewer than k vertices are removed. The connectivity $\kappa(G)$ of G is the maximum value of k for which G is k-connected.

If C is a cycle in G, we let \overrightarrow{C} denote the cycle C with a clockwise or anticlockwise orientation. For $u, v \in V(C)$ with a fixed chosen orientation for C, we let $\overrightarrow{C}[u, v]$ denote the consecutive vertices on C from u to v in the direction specified by \overrightarrow{C} . The same vertices, in reverse order, are given by $\overleftarrow{C}[v, u]$. Both $\overrightarrow{C}[u, v]$ and $\overleftarrow{C}[v, u]$ are considered as paths and as vertex sets in the sequel. Note that we do not exclude the possibility that u = v; in this case both $\overrightarrow{C}[u, v]$ and $\overleftarrow{C}[v, u]$ reduce to one vertex.

A cycle passing through all the vertices of a graph is called a Hamilton cycle. Similarly, a path passing through all the vertices of a graph is called a Hamilton path. A graph G is said to be Hamiltonian if G has a Hamilton cycle, traceable if G has a Hamilton path, and Hamilton-connected if every two vertices of G are connected by a Hamilton path. Already back in 1952, Dirac [3] gave the following minimum degree condition for a graph to be Hamiltonian.

Theorem 1 [3]. A graph G with $n \ge 3$ vertices is Hamiltonian if $\delta(G) \ge n/2$.

There exist many generalizations of Dirac's Theorem. In this note we refrain from providing more details. For more information on some of these generalizations, we refer the reader to [2, 4-7, 9, 10].

Motivated by Dirac's Theorem, in a paper of 2017 McDiarmid and Yolov [8] introduced a new graph parameter which they named the bipartite-hole-number.

Definition [8]. An (s, t)-bipartite-hole in a graph G consists of two disjoint sets of vertices S and T with |S| = s and |T| = t such that $E(S, T) = \emptyset$. The bipartite-hole-number $\tilde{\alpha}(G)$ is the least integer r that can be written as r = s+t-1 for some positive integers s and t such that G does not contain an (s, t)-bipartite-hole.

As stated in [8], an equivalent definition of $\tilde{\alpha}(G)$ is the maximum integer r such that G contains an (s, t)-bipartite-hole for every pair of nonnegative integers s and t with s + t = r.

In [8], the authors presented the following tight sufficient condition for Hamiltonicity in terms of the minimum degree and the bipartite hole number, improving Theorem 1.

Theorem 2 [8]. A graph G with at least three vertices is Hamiltonian if $\delta(G) \geq \widetilde{\alpha}(G)$.

As noted in [8], it is easy to check that a graph G with $\delta(G) \ge n/2$ has no $(1, \lfloor \frac{n}{2} \rfloor)$ -bipartite-hole, so for such a graph $\delta(G) \ge n/2 \ge \tilde{\alpha}(G)$. Motivated by this result, it is natural to consider possible counterparts of this result for other Hamiltonian properties.

In [8], the authors also presented the following result.

Theorem 3 [8]. Let $r \ge 0$ be an integer and let G be a graph with at least three vertices such that $\delta(G) \ge (r+1)\widetilde{\alpha}(G) + 3r$. Then G contains r+1 edge-disjoint Hamilton cycles.

The rest of this note is organized as follows. In Section 2, we will present our results, including the natural counterparts of Theorem 2 for traceable graphs and for Hamilton-connected graphs. In Section 3, we will present the proofs of our results.

2. MAIN RESULTS

We start with the following counterpart of Theorem 2 for traceable graphs.

Theorem 4. A graph G on at least three vertices is traceable if $\delta(G) \geq \widetilde{\alpha}(G) - 1$.

It is easy to come up with examples showing that the result is sharp. Consider, e.g., the nontraceable graph $G = K_{r,r+2}$, for which clearly $\delta(G) = r$ and $\tilde{\alpha}(G) = r+2$. Our next result is a counterpart of Theorem 3, providing a sufficient condition for the existence of many edge-disjoint Hamilton paths.

Theorem 5. Let $r \ge 0$ be an integer, and let G be a graph on at least three vertices with $\delta(G) \ge (r+1)\widetilde{\alpha}(G) + 3r - 1$. Then G contains r+1 edge-disjoint Hamilton paths, which have 2(r+1) distinct end vertices.

We believe that the above result is only sharp for r = 0, but we were not able to relax the condition either, and leave it as an open problem. Next, we present the analogue of Theorem 4 for Hamilton-connected graphs.

Theorem 6. A graph G on at least three vertices is Hamilton-connected if $\delta(G) \ge \widetilde{\alpha}(G) + 1$.

This result is also sharp, in the sense that there exist non-Hamilton-connected graphs G with $\delta(G) = \tilde{\alpha}(G) = r$ for any positive integer r. An obvious example is the graph $G = K_{r,r}$, satisfying $\delta(G) = \tilde{\alpha}(G) = r$. An analogue of Dirac's Theorem for Hamilton-connected graphs states that a graph G of order n is Hamilton-connected if $\delta(G) \geq \frac{n+1}{2}$. It is not difficult to show that Theorem 6 improves this result. For a graph satisfying $\delta(G) \geq \frac{n+1}{2}$, it is easy to check that there is no $\left(1, \left|\frac{n-1}{2}\right|\right)$ -bipartite-hole. Hence, for such a graph $\delta(G) \geq \frac{n+1}{2} \geq \tilde{\alpha}(G) + 1$.

The following generalization of Dirac's Theorem for Hamilton-connected graphs is due to Chvátal and Erdős.

Theorem 7 [2]. A graph G with at least three vertices is Hamilton-connected if $\kappa(G) \ge \alpha(G) + 1$.



Figure 1. Graph G.

We observe that the condition of Theorem 6 is very similar to that of Theorem 7. But comparing these two theorems, neither condition implies the other. We first show an example of a graph G meeting the condition of Theorem 6 but

720

not of Theorem 7. Let G be the graph on vertex set $V(G) = V(A) \cup V(B) \cup V(C)$, where $A = K_{\ell}$, $B = \overline{K_k}$, $C = K_k$, $\ell \leq k$, and all these subgraphs are mutually vertex-disjoint. Let the edge set of G be defined as $E(G) = E(A) \cup$ $E(C) \cup \{ab \mid a \in V(A), b \in V(B)\} \cup \{bc \mid b \in V(B), c \in V(C)\}$. Obviously, we have $\kappa(G) = k = \alpha(G)$, and if we take $\ell \geq 3$ and $k \geq \ell + 3$, we get $\delta(G) = \ell - 1 + k \geq \min\{2\ell + 1, k + 1\} + 1 = \tilde{\alpha}(G) + 1$. In the other direction, the graph G that is depicted in Figure 1 satisfies $\kappa(G) = 3$, $\alpha(G) = 2$ but $\delta(G) = \tilde{\alpha}(G) = 3$.

In the next section, we will present the details of our proofs of the above theorems.

3. The Proofs

Our proof of Theorem 4 is an easy consequence of Theorem 2 and the following observation.

Lemma 8 (Exercise 18.1.6 on Page 474 of [1]). Let G be a graph on at least two vertices. Then G is traceable if and only if $G \vee K_1$ is Hamiltonian.

Proof of Theorem 4. Suppose $H = G \vee K_1$ with vertex set $V(H) = V(G) \cup \{v\}$ and edge set $E(H) = E(G) \cup \{vx \mid x \in V(G)\}$. By the definition of the bipartite-hole-number, we know that $\widetilde{\alpha}(H) = \widetilde{\alpha}(G)$. Then $\delta(H) = \delta(G) + 1 \ge \widetilde{\alpha}(G) - 1 + 1 = \widetilde{\alpha}(G) = \widetilde{\alpha}(H)$. Using Theorem 2, we obtain that H is Hamiltonian. Then by Lemma 8, G is traceable.

Our proof of Theorem 5 is also based on Lemma 8, and makes use of Theorem 3.

Proof of Theorem 5. Let $H = G \vee K_1$ be defined as above. Similarly as in the above proof, we get $\delta(H) \ge (r+1)\widetilde{\alpha}(G) + 3r - 1 + 1 = (r+1)\widetilde{\alpha}(H) + 3r$. By Theorem 3, H has r + 1 edge-disjoint Hamilton cycles. Using Lemma 8, we conclude that G has r + 1 edge-disjoint Hamilton paths, and that these paths have 2(r+1) distinct end vertices.

At the end of this note, we present our proof of Theorem 6.

Proof of Theorem 6. If $\tilde{\alpha}(G) = 1$, then G is complete, and so G is Hamiltonconnected. Hence we may suppose that $\tilde{\alpha}(G) \geq 2$ and G is not Hamiltonconnected. Then there exist two vertices u and v such that there is no Hamilton path connecting them. By Theorem 2, we know G is Hamiltonian. Let C be a Hamilton cycle in G, and let |V(C)| = n. Label the vertices in V(C) with $[n] = \{1, 2, \ldots, n\}$ in order according to the clockwise direction, where u = n and v = k for some $k \notin \{1, n - 1, n\}$. For a set $S \subseteq V(C)$, denote by S^+ the set of immediate successors x^+ on C of elements x in S, and denote by S^- the set of immediate predecessors x^- .

Let $1 \leq s \leq t$ be such that $\tilde{\alpha}(G) + 1 = s + t$ and G has no (s, t)-bipartite-hole. Since $\tilde{\alpha}(G) \geq 2$, we have $1 \leq s \leq \frac{\tilde{\alpha}(G)+1}{2} < \tilde{\alpha}(G)$, and hence

$$|N(1) \cap \{1,2\})| = 1 \le s \le \delta(G) - 2 \le |N(1) \cap (2,n)| = d(1) - 2.$$

Therefore we can choose $\ell \in (1, n)$ such that $|N(1) \cap (1, \ell)| = s$. We choose the smallest ℓ with this property and note that this choice implies $1\ell \in E(G)$.

We know that 1 is not adjacent to k+1 since there is no Hamilton path from n to k. Hence, we have $\ell \in (1, k]$ or $\ell \in (k+1, n)$. Next, we consider these two cases.

Case 1. $\ell \in (1, k]$. We describe five situations (referring to Figure 2) in which there is a Hamilton path connecting n and k, denoted as an (n, k)-H-path in the remainder of the proof.



Figure 2. Situations (a)–(e).

(a) If for some $i \in (1, \ell]$ we have $i \in N(1)$ and $i^- \in N(k+1)$, then $\overleftarrow{C}[n, k+1]$ $\overleftarrow{C}[i^-, 1]\overrightarrow{C}[i, k]$ is an (n, k)-H-path.

(b) If for some $i \in (1, \ell]$ and $j \in (\ell, k]$ we have $i \in N(1), j \in N(k+1)$ and $i^-j^+ \in E(G)$, then $\overline{C}[n, k+1]\overline{C}[j, i]\overline{C}[1, i^-]\overline{C}[j^+, k]$ is an (n, k)-H-path. In the particular case that j = k, then $\overline{C}[n, k+1]\overline{C}[i^-, 1]\overline{C}[i, k]$ is an (n, k)-H-path.

(c) If for some $i \in (\underline{1}, \ell]$ and $j \in (k + 1, n]$ we have $i \in N(1), j \in N(k + 1)$ and $i^-j^- \in E(G)$, then $\overleftarrow{C}[n, j]\overrightarrow{C}[k + 1, j^-]\overrightarrow{C}[i^-, 1]\overrightarrow{C}[i, k]$ is an (n, k)-H-path.

(d) If for some $i \in (\ell, k]$ and $j \in (1, \ell]$ we have $i \in N(1), j \in N(k+1)$ and $i^-j^- \in E(G)$, then $\overleftarrow{C}[n, k+1]\overrightarrow{C}[j, i^-]\overleftarrow{C}[j^-, 1]\overrightarrow{C}[i, k]$ is an (n, k)-H-path.

(e) If for some $i \in (k+1, n]$ and $j \in (1, \ell]$ we have $i \in N(1), j \in N(k+1)$ and $i^-j^- \in E(G)$, then $\overrightarrow{C}[n, i]\overrightarrow{C}[1, j^-]\overleftarrow{C}[i^-, k+1]\overrightarrow{C}[j, k]$ is an (n, k)-H-path.

We shall show that at least one of these situations must occur. Suppose for a contradiction that this is not the case. Then for every $\ell \in (1, k]$

(1)
$$E[(N(1) \cap (1, \ell])^-, (N(k+1) \cap (\ell, k])^+ \cup (N(k+1) \cap (k+1, n])^-] = \emptyset,$$

since (a), (b) and (c) do not occur; and

(2)
$$E[(N(1) \cap (\ell, k])^{-} \cup (N(1) \cap (k+1, n])^{-}, (N(k+1) \cap (1, \ell])^{-}] = \emptyset,$$

since (d) and (e) do not occur.

Then equation (1) implies $|(N(k+1) \cap (\ell, k])^+ \cup (N(k+1) \cap (k+1, n])^-| < t$. Since the two sets $(N(k+1) \cap (\ell, k])^+$ and $(N(k+1) \cap (k+1, n])^-$ both contain the vertex k+1, we have

$$|(N(k+1) \cap (\ell,k])^+ \cap (N(k+1) \cap (k+1,n])^-| = |\{k+1\}| = 1$$

when $\ell < k$, and

$$|(N(k+1) \cap (\ell, k])^+ \cap (N(k+1) \cap (k+1, n])^-| = |\emptyset| = 0$$

when $\ell = k$. Then

$$\begin{split} d(k+1) &= |N(k+1) \cap (1,\ell]| + |N(k+1) \cap (\ell,k]| + |N(k+1) \cap (k+1,n]| \\ &= |N(k+1) \cap (1,\ell]| + |(N(k+1) \cap (\ell,k])^+| + |(N(k+1) \cap (k+1,n])^-| \\ &= |N(k+1) \cap (1,\ell]| + |(N(k+1) \cap (\ell,k])^+ \cup (N(k+1) \cap (k+1,n])^-| \\ &+ |(N(k+1) \cap (\ell,k])^+ \cap (N(k+1) \cap (k+1,n])^-| \\ &\geq \delta(G). \end{split}$$

Now we have $|N(k+1) \cap (1,\ell]| > \delta(G) - t - 1 \ge \widetilde{\alpha}(G) + 1 - t - 1 = s - 1$, i.e., $|N(k+1) \cap (1,\ell]| \ge s$ when $\ell < k$, and $|N(k+1) \cap (1,\ell]| > \delta(G) - t \ge \widetilde{\alpha}(G) + 1 - t = s$, i.e., $|N(k+1) \cap (1,\ell]| \ge s + 1$ when $\ell = k$. No matter whether $\ell < k$ or $\ell = k$, equation (2) implies $|(N(1) \cap (\ell,k])^- \cup (N(1) \cap (k+1,n])^-| < t$. It is obvious that $(N(1) \cap (\ell,k])^-$ and $(N(1) \cap (k+1,n])^-$ are disjoint. Hence

$$\begin{split} \delta(G) &\leq d(1) = |N(1) \cap (1,\ell]| + |N(1) \cap (\ell,k]| + |N(1) \cap (k+1,n]| \\ &= |N(1) \cap [1,\ell]| + |(N(1) \cap (\ell,k])^-| + |(N(1) \cap (k+1,n])^-| \\ &= |N(1) \cap [1,\ell]| + |(N(1) \cap (\ell,k])^- \cup (N(1) \cap (k+1,n])^-| \\ &< s+t = \widetilde{\alpha}(G) + 1 \leq \delta(G), \end{split}$$

a contradiction.

Case 2. $\ell \in (k+1, n)$. Here, we describe four situations (referring to Figure 3) in which there is an (n, k)-H-path. Recall that $1\ell \in E(G)$.



Figure 3. Situations (f)-(t).

(f) If for some $i \in (1, k]$ and $j \in (\ell, n]$ we have $i \in N(1), j \in N(k+1)$ and $i^-j^- \in E(G)$, then $\overleftarrow{C}[n, j]\overrightarrow{C}[k+1, j^-]\overrightarrow{C}[i^-, 1]\overrightarrow{C}[i, k]$ is an (n, k)-H-path.

(g) If for some $i \in (k+1, \ell]$ and $j \in (\ell, n]$ we have $i \in N(1), j \in N(k+1)$ and $i^-j^- \in E(G)$, then $\overrightarrow{C}[n, j]\overrightarrow{C}[k+1, i^-]\overleftarrow{C}[j^-, i]\overrightarrow{C}[1, k]$ is an (n, k)-H-path.

(h) If for some $i \in [\ell, n)$ and $j \in [1, k]$ we have $i \in N(1), j \in N(k+1)$ and $i^+j^- \in E(G)$, then $\overleftarrow{C}[n, i^+]\overleftarrow{C}[j^-, 1]\overleftarrow{C}[i, k+1]\overrightarrow{C}[j, k]$ is an (n, k)-H-path.

(t) If for some $i \in [\ell, n)$ and $j \in (k + 1, \ell]$ we have $i \in N(1), j \in N(k + 1)$ and $i^+j^- \in E(G)$, then $\overleftarrow{C}[n, i^+]\overleftarrow{C}[j^-, k + 1]\overrightarrow{C}[j, i]\overrightarrow{C}[1, k]$ is an (n, k)-H-path.

We shall show that at least one of these situations must occur. Suppose for a contradiction that this is not the case. Then for every $\ell \in (1, k)$

(3)
$$E[(N(1) \cap (1, \ell])^-, (N(k+1) \cap (\ell, n])^-] = \emptyset,$$

since (f) and (g) do not occur; and

(4)
$$E[(N(1) \cap [\ell, n))^+, (N(k+1) \cap [1, \ell])^-] = \emptyset,$$

since (h) and (t) do not occur.

Then equation (3) implies $|N(k+1) \cap (\ell, n]| = |(N(k+1) \cap (\ell, n])^-| < t$. Then

$$\begin{split} |(N(k+1)\cap [1,\ell])^-| &= |(N(k+1)\cap [1,\ell])| \ge \delta(G) - |(N(k+1)\cap (\ell,n])| \\ &> \delta(G) - t \ge \widetilde{\alpha}(G) + 1 - t = s + t - t = s. \end{split}$$

Now we have $|(N(k+1) \cap [1,\ell])^-| \ge s+1$. Then equation (4) implies that $|N(1) \cap [\ell,n)| = |(N(1) \cap [\ell,n))^+| < t$. Therefore

$$\begin{split} \delta(G) &\leq d(1) = |N(1) \cap (1,\ell]| + |N(1) \cap [\ell,n)| + |\{n\}| - |\{\ell\}| \\ &\leq s+t-1 = \widetilde{\alpha}(G) + 1 - 1 \leq \delta(G) - 1, \end{split}$$

a contradiction.

This completes the proof of Theorem 6.

References

- J.A. Bondy and U.S.R. Murty, Graph Theory, in: Grad. Texts in Math. 244 (Springer, New York, 2008).
- [2] V. Chvátal and P. Erdős, A note on Hamiltonian circuits, Discrete Math. 2 (1972) 111–113. https://doi.org/10.1016/0012-365X(72)90079-9
- [3] G.A. Dirac, Some theorems on abstract graphs, Proc. Lond. Math. Soc. (3) 2 (1952) 69–81. https://doi.org/10.1112/plms/s3-2.1.69
- [4] G.-H. Fan, New sufficient conditions for cycles in graphs, J. Combin. Theory Ser. B 37 (1984) 221–227. https://doi.org/10.1016/0095-8956(84)90054-6
- R.J. Faudree, R.J. Gould, M.S. Jacobson and R.H. Schelp, Neighborhood unions and hamiltonian properties in graphs, J. Combin. Theory Ser. B 47 (1989) 1–9. https://doi.org/10.1016/0095-8956(89)90060-9
- [6] R. Gould, Advances on the Hamiltonian problem—A survey, Graphs Combin. 19 (2003) 7–52. https://doi.org/10.1007/s00373-002-0492-x
- H. Li, Generalizations of Dirac's theorem in Hamiltonian graph theory—A survey, Discrete Math. 313 (2013) 2034–2053. https://doi.org/10.1016/j.disc.2012.11.025
- [8] C. McDiarmid and N. Yolov, Hamilton cycles, minimum degree, and bipartite holes, J. Graph Theory 86 (2017) 277–285. https://doi.org/10.1002/jgt.22114

- [9] O. Ore, Note on Hamilton circuits, Amer. Math. Monthly 67 (1960) 55. https://doi.org/10.2307/2308928
- R.H. Shi, 2-neighborhoods and hamiltonian conditions, J. Graph Theory 16 (1992) 267–271. https://doi.org/10.1002/jgt.3190160310

Received 7 May 2021 Revised 8 July 2022 Accepted 8 July 2022 Available online 22 July 2022

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License https://creativecommons.org/licenses/by-nc-nd/4.0/