

COLORING SQUARES OF PLANAR GRAPHS WITH SMALL MAXIMUM DEGREE*

MATEUSZ KRZYZIŃSKI¹, PAWEŁ RZAŻEWSKI^{1,2}

AND

SZYMON TUR¹

¹*Faculty of Mathematics and Information Science
Warsaw University of Technology
Warsaw, Poland*

²*Institute of Informatics, University of Warsaw
Warsaw, Poland*

e-mail: mateusz.krzyzinski.stud@pw.edu.pl
pawel.rzazewski@pw.edu.pl
szymon.tur2.stud@pw.edu.pl

Abstract

For a graph G , by $\chi_2(G)$ we denote the minimum integer k , such that there is a k -coloring of the vertices of G in which vertices at distance at most 2 receive distinct colors. Equivalently, $\chi_2(G)$ is the chromatic number of the square of G . In 1977 Wegner conjectured that if G is planar and has maximum degree Δ , then $\chi_2(G) \leq 7$ if $\Delta \leq 3$, $\chi_2(G) \leq \Delta + 5$ if $4 \leq \Delta \leq 7$, and $\lfloor 3\Delta/2 \rfloor + 1$ if $\Delta \geq 8$. Despite extensive work, the known upper bounds are quite far from the conjectured ones, especially for small values of Δ . In this work we show that for every planar graph G with maximum degree Δ it holds that $\chi_2(G) \leq 3\Delta + 4$. This result provides the best known upper bound for $6 \leq \Delta \leq 14$.

Keywords: Wegner's conjecture, coloring squares of planar graphs, discharging, graph coloring.

2020 Mathematics Subject Classification: 05C15.

*MK and ST were supported by the “Szkola Orłów” (“School of Eagles”) project, co-financed by the European Social Fund under the Knowledge-Education-Development Operational Programme, Axis III, Higher Education For The Economy And Development, measure 3.1, Competences In Higher Education. PRz was supported by the project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme Grant Agreement 714704.



European Research Council
Established by the European Commission



1. INTRODUCTION

Graph coloring is undoubtedly among the best studied problems in graph theory. The origins of the research on graph colorings date back to 19th century and are related to the question whether all planar graphs can be properly colored with four colors. The affirmative answer to this question, i.e., the celebrated four color theorem, remains one of the most famous results in graph theory [4, 5, 18]. The study of the restrictions and generalizations of the problem [2, 10–13, 19] led to many exciting results and much better understanding of the structure of planar graphs.

Other variants of coloring planar graphs were also considered. Already in 1977, Wegner [22] studied the problem of coloring graphs in such a way that the vertices with the same color must be at distance more than d , where d is a fixed integer. Such a coloring is called *distance- d coloring* and the minimum number of colors in a distance- d coloring of a graph G is denoted by $\chi_d(G)$. Note that for $d = 1$ we obtain the classic graph coloring problem. The next case that has received the most attention is $d = 2$. The problem of distance-2 coloring is also known as *$L(1, 1)$ -labeling* [8]. Let us give a brief overview of the known results on distance-2 coloring of planar graphs. In what follows G is a planar graph with maximum degree Δ .

First, observe that, in contrast to the classic coloring, there is no universal constant c such that $\chi_2(G) \leq c$ for all planar graphs G . Indeed, for the n -vertex star $K_{1,n-1}$ it holds that $\chi_2(K_{1,n-1}) = n$. This implies that every graph G satisfies $\chi_2(G) \geq \Delta + 1$. On the other hand, as every planar graph has a vertex of degree at most 5, a simple greedy algorithm yields the bound $\chi_2(G) \leq 5\Delta + 1$. Thus $\chi_2(G)$ is bounded by a linear function of Δ .

Wegner [22] was probably the first who studied this dependence. Among other results, he showed that if G is planar and has maximum degree at most 3, then $\chi_2(G) \leq 8$. He also presented some families of planar graphs which require a large number of colors in any distance-2 coloring and conjectured that the lower bounds given by these families are actually tight. This problem is known as Wegner's conjecture.

Wegner's Conjecture. *Every planar graph G with maximum degree Δ satisfies*

$$\chi_2(G) \leq \begin{cases} 7 & \text{if } \Delta \leq 3, \\ \Delta + 5 & \text{if } 4 \leq \Delta \leq 7, \\ \lfloor \frac{3\Delta}{2} \rfloor + 1 & \text{if } \Delta \geq 8. \end{cases}$$

The problem of bounding $\chi_2(G)$ received a considerable attention. However, despite many partial results, the only case for which we know tight bound is $\Delta = 3$: Thomassen [20] confirmed the conjecture by showing that seven colors

always suffice. For $\Delta \geq 4$ the conjecture is wide open; we summarize the known bounds in Table 1.

Authors	Restriction	Result
Thomassen [20]	$\Delta \leq 3$	$\chi_2(G) \leq 7$
Jonas [15]	$\Delta \geq 7$	$\chi_2(G) \leq 8\Delta - 22$
Wong [23]	$\Delta \geq 7$	$\chi_2(G) \leq 3\Delta + 5$
Madaras and Marcinová [16]	$\Delta \geq 12$	$\chi_2(G) \leq 2\Delta + 18$
Borodin <i>et al.</i> [6]	$\Delta \leq 20$	$\chi_2(G) \leq 59$
	$21 \leq \Delta \leq 46$	$\chi_2(G) \leq \Delta + 39$
	$\Delta \geq 47$	$\chi_2(G) \leq \lceil \frac{9\Delta}{5} \rceil + 1$
van den Heuvel and McGuinness [21]	$\Delta \geq 5$	$\chi_2(G) \leq 9\Delta - 19$
		$\chi_2(G) \leq 2\Delta + 25$
Agnarsson and Halldórsson [1]	$\Delta \geq 749$	$\chi_2(G) \leq \lfloor \frac{9\Delta}{5} \rfloor + 2$
Molloy and Salavatipour [17]	$\Delta \geq 241$	$\chi_2(G) \leq \lceil \frac{5\Delta}{3} \rceil + 25$
		$\chi_2(G) \leq \lceil \frac{5\Delta}{3} \rceil + 78$
Zhu and Bu [24]	$\Delta \leq 5$	$\chi_2(G) \leq 20$
	$\Delta \geq 6$	$\chi_2(G) \leq 5\Delta - 7$
Cranston and Rabern [9]	$\Delta \geq 3$	$\chi_2(G) \leq \Delta^2 - 1$
Bousquet <i>et al.</i> [7]	$\Delta = 4$	$\chi_2(G) \leq 12$
Amini <i>et al.</i> [3]		$\chi_2(G) \leq \frac{3}{2}\Delta + o(\Delta)$
Havet <i>et al.</i> [14]		$\chi_2(G) \leq \frac{3}{2}\Delta(1 + o(1))$

Table 1. The progress on Wegner's conjecture.

Let us highlight that the currently best known bound asymptotically is $\frac{3}{2}\Delta(1 + o(1))$, independently showed by Amini *et al.* [3] and Havet *et al.* [14], and the best known bound of type $c\Delta + \mathcal{O}(1)$, where c is a constant, is $\lceil \frac{5\Delta}{3} \rceil + \mathcal{O}(1)$ by Molloy and Salavatipour [17]. However, since the additive constant is large, this bound

is very far from the conjectured one for small values of Δ . Thus some attention has been put on refining the bounds for graphs with small maximum degree, see Figure 1. We continue this line of research and show the following result.




Figure 1. Dependence of $\chi_2(G)$ on Δ for $3 \leq \Delta \leq 23$.

Theorem 1. *Every planar graph G with maximum degree $\Delta \geq 6$ satisfies $\chi_2(G) \leq 3\Delta + 4$.*

We point out that Theorem 1 provides the best known upper bound for the cases $6 \leq \Delta \leq 14$.

The proof of Theorem 1 uses the discharging method. We consider a minimal counterexample G and we fix its plane embedding. In Section 3.1 we show that G cannot contain certain subgraphs, as this would contradict the minimality of G . Then, in Section 3.2, we distribute some integer values, called *charges*, to the vertices and faces of G in a way that the total charge is negative. Next, we apply six *discharging rules* to transfer charges between the vertices and faces of G . Eventually, we analyze the final charges and find out that every vertex and every face has nonnegative charge. As in the discharging phase no charge is created nor lost, this is a contradiction. Thus the counterexample to Theorem 1 cannot exist.

2. PRELIMINARIES

All graphs considered in the paper are simple and finite. For a graph G , by $V(G)$ and $E(G)$ we denote, respectively, the vertex set and the edge set of G . Furthermore, if G is planar and given along with a fixed plane embedding, then $F(G)$ denotes the set of faces of G .

For two vertices v and u , by $\text{dist}_G(v, u)$ we denote the *distance* between these vertices, i.e., the number of edges on a shortest u - v path in G . For a vertex v , by $N_G(v)$ we denote its *neighborhood*, i.e., the set of all vertices adjacent to v , and by $\deg_G(v)$ we denote the *degree* of a vertex v , i.e., $|N_G(v)|$. The maximum and the minimum degree of G are denoted by, respectively, $\Delta(G)$ and $\delta(G)$.

For a vertex $v \in V(G)$, by $G - v$ we denote the graph obtained from G by removing v with all incident edges. For $u, v \in V(G)$, by $G + uv$ we denote the graph with vertex set $V(G)$ and edge set $E(G) \cup \{uv\}$. Note that if $uv \in E(G)$, then $G = G + uv$. In other words, we never create multiple edges.

Each face f is bounded by a closed walk called a *boundary*. We write $f = [v_1, v_2, \dots, v_k]$ to denote the cyclic ordering of vertices along the boundary of f . Note that if G is 2-connected, then the boundary of each face is a cycle; we will always work in this setting.

We say that a vertex v and a face f are *incident* if v lies on the boundary of f . By $\deg_G(f)$ we denote the *degree* of a face f , i.e., the number of vertices incident to f .

If the graph G is clear from the context, we drop the subscript in the notation above.

For an integer d , a vertex v is said to be a d -vertex (respectively, a d^+ -vertex, a d^- -vertex) if its degree is exactly d (respectively, at least d , at most d). Similarly, a face f is said to be a d -face (respectively, a d^+ -face, a d^- -face) if its degree is exactly d (respectively, at least d , at most d).

Often we will consider a situation where some subset of vertices of a graph G is colored. For an uncolored vertex v , we say that a color is *blocked* if it appears on a vertex within distance at most 2 from v . A color that is not blocked is *free*.

3. MAIN PROOF

By contradiction, suppose that Theorem 1 does not hold and let G be a minimum counterexample. Thus, for any planar graph G' , if $|V(G')| + |E(G')| < |V(G)| + |E(G)|$ and $\Delta(G') \leq \Delta$, then $\chi_2(G') \leq 3\Delta + 4$.

We can assume that G is connected, as the coloring of G can be obtained by coloring each connected component independently and each of them is smaller than G .

Fix some plane embedding of G . Whenever we refer to faces of G , we mean the faces of this fixed plane embedding.

3.1. Forbidden configurations

In this section we present a series of technical claims in which we analyze the structure of the graph G .

Claim 2. G is 2-connected.

Proof. Assume that G has a cutvertex v and let C be the vertex set of one connected component of $G - v$. Define $G'_1 = G[C \cup \{v\}]$ and $G'_2 = G - C$. By the minimality of G , for each $i \in \{1, 2\}$ there is a distance-2 $(3\Delta + 4)$ -coloring φ_i of G'_i . We can permute the colors in φ_2 so that (i) $\varphi_1(v) = \varphi_2(v)$ and (ii) $\varphi_1(N_{G_1}(v)) \cap \varphi_2(N_{G_2}(v)) = \emptyset$. The union of these colorings is a distance-2 $(3\Delta + 4)$ -coloring of G , a contradiction. ■

So by Claim 2 from now on we can assume that the boundary of each face is a simple cycle with at least three edges.

The proofs of the next few claims follow the same outline. First, assume that G contains some configuration that we want to exclude. We modify G by removing a single vertex v and possibly adding some new edges, in order to obtain a graph G' with the following properties:

- (i) $|V(G')| + |E(G')| \leq |V(G)| + |E(G)|$,
- (ii) $\Delta(G') \leq \Delta$,
- (iii) G' is planar (and its plane embedding can be easily obtained from the plane embedding of G),
- (iv) all pairs of vertices in $V(G) \setminus \{v\}$ that are at distance at most 2 in G are at distance at most 2 in G' .

By properties (i), (ii), and (iii) and the minimality of G we observe that G' has a distance-2 $(3\Delta + 4)$ -coloring φ . By property (iv), we can safely color all vertices of $V(G) \setminus \{v\}$ according to φ . To obtain a distance-2 $(3\Delta + 4)$ -coloring for G we only need to find a color for v . We do this by ensuring that the number of colors that are blocked for v is strictly less than $3\Delta + 4$. Thus there is a free color for v , as G admits a distance-2 $(3\Delta + 4)$ -coloring, a contradiction.

For brevity, in the proofs we only say how to define G' and compute the number of colors that are blocked for v . In particular, we will not explicitly check properties (i)–(iv), as verifying them is straightforward.

Claim 3. $\delta(G) \geq 3$.

Proof. First suppose that $\deg(v) = 1$. We set $G' = G - v$ and observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $\Delta < 3\Delta + 4$ colors for v .

Now, assume that $\deg(v) = 2$, let $N(v) = \{v_1, v_2\}$, see Figure 2. Let $G' = G - v + v_1v_2$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $2\Delta < 3\Delta + 4$ colors for v . ■




Figure 2. Forbidden configuration in Claim 3.

Claim 4. G has no two adjacent 3-vertices.

Proof. Suppose u and v are adjacent 3-vertices and let $N(v) = \{u, v_1, v_2\}$, see Figure 3. Let $G' = G - v + uv_1 + uv_2$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $2\Delta + 3 < 3\Delta + 4$ colors for v . ■



Figure 3. Forbidden configuration in Claim 4.

Claim 5. G has no 3-vertex incident to a 3-face.

Proof. Suppose that G has a 3-vertex v incident to a 3-face $[v, v_1, v_2]$ and let $N(v) = \{v_1, v_2, v_3\}$, see Figure 4. Let $G' = G - v + v_1v_3$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $\Delta + 2(\Delta - 1) = 3\Delta - 2 < 3\Delta + 4$ colors for v . ■




Figure 4. Forbidden configuration in Claim 5.

Claim 6. *G has no 3-vertex incident to two 4-faces.*

Proof. Suppose that G has a 3-vertex v incident to two 4-faces: $[v, v_1, u, v_2]$ and $[v, v_3, w, v_2]$, see Figure 5. Let $G' = G - v + v_1v_3$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $2(\Delta - 1) + \Delta = 3\Delta - 2 < 3\Delta + 4$ colors for v . ■




Figure 5. Forbidden configuration in Claim 6.

Claim 7. *If a 4-vertex of G is incident to a 3-face, then the other two vertices on that 3-face are 6^+ -vertices.*

Proof. Suppose that G has a 4-vertex v incident to a 3-face $[v, v_1, v_2]$, where $\deg(v_1) \leq 5$ and $N(v) = \{v_1, v_2, v_3, v_4\}$, see Figure 6. Let $G' = G - v + v_1v_3 + v_1v_4$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $2\Delta + (\Delta - 1) + 4 = 3\Delta + 3 < 3\Delta + 4$ colors for v . ■




Figure 6. Forbidden configuration in Claim 7.

Claim 8. *If G has two 3-faces that share an edge uv on their boundaries and $\deg(u) = 4$, then $\deg(v) \geq 8$.*

Proof. Suppose that G has a 4-vertex u incident to two adjacent 3-faces: $[v, v_1, u]$ and $[v, v_2, u]$, where $\deg(v) \leq 7$ and $N(u) = \{v, v_1, v_2, v_3\}$; see Figure 7. Let $G' = G - u + vv_3$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $\Delta + 2(\Delta - 1) + 5 = 3\Delta + 3 < 3\Delta + 4$ colors for u . ■





Figure 7. Forbidden configuration in Claim 8.

Claim 8 implies two statements that will be directly used in our proof. Let us start with some definitions. A 3-face is *weird* if it is incident to a 4-vertex and two 6^+ -vertices. Consider a vertex v and let f_0, f_1, \dots, f_{d-1} be the sequence of faces incident to v in the cyclic ordering around v in our fixed plane embedding of G . Arithmetic operations on indices will be performed modulo d . A *fan centered at v* is a sequence $\mathbf{f} = f_i, f_{i+1}, \dots, f_{i+p}$, $p \in \{0, 1, \dots, d-2\}$, such that (i) each element of \mathbf{f} is a 3-face, and (ii) f_{i-1} and f_{i+p+1} are not 3-faces (it is possible that $f_{i-1} = f_{i+p+1}$); see Figure 8. The faces f_i and f_{i+p} are *outer faces* of \mathbf{f} . By the vertices of \mathbf{f} we mean the vertices incident to the faces of \mathbf{f} , except for v . The vertices of f_i and f_{i+p} that are not shared with other elements of \mathbf{f} are *outer vertices* of \mathbf{f} .


 Figure 8. Fans centered at a vertex v are shaded and their outer faces are darker. Outer vertices are marked with squares.

Claim 9. *If v is a 7^- -vertex, $d(v) > 4$ and all faces incident to v are 3-faces, then v is not incident to a weird 3-face.*

Proof. By Claim 8 we observe that v is adjacent to no 4-vertex and thus is not incident to a weird 3-face. ■

Claim 10. *Let v be a 7^- -vertex, $d(v) > 4$ and let \mathbf{f} be a fan centered at v . If \mathbf{f} contains a weird 3-face f , then f is an outer face of \mathbf{f} and the 4-vertex of f is an outer vertex of \mathbf{f} .*

Proof. Observe that if the 4-vertex v of f is not an outer vertex of \mathbf{f} , then the edge uv belongs to the boundaries of two 3-faces, which contradicts Claim 8. ■

Claim 11. *If a 5-vertex of G is incident to five 3-faces, then it has at least four 7^+ -vertices as neighbors.*

Proof. Suppose that G has a 5-vertex v incident to five 3-faces, see Figure 9. Let v_1, v_2 be distinct 6^- -vertices in $N(v)$. Let $G' = G - v$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $2 \cdot 4 + 3(\Delta - 2) = 3\Delta + 2 < 3\Delta + 4$ colors for v . ■



Figure 9. Forbidden configuration in Claim 11.

Claim 12. *Let v be a 5-vertex of G adjacent to exactly one 4^+ -face, and let v_1 and v_2 be the neighbors of v on the boundary of this 4^+ -face. Then either both v_1 and v_2 are 6^+ -vertices, or at least one of them is a 7^+ -vertex.*

Proof. Let v, v_1, v_2 be as in the assumptions of the claim (see Figure 10) and suppose that the statement does not hold. In particular, $\deg(v_1) + \deg(v_2) \leq 11$. Let $G' = G - v + v_1v_2$. We observe that a distance-2 $(3\Delta + 4)$ -coloring of G' blocks at most $(\deg(v_1) + \deg(v_2) - 2) + 3(\Delta - 2) \leq 3\Delta + 3 < 3\Delta + 4$ colors for v . ■

In the next claims we will need the following definitions. For a vertex v , a vertex $u \in N(v)$ is a *bad neighbor* of v if (i) $\deg(u) = 5$ and (ii) u is incident to four 3-faces and one 4^+ -face f , and (iii) v and u are consecutive vertices of the boundary of f . A vertex $u \in N(v)$ is a *very bad neighbor* of v if $\deg(u) = 5$ and u is incident to five 3-faces.

Claim 13. *The very bad neighbors of any vertex are pairwise nonadjacent.*




Figure 10. Forbidden configuration in Claim 12.

Proof. Assume that some vertex has two adjacent very bad neighbors v_1, v_2 , see Figure 11. Let $G' = G - v_1v_2$. By the minimality of G , we observe that G' admits a distance-2 $(3\Delta + 4)$ -coloring φ . Finally, note that φ is also a distance-2 coloring of G , a contradiction. ■



Figure 11. Forbidden configuration in Claim 13.

The proofs of Claims 14 and 15 are again similar to each other. In both of them the forbidden configuration involves two adjacent vertices v_1 and v_2 . We assume that such vertices exist in G and obtain a new graph G' by removing the edge v_1v_2 . Note that all pairs of vertices from $V(G) \setminus \{v_1, v_2\}$ that are at distance at most 2 in G remain so in G' . By the minimality of G , we observe that G' admits a distance-2 $(3\Delta + 4)$ -coloring. However, after restoring the edge v_1v_2 the colors assigned to v_1 and v_2 might be in conflict with each other and with the colors of some other vertices. Thus we erase the colors of v_1 and v_2 , and recolor these vertices in a greedy way. Again, we ensure that this is possible by counting the number of blocked colors.

Claim 14. *If v is an 8^- -vertex, v_1 is its very bad neighbor and v_2 is its bad neighbor, then v_1 and v_2 are nonadjacent.*

Proof. By contradiction, suppose that G has three vertices v, v_1, v_2 as in the assumption and v_1 is adjacent to v_2 , see Figure 12. Let $G' = G - v_1v_2$ and

consider a distance-2 $(3\Delta + 4)$ -coloring of G' with colors of v_1 and v_2 erased. We observe that the number of colors that are blocked for v_1 is at most $2(\Delta - 2) + (\Delta - 3) + 3 + 6 = 3\Delta + 2 < 3\Delta + 4$. Then the number of colors that are blocked for v_2 is at most $(\Delta - 1) + (\Delta - 2) + (\Delta - 3) + 3 + 6 = 3\Delta + 3 < 3\Delta + 4$. ■




Figure 12. Forbidden configuration in Claim 14.

Claim 15. *If v is a 7-vertex and v_1, v_2 are its bad neighbors, then v_1 and v_2 are nonadjacent.*

Proof. By contradiction, suppose that G has vertices v, v_1, v_2 as in the assumption of the claim and v_1 is adjacent to v_2 , see Figure 13. Let $G' = G - v_1v_2$ and consider a distance-2 $(3\Delta + 4)$ -coloring of G' with colors of v_1 and v_2 erased. Note that the number of colors blocked for each of v_1, v_2 is at most $(\Delta - 1) + (\Delta - 2) + (\Delta - 3) + 3 + 6 = 3\Delta + 3 < 3\Delta + 4$. ■




Figure 13. Forbidden configuration in Claim 15.

3.2. Discharging

We give an initial charge of $\mathfrak{w}(x) = \deg(x) - 4$ to every $x \in V(G) \cup F(G)$. Using the Euler's formula $|V(G)| - |E(G)| + |F(G)| = 2$ and the handshaking lemma

$2|E(G)| = \sum_{v \in V(G)} \deg(v) = \sum_{f \in F(G)} \deg(f)$, we derive the equality

$$(1) \quad \sum_{x \in V(G) \cup F(G)} \mathfrak{w}(x) = \sum_{x \in V(G) \cup F(G)} (\deg(x) - 4) = -8.$$

The charges will be transferred between the elements of $V(G) \cup F(G)$ according to six discharging rules. As we will see at the end, after the application of these rules, each vertex and face will have a non-negative charge. Thus the total charge must be non-negative, which contradicts (1). This shows that a counterexample to Theorem 1 cannot exist.

Discharging rules

We apply the following discharging rules.

- R1** Every 5^+ -vertex sends $\frac{1}{3}$ to each incident 3-face.
- R2** Every 6^+ -vertex sends additional $\frac{1}{6}$ to each incident weird 3-face.
- R3** Every 5^+ -face sends $\frac{1}{2}$ to each incident 3-vertex.
- R4** Every 6-vertex sends $\frac{1}{6}$ to each bad neighbor.
- R5** Every 7^+ -vertex sends $\frac{1}{3}$ to each bad neighbor.
- R6** Every 7^+ -vertex sends $\frac{1}{6}$ to each very bad neighbor.

Final charges

For $x \in V(G) \cup V(F)$, let $\mathfrak{w}'(x)$ be the final charge (after applying discharging rules). We aim to show that $\mathfrak{w}'(x) \geq 0$ for all $x \in V(G) \cup F(G)$.

Let us start by bounding $\mathfrak{w}'(v)$ for $v \in V(G)$. As by Claim 3 we have $\deg(v) \geq 3$. The analysis is split into cases depending on the degree of v . Note that according to discharging rules, only 3-vertices and 5-vertices which are bad or very bad neighbors may receive any charge.

Case $\deg(v) = 3$. By Claims 5 and 6, v is incident to at least two 5^+ -faces. Note that v does not lose any charge and by **R3** it receives charge at least $2 \cdot \frac{1}{2} = 1$. Consequently $\mathfrak{w}'(v) \geq \mathfrak{w}(v) + 1 = (3 - 4) + 1 = 0$.

Case $\deg(v) = 4$. As v does not lose nor receive any charge, we have $\mathfrak{w}'(v) = \mathfrak{w}(v) = (4 - 4) = 0$.

Case $\deg(v) = 5$. Recall that v loses charge due to **R1** and might receive charge by **R4**, **R5**, and **R6**. Consider the following subcases.

- If v is incident to at most three 3-faces, then the charge lost by v due to **R1** is at most $3 \cdot \frac{1}{3} = 1$. Thus, $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - 1 = (5 - 4) - 1 = 0$.

- If v is incident to four 3-faces, then due to **R1** the charge lost is $4 \cdot \frac{1}{3} = \frac{4}{3}$. By Claim 12, rules **R4** and **R5** make v receive charge at least $\min\{2 \cdot \frac{1}{6}, \frac{1}{3}\} = \frac{1}{3}$. Thus, $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - \frac{4}{3} + \frac{1}{3} = (5 - 4) - 1 = 0$.
- If v is incident to five 3-faces, then v loses charge $5 \cdot \frac{1}{3} = \frac{5}{3}$ by **R1**. By Claim 11, the application of **R6** makes v receive charge at least $4 \cdot \frac{1}{6} = \frac{2}{3}$. Thus, $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - \frac{5}{3} + \frac{2}{3} = (5 - 4) - 1 = 0$.

Case $\deg(v) = 6$. Recall that v never receives any charge and loses charge due to **R1**, **R2**, and **R4**. Consider the following subcases.

- If v is incident to six 3-faces, then v has no bad neighbors. Furthermore, by Claim 9, v is not incident to a weird 3-face. Therefore v loses charge only due to **R1** and the total charge lost is $6 \cdot \frac{1}{3} = 2$, so $\mathfrak{w}'(v) = \mathfrak{w}(v) - 2 = (6 - 4) - 2 = 0$.
- Now consider the case that v is incident to at most five 3-faces. Let \mathbf{F} be the set of fans centered at v . For each $\mathbf{f} \in \mathbf{F}$, there are $|\mathbf{f}| + 1$ neighbors of v that are incident to a face from \mathbf{f} . Furthermore, each neighbor of v is incident to a face of at most one fan. Thus, $\sum_{\mathbf{f} \in \mathbf{F}} (|\mathbf{f}| + 1) \leq \deg(v) = 6$.

If a fan $\mathbf{f} \in \mathbf{F}$ contains a weird 3-face f , then, by Claim 10, the face f can only be an outer face of \mathbf{f} . Furthermore, the 4-vertex from the boundary of f must be an outer vertex of \mathbf{f} . Moreover, every bad neighbor of v is an outer vertex of some fan in \mathbf{F} . Thus the total number of bad neighbors and weird 3-faces incident to v is bounded by $2|\mathbf{F}|$. Summing up, by **R1**, **R2**, and **R4**, the total charge lost by v is at most

$$\frac{1}{3} \sum_{\mathbf{f} \in \mathbf{F}} |\mathbf{f}| + \frac{1}{6} \cdot 2 \cdot |\mathbf{F}| = \frac{1}{3} \sum_{\mathbf{f} \in \mathbf{F}} |\mathbf{f}| + \frac{1}{3} \sum_{\mathbf{f} \in \mathbf{F}} 1 = \frac{1}{3} \sum_{\mathbf{f} \in \mathbf{F}} (|\mathbf{f}| + 1) \leq \frac{\deg(v)}{3} = 2.$$

Consequently, $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - 2 = (6 - 4) - 2 = 0$.

Case $\deg(v) = 7$. Recall that v never receives any charge and loses charge due to **R1**, **R2**, **R5**, and **R6**. Consider the following subcases.

- If v is incident to seven 3-faces, then v has no bad neighbor and, by Claim 13, v has at most three very bad neighbors. Furthermore, by Claim 9, v is not incident to a weird 3-face. Thus, by **R1** and **R6**, the total charge lost by v is at most $7 \cdot \frac{1}{3} + 3 \cdot \frac{1}{6} = \frac{17}{6}$ and thus $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - \frac{17}{6} = (7 - 4) - \frac{17}{6} = \frac{1}{6} \geq 0$.
- Now consider the case that v is incident to at most six 3-faces. Let \mathbf{F} be the set of fans centered at v . Recall that any two elements of \mathbf{F} are separated by at least one 4^+ -face incident to v . We proceed similarly as for the case of 6-vertices.

Consider a fan $\mathbf{f} \in \mathbf{F}$. By Claim 10, if \mathbf{f} contains a weird 3-face f , then f must be an outer face of \mathbf{f} and the 4-vertex on the boundary of f must be an outer vertex of \mathbf{f} . By Claims 13, 14 and 15, bad and very bad neighbors of v are pairwise nonadjacent. Moreover, every bad neighbor which is a vertex of \mathbf{f} must be an outer vertex of \mathbf{f} .

Let $w(\mathbf{f})$ be total charge sent by v to the vertices and faces of \mathbf{f} . The contribution of **R1** to $w(\mathbf{f})$ is exactly $\frac{1}{3} \cdot |\mathbf{f}|$.

If $|\mathbf{f}| = 1$, then the possible cases are as follows: (a) the unique face in \mathbf{f} is weird and none of the vertices of \mathbf{f} is a bad or a very bad neighbor of v , (b) the unique face in \mathbf{f} is not weird, one of the vertices of \mathbf{f} is a bad neighbor of v , and the other one is not bad nor very bad, (c) the unique face in \mathbf{f} is not weird, one of the vertices of \mathbf{f} is a very bad neighbor of v , and the other one is not bad nor very bad, (d) the unique face in \mathbf{f} is not weird and none of the vertices of \mathbf{f} is a bad or a very bad neighbor of v . Summing up, the total contribution of **R2**, **R5**, and **R6** to $w(\mathbf{f})$ is at most $\frac{1}{3}$ (this happens in case (b) above).

Now consider the case that $|\mathbf{f}| > 1$. The total charge sent to the outer faces of \mathbf{f} and their vertices is at most $\frac{2}{3}$; it happens if each of outer vertices is a bad neighbor. Now let us consider the vertices u of \mathbf{f} that are not incident to the outer faces, note that there are exactly $(|\mathbf{f}| + 1) - 4 = |\mathbf{f}| - 3$ of them. Due to **R6**, the vertex v sends charge $\frac{1}{6}$ to u if u is a very bad neighbor of v . Since very bad neighbors are pairwise nonadjacent, we conclude that the number of very bad neighbors that are not the vertices of outer faces of \mathbf{f} is at most $\left\lceil \frac{|\mathbf{f}| - 3}{2} \right\rceil$.

Summing up, we obtain that $w(\mathbf{f}) \leq \alpha(|\mathbf{f}|)$, where

$$\alpha(t) = \begin{cases} \frac{4}{6} & \text{if } t = 1, \\ \frac{8}{6} & \text{if } t = 2, \\ \frac{10}{6} & \text{if } t = 3, \\ \frac{13}{6} & \text{if } t = 4, \\ \frac{15}{6} & \text{if } t = 5, \\ \frac{18}{6} & \text{if } t = 6. \end{cases}$$

A straightforward case analysis shows that $\sum_{\mathbf{f} \in \mathbf{F}} \alpha(|\mathbf{f}|) \leq 3$. Thus we obtain that $\mathbf{w}'(v) = \mathbf{w}(v) - \sum_{\mathbf{f} \in \mathbf{F}} w(\mathbf{f}) \geq (7 - 4) - \sum_{\mathbf{f} \in \mathbf{F}} \alpha(|\mathbf{f}|) \geq 3 - 3 = 0$.

Case $\deg(v) \geq 8$. Similarly to the previous case, v never receives any charge and loses charge due to **R1**, **R2**, **R5**, and **R6**. Consider the following subcases.

- If all faces incident to v are 3-faces, then v has no bad neighbor and thus **R5** does not apply. Recall that if v is incident to a weird 3-face, then no vertex incident to that face is a very bad neighbor of v . So the total charge lost by v due to **R1**, **R2**, and **R6** is at most $\frac{1}{3} \cdot \deg(v) + \frac{1}{6} \cdot \deg(v) = \frac{1}{2} \cdot \deg(v)$. This gives us the final charge $\mathbf{w}'(v) \geq \mathbf{w}(v) - \frac{1}{2} \deg(v) = (\deg(v) - 4) - \frac{1}{2} \deg(v) \geq 0$ as $\deg(v) \geq 8$.
- Assume that v is incident to some 4^+ face. Let \mathbf{F} be the set of fans centered at v and consider some $\mathbf{f} \in \mathbf{F}$. Again, let $w(\mathbf{f})$ be the total charge sent by v to vertices and faces of \mathbf{f} . The contribution of **R1** to $w(\mathbf{f})$ is $\frac{1}{3}|\mathbf{f}|$. Denoting by p the number of weird 3-faces in \mathbf{f} , we obtain that the contribution of **R2** to $w(\mathbf{f})$ is $\frac{1}{6}p$. Recall that bad and very bad neighbors of v are 5-vertices. By Claim 7,

the number of 5-vertices in \mathbf{f} is at most $(|\mathbf{f}| + 1) - (p + 1) = |\mathbf{f}| - p$. Recall that if \mathbf{f} contains a bad neighbor u of v , then u must be an outer vertex of \mathbf{f} , so at most 2 of these 5-vertices are bad neighbors of v . Thus the total contribution of **R5** and **R6** to $w(\mathbf{f})$ is at most $\frac{1}{3} \cdot 2 + \frac{1}{6} \cdot (|\mathbf{f}| - p - 2)$. Summing up, we obtain $w(\mathbf{f}) \leq \frac{1}{3}|\mathbf{f}| + \frac{1}{6}p + \frac{1}{3} \cdot 2 + \frac{1}{6} \cdot (|\mathbf{f}| - p - 2) = \frac{1}{2}|\mathbf{f}| + \frac{1}{3} \leq \frac{1}{2}(|\mathbf{f}| + 1)$. So the final charge is $\mathfrak{w}'(v) \geq \mathfrak{w}(v) - \sum_{\mathbf{f} \in \mathbf{F}} w(\mathbf{f}) \geq (\deg(v) - 4) - \sum_{\mathbf{f} \in \mathbf{F}} \frac{1}{2}(|\mathbf{f}| + 1) = (\deg(v) - 4) - \frac{1}{2}\deg(v) \geq 0$ as $\deg(v) \geq 8$.

Now let us consider the values of $\mathfrak{w}'(f)$ for $f \in F(G)$. Again, we consider the cases.

Case $\deg(f) = 3$. Recall that f receives charge due to **R1** and possibly **R2** and never loses any charge. By Claims 5 and 7, f is incident to three 5^+ -vertices or at least two 6^+ -vertices. By **R1** and **R2**, f receives charge at least $\min \{3 \cdot \frac{1}{3}, 2 \cdot (\frac{1}{3} + \frac{1}{6})\} = 1$, thus the final charge is $\mathfrak{w}'(f) \geq \mathfrak{w}(f) + 1 = (3 - 4) + 1 = 0$.

Case $\deg(f) = 4$. Recall that f never sends nor receives any charge, so $\mathfrak{w}'(f) = \mathfrak{w}(v) = 4 - 4 = 0$.

Case $\deg(f) \geq 5$. Recall that f never receives any charge and loses charge due to **R3**. By Claim 4, the number of 3-vertices incident to f is at most $\left\lfloor \frac{\deg(f)}{2} \right\rfloor$. So the total charge lost by f is at most $\frac{1}{2} \cdot \left\lfloor \frac{\deg(f)}{2} \right\rfloor$ and the final charge is

$$\begin{aligned} \mathfrak{w}'(f) &\geq \mathfrak{w}(f) - \frac{1}{2} \left\lfloor \frac{\deg(f)}{2} \right\rfloor \geq \\ &\geq \begin{cases} (5 - 4) - \frac{1}{2} \cdot 2 = 0 & \text{if } \deg(f) = 5, \\ (\deg(f) - 4) - \frac{\deg(f)}{4} = \frac{3\deg(f)}{4} - 4 \geq \frac{18}{4} - 4 \geq 0 & \text{if } \deg(f) \geq 6. \end{cases} \end{aligned}$$

Summing up, we showed that for every $x \in V(G) \cup E(G)$ it holds that $\mathfrak{w}'(x) \geq 0$. As the application of discharging rules did not create any new charge, we have

$$0 \leq \sum_{x \in V(G) \cup F(G)} \mathfrak{w}'(x) = \sum_{x \in V(G) \cup F(G)} \mathfrak{w}(x),$$

which contradicts (1). Thus the hypothetical counterexample to Theorem 1 cannot exist. This completes the proof.

4. CONCLUSION

Let us recall that our Theorem 1 is the best known bound for the cases $6 \leq \Delta \leq 14$. However, it does not confirm the Wegner's conjecture for either of them. Hence the natural question arises of how the bounds can be improved.

Let us point out that all forbidden configurations appearing in our proof are quite small and can be easily analyzed by hand. However, in proofs using the discharging method, it is typical to have computer-assisted proofs that consider hundreds of configurations [4, 5, 18]. We believe that considering larger configurations, possibly with a help of a computer, could lead to a significant improvement of the upper bound.

Acknowledgment

The authors are sincerely grateful to Marthe Bonamy for introducing us to the problem.

REFERENCES

- [1] G. Agnarsson and M.M. Halldórsson, *Coloring powers of planar graphs*, SIAM J. Discrete Math. **16** (2003) 651–662.
<https://doi.org/10.1137/S0895480100367950>
- [2] N. Alon and M. Tarsi, *Colorings and orientations of graphs*, Combinatorica **12** (1992) 125–134.
<https://doi.org/10.1007/bf01204715>
- [3] O. Amini, L. Esperet and J. van den Heuvel, *A unified approach to distance-two colouring of graphs on surfaces*, Combinatorica **33** (2013) 253–296.
<https://doi.org/10.1007/s00493-013-2573-2>
- [4] K. Appel and W. Haken, *Every planar map is four colorable. Part I: Discharging*, Illinois J. Math. **21** (1977) 429–490.
<https://doi.org/10.1215/ijm/1256049011>
- [5] K. Appel, W. Haken and J. Koch, *Every planar map is four colorable. Part II: Reducibility*, Illinois J. Math. **21** (1977) 491–567.
<https://doi.org/10.1215/ijm/1256049012>
- [6] O.V. Borodin, H. Broersma, A.N. Glebov and J. van den Heuvel, *Stars and bunches in planar graphs. Part II: General planar graphs and colourings*, CDAM Research Report Series, LSE-CDAM-2002-05 (2002).
- [7] N. Bousquet, L. de Meyer, Q. Deschamps and T. Pierron, *Square coloring planar graphs with automatic discharging* (2022).
arXiv:2204.05791
- [8] T. Calamoneri, *The $L(h, k)$ -labelling problem: An updated survey and annotated bibliography*, Comput. J. **54** (2011) 1344–1371.
<https://doi.org/10.1093/comjnl/bxr037>
- [9] D.W. Cranston and L. Rabern, *Painting squares in $\Delta^2 - 1$ shades*, Electron. J. Combin. **23(2)** (2016) #P2.50.
<https://doi.org/10.37236/4978>
- [10] Z. Dvořák and L. Postle, *Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8*, J. Combin. Theory Ser. B **129** (2018) 38–54.
<https://doi.org/10.1016/j.jctb.2017.09.001>

- [11] Z. Dvořák, D. Král' and R. Thomas, *Three-coloring triangle-free graphs on surfaces V. Coloring planar graphs with distant anomalies*, J. Combin. Theory Ser. B **150** (2020) 244–269.
<https://doi.org/10.1016/j.jctb.2020.04.006>
- [12] H. Grötzsch, *Zur Theorie der diskreten Gebilde, VII: Ein Dreifarbensatz für dreikreisfreie Netze auf der Kugel*, Wiss. Z. Martin-Luther-Univ. Halle-Wittenberg, Math.-Naturwiss. Reihe **8** (1959) 109–120.
- [13] J. Grytczuk and X. Zhu, *The Alon-Tarsi number of a planar graph minus a matching*, J. Combin. Theory, Ser. B **145** (2020) 511–520.
<https://doi.org/10.1016/j.jctb.2020.02.005>
- [14] F. Havet, J. van den Heuvel, C. McDiarmid and B.A. Reed, *List colouring squares of planar graphs*, Electron. Notes Discrete Math. **29** (2007) 515–519.
<https://doi.org/10.1016/j.endm.2007.07.079>
- [15] K. Jonas, *Graph Coloring Analogues with a Condition at Distance Two: $L(2, 1)$ -Labellings and List λ -Labellings*, PhD Thesis (University of South Carolina, 1993).
- [16] T. Madaras and A. Marcinová, *On the structural result on normal plane maps*, Discuss. Math. Graph Theory **22** (2002) 293–303.
<https://doi.org/10.7151/dmgt.1176>
- [17] M. Molloy and M.R. Salavatipour, *A bound on the chromatic number of the square of a planar graph*, J. Combin. Theory Ser. B **94** (2005) 189–213.
<https://doi.org/10.1016/j.jctb.2004.12.005>
- [18] N. Robertson, D.P. Sanders, P. Seymour and R. Thomas, *The four-colour theorem*, J. Combin. Theory Ser. B **70** (1997) 2–44.
<https://doi.org/10.1006/jctb.1997.1750>
- [19] C. Thomassen, *Every planar graph is 5-choosable*, J. Combin. Theory Ser. B **62** (1994) 180–181.
<https://doi.org/10.1006/jctb.1994.1062>
- [20] C. Thomassen, *Applications of Tutte Cycles* (Technical Report, Technical University of Denmark, 2001).
- [21] J. van den Heuvel and S. McGuinness, *Coloring the square of a planar graph*, J. Graph Theory **42** (2003) 110–124.
<https://doi.org/10.1002/jgt.10077>
- [22] G. Wegner, *Graphs with Given Diameter and a Coloring Problem* (Technical Report, Technical University of Dortmund, 1977).
<https://doi.org/10.17877/DE290R-16496>
- [23] S.A. Wong, *Colouring Graphs with Respect to Distance*, Master's Thesis (University of Waterloo, 1996).

- [24] J. Zhu and Y. Bu, *Minimum 2-distance coloring of planar graphs and channel assignment*, J. Comb. Optim. **36** (2018) 55–64.
<https://doi.org/10.1007/s10878-018-0285-7>

Received 3 June 2022

Revised 14 September 2022

Accepted 15 September 2022

Available online 30 September 2022