

A STUDY OF A COMBINATION OF DISTANCE DOMINATION AND RESOLVABILITY IN GRAPHS

DWI AGUSTIN RETNOWARDANI

MOHAMMAD IMAM UTOYO

Mathematics Department, University of Airlangga
Surabaya 60115, Indonesia

e-mail: 2i.agustin@ikipjember.ac.id
m.i.utoyo@fst.unair.ac.id

DAFIK

Mathematics Education Department, University of Jember
Jember 68121, Indonesia

e-mail: d.dafik@unej.ac.id

LILIEK SUSILOWATI

Mathematics Department, University of Airlangga
Surabaya 60115, Indonesia

e-mail: lilie-k-s@fst.unair.ac.id

AND

KAMAL DLIYOU

National School of Applied Sciences (ENSA)
Ibn Zohr University
B.P. 1136, Agadir, Morocco

e-mail: dlioukamal@gmail.com

Abstract

For $k \geq 1$, in a graph $G = (V, E)$, a set of vertices D is a distance k -dominating set of G , if any vertex in $V \setminus D$ is at distance at most k from some vertex in D . The minimum cardinality of a distance k -dominating set of G is the distance k -domination number, denoted by $\gamma_k(G)$. An ordered set of vertices $W = \{w_1, w_2, \dots, w_r\}$ is a resolving set of G , if for any two distinct vertices x and y in $V \setminus W$, there exists $1 \leq i \leq r$ such that

$d_G(x, w_i) \neq d_G(y, w_i)$. The minimum cardinality of a resolving set of G is the metric dimension of the graph G , denoted by $\dim(G)$. In this paper, we introduce the distance k -resolving dominating set which is a subset of V that is both a distance k -dominating set and a resolving set of G . The minimum cardinality of a distance k -resolving dominating set of G is called the distance k -resolving domination number and is denoted by $\gamma_k^r(G)$. We give several bounds for $\gamma_k^r(G)$, some in terms of the metric dimension $\dim(G)$ and the distance k -domination number $\gamma_k(G)$. We determine $\gamma_k^r(G)$ when G is a path or a cycle. Afterwards, we characterize the connected graphs of order n having $\gamma_k^r(G)$ equal to 1, $n - 2$, and $n - 1$, for $k \geq 2$. Then, we construct graphs realizing all the possible triples $(\dim(G), \gamma_k(G), \gamma_k^r(G))$, for all $k \geq 2$. Later, we determine the maximum order of a graph G having distance k -resolving domination number $\gamma_k^r(G) = \gamma_k^r \geq 1$, we provide graphs achieving this maximum order for any positive integers k and γ_k^r . Then, we establish Nordhaus-Gaddum bounds for $\gamma_k^r(G)$, for $k \geq 2$. Finally, we give relations between $\gamma_k^r(G)$ and the k -truncated metric dimension of graphs and give some directions for future work.

Keywords: resolving set, metric dimension, distance k -domination, distance k -resolving domination.

2020 Mathematics Subject Classification: 05C12, 05C69.

1. INTRODUCTION

In this paper, we study finite, simple, and undirected graphs. For graph terminology, we refer to [9].

In 1975, Meir and Moon [28] studied a combination of two concepts distance and domination in graphs. For $k \geq 1$, we call a *distance k -dominating set* in a graph $G = (V, E)$, a subset D of the vertex set V such that for any vertex $v \in V \setminus D$, we have $d_G(v, D) = \min\{d_G(v, x) : x \in D\} \leq k$, where $d_G(v, x)$ is the distance in G between the vertex v and x . The minimum cardinality overall distance k -dominating sets of G , is the *distance k -domination number* and is denoted by $\gamma_k(G)$. When $k = 1$, the distance 1-domination number is the well-known domination number of the graph denoted by $\gamma(G)$. Distance k -dominating sets find multiple applications in problems involving graphs like communication networks [31], geometric problems [26], facility location problems [19]. Results about this well-studied concept can be found surveyed in a recent book chapter [18].

Another concept associated with distance in graphs is resolvability and the metric dimension of graphs, introduced by Harary and Melter [17] and Slater [30]. Let $W = \{w_1, w_2, \dots, w_r\}$ be an ordered set of vertices in a graph G , the *metric representation* of v with respect to W is the r -vector $c(v|W) = (d_G(v, w_1), d_G(v, w_2), \dots, d_G(v, w_r))$. The set W is a *resolving set* of G , if for

every two distinct vertices $v, u \in V \setminus W$, $c(v|W) \neq c(u|W)$. The minimum cardinality of a resolving set of G is the *metric dimension* of G , and is denoted $\dim(G)$. Due to their important role from both a theoretical and a practical point of view, resolving sets and the metric dimension of graphs attracted attention these past years (see surveys [2, 32]). Resolving sets find many applications in several areas like network verification [3], robot navigation [25], pharmaceutical chemistry [8], coin weighing problems, Mastermind game (see references in [6, 23]) and more.

The problems of finding $\gamma_k(G)$ and $\dim(G)$ are both NP-Hard problems in general, see respectively [7] and [25].

To join the utility of resolving sets and distance k -dominating sets, we study a set satisfying the two properties.

Definition 1.1. A distance k -resolving dominating set is a set $S \subseteq V$, where S is both a resolving set and a distance k -dominating set of G . The distance k -resolving domination number, denoted by $\gamma_k^r(G)$, is the minimum cardinality of a distance k -resolving dominating set of G , i.e., $\gamma_k^r(G) = \min\{|S| : S \text{ is a distance } k\text{-resolving dominating set of } G\}$.

A situation where the uses of resolving sets and distance k -dominating sets are both needed could represent a possible application of distance k -resolving dominating sets. For example, consider a network of vehicles. We want to identify the position of each vehicle, where the detection range within the network is not limited, and every position must be within a specific distance of a station that provides a service, such as energy supply or maintenance. The distance k -resolving dominating sets are required.

Resolving sets that satisfy additional properties are known and studied. For example, independent resolving set [11], is a resolving set that is also an independent set. Connected resolving set [29], is a resolving set that is also a connected set. For $k = 1$, the distance 1-resolving dominating set is a resolving set that is also a dominating set, the minimum cardinality of such set was first studied under the name of resolving domination number in [4], while it appeared as metric-location-domination number in [20]. More studies were done about that case relating it with other graph parameters, see for example [5, 16, 22]. Here we use the name resolving domination number and denote by $\gamma^r(G)$.

For $k \geq 1$ and $v, u \in V$, let $d_k(v, u) = \min\{d_G(v, u), k + 1\}$. A variation of the metric dimension that could be related to $\gamma_k^r(G)$ is the *k -truncated metric dimension*, $\dim_k(G)$, defined as the minimum cardinality of a *k -truncated resolving set* of G , which is a set $W \subseteq V$ verifying for any two distinct vertices $v, u \in V$, there exists a vertex x in W such that $d_k(v, x) \neq d_k(u, x)$. The k -truncated metric dimension was first studied when $k = 1$ in [24], also called adjacency dimension, where it was used to investigate the metric dimension of lexicographic product of graphs. For $k \geq 1$, the k -truncated metric dimension coincides with

the $(1, k + 1)$ -metric dimension of graphs in [13]. Results on $\dim_k(G)$ can be found in [14, 15, 33].

In Section 2, we give sharp bounds for $\gamma_k^r(G)$ in terms of the metric dimension, the distance k -domination number, the order, the diameter, the radius, and the girth of the graph. Also, we give the distance k -resolving domination number of the families of paths and cycles. In Section 3, for all $k \geq 1$, we show that $\gamma_k^r(G)$ is equal to 1 if and only if G is a path of order at most $k + 1$. For $k \geq 2$, we show an equivalence between $\gamma_k^r(G)$ and $\dim(G)$, which we use to characterize all graphs of order n having $\gamma_k^r(G)$ equal to $n - 1$ and $n - 2$. In Section 4, we determine all the realizable triples of positive integers (β, γ, α) by a graph G having $\dim(G) = \beta$, $\gamma_k(G) = \gamma$, and $\gamma_k^r(G) = \alpha$ when $k \geq 2$, in particular the graphs we construct realizing these values are all trees. In Section 5, for all $k \geq 1$, we show that a graph G having distance k -resolving domination number $\gamma_k^r(G) = \gamma_k^r \geq 1$, has a maximum order of $\gamma_k^r + \gamma_k^r \sum_{p=1}^k (2p + 1)^{\gamma_k^r - 1}$. Also, we construct graphs attaining this maximum order for any arbitrary positive integers k and γ_k^r . Section 6 is devoted to Nordhaus-Gaddum bounds for the distance k -resolving domination number of graphs for $k \geq 2$. Finally, in Section 7, we discuss relations between $\gamma_k^r(G)$ and $\dim_k(G)$, we then conclude with some open questions.

2. PRELIMINARY RESULTS AND BOUNDS FOR $\gamma_k^r(G)$

Every superset of a distance k -dominating set is a distance k -dominating set. It is true also for resolving sets. This means that every superset of a distance k -resolving dominating set is also a distance k -resolving dominating set. We give the following bounds that extend bounds given for k equal to 1 and 2, in [5] and [34] respectively to all $k \geq 1$.

Proposition 2.1. *Let G be a connected graph of order $n \geq 2$. For $k \geq 1$, we have*

$$\max\{\gamma_k(G), \dim(G)\} \leq \gamma_k^r(G) \leq \min\{\gamma_k(G) + \dim(G), n - 1\}.$$

Proof. Let S be a minimum distance k -resolving dominating set of G . Since S is both a resolving set and a distance k -dominating set, then $\dim(G) \leq |S|$, and $\gamma_k(G) \leq |S|$. Thus $\max\{\gamma_k(G), \dim(G)\} \leq \gamma_k^r(G)$.

Let D and W be respectively a minimum distance k -dominating set and a minimum resolving set of G . The set $S = D \cup W$ is a distance k -resolving dominating set of cardinality $|S| = \gamma_k(G) + \dim(G)$. Also, any subset of V of cardinality $n - 1$ is both a resolving set and a distance k -dominating set. Then we have $\gamma_k^r(G) \leq \min\{\gamma_k(G) + \dim(G), n - 1\}$. ■

For any two positive integer k and k' such that $k \geq k' \geq 1$, every distance k' -dominating set is a distance k -dominating set. Therefore any distance k' -resolving

dominating set is a distance k -resolving dominating set.

Observation 2.2. For $k \geq k' \geq 1$, if G is a connected graph, then we have $\dim(G) \leq \gamma_k^r(G) \leq \gamma_{k'}^r(G) \leq \gamma^r(G)$.

The *eccentricity* of a vertex v in G is the maximum distance between v and any other vertex in G . The maximum and minimum eccentricity in G are respectively the *diameter* and the *radius* of G denoted respectively $\text{diam}(G)$ and $\text{rad}(G)$.

Lemma 2.3. Let G be a connected graph. For $k \geq \text{diam}(G)$, $\gamma_k^r(G) = \dim(G)$.

Proof. If $k \geq \text{diam}(G)$, then any non-empty set of vertices in V is a distance k -dominating set. Hence any resolving set is also a distance k -dominating set of G . Therefore, $\gamma_k^r(G) \leq \dim(G)$. From Proposition 2.1 it follows that $\gamma_k^r(G) = \dim(G)$. ■

Let P_n denote the path graph with $V(P_n) = \{1, 2, \dots, n\}$ and $E(P_n) = \{i(i+1) : 1 \leq i \leq n-1\}$. It is proved that $\dim(P_n) = 1$ [8], and for $k \geq 1$, $\gamma_k(P_n) = \lceil \frac{n}{2k+1} \rceil$ [12]. The values of the distance k -resolving domination number of P_n for k equal to 1 and 2 are given respectively in [4] and [34]. In the following we give $\gamma_k^r(P_n)$ for all $k \geq 1$.

Proposition 2.4. For $k \geq 1$ and $n \geq 2$,

$$\gamma_k^r(P_n) = \begin{cases} 1, & \text{if } k \geq n-1, \\ 2, & \text{if } \lfloor \frac{n}{2} \rfloor \leq k \leq n-2, \\ \lceil \frac{n}{2k+1} \rceil, & \text{if } 1 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1. \end{cases}$$

Proof. In [8], we have $\dim(P_n) = 1$. So by Proposition 2.1, $\gamma_k(P_n) \leq \gamma_k^r(P_n) \leq \gamma_k(P_n) + 1$. Also, for $1 \leq i, j \leq n$, with $i \neq j$, we have $d_{P_n}(i, j) = |i - j|$. Then any resolving set of cardinality 1 must be $\{1\}$ or $\{n\}$.

- For $k \geq n-1$, since $\text{diam}(P_n) = n-1$, it follows from Lemma 2.3 that $\gamma_k^r(P_n) = \dim(P_n) = 1$.
- For $\lfloor \frac{n}{2} \rfloor \leq k \leq n-2$, based on [12] $\gamma_k(P_n) = 1$, then $\gamma_k^r(P_n)$ is equal to 1 or 2. It is clear that an end-vertex is not distance k -dominating. Thus, $\gamma_k^r(P_n) = \gamma_k(P_n) + 1 = 2$.
- For $1 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$, in [12] we have $\gamma_k(P_n) = \lceil \frac{n}{2k+1} \rceil \geq 2$. Also, any set S consisting of two or more distinct vertices in $V(P_n)$ is a resolving set of P_n . Thus, $\gamma_k^r(P_n) = \gamma_k(P_n) = \lceil \frac{n}{2k+1} \rceil$. ■

The path P_n is a graph achieving the bounds in Proposition 2.1. For $k \geq n-1$, we have $\gamma_k^r(P_n) = \dim(P_n)$. For $1 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$, $\gamma_k^r(P_n) = \gamma_k(P_n)$, and for $\lfloor \frac{n}{2} \rfloor \leq k \leq n-2$, $\gamma_k^r(P_n) = \gamma_k(P_n) + \dim(P_n)$.

Let C_n denote the cycle graph with $n \geq 3$, where $V(C_n) = \{0, 1, \dots, n-1\}$ and $E(C_n) = \{i(i+1) \pmod n : 0 \leq i \leq n-1\}$. We have $\dim(C_n) = 2$ [10], and for $k \geq 1$, $\gamma_k(C_n) = \lceil \frac{n}{2k+1} \rceil$ [12].

Proposition 2.5. For $k \geq 1$ and $n \geq 3$,

$$\gamma_k^r(C_n) = \begin{cases} 2, & \text{if } 4k+1 \geq n, \\ 3, & \text{if } 4k+2 = n, \\ \lceil \frac{n}{2k+1} \rceil, & \text{if } 4k+3 \leq n. \end{cases}$$

Proof. We have $d_{C_n}(i, j) = \min\{|i-j|, n-|i-j|\}$.

Claim 2.6. For $n \geq 2k+2$ and $n \neq 4k+2$, the set of vertices $W = \{0, 2k+1\}$ is a resolving set of C_n .

Proof. Let $i, j \in V(C_n) \setminus W$, with $i \neq j$. If $d_{C_n}(i, 0) \neq d_{C_n}(j, 0)$, then S is a resolving set. We suppose that $d_{C_n}(i, 0) = d_{C_n}(j, 0)$, then either $d_{C_n}(i, 0) = i$ and $d_{C_n}(j, 0) = n-j$ or $d_{C_n}(i, 0) = n-i$ and $d_{C_n}(j, 0) = j$. Without loss of generality we suppose that $d_{C_n}(i, 0) = i$ and $d_{C_n}(j, 0) = n-j$, which means that $i+j = n$. If $d_{C_n}(i, 2k+1) = d_{C_n}(j, 2k+1)$, then $\min\{|2k+1-i|, n-|2k+1-i|\} = \min\{|2k+1-j|, n-|2k+1-j|\}$. Since $\min\{x, y\} = \frac{x+y-|x-y|}{2}$, it follows that $|n-2(|2k+1-i|)| = |n-2(|2k+1-j|)|$.

We suppose that $n-2|2k+1-i| = n-2|2k+1-j|$, which means that $|2k+1-i| = |2k+1-j|$. Since $i \neq j$, necessarily $2k+1-i = j-2k-1$. It follows that $i+j = 4k+2 = n$, a contradiction since $n \neq 4k+2$.

Otherwise if $n-2|2k+1-i| = 2|2k+1-j|-n$, then $n = |2k+1-i| + |2k+1-j|$. If $|2k+1-i| = 2k+1-i$ and $|2k+1-j| = 2k+1-j$, then $n = 2k+1-i+2k+1-j$. Assuming that $i+j = n$, it means that $n = 2k+1$, a contradiction.

Now if $|2k+1-i| = i-(2k+1)$ and $|2k+1-j| = j-(2k+1)$, then $n = i+j-2(2k+1)$. Since $i+j = n$, it means that $k=0$, a contradiction.

Finally if $|2k+1-i| = i-(2k+1)$ or $|2k+1-j| = j-(2k+1)$, we suppose that $|2k+1-i| = i-(2k+1)$ and $|2k+1-j| = 2k+1-j$. Then we get that $n = i-j$, again a contradiction.

It follows that $d_{C_n}(i, 2k+1) \neq d_{C_n}(j, 2k+1)$. So for $i, j \in V(C_n) \setminus W$, if $i \neq j$, then $c(i|W) \neq c(j|W)$. \square

• If $2k+1 \geq n$, then $k \geq \text{diam}(C_n)$. By Lemma 2.3, $\gamma_k^r(C_n) = \dim(C_n)$. Since $\dim(C_n) = 2$, we have $\gamma_k^r(C_n) = 2$.

If $4k+1 \geq n \geq 2k+2$, we have $\gamma_k^r(C_n) \geq \dim(C_n) = 2$. From Claim 2.6, the set $\{0, 2k+1\}$ is a resolving set of C_n , it is also a distance k -dominating set of C_n for $4k+1 \geq n \geq 2k+2$. Therefore $\gamma_k^r(C_n) = 2$.

• If $4k+2 = n$, based on [12] we have $\gamma_k(C_{4k+2}) = 2$, then by Proposition 2.1, $\gamma_k^r(C_{4k+2}) \geq 2$. By using contradiction we suppose that $\gamma_k^r(C_{4k+2}) = 2$, and let

S be a distance k -resolving dominating set of cardinality 2. Since all the vertices have degree 2, if a vertex i is in a distance k -dominating set of cardinality 2, then the set contains necessarily $i + 2k + 1 \pmod{n}$. Since the cycle C_n is vertex-transitive, we suppose without loss of generality that $S = \{0, 2k + 1\}$. If we take the vertices 1 and $4k + 1$, then clearly $c(1|S) = c(4k + 1|S)$. It follows that S is not a resolving set of C_{4k+2} . Hence $\gamma_k^r(C_{4k+2}) > 2$.

Now, let us consider the set $S = \{0, 1, 2k + 1\}$, we will show first that $\{0, 1\} \subset S$ is a resolving set of C_{4k+2} . For $i \in V(C_n) \setminus S$, we have $c(i|\{0, 1\}) = (\min\{i, n - i\}, \min\{i - 1, n - i + 1\})$. For $i, j \in V(C_n) \setminus S$, if $c(i|\{0, 1\}) = c(j|\{0, 1\})$, it means that $\min\{i, n - i\} = \min\{j, n - j\}$ and $\min\{i - 1, n - i + 1\} = \min\{j - 1, n - j + 1\}$. Since $\min\{x, y\} = \frac{x+y-|x-y|}{2}$, it follows that $|n - 2i| = |n - 2j|$ and $|n - 2(i - 1)| = |n - 2(j - 1)|$. Assuming that $i \neq j$, then necessarily $n - 2i = 2j - n$ and $n - 2(i - 1) = 2(j - 1) - n$, which is impossible. Then if $i \neq j$, we have $c(i|\{0, 1\}) \neq c(j|\{0, 1\})$. Therefore $\{0, 1\}$ is a resolving set of C_{4k+2} .

Since $\{0, 2k + 1\}$ is a distance k -dominating set of C_{4k+2} , it follows that $S = \{0, 1, 2k + 1\}$ is a distance k -resolving dominating set of C_{4k+2} . Therefore $\gamma_k^r(C_{4k+2}) = 3$.

• If $4k + 3 \leq n$, in [12] we have $\gamma_k(C_n) = \lceil \frac{n}{2k+1} \rceil$. Let us consider the set $S = \{i(2k + 1) : 0 \leq i \leq \lceil \frac{n}{2k+1} \rceil - 1\}$, we have $|S| = \lceil \frac{n}{2k+1} \rceil$. Claim 2.6 shows that the set $\{0, 2k + 1\} \subset S$ is a resolving set of C_n . Also, it is easy to see that the set S is a distance k -dominating set of C_n . It follows that $\gamma_k^r(C_n) = \lceil \frac{n}{2k+1} \rceil$. ■

Proposition 2.7. For $k \geq 1$, let G be a connected graph such that $\text{rad}(G) \leq k$ or $\text{diam}(G) = k + 1$. Then we have $\dim(G) \leq \gamma_k^r(G) \leq \dim(G) + 1$.

Proof. Let G be a connected graph with $\text{rad}(G) \leq k$. This means that $\gamma_k(G) = 1$. Then by Proposition 2.1, we have $\dim(G) \leq \gamma_k^r(G) \leq \dim(G) + 1$.

If $\text{diam}(G) = k + 1$, let $W \subset V$ be a minimum resolving set of G . Let $v \in V \setminus \text{dom}_k(W)$, where $\text{dom}_k(W) = \{v \in V : d_G(v, W) \leq k\}$. Then v must be at distance greater or equal to $k + 1$ from all the vertices of W . Since $\text{diam}(G) = k + 1$, the only possible metric representation with respect to W of a vertex v such that $d_G(v, W) \geq k + 1$, is a vector having $k + 1$ as a value in all its coordinates. Since W is a resolving set, then there is at most one such vertex in G . Hence, $\dim(G) \leq \gamma_k^r(G) \leq \dim(G) + 1$. ■

For all $k \geq 1$ both bounds in Proposition 2.7 can be achieved. For $\lfloor \frac{n}{2} \rfloor \leq k \leq n - 2$, the graph P_n has $\text{rad}(P_n) \leq k$, from Proposition 2.4, $\gamma_k^r(P_n) = \dim(P_n) + 1$. From Lemma 2.3, if $\text{rad}(G) \leq \text{diam}(G) \leq k$, then for any G we have $\gamma_k^r(G) = \dim(G)$. The cycle graphs C_{2k+2} or C_{2k+3} according to Proposition 2.5 are examples of graphs with $\text{diam}(G) = k + 1$ having $\gamma_k^r(G) = \dim(G)$. Also from Proposition 2.4, the path P_{k+2} is a graph of $\text{diam}(G) = k + 1$ having $\gamma_k^r(G) = \dim(G) + 1$.

Lemma 2.8 [21]. *For $k \geq 1$, let G be a connected graph of order $n \geq k + 1$ and diameter $\text{diam}(G) \geq k$. Then there exists a minimum distance k -dominating set D of G satisfying for every vertex $v \in D$ there is a vertex $x \in V \setminus D$ such that $d_G(v, x) = k$ and $N_k(x) \cap D = \{v\}$.*

The following upper bound proved for $\dim(G)$ in [4] is true also for $\gamma_k^r(G)$, the proofs are similar.

Proposition 2.9. *For $k \geq 1$, let G be a connected graph of order $n \geq k + 1$ with $\text{diam}(G) \geq k$. Then $\gamma_k^r(G) \leq n - k\gamma_k(G)$, and this upper bound is achieved for any positive integers k and $\gamma_k(G)$.*


Proof. Suppose that $\gamma_k(G) = \gamma$. Based on Lemma 2.8, let us consider $D = \{1, 2, \dots, \gamma\}$ a minimum distance k -dominating set such that for all $1 \leq i \leq \gamma$, there exists a vertex $w_{i,k}$ verifying that $d_G(i, w_{i,k}) = k$, and for $j \neq i$, $d_G(j, w_{i,k}) > k$. Now let $P_i = iw_{i,1}w_{i,2} \cdots w_{i,k}$ be a shortest $(i, w_{i,k})$ -path. We can see that for $1 \leq p \leq k$, we have $d_G(i, w_{i,p}) = p$ and $d_G(j, w_{i,p}) > p$. For any two different vertices $w_{i,p}$, $w_{j,q}$, with $1 \leq i, j \leq \gamma$ and $1 \leq p, q \leq k$, we will check the vector of distances with respect to the set D , we discuss the following two cases.

- (i) If $i \neq j$, we suppose without loss of generality that $q \geq p$. We have $d_G(i, w_{i,p}) = p$ and $d_G(i, w_{j,q}) \geq q + 1 > p$.
- (ii) If $i = j$ and $p \neq q$, we have $d_G(i, w_{i,p}) = p$ and $d_G(i, w_{i,q}) = q \neq p$.

It follows that the set D resolves all the vertices $w_{i,p}$, where $1 \leq i \leq \gamma$, and $1 \leq p \leq k$. Then the set $S = V \setminus \bigcup_{i=1}^{\gamma} \{w_{i,j}\}_{j=1}^k$ is both a distance k -dominating set and a resolving set. Hence $\gamma_k^r(G) \leq |S| = |V \setminus \bigcup_{i=1}^{\gamma} \{w_{i,j}\}_{j=1}^k| = n - k\gamma = n - k\gamma_k(G)$.

The family of trees $\{T_\gamma : \gamma \geq 1\}$ illustrated as an example in Figure 1 has $\gamma_k^r(T_\gamma) = n - k\gamma$, for $k, \gamma \geq 1$, where $\gamma_k(T_\gamma) = \gamma$. We have any distance k -dominating set in T_γ must contain at least one vertex in each branch $iw_{i,1} \cdots w_{i,k}$, with $1 \leq i \leq \gamma$. Also, the set of vertices $\{1, 2, \dots, \gamma\}$ is a distance k -dominating set of T_γ . Then clearly $\gamma_k(T_\gamma) = \gamma$. We can check as above that the set of vertices $\{1, 2, \dots, \gamma\}$ is a resolving set of T_γ . It follows from Proposition 2.1 that it is a minimum distance k -resolving dominating set of T_γ of cardinality $n - k\gamma = n - k\gamma_k(T_\gamma)$. ■

For a connected graph G of order n and diameter d , we have $\dim(G) \leq n - d$ [8]. The graphs achieving equality are characterized in [23]. This type of bound involving the order and the diameter of the graph was provided for the resolving domination number in [5]. We give a general upper bound for all $k \geq 1$.

Figure 1. Tree graph T_γ having $\gamma_k^r(T_\gamma) = n - k\gamma_k(T_\gamma)$.

Proposition 2.10. For $k \geq 1$, let G be a connected graph of order n and diameter d . Then

$$\gamma_k^r(G) \leq \begin{cases} n - d, & \text{if } d \leq k, \\ n - d + 1, & \text{if } k + 1 \leq d \leq 2k, \\ n - d + \lfloor \frac{d}{2k+1} \rfloor, & \text{if } d \geq 2k + 1. \end{cases}$$

These bounds are sharp.

Proof. Let $P = (0, 1, \dots, d)$ be a diametral path in G , i.e., P is a shortest path of length d . For any two vertices i and j in P , we have $d_G(i, j) = |i - j|$.

If $d \leq k$, then by Lemma 2.3, $\gamma_k^r(G) = \dim(G)$. Based on [8], we have $\gamma_k^r(G) \leq n - d$.

If $k + 1 \leq d \leq 2k$, we consider the set of vertices $\{k, d\}$. For $0 \leq l, m \leq d - 1$, with $l \neq m$, we have $d_G(l, d) = |l - d| \neq |m - d| = d_G(m, d)$. Also, for any $0 \leq l \leq d$, we have $d_G(l, k) = |l - k| \leq k$. This means that the set $\{k, d\}$ is resolving and distance k -dominating of the vertices $i \notin \{k, d\}$. Now, let $S' = V \setminus \{i : i \notin \{k, d\}\}$. Then S' is a distance k -resolving dominating set of G . Hence, $\gamma_k^r(G) \leq |S'| = n - d + 1$.

If $d \geq 2k + 1$, let us consider the set of vertices $S = \{k, k + (2k + 1), \dots, k + j(2k + 1), \dots, \min\{k + \lfloor \frac{d}{2k+1} \rfloor(2k + 1), d\}\}$. Let l be a vertex in $P \setminus S$. If $\min\{k + \lfloor \frac{d}{2k+1} \rfloor(2k + 1), d\} = k + \lfloor \frac{d}{2k+1} \rfloor(2k + 1)$, then either $k + \lfloor \frac{d}{2k+1} \rfloor(2k + 1) < l \leq d$ or there exists $1 \leq i \leq \lfloor \frac{d}{2k+1} \rfloor$ such that $k + (i - 1)(2k + 1) < l < k + i(2k + 1)$, or $0 \leq l < k$. In all those cases there exists a vertex in S at distance less or equal to k from l . The same can be observed when $\min\{k + \lfloor \frac{d}{2k+1} \rfloor(2k + 1), d\} = d$. Furthermore, since $|S| \geq 2$ and for $0 \leq i, j \leq d$, $d_G(i, j) = |i - j|$, it is straightforward that S resolves the vertices in $P \setminus S$.

If we consider the set $S' = V \setminus \{P \setminus S\}$, then S' is a distance k -resolving dominating set of the graph G . Hence, $\gamma_k^r(G) \leq |S'| = n - d + \lfloor \frac{d}{2k+1} \rfloor$.

The graph path P_n has diameter $n - 1$. From Proposition 2.4 it is a graph achieving the upper bound $n - d$ for $n \leq k + 1$. It achieves the upper bound $n - d + 1$ when $k + 2 \leq n \leq 2k + 1$. The path graph P_n also achieves the upper bound $n - d + \lfloor \frac{d}{2k+1} \rfloor$ when $n \geq 2k + 2$. ■

If $k = 1$, for a connected graph of diameter $d \geq 3$, the upper bound in Proposition 2.10 is precisely the bound given in terms of the order and the diameter in [5].

The girth of the graph is the length of a shortest cycle in the graph. The following lower bounds proved in [12] for $\gamma_k(G)$ holds also for $\gamma_k^r(G)$ and they are achieved.

Proposition 2.11. *For $k \geq 1$, let G be a connected graph having diameter d , radius r , and girth g . Then we have*

- (1) $\gamma_k^r(G) \geq \frac{d+1}{2k+1}$;
- (2) $\gamma_k^r(G) \geq \frac{2r}{2k+1}$;
- (3) $\gamma_k^r(G) \geq \frac{g}{2k+1}$, if $g < \infty$.

These bounds are sharp.

Proof. In [12], it is shown that if G is a connected graph of diameter d , then $\gamma_k(G) \geq \frac{d+1}{2k+1}$. In the same paper we have if G has radius r , then $\gamma_k(G) \geq \frac{2r}{2k+1}$. Also in [12], for a connected graph of girth $g < \infty$, we have $\gamma_k(G) \geq \frac{g}{2k+1}$. Since $\gamma_k^r(G) \geq \gamma_k(G)$, the above lower bounds for $\gamma_k(G)$ are true also for $\gamma_k^r(G)$.

Some graphs in Proposition 2.4 and 2.5 are examples of graphs attaining these bounds. In (1) consider the path graph of order $n = p(2k + 1)$ for $p \geq 2$, since $d = n - 1$, we get that $\gamma_k^r(G) = \frac{d+1}{2k+1}$. In (2) consider the path graph of order $n = 2p(2k + 1)$. We have $r = p(2k + 1)$, then from proposition 2.4, $\gamma_k^r(G) = \frac{2r}{2k+1}$. In (3) take a cycle graph of order $n = p(2k + 1)$ for $p \geq 3$, since $g = n$, then this is a graph having $\gamma_k^r(G) = \frac{g}{2k+1}$. ■

3. GRAPHS WITH $\gamma_k^r(G)$ EQUAL TO 1, $n - 2$, AND $n - 1$

Further, let K_n denote the complete graph on n vertices, and let $K_{s,t}$ with $s, t \geq 1$ denote the complete bipartite graph. For two graphs G_1 and G_2 the disjoint union of G_1 and G_2 , denoted by $G_1 \cup G_2$, is the graph with vertex set $V(G_1 \cup G_2) = V(G_1) \cup V(G_2)$ and edge set $E(G_1 \cup G_2) = E(G_1) \cup E(G_2)$. The join graph of G_1 and G_2 , denoted by $G_1 + G_2$, is the graph obtained from $G_1 \cup G_2$ by joining each vertex from $V(G_1)$ to each vertex in $V(G_2)$. We denote by \overline{G} the complement graph of G .

Theorem 3.1 [8]. *For a connected graph G of order $n \geq 2$, we have the following.*

- $\dim(G) = 1$ if and only if $G \cong P_n$.
- If $n \geq 4$, then $\dim(G) = n - 2$ if and only if $G \in \{K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$.
- $\dim(G) = n - 1$ if and only if $G \cong K_n$.

In a connected graph G of order $n \geq k + 1$, any subset of V of order greater or equal to $n - k$ is a distance k -dominating set.

Lemma 3.2. *Let $k \geq 2$. For $1 \leq i \leq k$, if G is a connected graph of order $n \geq i + 2$ that is not a path graph, then $\gamma_k^r(G) = n - i$ if and only if $\dim(G) = n - i$.*

Proof. For all $1 \leq i \leq k$, if $\dim(G) = n - i$, any subset of V of cardinality $n - i \geq n - k$ is a distance k -dominating set. Then a resolving set of cardinality $\dim(G) = n - i$ is also a distance k -dominating set. Therefore $\gamma_k^r(G) = \dim(G) = n - i$.

Conversely, if $\gamma_k^r(G) = n - i$, by Proposition 2.1, we have $\dim(G) \leq n - i$. If $n = i + 2$, then $\gamma_k^r(G) = n - i = 2$. It follows that $\dim(G)$ is equal to 1 or 2. Based on Theorem 3.1, the only graphs with $\dim(G) = 1$ are path graphs, it follows that $\dim(G) = 2$.

If $n \geq i + 3$, we suppose that $\dim(G) < n - i$. If $i \leq k - 1$, then a resolving set of cardinality $n - (i + 1) \geq n - k$ is also a distance k -dominating set. Thus $\gamma_k^r(G) \leq n - (i + 1)$, which is impossible. Now if $i = k$, let $W \subseteq V$ be a resolving set of cardinality $n - (k + 1)$, and let us denote $1, 2, \dots, k + 1$ the vertices in $V \setminus W$. Assuming that $\gamma_k^r(G) = n - k$, then there is at least one vertex v in $V \setminus W$ such that $d_G(v, W) = k + 1$. Let $w \in W$ be such that $d_G(v, w) = d_G(v, W) = k + 1$, and let Q be a shortest (v, w) -path. Since $d_G(v, w) = d_G(v, W)$, and G is a connected graph, the only vertex in $W \cap Q$ is w . We have $|Q| = k + 2$ and $|W| = n - (k + 1)$, which means that the subgraph induced by the vertices $1, 2, \dots, k + 1$ and w is the path Q . Without loss of generality, we suppose that the path Q is $(k + 1)k \cdots 1w$. Now, let $S = (W \setminus \{w\}) \cup \{k\}$. We have $d_G(k, k + 1) = d_G(k, k - 1) = 1$, $d_G(k, w) = k \geq 2$, and if $k \geq 3$, for $1 \leq j \leq k - 2$, we have $d_G(k, j) = k - j \geq 2$. Also $d_G(k + 1, S \setminus \{k\}) \geq k + 1$, since G is a connected graph and $n \geq k + 3$, then there exists a vertex $u \in S \setminus \{k\}$ such that either 1 and w are adjacent to u . This means that $d_G(k - 1, u) \leq k$. It follows that S is a resolving set of G . Since $d_G(k, i) \leq k$, for $1 \leq i \leq k + 1$, $i \neq k$, and $d_G(k, w) = k$, it means that the set S is also a distance k -dominating set of G . Hence $\gamma_k^r(G) \leq |S| = n - (k + 1)$, a contradiction. Therefore $\dim(G) = n - k$. ■

By combining Theorem 3.1 and Lemma 3.2 with Proposition 2.4, we give the following characterizations.

Theorem 3.3. *For any graph G of order $n \geq 2$, the following statements hold.*

- (a) *For all $k \geq 1$, $\gamma_k^r(G) = 1$ if and only if $G \in \{P_i\}_{i=2}^{k+1}$.*

- (b) If G is a connected graph of order $n \geq 4$, $\gamma_2^r(G) = n - 2$ if and only if $G \in \{P_4, K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$. For all $k \geq 3$, $\gamma_k^r(G) = n - 2$ if and only if $G \in \{K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$.
- (c) If G is a connected graph, for any $k \geq 2$, $\gamma_k^r(G) = n - 1$ if and only if $G \cong K_n$.

Proof. (a) For $k \geq 1$, if $\gamma_k^r(G) = 1$, then G is a connected graph and from Proposition 2.1, $\dim(G) = 1$. The equivalence is completed by Theorem 3.1 and Proposition 2.4.

(b) If G is a connected graph of order $n \geq 4$ different from a path graph, then by Lemma 3.2 we have $\gamma_k^r(G) = n - 2$ if and only if $\dim(G) = n - 2$. Which means by Theorem 3.1 that it is equivalent to $G \in \{K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$. From Proposition 2.4, we have $\gamma_k^r(P_n) = n - 2$, it occurs only when $k = 2$ and $n = 4$. Then $\gamma_2^r(G) = n - 2$ if and only if $G \in \{P_4, K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$. Also, for $k \geq 3$, $\gamma_k^r(G) = n - 2$ if and only if $G \in \{K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$.

(c) The only connected graphs of order 2 and 3 are respectively K_2 and P_3 or K_3 . For $k \geq 2$, from Proposition 2.4 and Theorem 3.1, we have $\gamma_k^r(K_2) = 1$, $\gamma_k^r(P_3) = 1$, and $\gamma_k^r(K_3) = 2$. If G has order $n \geq 4$, then by Lemma 3.2 and Theorem 3.1, we have $\gamma_k^r(G) = n - 1$ if and only if $G \cong K_n$. ■

For $k = 1$, we have $\gamma^r(G) = n - 1$ if and only if $G \in \{K_{1,n-1}, K_n\}$ [4, 20]. The graphs having $\gamma^r(G)$ equal to 2 and $n - 2$ are fully determined in [5] and [20], respectively.

4. REALIZABLE VALUES FOR $\dim(G)$, $\gamma_k(G)$, AND $\gamma_k^r(G)$.

In Proposition 2.1, we have $\max\{\gamma_k(G), \dim(G)\} \leq \gamma_k^r(G) \leq \gamma_k(G) + \dim(G)$. For $k = 1$, in [5] it is shown that for any three positive integers β , γ , and α , verifying that $\max\{\gamma, \beta\} \leq \alpha \leq \gamma + \beta$, and $(\beta, \gamma, \alpha) \notin \{(1, \gamma, \gamma + 1) : \gamma \geq 2\}$, there is always a graph G having $\dim(G) = \beta$, $\gamma(G) = \gamma$, and $\gamma^r(G) = \alpha$. We give a similar result for $\dim(G)$, $\gamma_k(G)$, and $\gamma_k^r(G)$, for all $k \geq 2$.

The graph families we provide in Theorem 4.2 are all trees. To determine $\gamma_k^r(G)$ of some of these graphs, we will need the next formula for the metric dimension of trees that appeared in [8, 17, 30]. We will recall some terminology given in [8]. In a tree T for $v \in V$, if the degree $\deg(v) \geq 3$, then v is called a *major vertex*. A leaf l , i.e., a vertex of degree one, in T is a *terminal vertex* of a major vertex v , if v is the closest major vertex in terms of distance to l , i.e., for u a major vertex in T different from v , we have $d_T(v, l) < d_T(u, l)$. If v is a major vertex having at least one terminal vertex, then v is called an *exterior*

major vertex. Let $L(T)$ and $EX(T)$ denote respectively the number of leaves and the number of exterior major vertices in a tree T .

Theorem 4.1 [8, 17, 30]. *If T is a tree that is not a path graph, then $\dim(T) = L(T) - EX(T)$. Also, any resolving set of T must contain at least one vertex from each branch at an exterior major vertex containing its terminal vertices with at most one exception.*

Theorem 4.2. *For any three positive integers β , γ , and α such that $\max\{\gamma, \beta\} \leq \alpha \leq \gamma + \beta$ and $(\beta, \gamma, \alpha) \notin \{(1, \gamma, \gamma + 1) : \gamma \geq 2\}$, and for all $k \geq 2$, there always exists a tree graph T having $\dim(T) = \beta$, $\gamma_k(T) = \gamma$, and $\gamma_k^r(T) = \alpha$. There is no graph realizing the triples $\{(1, \gamma, \gamma + 1) : \gamma \geq 2\}$.*

Proof. Let $\beta, \gamma, \alpha \geq 1$ be such that $\max\{\gamma, \beta\} \leq \alpha \leq \gamma + \beta$. We discuss the possible values for the triple $(\dim(G), \gamma_k(G), \gamma_k^r(G)) = (\beta, \gamma, \alpha)$, according to the following cases.

- If $\beta = 1$, then $\gamma \leq \alpha \leq \gamma + 1$. Also by Theorem 3.1 we have the path graphs are the only graphs having the metric dimension equal to 1. For $k \geq 2$, in a path graph any subset of vertices of order greater or equal to 2 is a resolving set. Then if $\gamma \geq 2$, we have $\alpha = \gamma$. This means that the triple $(1, \gamma, \gamma + 1)$ is not realizable by any graph for $\gamma \geq 2$. Also, according to Proposition 2.4 the path graphs realizes the following cases. (i) If $k + 1 \geq n$, then we have $\gamma = \beta = \alpha = 1$. (ii) If $\lfloor \frac{n}{2} \rfloor \leq k \leq n - 2$, then $\gamma = \beta = 1$ and $\alpha = 2 = \gamma + \beta$. (iii) If $1 \leq k \leq \lfloor \frac{n}{2} \rfloor - 1$, then $\beta = 1 < \gamma = \alpha = \lceil \frac{n}{2k+1} \rceil \geq 2$.
- If $\gamma = 1$, for any $\beta \geq 2$, then we have $\beta \leq \alpha \leq \beta + 1$. The star graph $K_{1, \beta+1}$ has $\gamma_k(K_{1, \beta+1}) = 1$, and from Theorem 3.1 and Theorem 3.3 we have $\gamma_k^r(K_{1, \beta+1}) = \dim(K_{1, \beta+1}) = \beta$, for any $k \geq 2$. This means that for $k \geq 2$, the triple $(\beta, 1, \beta)$ is realized for all $\beta \geq 2$. For the case of the triple $(\beta, 1, \beta + 1)$, we consider the spider tree graph, denoted by $S_{\beta+1, k}$, having one vertex v_0 of degree $\beta + 1$ with $\beta + 1$ leaves l_i , $1 \leq i \leq \beta + 1$, at distance k from v_0 . Note that all the vertices of $S_{\beta+1, k}$ are of degree less or equal to 2 except v_0 . Clearly $\gamma_k(S_{\beta+1, k}) = 1$, and based on Theorem 4.1, we have $\dim(S_{\beta+1, k}) = \beta$. Also any resolving set must contain at least one vertex in all but one of the (v_0, l_i) -paths, where $1 \leq i \leq \beta + 1$. By using contradiction, we suppose that $\gamma_k^r(S_{\beta+1, k}) = \beta$. From Theorem 4.1, we consider that a minimum distance k -resolving dominating set W of $S_{\beta+1, k}$ having cardinality β contains one vertex in any of the (v_0, l_i) -paths, with $1 \leq i \leq \beta$. We have the vertex $l_{\beta+1}$ is at distance greater than k from the vertices in W . This means that W is not a distance k -dominating set, a contradiction. Hence, $\gamma_k^r(S_{\beta+1, k}) = \beta + 1$.
- If $\beta \geq 2$ and $\gamma \geq 2$, with $\max(\gamma, \beta) \leq \alpha \leq \gamma + \beta$, then the realizable values for the triple (β, γ, α) are considered depending on the following five subcases.

(i) If $2 \leq \beta = \gamma < \alpha$, then the trees $T^1 = \{T_{k,m,l}^1 : m \geq 0, l \geq 1, k \geq 2\}$ in Figure 2 illustrate graphs realizing this case.

Claim 4.3. We have $\gamma_k(T_{k,m,l}^1) = \dim(T_{k,m,l}^1) = m + l$, and $\gamma_k^r(T_{k,m,l}^1) = m + 2l$.

Proof. Suppose that $\gamma_k(T_{k,m,l}^1) = \gamma$, $\dim(T_{k,m,l}^1) = \beta$, and $\gamma_k^r(T_{k,m,l}^1) = \alpha$. It is clear that $\{v_i\}_{i=1}^m \cup \{w_i\}_{i=1}^l$ is a minimum distance k -dominating set. Then $\gamma = m + l$. Based on Theorem 4.1, we have $\beta = m + l$, and for each $1 \leq i \leq l$, a resolving set must contain one vertex from the set of vertices $\{v_{i,j}\}_{j=0}^k$. Also, for each $1 \leq i \leq m$, a resolving set must contain one vertex from the set of vertices $\{w_{i,j}, w'_{i,j}\}_{j=0}^k$. Now, let S be a minimum distance k -resolving dominating set of cardinality α . We suppose without loss of generality, that S contain a vertex from each $\{v_{i,j}\}_{j=0}^k$ with $1 \leq i \leq m$, and one vertex from each $\{w_{i,j}\}_{j=0}^k$ with $1 \leq i \leq l$. Since $d_G(w_i, w'_{i,k}) = k$, and for $x \notin \{w_i, w'_{i,j}\}$ we have $d_G(x, w'_{i,k}) > k$. Then to be a distance k -dominating set, S must contain for each $1 \leq i \leq l$, at least w_i or a vertex in $\{w'_{i,j}\}_{j=0}^k$. Hence $\alpha \geq m + 2l$. It is easy to check that the set of vertices $\{v_{i,1}\}_{i=1}^m \cup \{w_{i,k}\}_{i=1}^l \cup \{w_i\}_{i=1}^l$ is a distance k -resolving dominating set. Thus $\alpha \leq m + 2l$. It follows that $\alpha = m + 2l$. \square






Figure 2. Tree $T_{k,m,l}^1$.

The proofs for the remaining cases use similar arguments as in the proof of Claim 4.3. In the following, we only provide examples of minimum distance k -dominating sets, minimum resolving sets, and minimum distance k -resolving dominating sets for each family of trees.

(ii) If $2 \leq \gamma \leq \beta = \alpha$, then the family of trees $T^2 = \{T_{k,m,l}^2 : m \geq 1, l \geq 1, k \geq 2\}$ represented in Figure 3 realizes this case. The set of vertices $\{v_i\}_{i=1}^m \cup \{w\}$ is a minimum distance k -dominating set of cardinality $m + 1$. Also, the set of vertices $\{v_{i,1}\}_{i=1}^m \cup \{w_{i,1}\}_{i=1}^l$ is both a minimum resolving set and a minimum distance k -resolving dominating set of cardinality $m + l$.

Figure 3. Tree $T_{k,m,l}^2$.

(iii) If $2 \leq \beta < \gamma = \alpha$, then the family of trees $T^3 = \{T_{k,m,l}^3 : m \geq 1, l \geq 1, k \geq 2\}$ represented in Figure 4 realizes this case. From Theorem 4.1, we have $\dim(T_{k,m,l}^3) = m + 1$. Also, the set $\{v_i\}_{i=1}^m \cup \{w_i\}_{i=1}^l \cup \{u\}$ is a minimum distance k -dominating set of $T_{k,m,l}^3$ of cardinality $m + l + 1$. Finally, the set $\{v_{i,1}\}_{i=1}^m \cup \{w_{i,1}\}_{i=1}^l \cup \{u_1\}$ is a distance k -resolving dominating set of cardinality $m + l + 1$. It follows that $\gamma_k^T(T_{k,m,l}^3) = \gamma_k(T_{k,m,l}^3) = m + l + 1$.

Figure 4. Trees $T_{k,m,l}^3$.

(iv) If $2 \leq \gamma < \beta < \alpha$, then the family of trees $T^4 = \{T_{k,m,l,r}^4 : m \geq 0, l \geq 0, r \geq 3, k \geq 2\}$ represented in Figure 5 illustrates graphs realizing this case, where $(m, l) \neq (0, 0)$. The set of vertices $\{v_{i,k}\}_{i=1}^m \cup \{w_{i,k}\}_{i=1}^l \cup \{u_{i,k}\}_{i=1}^{r-1}$ is a minimum resolving set of cardinality $m + l + r - 1$. The set of vertices $\{v_i\}_{i=1}^m \cup \{w_i\}_{i=1}^l \cup \{u\}$ is a minimum distance k -dominating set of cardinality $m + l + 1$. The set of vertices $\{v_{i,1}\}_{i=1}^m \cup \{w_{i,k}, w_i\}_{i=1}^l \cup \{u_{i,k}\}_{i=1}^{r-1} \cup \{u\}$ is a minimum distance k -resolving dominating set of cardinality $m + 2l + r$.

(v) If $2 \leq \beta < \gamma < \alpha$, then Figure 6 illustrates a family of trees $T^5 = \{T_{k,m,l,r}^5 : m \geq 0, l \geq 0, r \geq 2, k \geq 2\}$ realizing this case, where $(m, l) \neq (0, 0)$. The set of vertices $\{v_{i,k}\}_{i=1}^m \cup \{w_{i,k}\}_{i=1}^l \cup \{u_{r,k}\}$ is a minimum resolving set of cardinality




Figure 5. Tree $T_{k,m,l,r}^4$.

$m + l + 1$. The set of vertices $\{v_i\}_{i=1}^m \cup \{w_i\}_{i=1}^l \cup \{u_i\}_{i=1}^r$ is a minimum distance k -dominating set of cardinality $m + l + r$. The set of vertices $\{v_{i,1}\}_{i=1}^m \cup \{w_{i,k}\}_{i=1}^l \cup \{w_i\}_{i=1}^l \cup \{u_i\}_{i=1}^r \cup \{u_{r,k}\}$ is a minimum distance k -resolving dominating set of cardinality $m + 2l + r + 1$.




Figure 6. Tree $T_{k,m,l,r}^5$. ■

5. MAXIMUM ORDER GRAPHS

The maximum order n of a graph G having diameter d and metric dimension $\dim(G) = \beta$, was shown to be $\beta + d^\beta$ [8, 25]. This was proved by considering the maximum possible number of distinct metric representations with respect to a minimum resolving set. But this maximum order is only achieved when $d \leq 3$ or $\beta = 1$. Later, Hernando *et al.* [23] proved a stronger result by showing that

$$n \leq (\lfloor \frac{2d}{3} \rfloor + 1)^\beta + \beta \sum_{i=1}^{\lceil \frac{d}{3} \rceil} (2i - 1)^{\beta-1},$$

where the maximum order is achieved for any arbitrary positive integers d and β .

Cáceres *et al.* [5] showed that for a graph G of order n having $\gamma^r(G) = \gamma^r$, then $n \leq \gamma^r + \gamma^r \cdot 3^{\gamma^r - 1}$. They also provided graphs achieving this maximum order. Next, we generalize this result for $\gamma_k^r(G)$ for all $k \geq 1$.

Theorem 5.1. *For $k \geq 1$, the maximum order of a connected graph G having distance k -resolving domination number γ_k^r is $\gamma_k^r + \gamma_k^r \sum_{p=1}^k (2p+1)^{\gamma_k^r - 1}$. This maximum order is achieved for any $k, \gamma_k^r \geq 1$.*

Proof. Let G be a graph of order n and let S be a minimum distance k -resolving dominating set of G . For any vertex $x \in V \setminus S$, let us consider v_i a vertex in S such that $d_G(x, v_i) = p \leq k$. If $\gamma_k^r(G) = \gamma_k^r \geq 2$, for any vertex v_j from S different from v_i , the triangle inequality gives $|d_G(x, v_j) - d_G(v_i, v_j)| \leq d_G(x, v_i) = p$. It follows that the metric representation of x with respect to S has the coordinate corresponding to v_i equal to p and for the other coordinates there are at most $2p+1$ possible values in each of the other $\gamma_k^r - 1$ coordinates. Therefore, there are at most $(2p+1)^{\gamma_k^r - 1}$ possible metric representations of x with respect to the set S . Since $1 \leq p \leq k$, there are at most $\sum_{p=1}^k (2p+1)^{\gamma_k^r - 1}$ distinct metric representations for the vertices at distance less or equal to k from v_i . Since $|S| = \gamma_k^r$, we have $n \leq \gamma_k^r + \gamma_k^r \sum_{p=1}^k (2p+1)^{\gamma_k^r - 1}$.

Let k and γ_k^r be two arbitrary positive integers, we will prove that there exists a graph having distance k -resolving domination number γ_k^r and order $\gamma_k^r + \gamma_k^r \sum_{p=1}^k (2p+1)^{\gamma_k^r - 1}$.

If $\gamma_k^r = 1$, then from Theorem 3.3 the graph G is a path graph of maximum order $k+1$, which coincides with the maximum order bound. If $\gamma_k^r = r \geq 2$, we consider the following subsets of \mathbb{Z}^r ,

$$Q_0 = \{(0, 2k+1, 2k+1, \dots, 2k+1), (2k+1, 0, 2k+1, \dots, 2k+1), \dots, (2k+1, 2k+1, \dots, 2k+1, 0)\}.$$

For all $1 \leq i \leq r$,

$$Q_i = \{(q_1, q_2, \dots, q_r) : 1 \leq q_i \leq k, \text{ and for } j \neq i, 2k - q_i + 1 \leq q_j \leq 2k + q_i + 1\}.$$

Let G_r be the graph whose vertex set is $V(G_r) = \bigcup_{i=0}^r Q_i$. For which two vertices $q = (q_1, q_2, \dots, q_r)$ and $q' = (q'_1, q'_2, \dots, q'_r)$ are adjacent if and only if $|q_j - q'_j| \leq 1$ for each $1 \leq j \leq r$.

Claim 5.2. *The graph G_r is a connected graph.*

Proof. If $q_{i,0}, q_{j,0} \in Q_0$, where $q_{i,0}$ has the i -th element equal to 0 and $q_{j,0}$ has the j -th element equal to 0, we construct a $(q_{i,0}, q_{j,0})$ -path as following,

$$\begin{aligned}
& (2k+1, \dots, 0, 2k+1, \dots, 2k+1, \dots, 2k+1)(2k+1, \dots, 1, 2k+1, \dots, 2k, 2k+1, \dots, 2k+1) \\
& (2k+1, \dots, 2, 2k+1, \dots, 2k-1, 2k+1, \dots, 2k+1) \dots \dots \\
& (2k+1, \dots, k, 2k+1, \dots, k+1, 2k+1, \dots, 2k+1)(2k+1, \dots, k+1, 2k+1, \dots, k, 2k+1, \dots, 2k+1) \\
& (2k+1, \dots, k+2, 2k+1, \dots, k-1, 2k+1, \dots, 2k+1) \dots \dots \\
& (2k+1, \dots, 2k, 2k+1, \dots, 1, 2k+1, \dots, 2k+1)(2k+1, \dots, 2k+1, 2k+1, \dots, 0, 2k+1, \dots, 2k+1).
\end{aligned}$$

Also, for each $1 \leq i \leq r$, if $q = (q_1, q_2, \dots, q_r) \in Q_i$, it is easy to see from the definition of the adjacency in G_r , that there is a $(q, q_{i,0})$ -path. Hence, the graph G_r is a connected graph. \square

For $1 \leq i \leq r$ and $q \in V(G_r) \setminus Q_0$, we denote $L_i(q) = (f_i(q_1), f_i(q_2), \dots, f_i(q_r))$, where f_i is an integer-valued function defined as following.

If $q = (q_1, q_2, \dots, q_r) \in Q_s$, with $s \neq i$.

- For $j \notin \{s, i\}$, $f_i(q_j) = \begin{cases} q_j, & \text{if } q_j = 2k+1, \\ q_j - 1, & \text{if } q_j > 2k+1, \\ q_j + 1, & \text{if } q_j < 2k+1. \end{cases}$
- $f_i(q_s) = \begin{cases} q_s, & \text{if } q_s = k, \\ q_s + 1, & \text{if } q_s < k \text{ or } q_i = k+1. \end{cases}$
- $f_i(q_i) = q_i - 1$.

If $q = (q_1, q_2, \dots, q_r) \in Q_i$.

- For $j \neq i$, $f_i(q_j) = \begin{cases} q_j, & \text{if } q_j = 2k+1, \\ q_j - 1, & \text{if } q_j > 2k+1, \\ q_j + 1, & \text{if } q_j < 2k+1. \end{cases}$
- $f_i(q_i) = q_i - 1$.

For $t \geq 1$, we define $L_i^t(q)$ with $L_i^1(q) = L_i(q)$. For $t \geq 2$, $L_i^t(q) = L_i(L_i^{t-1}(q)) = (f_i^t(q_1), f_i^t(q_2), \dots, f_i^t(q_r))$, where f_i^t is the t -th iterated function of f_i , i.e., $f_i^t = \underbrace{f_i \circ f_i \circ \dots \circ f_i}_{t \text{ times}}$.

Claim 5.3. For all $1 \leq i \leq r$, for any vertex $q = (q_1, q_2, \dots, q_r) \in V(G_r) \setminus Q_0$, we have $L_i(q) \in V(G_r)$. Also, $L_i(q)$ is adjacent in G_r to q , and $L_i^{q_i}(q) = q_{0,i}$.

Proof. Let $q = (q_1, q_2, \dots, q_r) \in V(G_r) \setminus Q_0$. For $1 \leq i \leq r$, we have $L_i(q) = (f_i(q_1), f_i(q_2), \dots, f_i(q_r))$. If $q \in Q_s$, where $s \neq i$, for $j \neq s$, we have $2k - q_s + 1 \leq q_j \leq 2k + q_s + 1$ and $1 \leq q_s \leq k$. We discuss the membership of $L_i(q)$ according to the following cases.

- (i) If $q_s < k$, then we have $f_i(q_s) = q_s + 1 \leq k$, $f_i(q_i) = q_i - 1 \geq 2k - q_s$, and for $j \notin \{i, s\}$, $2k - q_s + 2 \leq f_i(q_j) \leq 2k + q_s$. So $L_i(q) = (f_i(q_1), f_i(q_2), \dots, f_i(q_r)) \in Q_s$.

- (ii) If $q_s = k$ and $q_i > k + 1$, then $f_i(q_i) = q_i - 1 \geq k + 1$, $f_i(q_s) = k$, and for $j \notin \{i, s\}$, $k + 1 \leq f_i(q_j) \leq 3k + 1$. So $L_i(q) = (f_i(q_1), f_i(q_2), \dots, f_i(q_r)) \in Q_s$.
- (iii) If $q_i = k + 1$, then $q_s = k$. It follows that $f_i(q_i) = k$, $f_i(q_s) = k + 1$, and for $j \notin \{i, s\}$, $k + 1 \leq f_i(q_j) \leq 3k + 1$. Therefore, $L_i(q) \in Q_i$.

Now, if $q \in Q_i$, from the definition of f it is easy to see that $L_i(q) \in Q_i$. Hence, for any vertex $q \in V(G_r) \setminus Q_0$, we have $L_i(q) \in V(G_r)$. Moreover, for $q \in V(G_r) \setminus Q_0$, and all $1 \leq i, j \leq r$, we have $|f_i(q_j) - q_j| \leq 1$, $f_i^{q_i}(q_i) = 0$, and for $j \neq i$, $f_i^{q_i}(q_j) = 2k + 1$. Thus, $L_i(q)q \in E(G_r)$, and $L_i^{q_i}(q) = q_{0,i}$. \square

Claim 5.4. For all $1 \leq i \leq r$, for any vertex $q = (q_1, q_2, \dots, q_r) \in V(G_r) \setminus Q_0$, $d_{G^r}(q, q_{0,i}) = q_i$.

Proof. Based on Claim 5.3 for $1 \leq i \leq r$, we have $qL_i(q)L_i^2(q) \cdots L_i^{q_i}(q) = q_{0,i}$ is a $(q, q_{0,i})$ -path in G_r of length q_i . Hence $d_{G^r}(q, q_{0,i}) \leq q_i$. Since $q_{0,i}$ and q are vertices having respectively 0 and q_i at the i -th coordinate and any two vertices in G_r can be adjacent only if the difference between the respective coordinates is at most 1, it follows that $d_{G^r}(q, q_{0,i}) \geq q_i$. Therefore, $d_{G^r}(q, q_{0,i}) = q_i$. \square

From above we can conclude that for any two different vertices q and q' in $V(G_r) \setminus Q_0$, there exists $1 \leq i \leq r$ such that $d_{G^r}(q, q_{0,i}) = q_i \neq d_{G^r}(q', q_{0,i}) = q'_i$. It follows that the set of vertices Q_0 is a resolving set of G_r . Also, for all $1 \leq i \leq r$, and any vertex $q \in Q_i$, $d_{G^r}(q, q_{0,i}) = q_i \leq k$. Hence, the set Q_0 is as well a distance k -dominating set of G_r . Hence, $\gamma_k^r(G_r) \leq |Q_0| = r$.

Suppose that $\gamma_k^r(G_r) \leq r - 1$. We have the order of the graph G_r is $|G_r| = r + r \sum_{p=1}^k (2p+1)^{r-1}$. Also the maximum order of a graph having $\gamma_k^r(G_r) \leq r - 1$ was previously proved to be less or equal to $\gamma_k^r(G_r) + \gamma_k^r(G_r) \sum_{p=1}^k (2p+1)^{\gamma_k^r(G_r)-1} \leq (r-1) + (r-1) \sum_{p=1}^k (2p+1)^{r-2}$, it is a contradiction. Therefore, $\gamma_k^r(G_r) = r$. \blacksquare

For $k = 1$, the maximum order in Theorem 5.1 is precisely the maximum order given in [5].

6. NORDHAUS-GADDUM TYPE BOUNDS

Nordhaus-Gaddum bounds are sharp bounds on the sum or the product of a parameter of a graph G and its complement \overline{G} . The survey [1] contains a bibliography of these types of bounds for some graph parameters. Hernando *et al.* [22] found Nordhaus-Gaddum type of bounds for the metric dimension and the resolving domination number. We provide those bounds for the distance k -resolving domination number for $k \geq 2$.

Theorem 6.1. For any graph G of order $n \geq 2$, we have the following.

- If $k = 2$, then

$$3 \leq \gamma_2^r(G) + \gamma_2^r(\overline{G}) \leq 2n - 1 \text{ and } 2 \leq \gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) \leq n(n - 1).$$

The lower bounds are attained if and only if $G \in \{K_2, \overline{K}_2, P_3, \overline{P}_3\}$.

The upper bounds are attained if and only if $G \in \{K_n, \overline{K}_n\}$.

- If $k \geq 3$, then

$$2 \leq \gamma_k^r(G) + \gamma_k^r(\overline{G}) \leq 2n - 1 \text{ and } 1 \leq \gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \leq n(n - 1).$$

The lower bounds are attained if and only if $G \cong P_4$.

The upper bounds are attained if and only if $G \in \{K_n, \overline{K}_n\}$.

Proof. If $k = 2$, then we have from Theorem 3.3 (a), $\gamma_2^r(G) = 1$ if and only if G is K_2 or P_3 . Also, for any other graph G , we have $\gamma_2^r(G) \geq 2$. This means that $\gamma_2^r(G) + \gamma_2^r(\overline{G}) \geq 3$ and $\gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) \geq 2$. Since $\gamma_2^r(\overline{K}_2) = 2$ and $\gamma_2^r(\overline{P}_3) = 2$, we can conclude that these lower bounds are attained if and only if $G \in \{K_2, \overline{K}_2, P_3, \overline{P}_3\}$.

If $k \geq 3$, then based on Theorem 3.3 (a), we have $\gamma_k^r(G) = 1$ if and only if $G \in \{P_2, P_3, \dots, P_{k+1}\}$. The graph P_4 is a self-complementary graph, i.e., $\overline{P}_4 \cong P_4$, we have $\gamma_k^r(\overline{P}_4) = \gamma_k^r(P_4) = 1$. Also, P_4 is the only graph whose complement is also a path and has a distance k -resolving domination number equal to 1. Therefore $\gamma_k^r(G) + \gamma_k^r(\overline{G}) \geq 2$ and $\gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \geq 1$, also these lower bounds are achieved if and only if G is P_4 .

Otherwise, for $k \geq 2$, we have $\gamma_k^r(G) = n$ if and only if G is the empty graph on n vertices \overline{K}_n , whose complement graph is the complete graph K_n . According to Theorem 3.3 (c), we have $\gamma_k^r(K_n) = n - 1$. Therefore, for any graph G of order $n \geq 2$, for $k \geq 2$, we have $\gamma_k^r(G) + \gamma_k^r(\overline{G}) \leq 2n - 1$ and $\gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \leq n(n - 1)$. Moreover, these upper bounds are achieved if and only if $G \in \{K_n, \overline{K}_n\}$. ■

Let G be a connected graph with $V(G) = \{1, 2, \dots, n\}$. The graph $G[H^i]$ is the graph obtained from G by replacing the vertex i with a graph H and joining each vertex of H to every vertex adjacent to i in G . Let H_1 and H_2 be two graphs, the graph $G[H_1^i, H_2^j]$ is the graph obtained from G by replacing the vertex i (respectively, j) with the graph H_1 (respectively, H_2) and joining each vertex of H_1 (respectively, H_2) to every vertex adjacent to i (respectively, j) in G . If i and j are adjacent in G , join every vertex of H_1 to every vertex of H_2 . The Bull graph B is the graph with vertex set $V(B) = \{1, 2, 3, 4, 5\}$ and edge set $E(B) = \{12, 13, 23, 14, 25\}$. The graph B is a self-complementary graph, i.e., $\overline{B} \cong B$.

Theorem 6.2. If G and \overline{G} are both connected graphs of order $n \geq 4$, we have the following.

- If $k = 2$, then

$$4 \leq \gamma_2^r(G) + \gamma_2^r(\overline{G}) \leq 2n - 4 \text{ and } 4 \leq \gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) \leq (n - 2)^2.$$

The upper bounds are attained if and only if $G \cong P_4$.

- If $k \geq 3$, then

$$2 \leq \gamma_k^r(G) + \gamma_k^r(\overline{G}) \leq 2n - 6 \text{ and } 1 \leq \gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \leq (n - 3)^2.$$

The lower bounds are attained if and only if $G \cong P_4$.

The upper bounds are attained if and only if $G \in \{P_4, C_5, B\} \cup \{P_4[K_{n-3}^1], P_4[\overline{K}_{n-3}^1], P_4[K_{n-3}^2], P_4[\overline{K}_{n-3}^2]\} \cup \{P_4[K_r^1, K_{n-r-2}^2] : 1 \leq r \leq n - 3\} \cup \{P_4[\overline{K}_r^1, \overline{K}_{n-r-2}^3] : 1 \leq r \leq n - 3\}$.

Proof. For $k = 2$, let G be a graph such that G and \overline{G} are connected graphs. From Theorem 3.3 (a), $\gamma_2^r(G) = 1$ if and only if G is either K_2 or P_3 . Then both G and \overline{G} have distance 2-resolving domination number greater or equal to 2. Hence, $\gamma_2^r(G) + \gamma_2^r(\overline{G}) \geq 4$ and $\gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) \geq 4$. Also based on Proposition 2.9 we have $\gamma_2^r(P_4) = \gamma_2^r(\overline{P}_4) = 2$, then the lower bounds are sharp.

Otherwise, we have from Theorem 3.3 (c), K_n is the only connected graph with distance 2-resolving domination number equal to $n - 1$. Since the complement of the complete graph is disconnected, it follows that $\gamma_2^r(G) \leq n - 2$. Moreover, from Theorem 3.3 (b), for $n \geq 4$, $\gamma_2^r(G) = n - 2$ if and only if G is either P_4 , $K_{s,t}(s, t \geq 1)$, $K_s + \overline{K}_t(s \geq 1, t \geq 2)$, or $K_s + (K_1 \cup K_t)(s, t \geq 1)$. The only graph from these graphs whose complement graph is also connected is the path P_4 . Since P_4 is self-complementary, we can conclude that $\gamma_2^r(G) + \gamma_2^r(\overline{G}) \leq 2n - 4$ and $\gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) \leq (n - 2)^2$, where the equality holds if and only if $G \cong P_4$.

For $k \geq 3$, we have $\gamma_k^r(P_4) = 1$. The graph P_4 is self-complementary and is the only graph in Theorem 3.3 (a) whose complement is a path graph having $\gamma_k^r(\overline{G}) = 1$. Then $\gamma_k^r(G) + \gamma_k^r(\overline{G}) \geq 2$ and $\gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \geq 1$, and these lower bounds are achieved if and only if G is P_4 .

Otherwise, we have from Theorem 3.3 (c), $\gamma_k^r(G) = n - 1$ if and only if G is a complete graph. It follows that $\gamma_k^r(G) \leq n - 2$. Furthermore, in Theorem 3.3 (b), $\gamma_k^r(G) = n - 2$ if and only if G is either $K_{s,t}(s, t \geq 1)$, $K_s + \overline{K}_t(s \geq 1, t \geq 2)$, or $K_s + (K_1 \cup K_t)(s, t \geq 1)$. Since the complements of these graphs are all disconnected, it follows that $\gamma_k^r(G) \leq n - 3$ and $\gamma_k^r(\overline{G}) \leq n - 3$. Therefore, for $k \geq 3$, $\gamma_k^r(G) + \gamma_k^r(\overline{G}) \leq 2n - 6$ and $\gamma_k^r(G) \cdot \gamma_k^r(\overline{G}) \leq (n - 3)^2$. The only connected graph of order 4 whose complement graph is also a connected graph is P_4 , we have $\gamma_k^r(P_4) = \gamma_k^r(\overline{P}_4) = 1$. Also for $n \geq 5$, based on Lemma 3.2, we have $\gamma_k^r(G) = \gamma_k^r(\overline{G}) = n - 3$ if and only if $\dim(G) = \dim(\overline{G}) = n - 3$. It follows that $\gamma_k^r(G) + \gamma_k^r(\overline{G}) = 2n - 6$ if and only if $\dim(G) + \dim(\overline{G}) = 2n - 6$. In [22], if G and \overline{G} are both connected graphs, we have $\dim(G) + \dim(\overline{G}) = 2n - 6$ if and only if

$G \in \{P_4, C_5, B\} \cup \{P_4[K_{n-3}^1], P_4[\overline{K}_{n-3}^1], P_4[K_{n-3}^2], P_4[\overline{K}_{n-3}^2]\} \cup \{P_4[K_r^1, K_{n-r-2}^2] : 1 \leq r \leq n-3\} \cup \{P_4[\overline{K}_r^1, \overline{K}_{n-r-2}^3] : 1 \leq r \leq n-3\}$. ■

7. SOME RELATIONS BETWEEN $\gamma_k^r(G)$ AND $\dim_k(G)$

For $k \geq 1$, let W be a k -truncated resolving set of a graph G . For any two distinct vertices $v, u \in V$, there exists a vertex x in W such that $d_k(v, x) = \min\{d_G(v, x), k+1\} \neq d_k(u, x) = \min\{d_G(u, x), k+1\}$. We have W is a resolving set of G . Also, at least one of u and v is at distance at most k from x . Based on this observation we get the following upper bound for $\gamma_k^r(G)$ in terms of $\dim_k(G)$.

Proposition 7.1. *For $k \geq 1$, let G be a connected graph. Then we have $\gamma_k^r(G) \leq \dim_k(G) + 1$.*

Proof. Let W be a minimum k -truncated resolving set of G . Then there is at most one vertex v in V such that $d_G(v, W) > k$. Otherwise, if v and u are two distinct vertices at distance greater than k from W , then $d_k(v, x) = d_k(u, x) = k+1$, for every $x \in W$. Now, suppose that there exists a vertex v such that $d_G(v, W) > k$, then the set $W \cup \{v\}$ is a distance k -dominating set of G . Since W is a resolving set of G , we have $W \cup \{v\}$ is a distance k -resolving dominating set of G . Thus, $\gamma_k^r(G) \leq |W| + 1 = \dim_k(G) + 1$. ■

If there exists a minimum k -truncated resolving set W of a connected graph G such that $d_G(v, W) \leq k$ for any $v \in V$, then necessarily $\gamma_k^r(G) \leq \dim_k(G)$.

In the following, we show that every k -truncated resolving set is a distance $(k+1)$ -resolving dominating set.

Proposition 7.2. *For $k \geq 1$, let G be a connected graph. Then we have $\gamma_{k+1}^r(G) \leq \dim_k(G)$.*

Proof. Let W be a minimum k -truncated resolving set of G . Suppose that there is a vertex v in V such that $d_G(v, W) \geq k+2$. Let u be a vertex adjacent to v . Then necessarily $d_G(u, W) \geq k+1$, otherwise $d_G(v, W) \leq k+1$. This means that $d_k(v, x) = d_k(u, x) = k+1$, for all $x \in W$, a contradiction. Therefore $d_G(v, W) \leq k+1$, for any vertex v in V . Thus W is a distance $(k+1)$ -resolving dominating set. Hence $\gamma_{k+1}^r(G) \leq |W| = \dim_k(G)$. ■

For $k \geq 1$, for a connected graph G of order n , we have $1 \leq \dim_k(G) \leq n-1$. A characterization of connected graphs of order n having $\dim_k(G) \in \{1, n-2, n-1\}$ is given in the following.

Theorem 7.3. *For a connected graph G of order $n \geq 2$, the following statements hold.*

- (a) [13] For $k \geq 1$, $\dim_k(G) = 1$ if and only if $G \in \{P_i\}_{i=2}^{k+2}$.
- (b) [14] For $n \geq 4$, $\dim_1(G) = n - 2$ if and only if $G \in \{P_4, K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$. For $k \geq 2$ and $n \geq 4$, $\dim_k(G) = n - 2$ if and only if $G \in \{K_{s,t}(s, t \geq 1), K_s + \overline{K}_t(s \geq 1, t \geq 2), K_s + (K_1 \cup K_t)(s, t \geq 1)\}$.
- (c) [14] For $k \geq 1$, $\dim_k(G) = n - 1$ if and only if $G \cong K_n$.

In Theorems 3.3 and 7.3, if G is a connected graph of order $n \geq 2$, we can see that for $k \geq 1$ and $r \in \{1, n - 2, n - 1\}$, we have $\dim_k(G) = r$ if and only if $\gamma_{k+1}^r(G) = r$.

Proposition 7.4. For $k \geq 1$, and any positive integers $\beta \geq 1$ and $\beta \leq \gamma \leq \beta + 1$, there exists a connected graph G having $\dim_k(G) = \beta$ and $\gamma_k^r(G) = \gamma$. For $\beta \geq 2$, the pair $(\beta, 1)$ is not realizable.

Proof. Let $\beta \geq 1$ and $\beta \leq \gamma \leq \beta + 1$. If $\gamma = \beta \geq 1$, we have $\gamma^r(K_{\beta+1}) = \beta$ [4]. By Theorem 3.3 and Theorem 7.3, for $k \geq 1$, $\dim_k(K_{\beta+1}) = \gamma_k^r(K_{\beta+1}) = \beta$.

If $\gamma = \beta + 1$, then for $k \geq 1$, if $\beta = 1$, according to Theorem 7.3 and Proposition 2.4, the path graph P_{k+2} has $\dim_k(P_{k+2}) = 1$ and $\gamma_k^r(P_{k+2}) = 2$. Now let $k \geq 1$ and $\beta \geq 2$. Let $S_{\beta+1,k}$ be the spider tree graph considered in the proof of Theorem 4.2 having one vertex v_0 of degree $\beta + 1$ and $\beta + 1$ leaves at distance k from v_0 . As shown previously in Theorem 4.2, we have $\dim(S_{\beta+1,k}) = \beta$ and $\gamma_k^r(S_{\beta+1,k}) = \beta + 1$. Let $v_1, v_2, \dots, v_{\beta+1}$ be the neighbors of v_0 in $S_{\beta+1,k}$ and let $W = \{v_1, v_2, \dots, v_{\beta+1}\}$. It is easy to check that W is a k -truncated resolving set of $S_{\beta+1,k}$. Therefore, $\dim_k(S_{\beta+1,k}) \leq |W| = \beta$. Since $\dim_k(S_{\beta+1,k}) \geq \dim(S_{\beta+1,k}) = \beta$, it follows that $\dim_k(S_{\beta+1,k}) = \beta$.

From Theorem 3.3, we have $\gamma_k^r(G) = 1$ if and only if G is a path graph of order at most $k + 1$. If $n \leq k + 1$, in Theorem 7.3, we have $\dim_k(P_n) = 1$. Therefore, there is no connected graph G having $\gamma_k^r(G) = 1$ and $\dim_k(G) \geq 2$. ■

The case $\gamma = \beta + 1$, in Proposition 7.4, proves the sharpness of the upper bound in Proposition 7.1.

To provide examples of connected graphs having $\dim_k(G) > \gamma_k^r(G)$, we give the k -truncated metric dimension of path graphs which appeared in [14].

Theorem 7.5 [14]. For $k \geq 1$, we have

- $\dim_k(P_n) = 1$ for $2 \leq n \leq k + 2$;
- $\dim_k(P_n) = 2$ for $k + 3 \leq n \leq 3k + 3$;
- for $n \geq 3k + 4$, we have

$$\dim_k(P_n) = \begin{cases} \left\lfloor \frac{2n+3k-1}{3k+2} \right\rfloor, & \text{if } n \equiv 0, 1, \dots, k+2 \pmod{(3k+2)}, \\ \left\lfloor \frac{2n+4k-1}{3k+2} \right\rfloor, & \text{if } n \equiv k+3, \dots, \left\lceil \frac{3k+5}{2} \right\rceil - 1 \pmod{(3k+2)}, \\ \left\lfloor \frac{2n+3k-1}{3k+2} \right\rfloor, & \text{if } n \equiv \left\lceil \frac{3k+5}{2} \right\rceil, \dots, 3k+1 \pmod{(3k+2)}. \end{cases}$$

From Theorem 7.5 and Proposition 2.4, we can see, for example, that if G is a path graph of order $6k+3$, then $\dim_k(G) = 4 > \gamma_k^r(G) = 3$. Moreover, we remark that the difference $\dim_k(G) - \gamma_k^r(G)$ can be arbitrarily large.

Proposition 7.6. *Let $k \geq 1$. For any positive integer N there exists a connected graph G with $\dim_k(G) - \gamma_k^r(G) > N$.*

Proof. For $k \geq 1$, let G be a path graph of order $n = i(3k+2)$ where $i \geq 1$. Based on Theorem 7.5, we have $\dim_k(G) = 2i$. From Proposition 2.4, $\gamma_k^r(G) = \left\lceil \frac{i(3k+2)}{2k+1} \right\rceil < \frac{i(3k+2)}{2k+1} + 1 \leq \frac{5}{3}i + 1$. It follows that $\dim_k(G) - \gamma_k^r(G) > 2i - \frac{5}{3}i - 1 = \frac{1}{3}i - 1 \rightarrow \infty$ as $i \rightarrow \infty$. ■

The upper bound in Proposition 2.9 holds for $\dim_k(G)$ the proofs are similar.

Proposition 7.7. *For $k \geq 1$, let G be a connected graph of order $n \geq k+1$, with $\text{diam}(G) \geq k$. Then $\dim_k(G) \leq n - k\gamma_k(G)$.*

8. CONCLUDING REMARKS

The study of the distance k -resolving domination number could be extended to other graph families and operations on graphs not discussed here. For example for trees, a formula in [20] is provided to compute efficiently $\gamma^r(T)$ for any tree T . We ask if it would be possible also for $\gamma_k^r(T)$ when $k \geq 2$. Also, it would be interesting to investigate the following questions.

- Is there a characterization of graphs achieving the bounds in Proposition 2.1?
- For $k \geq 1$ and $2 \leq \gamma \leq n-3$, can we characterize the connected graphs G of order n having $\gamma_k^r(G) = \gamma$?

A characterization of connected graphs G with $\gamma_2^r(G) = 2$ will provide all the graphs having $\gamma_2^r(G) + \gamma_2^r(\overline{G}) = 4$ and $\gamma_2^r(G) \cdot \gamma_2^r(\overline{G}) = 4$ in Theorem 6.2, where G and \overline{G} are both connected graphs.

In view of the discussion in Section 7 the following questions naturally arise.

- What is a sharp upper bound for $\dim_k(G)$ in terms of $\gamma_k^r(G)$ and what can be said about the ratio $\frac{\dim_k(G)}{\gamma_k^r(G)}$ for a connected graph G ?

- Is there a characterization of graphs G having $\gamma_k^r(G) = \dim_k(G) + 1$ or $\gamma_{k+1}^r(G) = \dim_k(G)$?
- For which pair β, γ of positive integers with $\gamma < \beta$ does there exist a connected graph G such that $\dim_k(G) = \beta$ and $\gamma_k^r(G) = \gamma$?

For $k \geq 1$, we denote $N_k(v) = \{x \in V : 0 < d_G(v, x) \leq k\}$, the *open k -neighborhood* of a vertex v in V . The *k -locating-dominating set* defined as a set $X \subseteq V$, verifying for every $v, u \in V \setminus X$, we have $\emptyset \neq N_k(v) \cap X \neq N_k(u) \cap X \neq \emptyset$. The minimum cardinality of such set is called the *k -locating-domination number* denoted by $LD_k(G)$. Results about the k -locating-domination number can be found surveyed in [27]. Necessarily every k -locating-dominating set is a distance k -resolving dominating set, the opposite is not true. Therefore for all $k \geq 1$, we have $\gamma_k^r(G) \leq LD_k(G)$. For $k = 1$, in [5] it is shown that $LD_1(T) \leq 2\gamma^r(T) - 2$ for any tree T different from P_6 . In [16], it is proved that $LD_1(G) \leq (\gamma^r(G))^2$ for any graph G not containing C_4 or C_6 as a subgraph. Finding an upper bound for $LD_1(G)$ in terms of $\gamma^r(G)$ for graphs in general is still open, it is shown [16] that such an upper bound is at least exponential in terms of $\gamma^r(G)$. Is it possible to find upper bounds for $LD_k(G)$ in terms of $\gamma_k^r(G)$ when $k \geq 2$ for graphs?

Acknowledgement

We gratefully acknowledge Airlangga University, Indonesia, and Ibn Zohr University, Morocco, for their supervision and support in accomplishing this work. The authors are grateful to the reviewers for all of their careful and valuable comments, which contributed to improve the paper.

REFERENCES

- [1] M. Aouchiche and P. Hansen, *A survey of Nordhaus-Gaddum type relations*, Discrete Appl. Math. **161** (2013) 466–546.
<https://doi.org/10.1016/j.dam.2011.12.018>
- [2] R.F. Bailey and P.J. Cameron, *Base size, metric dimension and other invariants of groups and graphs*, Bull. Lond. Math. Soc. **43** (2011) 209–242.
<https://doi.org/10.1112/blms/bdq096>
- [3] Z. Beerliova, F. Eberhard, T. Eberhard, A. Hall, M. Hoffmann, M. Mihalák and L.S. Ram, *Network discovery and verification*, IEEE J. Sel. Areas Commun. **24** (2006) 2168–2181.
<https://doi.org/10.1109/JSAC.2006.884015>
- [4] R.C. Brigham, G. Chartrand, R.D. Dutton and P. Zhang, *Resolving domination in graphs*, Math. Bohem. **128** (2003) 25–36.
<https://doi.org/10.21136/MB.2003.133935>
- [5] J. Cáceres, C. Hernando, M. Mora, I.M. Pelayo and M.L. Puertas, *Locating-dominating codes: Bounds and extremal cardinalities*, Appl. Math. Comput. **220** (2013) 38–45.
<https://doi.org/10.1016/j.amc.2013.05.060>

- [6] J. Cáceres, C. Hernando, M. Mora, I.M. Pelayo, M.L. Puertas, C. Seara and D.R. Wood, *On the metric dimension of Cartesian products of graphs*, SIAM J. Discrete Math. **21** (2007) 423–441.
<https://doi.org/10.1137/050641867>
- [7] G.J. Chang and G.L. Nemhauser, *The k -domination and k -stability problem on graphs*, Tech. Rep. **540** (School of Operations Res. and Industrial Eng., Cornell Univ., 1982) 332–345.
- [8] G. Chartrand, L. Eroh, M.A. Jhonson and O.R. Oellermann, *Resolvability in graphs and the metric dimension of a graph*, Discrete Appl. Math. **105** (2000) 99–113.
[https://doi.org/10.1016/S0166-218X\(00\)00198-0](https://doi.org/10.1016/S0166-218X(00)00198-0)
- [9] G. Chartrand, L. Lesniak and P. Zhang, *Graphs & Digraphs*, 6th Ed. (Chapman & Hall, London, 2016).
<https://doi.org/10.1201/b19731>
- [10] G. Chartrand, C. Poisson and P. Zhang, *Resolvability and the upper dimension of graphs*, Comput. Math. Appl. **39** (2000) 19–28.
[https://doi.org/10.1016/S0898-1221\(00\)00126-7](https://doi.org/10.1016/S0898-1221(00)00126-7)
- [11] G. Chartrand, V. Saenpholphat and P. Zhang, *The independent resolving number of a graph*, Math. Bohem. **128** (2003) 379–393.
<https://doi.org/10.21136/MB.2003.134003>
- [12] R.R. Davila, C. Fast, M.A. Henning and F. Kenter, *Lower bounds on the distance domination number of a graph*, Contrib. Discrete Math. **12** (2017).
<https://doi.org/10.11575/cdm.v12i2.62487>
- [13] A. Estrada-Moreno, I.G. Yero and J.A. Rodríguez-Velázquez, *On the (k,t) -metric dimension of graphs*, Comput. J. **64** (2021) 707–720.
<https://doi.org/10.1093/comjnl/bxaa009>
- [14] R.M. Frongillo, J. Geneson, M.E. Lladser, R.C. Tillquist and E. Yi, *Truncated metric dimension for finite graphs*, Discrete Appl. Math. **320** (2022) 150–169.
<https://doi.org/10.1016/j.dam.2022.04.021>
- [15] J. Geneson and E. Yi, *The distance- k dimension of graphs* (2021).
arXiv:2106.08303v2
- [16] A. González, C. Hernando and M. Mora, *Metric-locating-dominating sets of graphs for constructing related subsets of vertices*, Appl. Math. Comput. **332** (2018) 449–456.
<https://doi.org/10.1016/j.amc.2018.03.053>
- [17] F. Harary and R.A. Melter, *On the metric dimension of a graph*, Ars Combin. **2** (1976) 191–195.
- [18] M.A. Henning, *Distance domination in graphs*, in: Topics in Domination in Graphs, T.W. Haynes, S.T. Hedetniemi and M.A. Henning (Ed(s)) (Springer, Cham, 2020) 205–250.
https://doi.org/10.1007/978-3-030-51117-3_7

- [19] M.A. Henning, *Distance domination in graphs*, in: Domination in Graphs: Advanced Topics, T.W. Haynes, S.T. Hedetniemi, and P.J. Slater (Ed(s)) (Marcel Dekker, Inc. New York, 1998) 321–349.
<https://doi.org/10.1201/9781315141428>
- [20] M.A. Henning and O.R. Oellermann, *Metric-locating-dominating sets in graphs*, Ars Combin. **73** (2004) 129–141.
- [21] M.A. Henning, O.R. Oellermann and H.C. Swart, *Relating pairs of distance domination parameters*, J. Combin. Math. Combin. Comput. **18** (1995) 233–244.
- [22] C. Hernando, M. Mora and I.M. Pelayo, *Nordhaus-Gaddum bound for locating domination*, European J. Combin. **36** (2014) 1–6.
<https://doi.org/10.1016/j.ejc.2013.04.009>
- [23] C. Hernando, M. Mora, I.M. Pelayo, C. Seara and D.R. Wood, *Extremal graph theory for metric dimension and diameter*, Electron. J. Combin. **17** (2010) #R30.
<https://doi.org/10.37236/302>
- [24] M. Jannesari and B. Omoomi, *The metric dimension of the lexicographic product of graphs*, Discrete Math **312** (2012) 3349–3356.
<https://doi.org/10.1016/j.disc.2012.07.025>
- [25] S. Khuller, B. Raghavachari and A. Rosenfeld, *Landmarks in graphs*, Discrete Appl. Math. **70** (1996) 217–229.
[https://doi.org/10.1016/0166-218X\(95\)00106-2](https://doi.org/10.1016/0166-218X(95)00106-2)
- [26] D. Lichtenstein, *Planar formulae and their uses*, SIAM J. Comput. **11** (1982) 329–343.
<https://doi.org/https://doi.org/10.1137/0211025>
- [27] A. Lobstein, O. Hudry and I. Charon, *Locating-domination and identification*, in: Topics in Domination in Graphs, T.W. Haynes, S.T. Hedetniemi and M.A. Henning (Ed(s)) (Springer, Cham, 2020) 251–299.
https://doi.org/10.1007/978-3-030-51117-3_8
- [28] A. Meir and J.W. Moon, *Relations between packing and covering numbers of a tree*, Pacific J. Math. **61**(1) (1975) 225–233.
<https://doi.org/10.2140/pjm.1975.61.225>
- [29] V. Saenpholphat and P. Zhang, *Connected resolvability of graphs*, Czechoslovak Math. J. **53** (2003) 827–840.
<https://doi.org/10.1023/B:CMAJ.0000024524.43125.cd>
- [30] P.J. Slater, *Leaves of trees*, Congr. Numer. **14** (1975) 549–559.
- [31] P.J. Slater, *R-domination in graphs*, J. ACM **23** (1976) 446–450.
<https://doi.org/10.1145/321958.321964>
- [32] R.C. Tillquist, R.M. Frongillo and M.E. Lladser, *Getting the lay of the land in discrete space: A survey of metric dimension and its applications* (2021).
arXiv:2104.07201

- [33] R.C. Tillquist, R.M. Frongillo and M.E. Lladser, *Truncated metric dimension for finite graphs* (2021).
arXiv:2106.14314v1
- [34] D.A.R. Wardani, M.I. Utoyo, Dafik and K. Dliou, *The distance 2-resolving domination number of graphs*, J. Phys. Conf. Ser. **1836** (2021) 012017.
<https://doi.org/10.1088/1742-6596/1836/1/012017>

Received 11 August 2022

Revised 18 January 2023

Accepted 27 January 2023

Available online 15 February 2023