

TIGHT DESCRIPTION OF FACES IN TOROIDAL GRAPHS WITH MINIMUM DEGREE AT LEAST 4

OLEG V. BORODIN¹

Sobolev Institute of Mathematics Siberian Branch
Russian Academy of Sciences, Novosibirsk, 630090, Russia
e-mail: brdnoleg@math.nsc.ru

AND

ANNA O. IVANOVA²

Ammosov North-Eastern Federal University
Yakutsk, 677013, Russia
e-mail: shmgnanna@mail.ru

Abstract

The degree $d(x)$ of a vertex or face x in a graph G is the number of incident edges. A face $f = v_1 \cdots v_{d(f)}$ in a graph G on the plane or other orientable surface is of type (k_1, k_2, \dots) if $d(v_i) \leq k_i$ for each i . By δ we denote the minimum vertex-degree of G .

It follows from the classical theorem by Lebesgue (1940) that every plane triangulation with $\delta \geq 4$ has a 3-face of types $(4, 4, \infty)$, $(4, 5, 19)$, $(4, 6, 11)$, $(4, 7, 9)$, $(5, 5, 9)$, or $(5, 6, 7)$. In 1999, Jendrol' gave a similar description: “ $(4, 4, \infty)$, $(4, 5, 13)$, $(4, 6, 17)$, $(4, 7, 8)$, $(5, 5, 7)$, $(5, 6, 6)$ ” and conjectured that “ $(4, 4, \infty)$, $(4, 5, 10)$, $(4, 6, 15)$, $(4, 7, 7)$, $(5, 5, 7)$, $(5, 6, 6)$ ” holds. In 2002, Lebesgue's description was strengthened by Borodin to “ $(4, 4, \infty)$, $(4, 5, 17)$, $(4, 6, 11)$, $(4, 7, 8)$, $(5, 5, 8)$, $(5, 6, 6)$ ”. In 2014, we obtained the following tight description, which, in particular, disproves the above mentioned conjecture by Jendrol': “ $(4, 4, \infty)$, $(4, 5, 11)$, $(4, 6, 10)$, $(4, 7, 7)$, $(5, 5, 7)$, $(5, 6, 6)$ ”, and recently proved another tight description of faces in plane triangulations with $\delta \geq 4$: “ $(4, 4, \infty)$, $(4, 6, 10)$, $(4, 7, 7)$, $(5, 5, 8)$, $(5, 6, 7)$ ”.

It follows from Lebesgue's theorem of 1940 that every plane 3-connected quadrangulation has a face of one of the types $(3, 3, 3, \infty)$, $(3, 3, 4, 11)$, $(3, 3, 5, 7)$, $(3, 4, 4, 5)$. Recently, we improved this description to “ $(3, 3, 3, \infty)$,

¹The first author's work was supported by the Ministry of Science and Higher Education of the Russian Federation (project no. FWNF-2022-0017)

²The second author's work was supported by the Ministry of Science and Higher Education of the Russian Federation (Grant No. FSRG-2023-0025)

$(3, 3, 4, 9)$, $(3, 3, 5, 6)$, $(3, 4, 4, 5)$ ”, where all parameters except possibly 9 are best possible and 9 cannot go down below 8.

In 1995, Avgustinovich and Borodin proved the following tight description of the faces of torus quadrangulations with $\delta \geq 3$: “ $(3, 3, 3, \infty)$, $(3, 3, 4, 10)$, $(3, 3, 5, 7)$, $(3, 3, 6, 6)$, $(3, 4, 4, 6)$, $(4, 4, 4, 4)$ ”.

Recently, we proved that every triangulation with $\delta \geq 4$ of the torus has a face of one of the types $(4, 4, \infty)$, $(4, 6, 12)$, $(4, 8, 8)$, $(5, 5, 8)$, $(5, 6, 7)$, or $(6, 6, 6)$, which description is tight.

The purpose of this paper is to prove that every graph with $\delta \geq 4$ that admits a closed 2-cell embedding on the torus has a face of one of the types $(4, 4, 4, 4)$, $(4, 4, \infty)$, $(4, 5, 16)$, $(4, 6, 12)$, $(4, 8, 8)$, $(5, 5, 8)$, $(5, 6, 7)$, or $(6, 6, 6)$, where all parameters are best possible.

Keywords: plane graph, toroidal graph, degree, face, structure.

2020 Mathematics Subject Classification: 05C75.

1. INTRODUCTION

The *degree* $d(x)$ of a vertex or face x in a plane or torus graph G is the number of incident edges. A k -*vertex* and k -*face* is one of degree k , a k^+ -*vertex* has degree at least k , and so on. A face f in a graph G on the plane or torus is of *type* (k_1, k_2, \dots) , or a (k_1, k_2, \dots) -*face* if $d(v_i) \leq k_i$ for each i . By δ and w denote the minimum vertex degree and smallest degree-sum of faces in G , respectively.

We now recall some results on the structure of faces in plane graph with $\delta \geq 3$, beginning with the fundamental theorem of Lebesgue [19] from 1940.

Theorem 1 (Lebesgue [19]). *Every plane graph with $\delta \geq 3$ has a face of one of the following types:*

$$\begin{aligned} &(3, 6, \infty), (3, 7, 41), (3, 8, 23), (3, 9, 17), (3, 10, 14), (3, 11, 13), \\ &(4, 4, \infty), (4, 5, 19), (4, 6, 11), (4, 7, 9), (5, 5, 9), (5, 6, 7), \\ &(3, 3, 3, \infty), (3, 3, 4, 11), (3, 3, 5, 7), (3, 4, 4, 5), (3, 3, 3, 3, 5). \end{aligned}$$

The classical Theorem 1, along with other ideas in Lebesgue [19], has a lot of applications to plane graph coloring problems (the first example of such applications and recent surveys can be found in [4, 13, 17, 21]).

Some parameters of Lebesgue’s theorem were improved for several narrow classes of plane graphs. Back in 1963, Kotzig [18] proved that every plane triangulation with $\delta = 5$ satisfies $w \leq 18$ and conjectured that $w \leq 17$ holds. In 1989, Kotzig’s conjecture was confirmed by Borodin [2] in a more general form by proving that every such a graph has a $(5, 5, 7)$ -face or a $(5, 6, 6)$ -face, where all parameters are tight. This result also confirmed a conjecture of Grünbaum [14]

from 1975 on the cyclic 11-connectivity of 5-connected planar graphs, and it has been extended to several classes of plane graphs over the last decades; see, for example, recent surveys [7, 13, 17] and also [3–5, 15, 16, 20].

It follows from the classical theorem by Lebesgue [19] that every plane triangulation with $\delta \geq 4$ has a 3-face of types $(4, 4, \infty)$, $(4, 5, 19)$, $(4, 6, 11)$, $(4, 7, 9)$, $(5, 5, 9)$, or $(5, 6, 7)$. In 1999, Jendrol' [16] gave a similar description: “ $(4, 4, \infty)$, $(4, 5, 13)$, $(4, 6, 17)$, $(4, 7, 8)$, $(5, 5, 7)$, $(5, 6, 6)$ ” and conjectured that “ $(4, 4, \infty)$, $(4, 5, 10)$, $(4, 6, 15)$, $(4, 7, 7)$, $(5, 5, 7)$, $(5, 6, 6)$ ” holds. In 2002, Lebesgue's description was strengthened by Borodin [3] to “ $(4, 4, \infty)$, $(4, 5, 17)$, $(4, 6, 11)$, $(4, 7, 8)$, $(5, 5, 8)$, $(5, 6, 6)$ ”. In 2014, we obtained [6] the following tight description, which, in particular, disproves the above mentioned conjecture by Jendrol': “ $(4, 4, \infty)$, $(4, 5, 11)$, $(4, 6, 10)$, $(4, 7, 7)$, $(5, 5, 7)$, $(5, 6, 6)$ ”, and recently proved [11] another tight description of faces in plane triangulations with $\delta \geq 4$: “ $(4, 4, \infty)$, $(4, 6, 10)$, $(4, 7, 7)$, $(5, 5, 8)$, $(5, 6, 7)$ ”.

In particular, precise descriptions of the structure of faces were obtained for plane graphs with $\delta \geq 4$ (Borodin, Ivanova [5]) and for plane triangulations (Borodin, Ivanova, Kostochka [12]). It follows from Theorem 1 that every plane quadrangulation with $\delta \geq 3$ has a face of one of the types $(3, 3, 3, \infty)$, $(3, 3, 4, 11)$, $(3, 3, 5, 7)$, $(3, 4, 4, 5)$. Recently, we improved [8] this result to the following description: “ $(3, 3, 3, \infty)$, $(3, 3, 4, 9)$, $(3, 3, 5, 6)$, $(3, 4, 4, 5)$ ”, where all parameters except possibly 9 are best possible, while 9 cannot go down below 8.

In 1995, Avgustinovich and Borodin gave the following tight description of faces in quadrangulations of the torus.

Theorem 2 (Avgustinovich, Borodin [1]). *Every torus quadrangulation with $\delta \geq 3$ has a face of one of the following types: $(3, 3, 3, \infty)$, $(3, 3, 4, 10)$, $(3, 3, 5, 7)$, $(3, 3, 6, 6)$, $(3, 4, 4, 6)$, $(4, 4, 4, 4)$, where all parameters are best possible.*

Recently, we proved [9] that every toroidal triangulation with $\delta \geq 5$ has a face of one of the types $(5, 5, 8)$, $(5, 6, 7)$, or $(6, 6, 6)$, and later on we extended [10] this description to $\delta \geq 4$ as follows: “ $(4, 4, \infty)$, $(4, 6, 12)$, $(4, 8, 8)$, $(5, 5, 8)$, $(5, 6, 7)$, $(6, 6, 6)$ ”, which is also tight.

The purpose of our paper is to prove the following further extension of the result in [10].

Theorem 3. *Every graph with $\delta \geq 4$ that admits a closed 2-cell embedding on the torus has a face of one of the types:*

- (Ta) $(4, 4, 4, 4)$,
- (Tb) $(4, 4, \infty)$,
- (Tc) $(4, 5, 16)$,
- (Td) $(4, 6, 12)$,
- (Te) $(4, 8, 8)$,

(Tf) $(5, 5, 8)$,

(Tg) $(5, 6, 7)$, *or*

(Th) $(6, 6, 6)$,

where all parameters are best possible.

2. THE TIGHTNESS OF THEOREM 3

It is easy to construct a 4-regular quadrangulation of the torus; for example, by deleting all 4-vertices from a graph on Figure 4, so the item (Ta) in our description is necessary. Now to justify (Tc), it suffices to replace each face of such a quadrangulation by a construction in Figure 1.




Figure 1. All 3-faces are of type $(4, 5, 16)$, and there are no $(4, 4, 4, 4)$ -faces.

The tightness of (Tb) is confirmed by the double n -pyramid augmented by an edge joining its n -vertices. Figure 2 represents a bipartite torus graph with four 6-faces. Putting a vertex inside each its face and joining it with the six boundary vertices produces a 6-regular triangulation $T(6, 6, 6)$, which confirms the necessity and sharpness of (Th).

Next we put a vertex on every edge of $T(6, 6, 6)$, followed by putting a vertex $v(f)$ inside each 6-face f obtained and joining $v(f)$ with the six vertices of f ; this results in a triangulation $T(4, 6, 12)$ confirming the tightness of (Td).







Figure 2. A bipartite torus graph with four 6-faces ([10]).

Figure 3. Producing a torus graph with all faces of type $(4, 6, 12)$ ([10]).

The next two constructions confirm the tightness of (Te) and (Tf), respectively.

Figure 4. All faces are of type $(4, 8, 8)$ ([10]).Figure 5. A torus graph with all $(5, 5, 8)$ -faces ([9]).

Finally, replacing each 6-face in Figure 2 by the construction shown in Figure 6 produces a torus triangulation which confirms that the term (Th) in Theorem 3 is also best possible.




Figure 6. A replacement for each 6-face in Figure 1 which results in all faces of type $(5^+, 6^+, 7^+)$ ([9]).

3. PROVING THE EXISTENCE OF FACE-TYPES IN THEOREM 3

Suppose G is a counterexample to Theorem 3. Euler's formula $|V| - |E| + |F| = 0$ for G , where V and F are the sets of its vertices and faces, respectively, can be rewritten as follows.

$$(1) \quad \sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2d(f) - 6) = 0.$$

We assign a *charge* $\mu(v) = d(v) - 6$ to every vertex v and $\mu(f) = 2d(f) - 6$ to every face f of G , so that only 5^- -vertices have a negative charge. Using the properties of G as a counterexample, we define a local redistribution of charges, preserving their sum, such that the *new charge* $\mu'(x)$ satisfies $\mu'(x) \geq 0$ whenever $x \in V \cup F$ and there is at least one x in $V \cup F$ with $\mu'(x) > 0$. This will contradict the fact that the sum of the new charges is by (1) equal to 0.

In what follows, by “non- (k, l, m) !” we mean a short-hand for “since T has no (k, l, m) -faces”. The neighbors of a vertex or face x in a cyclic order are denoted by $v_1, v_2, \dots, v_{d(x)}$.

We use the following rules R1–R3 of discharging (see Figure 7).

R1. Every 4^+ -face gives $\frac{1}{2}$ to each incident vertex of degree 4 or 5.

R2. Each 4-vertex v in a face $f_1 = v_1 v v_2$ with $d(v_1) \leq d(v_2)$ (where $d(v_1) \geq 5$ due to non- $(4, 4, \infty)$!) receives the following charges from v_1, v_2 through f_1 .

(R2a) If $d(v_1) = 5$ or $d(v_1) = 6$, then $d(v_2) \geq 17$ due to non- $(4, 5, 16)$! or $d(v_2) \geq 13$ due to non- $(4, 6, 12)$!, respectively, and v_2 gives $\frac{1}{2}$ to v .

(R2b) Suppose $7 \leq d(v_1) \leq 8$; then $d(v_2) \geq 9$ by non- $(4, 8, 8)$!, and now v_2 gives $\frac{1}{3}$ to v , while v_1 gives $\frac{1}{6}$.

(R2c) If $d(v_1) \geq 9$, then each of v_1 and v_2 gives $\frac{1}{3}$ to v .

R3. Each 5-vertex v in a face $f_1 = v_1 v v_2$ receives the following charges from v_1, v_2 through f_1 .

(R3a) If $d(v_1) = 7$, then v_1 gives $\frac{1}{7}$.

(R3b) If $d(v_1) = 8$ (and hence $d(v_2) \geq 6$ due to non- $(5, 5, 8)$!), then v_1 gives $\frac{5}{24}$.

(R3c) If $9 \leq d(v_1) \leq 12$ and $d(v_2) \geq 6$, then v_1 gives $\frac{1}{3}$ to v .

(R3d) Suppose $d(v_2) \geq 6$; then v_1 gives to v :

(R3d1) $\frac{1}{3}$ if $13 \leq d(v_1) \leq 16$, or

(R3d2) $\frac{1}{2}$ if $d(v_1) \geq 17$.

(R3e) If $d(v_2) = 5$ (and hence $d(v_1) \geq 9$ by non- $(5, 5, 8)$!), then v_1 gives $\frac{1}{7}$ to v and, by symmetry, also to v_2 .

(R3f) If $d(v_2) = 4$ (and hence $d(v_1) \geq 17$ by non- $(4, 5, 16)$!), then v_1 gives $\frac{1}{8}$ to v .

We now check that $\mu'(x) \geq 0$ whenever $x \in V \cup F$ and at least one vertex or face has a strictly positive new charge μ' .




Figure 7. Rules of discharging.

First consider $f \in F$. If $d(f) = 3$, then $\mu'(f) = \mu(f) = 0$ as f does not participate in discharging. If $d(f) \geq 4$, then $\mu'(f) \geq 2d(f) - 6 - d(f) \times \frac{1}{2} = \frac{3(d(f)-4)}{2} \geq 0$ in view of R1. In particular, $\mu'(f) > 0$ when $d(f) \geq 5$.

From now on suppose $v \in V$. Here, our proof splits.

Case 1. $d(v) = 4$. Note that v receives the total of at least $\frac{1}{2} = \frac{1}{3} + \frac{1}{6}$ through each incident face by R1, R2 due to non- $(4, 4, \infty)!$, non- $(4, 5, 16)!$, non- $(4, 6, 12)!$, and non- $(4, 8, 8)!$. Thus we already have $\mu'(v) \geq 4 - 6 + 4 \times \frac{1}{2} = 0$. Furthermore, it is not difficult to see that in fact $\mu'(v) > 0$ if v participates at least once in R2c.

Case 2. $d(v) = 5$. We know that v receives $\frac{1}{2}$ from each 4^+ -face by R1. Through each 3-face, v receives $\frac{1}{8}$ by R3f and at least $\frac{1}{7}$ by R3a–R3e. It is easy to see that $\mu'(v) \geq 5 - 6 + 2 \times \frac{1}{2} + 3 \times \frac{1}{8} > 0$ if v is incident with at least two 4^+ -faces. If v belongs to the boundary of precisely one 4^+ -face, then $\mu'(v) \geq -1 + \frac{1}{2} + 4 \times \frac{1}{8} \geq 0$, with the equality only when each of its incident 3-faces participates in R3f. In particular, $\mu'(v) > 0$ unless v has at least two 17^+ -neighbors due to non-(4, 5, 16)!.

From now on suppose v is completely surrounded by 3-faces. Note that v has at most two 6^- -neighbors by non-(5, 6, 6)!, so we can assume that $d(v_1) \geq 7$, $d(v_3) \geq 7$ and $d(v_4) \geq 7$.

To ensure $\mu'(v) \geq 0$, it suffices, due to symmetry, to check that v receives the total of at least $\frac{1}{2}$ from v_1 , v_2 and v_3 through the faces $f_1 = v_1vv_2$, $f_2 = v_2vv_3$ and $f_3 = v_3vv_4$.

If $d(v_2) = 4$, then $d(v_1) \geq 17$ and $d(v_3) \geq 17$ by non-(4, 5, 16)!, so we are done already due to the contribution of $\frac{1}{2}$ from v_3 by R3d2.

For $d(v_2) = 5$, we have $d(v_1) \geq 9$ and $d(v_3) \geq 9$ by non-(5, 5, 8)!, so v receives $\frac{1}{3} + \frac{1}{7}$ from v_3 by R3c or R3d1 combined with R3e. Furthermore, v receives $\frac{1}{7}$ from v_1 by R3e. Thus v receives $\frac{1}{3} + 2 \times \frac{1}{7} > \frac{1}{2}$ from v_1 and v_3 through the faces f_1 , f_2 and f_3 .

Next suppose $d(v_2) = 6$; now $d(v_1) \geq 8$ and $d(v_3) \geq 8$ by non-(5, 6, 7)!. This means that v receives from v_3 either $2 \times \frac{5}{24}$ by R3b if $d(v_3) = 8$, or $2 \times \frac{1}{3}$ by R3c combined with R3d1 when $d(v_3) \geq 9$. From v_1 , our v receives through f_1 either $\frac{5}{24}$ by R3b if $d(v_1) = 8$, or $\frac{1}{3}$ by R3c when $9 \leq d(v_1) \leq 12$. Now if $d(v_1) \geq 13$, then v_1 sends to v through f_1 either $\frac{1}{3}$ by R3d1 or $\frac{1}{2}$ by R3d2. In total, v receives at least $3 \times \frac{5}{24} > \frac{1}{2}$ through the faces f_1 , f_2 and f_3 .

Finally, suppose $d(v_2) \geq 7$. Here, v receives from v_2 through the faces f_1 and f_2 either at least $2 \times \frac{1}{7}$ by R3a–R3c when $d(v_2) \leq 12$, or $2 \times \frac{1}{3}$ by R3d1 otherwise. By symmetry, the same donation occurs from v_3 through f_2 and f_3 , and so v receives at least $\frac{4}{7} > \frac{1}{2}$ from v_2 and v_3 through the faces f_1 , f_2 and f_3 , as desired.

In particular, we have proved that a 5-vertex v not only satisfies $\mu'(v) \geq 0$, but in fact $\mu'(v) > 0$ holds unless v is incident with precisely one 4^+ -face and has two or three 17^+ -neighbors along with three or two 4-neighbors, respectively.

Case 3. $d(v) = 6$. Since v does not participate in discharging, we have $\mu'(v) = 6 - 6 = 0$.

Case 4. $d(v) = 7$. Such a vertex v sends at most $\frac{1}{6}$ through each incident 3-face which is incident with a 5^- -vertex; namely, by R2b and R3a. If v is incident with at least one 4^+ -face, then $\mu'(v) \geq 7 - 6 - 6 \times \frac{1}{6} = 0$. Otherwise, due to non-(5, 5, 8)!, our v has at most three 5^- -neighbors, and hence again we have $\mu'(v) \geq 1 - 2 \times 3 \times \frac{1}{6} = 0$, as desired.

Case 5. $d(v) = 8$. Now v can only send at most $\frac{5}{24}$ through each incident 3-face by R2b or R3b, in view of non-(5, 5, 8)!, which means that $\mu'(v) \geq 8 - 6 - 8 \times \frac{5}{24} > 0$.

Case 6. $9 \leq d(v) \leq 12$. Note that v sends $\frac{1}{3}$ through each incident 3-face to a unique 5^- -vertex in the boundary of that face by R2b, R2c and R3c or $2 \times \frac{1}{7}$ through a face incident with two 5-vertices by R3e due to non-(4, 5, 16)!. This implies $\mu'(v) \geq d(v) - 6 - \frac{d(v)}{3} = \frac{2(d(v)-9)}{3} \geq 0$. In particular, $\mu'(v) > 0$ whenever $10 \leq d(v) \leq 12$.

Now suppose $d(v) = 9$. If v is incident with a 4^+ -face or has two consecutive 6^+ -neighbors in an incident 3-face, then $\mu'(v) \geq 9 - 6 - 8 \times \frac{1}{3} > 0$. Otherwise, v is completely surrounded by 3-faces and, by parity reasons, has two consecutive 5-neighbors, which implies $\mu'(v) \geq 3 - 8 \times \frac{1}{3} - \frac{2}{7} > 0$.

Case 7. $13 \leq d(v) \leq 16$. Such a v sends through each incident 3-face at most $\frac{1}{2}$ to a unique 5^- -vertex by R2, R3d1 or $2 \times \frac{1}{7}$ by R3e, since R3f is not applicable in view of non-(4, 5, 16)!. This implies that $\mu'(v) \geq d(v) - 6 - \frac{d(v)}{2} > \frac{d(v)-12}{2} > 0$.

Case 8. $d(v) \geq 17$. Here, v sends through each incident 3-face vv_1v_2 either $\frac{1}{2} + \frac{1}{8}$ by R2a combined with R3f if $d(v_1) = 4$ and $d(v_2) = 5$, or at most $\frac{1}{2}$ by R2, R3 otherwise, which yields $\mu'(v) \geq d(v) - 6 - \frac{5d(v)}{8} = \frac{3(d(v)-16)}{8} > 0$.

Remark 4. Thus we have proved $\mu'(v) \geq 0$ for every $v \in V$. Furthermore, as shown in *Cases 5–8*, a vertex v satisfies $\mu'(v) > 0$ if $d(v) \geq 8$.

To arrive at a final contradiction with (1), it suffices to show, assuming $\delta(G) \leq 7$, that in fact there is an $x \in V \cup F$ with $\mu'(x) > 0$, since this will imply

$$0 = \sum_{v \in V} (d(v) - 6) + \sum_{f \in F} (2d(f) - 6) = \sum_{x \in V \cup F} \mu'(x) > 0.$$

To this aim, first consider a vertex v . If $d(v) = 5$, then, as shown in *Case 2*, $\mu'(v) > 0$ unless v has a 17^+ -neighbor, which possibility is already excluded. Thus it remains to assume that G has no 5-vertices.

Next suppose $d(v) = 7$. It now follows from non-(4, 8, 8)! that v cannot send a positive charge through a 3-face by R2b due to the absence of 9^+ -vertices in G and since R3a is not applicable because of the absence of 5-vertices, which means that $\mu'(v) = \mu(v) = 1$. Thus it remains to assume, moreover, that G has no 7-vertices either.

Furthermore, now each 3-face of G must be incident with three 6^- -vertices, contrary to non-(6, 6, 6)!. This implies that G has no 3-faces.

Finally, as follows from an inequality two lines above *Case 1*, G cannot have 5^+ -faces since they have $\mu' > 0$. Hence all faces of G are 4-faces. However, each

4-face f must be incident with at least one 6-vertex by non- $(4, 4, 4, 4)!$, which 6-vertex does not receive charge from f by R1, and so $\mu'(f) \geq 2 \times 4 - 6 - 3 \times \frac{1}{2} > 0$.

This contradiction completes the proof of Theorem 3.

REFERENCES

- [1] S.V. Avgustinovich and O.V. Borodin, *Edge neighborhoods in normal maps*, Diskretn. Anal. Issled. Oper. **2(3)** (1989) 9–12, in Russian.
Translation in: Operations Research and Discrete Analysis, A.D. Korshunov (Eds), Math. Appl. (N.Y.) **391** (Springer Netherlands, 1997) 17–22.
https://doi.org/10.1007/978-94-011-5678-3_3
- [2] O.V. Borodin, *Solution of problems of Kotzig and Grünbaum concerning the isolation of cycles in planar graphs*, Mat. Zametki **46(5)** (1989) 9–12, in Russian.
Translation in: Math. Notes **46(5)** (1989) 835–837.
<https://doi.org/10.1007/BF01139613>
- [3] O.V. Borodin, *An improvement of Lebesgues theorem on the structure of minor faces of 3-polytopes*, Diskretn. Anal. Issled. Oper. **9(3)** (2002) 29–39, in Russian.
- [4] O.V. Borodin, *Colorings of plane graphs: A survey*, Discrete Math. **313** (2013) 517–539.
<https://doi.org/10.1016/j.disc.2012.11.011>
- [5] O.V. Borodin and A.O. Ivanova, *Describing 3-faces in normal plane maps with minimum degree 4*, Discrete Math. **313** (2013) 2841–2847.
<https://doi.org/10.1016/j.disc.2013.08.028>
- [6] O.V. Borodin and A.O. Ivanova, *Combinatorial structure of faces in triangulated 3-polytopes with minimum degree 4*, Sib. Math. J. **55** (2014) 12–18.
<https://doi.org/10.1134/S0037446614010030>
- [7] O.V. Borodin, A.O. Ivanova, *New results about the structure of plane graphs: A survey*, AIP Conference Proceedings **1907** (2017) 030051.
<https://doi.org/10.1063/1.5012673>
- [8] O.V. Borodin and A.O. Ivanova, *An improvement of Lebesgue’s description of edges in 3-polytopes and faces in plane quadrangulations*, Discrete Math. **342** (2019) 1820–1827.
<https://doi.org/10.1016/j.disc.2019.02.019>
- [9] O.V. Borodin and A.O. Ivanova, *Tight description of faces in torus triangulations with minimum degree 5*, Sib. Elektron. Mat. Izv. **18** (2021) 1475–1481.
<https://doi.org/10.33048/semi.2021.18.110>
- [10] O.V. Borodin and A.O. Ivanova, *Combinatorial structure of faces in triangulations on surfaces*, Sib. Math. J. **63** (2022) 662–669.
<https://doi.org/10.1134/S0037446622040061>
- [11] O.V. Borodin and A.O. Ivanova, *Another tight description of faces in plane triangulations with minimum degree 4*, Discrete Math. **345(9)** (2022) 112964.
<https://doi.org/10.1016/j.disc.2022.112964>

- [12] O.V. Borodin, A.O. Ivanova and A.V. Kostochka, *Describing faces in plane triangulations*, Discrete Math. **319** (2014) 47–61.
<https://doi.org/10.1016/j.disc.2013.11.021>
- [13] D.W. Cranston and D.B. West, *An introduction to the discharging method via graph coloring*, Discrete Math. **340** (2017) 766–793.
<https://doi.org/10.1016/j.disc.2016.11.022>
- [14] B. Grünbaum, *Polytopal graphs*, in: Studies in Graph Theory, Part II, D.R. Fulkerson (Ed(s)) (MMA Studies in Mathematics **12**, 1975) 201–224.
- [15] M. Hornák and S. Jendrol', *Unavoidable sets of face types for planar maps*, Discuss. Math. Graph Theory **16** (1996) 123–141.
<https://doi.org/10.7151/dmgt.1028>
- [16] S. Jendrol', *Triangles with restricted degrees of their boundary vertices in plane triangulations*, Discrete Math. **196** (1999) 177–196.
[https://doi.org/10.1016/S0012-365X\(98\)00172-1](https://doi.org/10.1016/S0012-365X(98)00172-1)
- [17] S. Jendrol' and H.-J. Voss, *Light subgraphs of graphs embedded in the plane – A survey*, Discrete Math. **313** (2013) 406–421.
<https://doi.org/10.1016/j.disc.2012.11.007>
- [18] A. Kotzig, *From the theory of Euler's polyhedrons*, Mat.-Fyz. Čas. **13** (1963) 20–34, in Russian.
- [19] H. Lebesgue, *Quelques conséquences simples de la formule d'Euler*, J. Math. Pures Appl. **19** (1940) 27–43.
- [20] B. Mohar, R. Škrekovski and H.-J. Voss, *Light subgraphs in planar graphs of minimum degree 4 and edge-degree 9*, J. Graph Theory **44** (2003) 261–295.
<https://doi.org/10.1002/jgt.10144>
- [21] O. Ore and M.D. Plummer, *Cyclic coloration of plane graphs*, in: Recent Progress in Combinatorics, W.T. Tutte (Ed(s)) (Academic Press, New York, 1969) 287–293.

Received 7 June 2023

Revised 26 October 2023

Accepted 27 October 2023

Available online 17 November 2023