

MAXIMISING 1'S THROUGH PROPER LABELLINGS

JULIEN BENSMAIL

Université Côte d'Azur, CNRS, Inria, I3S, France

e-mail: julien.bensmail.phd@gmail.com

Abstract

We investigate graph proper labellings, i.e., assignments of labels to the edges so that no two adjacent vertices are incident to the same sum of labels, with the additional requirement that label 1 must be assigned to as many edges as possible. The study of such objects is motivated by practical concerns, and by connections with other types of proper labellings in which other additional properties (such as minimising the sum of assigned labels, or minimising the use of label 3) must be met. We prove that maximising 1's is a problem on its own, in that it is not equivalent to any of these other labelling problems with optimisation concerns. We then provide labelling tools and techniques for designing proper labellings with many 1's. As a result, we prove that, for several graph classes, it is always possible to design proper labellings where label 1 is assigned to about half the edges.

Keywords: proper labelling, incident sum, label 1, 1-2-3 conjecture.

2020 Mathematics Subject Classification: 05C78; 05C15; 05C85.

1. INTRODUCTION

We deal with a variant of proper labellings of graphs, which are usually defined as follows. Let G be a graph. A k -labelling ℓ of G is an assignment $\ell : E(G) \rightarrow \{1, \dots, k\}$ of labels (from $\{1, \dots, k\}$) to the edges of G . For every vertex v of G , one can then compute the sum $\sigma(v)$ of labels incident to v by ℓ , being formally defined as $\sigma(v) = \sum_{u \in N(v)} \ell(vu)$. If no two adjacent vertices of G have the same sum by ℓ , or, in other words, if $\sigma(u) \neq \sigma(v)$ for every edge $uv \in E(G)$, then we say ℓ is *proper*. We denote by $\chi_S(G)$ the smallest $k \geq 1$, if any, such that G admits proper k -labellings. Actually, greedy labelling arguments can be invoked to prove formally that $\chi_S(G)$ is well defined if and only if G is *nice*, i.e., does not contain K_2 (the complete graph on two vertices) as a connected component.

Proper labellings have been intensively studied in the context of the so-called 1-2-3 Conjecture (stating that $\chi_S(G) \leq 3$ holds for every nice graph G) raised by Karoński, Łuczak, and Thomason [9] in 2004, whose resolution was recently claimed by Keusch [10]. We refer the interested reader to e.g. [2, 4, 11] for an overview of the investigations dedicated to proper labellings (and their variants) to date.

The origins behind these notions trace back to the 1980s, to considerations first raised by Chartrand *et al.* [7]. In the kind of problems they introduced there, a graph G is given, and the goal is to make it somewhat irregular, i.e., so that some of its vertices have pairwise different degrees, by replacing its edges with parallel edges, all the while keeping the resulting multigraph of relatively small size (in terms of number of edges). Such a problem can be formalised and studied through the lens of labellings satisfying certain distinction properties. As an illustration, consider proper labellings. Let G be a nice graph, given together with a proper k -labelling ℓ (for any $k \geq 1$). Let $M(G, \ell)$ be the multigraph obtained from G by considering every edge $e = uv$ of G in turn, and replacing e with $\ell(e) \geq 1$ parallel edges¹ (i.e., all joining u and v). Since ℓ is proper, note that $M(G, \ell)$ must be *locally irregular*, meaning that no two of its adjacent vertices have the same degree. Also, by the way $M(G, \ell)$ was constructed, G and $M(G, \ell)$ have the same structure, as no new adjacencies have been created. Now, in this formalism, the parameter $\chi_S(G)$ means there is a way to obtain such a locally irregular $M(G, \ell)$ from G by replacing every edge by at most $\chi_S(G)$ parallel edges. The 1-2-3 Conjecture, now, implies there is always a locally irregular $M(G, \ell)$ with size at most $3|E(G)|$. This was one of the main motivations from [7] for investigating labellings minimising the maximum value of an assigned label.

As raised in [4], at least in the context of proper labellings, minimising the maximum label value by ℓ is not a guarantee that $M(G, \ell)$ will necessarily be the smallest locally irregular multigraph overlying G . This led the authors to consider proper labellings minimising the sum of assigned labels, which we will introduce more formally in later Section 2. One intuitive way of minimising the label sum by a proper 3-labelling is through minimising the number of assigned 3's, which is a concern considered by other authors [2].

In this work, we introduce and study the sort of dual approach, consisting in maximising 1's by proper labellings. Our formal terminology is as follows. For a nice graph G , a labelling ℓ , and an $x \in \mathbb{N}^*$, we denote by $\text{nb}(\ell, x)$ the number of edges of G that are assigned label x by ℓ . For any $k \geq \chi_S(G)$, we denote by $\text{maxOne}(G, k)$ the maximum value of $\text{nb}(\ell, 1)$ over all proper k -labellings ℓ of G . Last, we denote by $\text{maxOne}(G)$ the maximum value of $\text{maxOne}(G, k)$ for some $k \geq 1$. Note that we always have $0 \leq \text{maxOne}(G) \leq |E(G)|$, where the upper

¹Hence why we consider k -labellings, assigning strictly positive labels.

bound is attained if and only if G is locally irregular. Also, as we will prove later through Theorem 5, there is no $k \geq 1$ such that $\maxOne(G) = \maxOne(G, k)$ for every nice graph G , which justifies to wonder about the more general parameter $\maxOne(G)$ (and not just about $\maxOne(G, 3)$, which could make sense due to the 1-2-3 Conjecture). From the motivation point of view and the connection between graphs G and associated overlying locally irregular multigraphs $M(G, \ell)$, considering proper labellings ℓ assigning many 1's also justifies in that the corresponding locally irregular multigraph $M(G, \ell)$ has many edges with multiplicity 1, and, thus, is even closer to the original graph G .

We organise this work as follows. In Section 2, we start by making more formal some of the connections we mentioned earlier, between proper labellings maximising 1's and other types of proper labellings fulfilling additional requirements. In Section 3, we investigate first properties of the parameter \maxOne , showing, for instance, that it sometimes requires to assign arbitrarily large labels, and that determining this parameter is quite different from determining other related parameters (such as the minimum label sum by a proper labelling). We also raise, in that section, our leading conjecture, Conjecture 10, stating, roughly, that, at worst, we should always be able to design a proper labelling assigning label 1 to about half of the edges, for any nice graph. We then prove this conjecture for several graph classes through Sections 4 and 5, along which the proofs and arguments we employ are of increasing complexity. In particular, in Section 5, we get to exploring both known and new labelling techniques and tools to establish lower bounds on the parameter \maxOne . We conclude with perspectives for further work on the topic in Section 6.

2. RELATED PARAMETERS AND THEIR INTERPLAY

In this section, we recall some other optimisation problems involving proper labellings from previous works. In particular, we make explicit their connections with the problem of maximising 1's, from which we can establish first properties of the parameter \maxOne .

2.1. Proper labellings with minimum number of 3's

In [2], the authors considered notions that are, to some extent, dual to those we introduce in the current work. Namely, they considered the existence of graphs requiring “lots” of 3's in all of their proper 3-labellings (which the 1-2-3 Conjecture is precisely about). Formally, this leads to the parameter $\minThree(G)$, which, for a given nice graph G , is defined as the smallest x such that, for all proper 3-labellings ℓ of G , we have $\text{nb}(\ell, 3) \leq x$.

While the parameters $\text{maxOne}(G)$ and $\text{minThree}(G)$ might seem dual ones at first glance, a fundamental difference lies in the fact that, for the latter parameter, we are focusing on proper 3-labellings only (while, for the former parameter, we are considering proper labellings assigning labels that can be arbitrarily large). One of the reasons why the latter parameter is defined this way is because, when assigning arbitrarily large labels, we can always make sure to avoid assigning label 3 (and, more generally, any given label).

Proposition 1. *Every nice graph admits proper labellings ℓ with $\text{nb}(\ell, 3) = 0$.*

Proof. Assume ℓ is a proper labelling of a graph G such that $\text{nb}(\ell, 3) > 0$. Let uv be an edge of G with $\ell(uv) = 3$. Note that changing the label assigned to uv cannot result in u and v being in conflict² (as $\ell(uv)$ contributes to both $\sigma(u)$ and $\sigma(v)$). In other words, when changing the label assigned to uv , to preserve properness we only need to make sure that u does not get in conflict with its at most $\Delta(G) - 1$ neighbours other than v , and similarly that v does not get in conflict with its at most $\Delta(G) - 1$ neighbours other than u . Thus, there must be a label in $\{1, \dots, 2\Delta(G)\} \setminus \{3\}$ that can be assigned to uv without introducing conflicts. Then we end up with a proper labelling ℓ' of G with $\text{nb}(\ell', 3) = \text{nb}(\ell, 3) - 1$, and by repeating these arguments while G has edges assigned label 3, eventually we get to a situation where G is labelled in a proper way and no edges are assigned label 3. ■

Due to these facts, the real parameter dual (in spirit) to $\text{minThree}(G)$ is actually $\text{maxOne}(G, 3)$, from which we deduce that the following relationship holds.

Observation 2. *If G is a nice graph, then $\text{maxOne}(G, 3) \leq |E(G)| - \text{minThree}(G)$.*

Indeed, by definition, every proper 3-labelling ℓ of a nice graph G must assign label 3 to at least $\text{minThree}(G)$ edges of G . This implies at most $|E(G)| - \text{minThree}(G)$ edges of G can be assigned label 1 or 2 by ℓ , which yields the claimed inequality.

As will be proved in later Theorem 5, there are nice graphs G for which the smallest $k \geq 1$ for which $\text{maxOne}(G) = \text{maxOne}(G, k)$ is arbitrarily large, and thus we can have $k > 3$. Hence, apart from consequences on the very particular case $\text{maxOne}(G, 3)$, we do not get much on $\text{maxOne}(G)$ from results from [2]. We will actually prove in Theorem 8 that maximising 1's and minimising 3's by proper 3-labellings are distant problems.

So that the reader can compare the results from the current work and those from [2], we now give a brief summary of the main contributions of that work. First, the authors proved that there is no absolute constant $x \geq 1$ such that every

²We say there is a *conflict* between two vertices u and v by a labelling if $\sigma(u) = \sigma(v)$.

nice connected graph admits a proper 3-labelling assigning label 3 to at most x edges. This led them to investigating classes of connected graphs requiring a large fraction of their edges be assigned label 3 by any proper 3-labelling. They then exhibited constructions of arbitrarily large connected graphs in which label 3 must be assigned to at least a tenth of the edges, and also provided other constructions, where the edge fraction is sometimes a bit smaller, providing graphs with specific properties (degree constraints, structural properties, *etc.*). These constructions and other arguments led the authors to conjecture that, perhaps, all nice connected graphs admit proper 3-labellings assigning label 3 to at most about a third of the edges, a conjecture they verified for several classes of nice connected graphs (dense enough graphs, graphs with large girth, cubic graphs, cacti, Halin graphs, *etc.*). Last, they also established general properties of the parameter minThree ; in particular, for any given nice connected graph G and fixed $x \geq 1$, it is NP-complete to decide whether $\text{minThree}(G) \leq x$.

2.2. Equitable proper labellings

The reason why maximising 1's and minimising 3's by proper 3-labellings might sound as dual problems, is because labels 1 and 3 seem sort of interchangeable. This is actually true in some contexts, such as in regular graphs, as noticed e.g. in [2]: in a nice regular graph, swapping 1's and 3's by a proper 3-labelling results in another proper 3-labelling.

While this phenomenon is not true in general, designing proper labellings where labels are assigned about the same number of times is caught by the notion of equitable proper labellings, first introduced in [1]. Formally, a proper labelling ℓ of a nice graph G is said *equitable* if $|\text{nb}(\ell, i) - \text{nb}(\ell, j)| \leq 1$ for any two labels i and j assigned by ℓ . We can then define $\overline{\chi_\Sigma}(G)$ as the smallest $k \geq 1$ such that equitable proper k -labellings of G exist.

These notions relate to our problem as follows.

Observation 3. *If a graph G admits an equitable proper k -labelling ℓ where $\text{nb}(\ell, 1) > 0$, then $\text{maxOne}(G, k) \geq \lfloor \frac{1}{k} |E(G)| \rfloor$ and, thus, $\text{maxOne}(G) \geq \lfloor \frac{1}{k} |E(G)| \rfloor$.*

Regarding the parameter maxOne , Observation 3 makes the most sense for graphs admitting equitable proper k -labellings where k is small. The results of [1] actually suggested that an equitable version of the 1-2-3 Conjecture (stating that $\overline{\chi_\Sigma}(G) \leq 3$ holds for most graphs G) might be true, which was formalised in [3]. If proved to hold, that conjecture would thus somewhat imply that $\text{maxOne}(G) \geq \lfloor \frac{1}{3} |E(G)| \rfloor$ holds for most graphs G .

For the record, the authors of [1] proved that $\overline{\chi_\Sigma}(G) \leq 2$ holds for any nice forest G , that $\overline{\chi_\Sigma}(G) = \chi_S(G) \leq 2$ holds for any nice complete bipartite graph G

different from $K_{3,3}$ (for which labels 1, 2, 3 are needed and sufficient in the equitable version), and that $\overline{\chi_\Sigma}(G) = \chi_S(G) = 3$ holds for any nice complete graph G different from K_4 (for which labels 1, 2, 3, 4 are needed and sufficient in the equitable setting). It turns out that, to date, K_4 is the only known connected graph G with $\overline{\chi_\Sigma}(G) \geq 4$. In [2], among other things, the authors established that there exist infinitely many graphs G with $2 = \chi_S(G) < \overline{\chi_\Sigma}(G) = 3$. Actually, deciding whether $\overline{\chi_\Sigma}(G) = 2$ holds for a given graph G with $\chi_S(G) = 2$ is an NP-complete problem. To date, the main open problem regarding the parameter $\overline{\chi_\Sigma}(G)$ is establishing a constant upper bound for any nice graph G . The best upper bound that has been established at this point is $\overline{\chi_\Sigma}(G) \leq |E(G)|$, which follows from investigations on a local version of the Antimagic Labelling Conjecture [5].

2.3. Proper labellings with minimum label sum

In [4], the authors considered a variant of proper labellings where the sum of assigned labels must be as small as possible. Formally, given a nice graph G and a labelling ℓ of G , one can define $\sigma(\ell)$ as the *sum* of the labels assigned by ℓ to the edges of G . Now, for any $k \geq 1$, we define $\text{minLabelSum}(G, k)$ as the smallest value of $\sigma(\ell)$ for a proper k -labelling ℓ of G , and $\text{minLabelSum}(G)$ as the minimum value of $\text{minLabelSum}(G, k)$ for some $k \geq 1$.

As will be proved later in Section 3, while this might seem natural, it turns out that, in general, minimising the label sum by a proper labelling does not necessarily require to maximise 1's, and *vice versa*. Still, there are contexts in which these two problems are equivalent, namely when very few different labels are assigned.

Observation 4. *For proper 2-labellings, minimising the label sum is equivalent to maximising 1's. Hence, for any nice graph G and proper 2-labelling ℓ with $\text{nb}(\ell, 1) + 2 \cdot \text{nb}(\ell, 2) = \text{minLabelSum}(G, 2)$, we have $\text{maxOne}(G, 2) = \text{nb}(\ell, 1)$ and, thus, $\text{maxOne}(G) \geq \text{nb}(\ell, 1)$.*

The main results from [4] deal mainly with a conjecture the authors raised, stating that we should have $\text{minLabelSum}(G) \leq 2|E(G)|$ for any nice connected graph G . Note that this would hold if one proved that any such G always admits a proper 3-labelling where the number of assigned 1's is always more than the number of assigned 3's, which shows that all variants we have mentioned up to this point are sort of all connected.

The authors of [4] determined the parameter minLabelSum for several graph classes, including nice complete bipartite graphs, nice complete graphs, and nice cycles. They also proved that determining $\text{minLabelSum}(G, 2)$ for a given graph G is NP-complete, which, due to Observation 4, implies that determining $\text{maxOne}(G, 2)$ for a given graph G is also NP-complete. Likewise, they proved that the same problem can be solved in polynomial time when G has bounded

treewidth, and this yields a similar result in our context. They also proved that, in general, the smallest $k \geq 1$ such that $\text{minLabelSum}(G, k) = \text{minLabelSum}(G)$ is not bounded by a constant, in the sense that, in order to minimise the label sum, it is sometimes necessary to assign arbitrarily large labels. Finally, they provided some upper bounds on $\text{minLabelSum}(G)$ for G lying in certain graph classes (nice connected bipartite graphs, nice trees, nice connected graphs with bounded chromatic number, *etc.*). In particular, they proved that every nice tree admits a proper 2-labelling where the number of assigned 1's is at least the number of assigned 2's; Observation 4 thus implies that we have $\text{maxOne}(T) \geq \text{maxOne}(T, 2) \geq \lceil \frac{1}{2}|E(T)| \rceil$ for every nice tree T .

3. GENERAL BEHAVIOUR OF maxOne

In this section, we give hints on the general behaviour of the parameter maxOne . In particular, we prove that maximising 1's by a proper labelling can require to assign arbitrarily large labels. We also prove that maximising 1's is not equivalent to one of the other related labelling problems we mentioned earlier (minimising the label sum, and minimising 3's in proper 3-labellings). All this leads us to raising a general conjecture, Conjecture 10, on the number of 1's we can assign in all nice graphs, by a proper labelling.

3.1. On the possible magnitude of maxOne

We start off by remarking that, in some contexts, maximising the number of assigned 1's might require to design proper k -labellings with k arbitrarily large.

Theorem 5. *For arbitrarily large values of k , there exist nice connected graphs G such that $\text{maxOne}(G) = \text{maxOne}(G, k)$, and $\text{maxOne}(G, i) < \text{maxOne}(G, k)$ for every $i < k$.*

Proof. Consider the following construction (see Figure 1). Start from an edge uv . Now, choose any $\alpha \geq 2$, and add α new vertices x_1, \dots, x_α adjacent to u , as well as $\alpha + 1$ new vertices $y_1, \dots, y_{\alpha+1}$ adjacent to v . Finally, add new degree-1 vertices adjacent to the x_i 's and y_i 's so that $d(x_1) = \alpha + 1$, $d(y_1) = \alpha + 2$, $d(y_2), \dots, d(y_{\alpha+1})$ are $\alpha + 3, \dots, 2\alpha + 2$, and $d(x_2), \dots, d(x_\alpha)$ are $2\alpha + 2, \dots, 3\alpha$. Let G denote the resulting graph (actually a tree).

Note that $d(u) = d(x_1) = \alpha + 1$ and $d(v) = d(y_1) = \alpha + 2$. Thus, G is not locally irregular, and $\text{maxOne}(G) < |E(G)|$ (while $|E(G)| = 1 + \sum_{i=\alpha+1}^{2\alpha+2} i + \sum_{i=2\alpha+2}^{3\alpha} i = 4\alpha^2 + 3\alpha + 3$). We claim that $\text{maxOne}(G) = |E(G)| - 1$, and that, by any proper labelling of G assigning label 1 to all but one edge, the unique label different from 1 must be assigned to uv . Note indeed that if the unique label different from 1 was assigned to an edge incident to one of the x_i 's,




Figure 1. Illustration of the construction in the proof of Theorem 5. Pendant edges incident to the x_i 's and y_i 's are not drawn. The number in each vertex indicates its actual degree in the whole graph.

then we would have $\sigma(v) = \sigma(y_1) = \alpha + 2$ and the labelling would actually not be proper. Likewise, we would have a similar conclusion if the unique label different from 1 was assigned to an edge incident to one of the y_i 's (we would get $\sigma(u) = \sigma(x_1) = \alpha + 1$). Thus, the unique label different from 1 must be assigned to uv .

Remark now that upon assigning some label $i > 1$ to uv and label 1 to all other edges of G , we get $\{\sigma(y_1), \dots, \sigma(y_{\alpha+1})\} = \{\alpha + 2, \dots, 2\alpha + 2\}$, while $d(v) = \alpha + 2$. Thus, so that we do not get a conflict involving v and one of the y_i 's, we must have $i > \alpha + 1$ so that $\sigma(v) > 2\alpha + 2$. Similarly, we get $\{\sigma(x_2), \dots, \sigma(x_\alpha)\} = \{2\alpha + 2, \dots, 3\alpha\}$. Thus, since $d(u) = \alpha + 1$, so that we do not get a conflict between u and the x_i 's, it must be that i does not lie in $\{\alpha + 2, \dots, 2\alpha\}$. Altogether, we thus deduce that we must have $i > 2\alpha$. The claim now follows from the fact that α can be chosen arbitrarily large. ■

3.2. Maximising 1's versus minimising the label sum

Recall that Theorem 5 is one of the main reasons why the two parameters `maxOne` and `minThree` are hardly comparable in general. The parameter `minLabelSum`, however, is subject to a similar behaviour, and one could thus wonder whether, in general, minimising the label sum by a proper labelling is somewhat equivalent to maximising 1's. While this is true in specific contexts (recall Observation 4), we prove this is not the case in general.

Theorem 6. *There are nice connected graphs for which any proper labelling maximising the number of 1's is arbitrarily far from minimising the label sum.*

Proof. This can be seen through the construction (illustrated in Figure 1) we provided in the proof of Theorem 5. For such a graph G (and some $\alpha \geq 2$),

recall that, by a proper labelling ℓ with $\text{nb}(\ell, 1)$ as large as possible, we have $\text{nb}(\ell, 1) = |E(G)| - 1$ and $\ell(uv) \geq 2\alpha + 1$; thus, such a proper labelling satisfies $\sigma(\ell) \geq |E(G)| + 2\alpha = 4\alpha^2 + 5\alpha + 3$. Meanwhile, it can be noticed that one can construct a proper 2-labelling ℓ' of G with $\sigma(\ell') = |E(G)| + 2 = 4\alpha^2 + 3\alpha + 5$. Indeed, to obtain such an ℓ' , it suffices to assign label 2 to any (pendant) edge incident to x_1 different from x_1u , label 2 to any (pendant) edge incident to y_1 different from y_1v , and label 1 to all other edges. This indeed results in ℓ' being proper, as we get $\sigma(z) = d(z)$ for all $z \in V(G)$ with $d(z) \geq 2$ and $z \notin \{x_1, y_1\}$, and the only pairs of adjacent vertices of G with the same degree are $\{u, x_1\}$ and $\{v, y_1\}$. In particular, recall as well that degree-1 vertices cannot be in conflict with their unique neighbour, upon assigning strictly positive labels. The full claim now follows from the fact that α can be chosen arbitrarily large in the construction of G . ■

Regarding the proof of Theorem 6, note that minimising the label sum by a proper labelling of the graphs G we construct can still be achieved through assigning label 1 to all but two edges of G , which is very close to maximising the number of 1's. For completeness, below we prove a similar result for the other direction; namely, we prove that there exist graphs in which minimising the label sum is far from maximising 1's.

Theorem 7. *There are nice connected graphs for which any proper labelling minimising the label sum is arbitrarily far from maximising the number of 1's.*

Proof. Let Q denote the graph (depicted in Figure 1(a)) constructed in the proof of Theorem 5 for $\alpha = 2$. Note that $|E(Q)| = 25$, and, by earlier arguments, we have $\text{minLabelSum}(Q) = \text{minLabelSum}(Q, 2) = |E(Q)| + 2 = 27$. Indeed, a proper labelling of Q assigning label 1 to all edges but one needs to assign a label with value at least 5 to uv , while we can produce a proper 2-labelling of Q that assigns label 2 to only two edges (and label 1 to the rest). Thus, by any proper labelling ℓ of Q minimising the label sum, we have $\text{nb}(\ell, 1) \leq |E(Q)| - 2 = 23$, while $\text{maxOne}(Q) = \text{maxOne}(Q, 5) = |E(Q)| - 1 = 24$.

Now choose any $q \geq 7$, and consider the following graph G with $25q$ edges. Start from q vertex-disjoint copies Q_1, \dots, Q_q of Q . For every $i \in \{1, \dots, q\}$, let z_i denote, in Q_i , any one of the two degree-1 vertices adjacent to x_1 (which, recall, has degree 3). Finally, identify all of z_1, \dots, z_q to a single vertex r to get our G . Note that G is, in some sense, a star with q branches, each of which is actually a copy of Q .

Since $d(r) = q \geq 7$ and all neighbours of r are of degree 3, note that, by any 2-labelling of G , we cannot have a conflict involving r and one of its neighbours. Also, due to its structure, designing a proper labelling of G with specific properties mainly falls down to designing proper labellings with the same properties in

Q_1, \dots, Q_q . In particular, we have

$$\min\text{LabelSum}(G) \leq \min\text{LabelSum}(G, 2) \leq q \cdot \min\text{LabelSum}(Q, 2) = 27q$$

and any proper 2-labelling ℓ with minimum label sum of G verifies $\text{nb}(\ell, 1) \leq 23q$. On the other hand, we have

$$\max\text{One}(G) \geq q \cdot \max\text{One}(Q) = 24q.$$

The desired conclusion then arises from the fact that q can be chosen arbitrarily large. \blacksquare

Remark that the construction used in the proof of Theorem 7 could also be used to prove Theorem 6. An important difference, however, is that, for graphs from the former construction, maximising 1's requires at least the use of label $1, \dots, 5$, while for those from the latter construction this requires only the use of labels 1 and 2. Thus, the two constructions achieve slightly different goals.

3.3. Maximising 1's versus minimising 3's

Due to Theorem 5 and our arguments from Section 2, maximising 1's and minimising 3's can only be compared in the context of proper 3-labellings. Still, we establish that, even in the context of proper 3-labellings, these two problems are sometimes quite different.

Theorem 8. *There are nice connected graphs for which any proper 3-labelling maximising the number of 1's is arbitrarily far from minimising the number of 3's, and vice versa.*

Proof. This can be established through a slight modification of the graphs we considered in the proofs of Theorems 6 and 7. Let Q be the graph with 45 edges depicted in Figure 2. Below, we deal with the vertices and edges of Q through the notation given in that figure.




Figure 2. The graph Q in the proof of Theorem 8. Pendant edges incident to some of the vertices are not drawn; the number in each vertex indicates its actual degree in the whole graph.

Note that Q is not locally irregular, since it has four pairs of adjacent vertices with the same degree (namely, $\{v_1, v_3\}$, $\{v_4, v_5\}$, $\{v_6, v_7\}$, and $\{v_8, v_9\}$). Also, due

to the structure of Q , and due to these pairs, we have $\maxOne(Q) \leq |E(Q)| - 2 = 43$, and, clearly, any proper labelling ℓ of Q assigning label 1 to all but two edges must assign a label different from 1 to v_3v_4 and v_7v_8 . Actually, to avoid any conflict between v_3 and v_1 and v_2 , and similarly between v_8 and v_9 and v_{10} , we must have $\ell(v_3v_4), \ell(v_7v_8) \geq 3$. Thus, the unique proper 3-labelling ℓ of Q maximising the number of assigned 1's assigns label 3 to v_3v_4 and v_7v_8 and label 1 to the rest (it can be checked that this is indeed proper). Thus, we have $\maxOne(Q, 3) = 43$, and this ℓ satisfies $\sigma(\ell) = |E(Q)| + 4 = 49$.

On the other hand, we have $\minLabelSum(Q) \leq \minLabelSum(Q, 2) \leq |E(Q)| + 3 = 48$. Indeed, one can obtain a proper 2-labelling ℓ of Q by assigning label 2 to any pendant edge incident to v_1 , to any pendant edge incident to v_9 , and to v_5v_6 (thus to only three edges), and label 1 to the other edges. Furthermore, this ℓ satisfies $\text{nb}(\ell, 1) = |E(Q)| - 3 = 42$.

Now choose any $q \geq 9$, and, just as in the proof of Theorem 7, consider the graph G with $45q$ edges obtained from q copies Q_1, \dots, Q_q of Q by identifying q degree-1 neighbours of the local copies of v_1 to a single vertex r (so that G is a kind of star with q branches each of which is a copy of Q , and all neighbours of r have degree 3). Once more, note that, by any proper 3-labelling of G , it cannot be that r is involved in a conflict; thus, any proper 3-labelling of G is, essentially, a combination of proper 3-labellings of Q_1, \dots, Q_q . Now, by earlier arguments, we have

$$\minLabelSum(G) \leq \minLabelSum(G, 2) \leq q \cdot \minLabelSum(Q, 2) = 48q,$$

and any proper 2-labelling ℓ with minimum label sum of G verifies $\text{nb}(\ell, 1) \leq 42q$. Meanwhile, we have

$$\maxOne(G) \geq \maxOne(G, 3) \geq q \cdot \maxOne(Q, 3) = 43q,$$

and any proper 3-labelling ℓ of G maximising 1's satisfies $\sigma(\ell) = 49q$. So, by considering large enough values of q , we get that, in G , a proper 3-labelling maximising 1's is arbitrarily far from minimising the label sum, and *vice versa*. We thus have our conclusion. \blacksquare

3.4. A conjecture

Due to the previous results, minimising the label sum and maximising 1's by a proper labelling can then be perceived as rather distant problems, and, in particular, problems we might raise regarding the parameter \maxOne do not have to be necessarily reminiscent of existing ones for the parameter \minLabelSum . In [4], the leading conjecture states that we should have $\minLabelSum(G) \leq 2|E(G)|$ for every nice graph G , formalising the intuition that we should always be able to design proper 3-labellings ℓ of G where $\text{nb}(\ell, 1)$ is about $\text{nb}(\ell, 3)$. Other

questions in [4] are on proper 2-labellings of nice bipartite graphs. That is, it is believed that there is some $c \geq 1$ such that, for every bipartite graph G with $\chi_S(G) \leq 2$, we should have $\text{minLabelSum}(G, 2) \leq \frac{3}{2}|E(G)| + c$, or, in other words, that we should always be able to assign labels 1 and 2 to about as many edges. An interesting point lies in the fact that those concerns involve assigning different labels about the same number of times, which sort of connects to equitable proper labellings as considered in [1].

To guide our upcoming investigations on the parameter maxOne , we first need to wonder about the existence of “bad” graphs for this parameter, namely nice graphs in which, by any proper labelling, we cannot assign label 1 to too many edges. Examining small graphs leads to considering K_3 , the complete graph on three vertices, which clearly verifies $\text{maxOne}(K_3) = \text{maxOne}(K_3, 3) = 1 = \frac{1}{3}|E(K_3)|$. This example is very particular, however, due to its very limited size ($\text{maxOne}(K_3)$ can also be written as $\lfloor \frac{1}{2}|E(K_3)| \rfloor$) and structure. Many other small graphs G actually verify $\text{maxOne}(G) = \frac{1}{2}|E(G)|$, which turns out to be a general phenomenon, as exposed by the next result.

Theorem 9. *There are arbitrarily large nice graphs G with $\text{maxOne}(G) \leq \frac{1}{2}|E(G)|$.*

Proof. Let us start with a straight observation. Let H be a graph containing a path $P = uv_1 \dots v_p w$ for some $p \equiv 3 \pmod{4}$ (thus with length 0 modulo 4), where the v_i ’s all have degree 2 in H . For any two adjacent vertices $v_i v_{i+1}$ of P , note that, by any proper labelling of H , so that $\sigma(v_i) \neq \sigma(v_{i+1})$, it must be that the two edges incident to v_i and v_{i+1} being at distance 2 (thus different from $v_i v_{i+1}$) get assigned distinct labels. In particular, if some edge of P is assigned label 1, then there is another edge of P at distance 2 that cannot be assigned label 1. Due to the fact that the length of P is a multiple of 4, we thus deduce that, by any proper labelling of H , at most half the edges of P can be assigned label 1.

Back to proving Theorem 9, consider any graph H , and let G be the (nice) graph obtained from H by considering every edge e of H in turn, and subdividing e $n_e \geq 3$ times, for some $n_e \equiv 3 \pmod{4}$. As a result, note that, in G , every edge e of H gets transformed into a path P_e whose length $n_e + 1$ is a multiple of 4. By the previous observation, by any proper labelling of G , it must be that, in any such path P_e , at most half the edges are assigned label 1, while the P_e ’s partition the edges of G , and G has even size. Thus, we have $\text{maxOne}(G) \leq \frac{1}{2}|E(G)|$, and the claim follows from the fact that the n_e ’s can be chosen arbitrarily large, as long as they are congruent to 3 modulo 4. ■

Note that Theorem 9 also adapts to various graph classes and properties, provided they comply with the subdivision operation we use. In particular, the result also holds for nice graphs with bounded maximum degree or arbitrarily large girth, for nice planar graphs, *etc.* Also, while subdividing all edges of H as described in the proof of Theorem 9 always results in G being bipartite, we

can obtain a very close result for nice non-bipartite graphs by considering a non-bipartite graph as H , and subdividing (as described above) all its edges but any one edge f belonging to an odd-length cycle. Doing so, as a result we get a G that has odd-length cycles (all going through f), such that $\maxOne(G)$ is, at best, just above $\frac{1}{2}|E(G)|$. Thus, Theorem 9 is not specific to nice bipartite graphs.

For these reasons, we believe the following might hold.

Conjecture 10. *If G is a nice connected graph, then*

$$\maxOne(G) \geq \left\lceil \frac{1}{2}|E(G)| \right\rceil - 1.$$

Again, Conjecture 10, if true, would probably not bring anything new regarding the main conjectures from [4] on the parameter \minLabelSum , since, as emphasised earlier, maximising 1's by a proper labelling can require to employ arbitrarily large labels, which might result in large label sum, or, to the least, larger than the best we can hope for. It is worth pointing out also that the bound in Conjecture 10 is expressed so that it fits with some of the results (Theorem 12, in particular) to be established in upcoming Section 4.

4. EASY CLASSES OF GRAPHS

In this section, we investigate the parameter \maxOne for easy graph classes, in which the structure is so predictable that the exact value can be determined through straight *ad hoc* arguments. More precisely, we consider nice paths, nice cycles, and nice complete bipartite graphs. In each case, Conjecture 10 is proved to hold. For transparency, let us mention that, here, the proof arguments we employ are close to arguments used in [4] to determine \minLabelSum for these graphs, the context being very close to that of Observation 4. It is also worth mentioning that, for these classes of graphs we consider, the proper labellings we design actually use labels $1, \dots, \chi_S(G)$ for a graph G . Thus, in other words, Theorem 5 would not restrict to these classes of graphs.

Theorem 11. *If P is a nice path, then*

- $\maxOne(P) = \maxOne(P, 1) = |E(P)|$ if $|E(P)| = 2$;
- $\maxOne(P) = \maxOne(P, 2) = \frac{1}{2}|E(P)|$ if $|E(P)| \equiv 0 \pmod{4}$;
- $\maxOne(P) = \maxOne(P, 2) = \frac{1}{2}|E(P)| + 1$ if $|E(P)| \equiv 2 \pmod{4}$ and $|E(P)| \geq 6$; and
- $\maxOne(P) = \maxOne(P, 2) = \left\lceil \frac{1}{2}|E(P)| \right\rceil$ otherwise.

Proof. If $|E(P)| = 2$, then P is locally irregular, and we end up with a proper labelling when assigning label 1 to all edges. This proves the first item.

Regarding the other items, note that, regardless of the length of P , if v_i, v_{i+1}, v_{i+2} , and v_{i+3} are four consecutive vertices, then, in order to have $\sigma(v_{i+1}) \neq \sigma(v_{i+2})$ by a labelling ℓ , we must have $\ell(v_i v_{i+1}) \neq \ell(v_{i+2} v_{i+3})$. Recall also that, by any labelling assigning strictly positive labels, a degree-1 vertex cannot be in conflict with its unique neighbour. From these arguments, it is easy to see that, in order to assign the most 1's by a proper labelling of P , one should assign 1's to pairs of subsequent edges, so that the pairs of subsequent edges preceding and succeeding it are not assigned label 1. More formally, if we denote by e_1, \dots, e_m the consecutive edges of P , then, assuming we assign label 1 to a pair $\{e_i, e_{i+1}\}$ of subsequent edges, the pairs $\{e_{i-2}, e_{i-1}\}$ and $\{e_{i+2}, e_{i+3}\}$ must be assigned other labels.

From these arguments, we deduce that, to assign label 1 to the most edges of P , one can assign label 1 to the edges in the pairs $\{e_1, e_2\}, \{e_5, e_6\}, \{e_9, e_{10}\}$, and so on (where, naturally, depending on the length of P , the last pair might contain only one edge). By the arguments above, the other edges, not part of these pairs, can be assigned e.g. label 2. In all cases, we end up with a proper 2-labelling, since no two edges at distance 2 are assigned the same label. Also, this labelling maximises the number of 1's by the previous arguments (as assigning more 1's would imply two edges at distance 2 are assigned label 1), and it can be checked that we have the claimed equalities, depending on $|E(P)|$. ■

Theorem 12. *If C is a nice cycle, then*

- $\maxOne(C) = \maxOne(C, 2) = \frac{1}{2}|E(C)|$ if $|E(C)| \equiv 0 \pmod{4}$;
- $\maxOne(C) = \maxOne(C, 3) = \frac{1}{2}|E(C)| - 1$ if $|E(C)| \equiv 2 \pmod{4}$;
- $\maxOne(C) = \maxOne(C, 3) = \lfloor \frac{1}{2}|E(C)| \rfloor$ otherwise.

Proof. This follows from similar reasons as for paths in the proof of Theorem 11. In particular, in any proper labelling of C , two edges at distance 2 must always be assigned distinct labels. Denoting by e_1, \dots, e_m the consecutive edges of C , finding a proper k -labelling of C is then equivalent to finding a proper k -vertex-colouring of G , the graph obtained from C by adding a vertex v_{e_i} for every edge e_i of C , and adding an edge joining two vertices v_{e_i} and v_{e_j} if and only if e_i and e_j are at distance 2 in C . Moreover, assigning the most 1's by a proper labelling of C is then equivalent to finding a proper vertex-colouring of G where a given one of the colours is assigned to the most vertices.

Observe, now, that G is either the disjoint union of two paths of length 1 (when $|E(C)| = 4$), the disjoint union of two even-length cycles (when $|E(C)|$ is a multiple of 4 at least 8), the disjoint union of two odd-length cycles (when $|E(C)| \equiv 2 \pmod{4}$), or an odd-length cycle (otherwise, when $|E(C)|$ is odd). The cardinality of the maximum independent sets in such easy structures are easy to deduce, and it can be noted that they lead to the equalities from the statement.

Also, any such maximum independent set can be converted to a proper 2-vertex-colouring or proper 3-vertex-colouring of G (depending on the length of C), which, in turn, converts to a proper 2-labelling or proper 3-labelling, respectively, of C , that maximises the number of 1's. For the record, it is indeed known that $\chi_S(C) = 2$ if $|E(C)| \equiv 0 \pmod{4}$, while we have $\chi_S(C) = 3$ otherwise (see e.g. [6]). ■

Theorem 13. *If $K_{n,m}$ is a nice complete bipartite graph with parts of size n and m , then*

- $\maxOne(K_{n,m}) = \maxOne(K_{n,m}, 1) = |E(K_{n,m})|$ if $n \neq m$; and
- $\maxOne(K_{n,m}) = \maxOne(K_{n,m}, 2) = |E(K_{n,m})| - \sqrt{|E(K_{n,m})|}$ otherwise.

Proof. If $n \neq m$, then $K_{n,m}$ is locally irregular, and the claim follows from the fact that a proper labelling results when assigning label 1 to all edges. Now, assume $n = m$ (with $n, m \geq 2$ as otherwise $K_{n,m}$ would not be nice). Since all vertices of $K_{n,m}$ have the same degree, so that we do not have a conflict between two (adjacent) vertices from different parts, it must be that at least n edges are not assigned label 1 (as, otherwise, we would deduce that, in both parts, there is a vertex with sum n). Thus, $\maxOne(K_{n,m}) \leq |E(K_{n,m})| - n$ (while $|E(K_{n,m})| = n^2$). Now, by choosing any vertex v of $K_{n,m}$, assigning label 2 to all edges incident to v , and label 1 to all other edges, we clearly obtain a proper labelling ℓ . Furthermore, $\text{nb}(\ell, 1) = |E(K_{n,m})| - n$, and the claimed equality thus holds. ■

5. GENERAL TECHNIQUES AND OTHER GRAPH CLASSES

We here come up with general tools and techniques for designing proper labellings assigning many 1's, which, depending on the graphs considered, can be of varying efficiency. Applying these, we get to determining the exact value of \maxOne for a few other graph classes, such as nice complete graphs and more generally dense enough nice graphs.

5.1. Special subgraphs

Let G be a graph. For a subset $F \subseteq E(G)$ of edges of G , we say that the corresponding subgraph H (i.e., with edge set F) of G is *special* if it fulfils the following properties.

- (A) If two of its isolated vertices, say u and v , are adjacent in G , then $d_G(u) \neq d_G(v)$;
- (B) no connected component of H with only one edge uv satisfies $d_G(u) = d_G(v)$.

The main idea behind special subgraphs is that if H is a special subgraph over edge set F of some graph G , then we can design a proper labelling of G assigning label 1 to all edges in $E(G) \setminus F$. This is due to the following lemma.

Lemma 14. *Let G be a nice graph, and H be a special subgraph of G defined over some edge set $F \subseteq E(G)$. Assume ℓ is a partial labelling of G where only the edges of $E(G) \setminus F$ are labelled, all with label 1. Then, it is possible to extend ℓ to a proper labelling of G by assigning labels to the edges of F (and preserving the labels of the edges in $E(G) \setminus F$).*

Proof. For every vertex v of G , let us denote by $s(v)$ the number of its incident edges in $E(G) \setminus F$. Then, by ℓ , we currently have $\sigma(v) = s(v)$. At this point, note also that only the isolated vertices of H have all their incident edges labelled, and, if u and v are two such adjacent vertices of G , then, by definition, we have $d_G(u) \neq d_G(v)$, and $\sigma(u) \neq \sigma(v)$.

We now explain how to label the edges of F . Start by considering every edge $uv \in F$ that forms a single-edge connected component of H . By definition, recall that $d_G(u) - d_H(u) \neq d_G(v) - d_H(v)$, so $s(u) \neq s(v)$, and, since uv is the only edge incident to u and v that remains to be labelled, note that, when assigning a label to uv , we only need to make sure u does not get in conflict with any of its neighbours other than v , and *vice versa*. By arguments used to prove Proposition 1, we can always find such a label (since we can assign arbitrarily large ones) we can properly assign to uv , to be done with it. We apply the same to every other such isolated edge of H .

It now remains to label all edges of the connected components of H with at least two edges. We treat every such connected component C in the following way. Let T be a spanning tree of C , and r be any vertex of T with degree at least 2 (in T). We start by assigning label 1 to all edges in $E(C) \setminus E(T)$. It now remains to label all edges of T . Note that every vertex of C is incident to an edge of T . To label these edges, we consider them in decreasing order e_m, \dots, e_1 of their distance to r in T ; in other words, we treat them in reverse order as the edges of T are traversed during a BFS algorithm from r .

We first explain how to deal with all edges e_m, \dots, e_3 , before explaining how to deal with e_2 and e_1 , as these edges require some extra care. Assume all edges e_m, \dots, e_{i+1} , for some $i \geq 3$, have been treated (and thus labelled), and now consider e_i . Assume $e_i = uv$, where, in T , u is closer to r than v is. Due to the ordering we have been considering, and because T is a tree, note that, upon labelling e_i , we will completely determine $\sigma(v)$ (while $\sigma(u)$ will be determined later on). Thus, the label assigned to e_i must be chosen so that v does not get in conflict with any other adjacent vertex of G whose all incident edges have already been assigned a label. Since there are at most $\Delta(G) - 1$ such vertices, there is thus a label in $\{1, \dots, \Delta(G)\}$ we can assign to e_i without raising any conflict involving v . Then we are done with e_i .

Assuming now we already dealt with all of e_m, \dots, e_3 , we now deal with e_2 . The point is that, once e_2 is treated, only one edge, e_1 , remains to be labelled, so, when labelling e_2 , we need to make sure that we do not create a conflicting situation that cannot be resolved upon labelling e_1 . Since, recall, we chose r so that $d_T(r) \geq 2$, it must be that e_2 and e_1 are both incident to r . Assume $e_2 = ru$ and $e_1 = rv$, where $u \neq v$. Upon labelling e_2 , note that we completely determine $\sigma(u)$, which must not be in conflict with any of its neighbours other than r in G . Also, note that, upon labelling e_2 , we would not be able to eventually (through labelling e_1) avoid $\sigma(r) = \sigma(v)$ if the resulting partial sums of r and v (i.e., the sums of the labels assigned to their incident edges that have been labelled at this point) were the same. So, upon labelling e_2 , we also need to guarantee r and v do not get the same resulting partial sum. Again, there is at least one label in $\{1, \dots, \Delta(G) + 1\}$ that can be properly assigned to e_2 here, so that we do not run into that issue.

It now remains to label $e_1 = rv$, which will fully determine $\sigma(r)$ and $\sigma(v)$. Recall that, when dealing with e_2 , we made sure that the current partial sums of r and v are different, which means that assigning any label to e_1 cannot result in r and v being in conflict. So, we only need to make sure that r does not get in conflict with its at most $\Delta(G) - 1$ neighbours other than v in G , and similarly for v and its other at most $\Delta(G) - 1$ neighbours in G . Again, by arguments used in the proof of Proposition 1, there is at least one proper label in $\{1, \dots, 2\Delta(G) - 1\}$ we can assign to e_1 , to be done with C . By then treating all other connected components with at least two edges of H this way, we end up with the desired proper labelling of G . ■

Regarding maximising 1's by a proper labelling, due to Lemma 14, we would definitely be more interested in graphs admitting special subgraphs over the least edges possible (so that the number of other edges, to which we would assign label 1, is maximised). Consequently, for any graph G , we define $\text{MSS}(G)$ as the number of edges of the largest special subgraph of G . The following now clearly holds.

Corollary 15. *For every nice graph G , we have $\text{maxOne}(G) \geq |E(G)| - \text{MSS}(G)$.*

Note that the parameter $\text{MSS}(G)$ is well defined for every nice graph G . In particular, it can be noted that any spanning tree of a nice connected graph forms a special subgraph. Finding the best special subgraph of a given nice graph, however, does not seem as an easy task in general, as, recall, we are also allowed to have isolated vertices in the subgraph (as long as they are not adjacent and have the same degree in the original graph), as well as isolated edges (as long as their two ends have different degrees in the original graph).

This being said, the fact that any spanning tree of a nice connected graph G forms a special subgraph implies the following side result.

Theorem 16. *If G is a nice connected graph, then $\maxOne(G) \geq |E(G)| - (|V(G)| - 1)$.*

Corollary 17. *Nice connected graphs G with $|E(G)| \geq 2(|V(G)| - 1)$ satisfy Conjecture 10.*

In particular, Corollary 17 implies Conjecture 10 for sufficiently dense nice graphs, such as graphs with minimum degree at least 4.

For specific graph classes, special subgraphs are actually crucial to understand the parameter \maxOne . We illustrate this through nice complete graphs, proving that, K_3 apart, they all comply with Conjecture 10. We indeed consider complete graphs on at least four vertices, as K_3 is a cycle and was thus treated earlier through Theorem 12.

Theorem 18. *If K_n is a complete graph with $n \geq 4$, then*

- $\maxOne(K_n) = |E(K_n)| - \frac{2}{3}(n - 1)$ if $n \equiv 1 \pmod{3}$;
- $\maxOne(K_n) = |E(K_n)| - (\frac{2}{3}(n - 2) + 1)$ if $n \equiv 2 \pmod{3}$; and
- $\maxOne(K_n) = |E(K_n)| - (\frac{2}{3}(n - 3) + 2)$ otherwise.

Proof. We apply the approach developed through Lemma 14 and Corollary 15. In the case of a complete graph K_n , note that a special subgraph can contain at most one isolated vertex, and that none of its connected components can contain a single edge. Furthermore, it is not too hard to see that, so that a special subgraph of a complete graph has the least edges, there must be exactly one isolated vertex, and all other vertices should, as much as possible, be part of connected components containing two edges only.

From these arguments, consider the subgraph H with edge set F of any complete graph K_n (with $n \geq 4$) being obtained as follows, where v_1, \dots, v_n denote the vertices of K_n .

- If $n \equiv 1 \pmod{3}$, then add v_2v_3 and v_3v_4 to F , then v_5v_6 and v_6v_7 , then v_7v_8 and v_8v_9 , and so on up to $v_{n-2}v_{n-1}$ and $v_{n-1}v_n$. Note that v_1 is the only isolated vertex of H , and that all other vertices are part of connected components being paths of length 2.
- If $n \equiv 2 \pmod{3}$, then perform the same as in the previous case up to adding $v_{n-3}v_{n-2}$ and $v_{n-2}v_{n-1}$ to F , and additionally add $v_{n-2}v_n$ to F . The difference with the previous case is that, here, one of the connected components of H , that containing v_n , is a star with three leaves.
- If $n \equiv 0 \pmod{3}$, then perform the same as previously up to adding $v_{n-4}v_{n-3}$ and $v_{n-3}v_{n-2}$ to F , and additionally add $v_{n-3}v_{n-1}$ and $v_{n-3}v_n$ to F . Here, one of the connected components of H , that containing v_{n-1} and v_n , is a star with four leaves.

It is important to mention that these procedures can indeed be achieved properly, since $n \geq 4$. Also, it is not too hard to see that $|F| = \text{MSS}(K_n)$, while $|F| = \frac{2}{3}(n-1)$ when $n \equiv 1 \pmod{3}$, $|F| = \frac{2}{3}(n-2) + 1$ when $n \equiv 2 \pmod{3}$, and $|F| = \frac{2}{3}(n-3) + 2$ when $n \equiv 0 \pmod{3}$. By Corollary 15, $\text{maxOne}(K_n)$ is thus at least the claimed values.

Towards a contradiction, assume now that for, say, some $n \equiv 1 \pmod{3}$ (the other cases can be treated similarly), there is a proper labelling assigning label 1 to more edges, that is, to all but at most $\frac{2}{3}(n-1) - 1$ edges. Denote by F the set of these edges. As mentioned earlier, by this labelling there can be at most one isolated vertex, while the other vertices must all be incident to at least one edge of F . Besides, for similar reasons as before, in $K_n[F]$, none of the connected components can contain only one edge. Now, again, the most efficient way for all vertices but at most one to be covered by such connected components in $K_n[F]$, is to have these connected components to be paths of length 2. Thus, we need $|F| \geq \frac{2}{3}(n-1)$, which is a contradiction, since $|F| \leq \frac{2}{3}(n-1) - 1$. ■

Let us emphasise that Theorem 18 deals with the general version of the parameter maxOne , while it is known, see e.g. [6], that $\chi_S(K_n) = 3$ for every $n \geq 3$. Thus, one could more specifically wonder about $\text{maxOne}(K_n, 3)$. We suspect this parameter might behave similarly as $\text{minLabelSum}(K_n)$, which is actually $\text{minLabelSum}(K_n, 3)$, as proved in [4].

It is also worth emphasising that deducing bounds on the parameter maxOne through special subgraphs makes more sense for somewhat dense nice graphs. In particular, in the case of sparse nice graphs, such as nice trees, this method is less likely to be efficient.

5.2. Irregularators

In the approach developed in the previous section, involving special subgraphs, the main point was to narrow the distinguishing part of designing a proper labelling down to the edges of a small subgraph, so that all other edges of the whole graph get sort of useless and can thus be assigned label 1 freely. We here explore a sort of opposite idea, relying, exploiting the structure of the graph, on assigning label 1 to the most edges possible so that the distinguishing part is satisfied for the most pairs of adjacent vertices possible, and then the remaining edges are assigned labels, so that a proper labelling results.

These ideas can be implemented through the concept of irregularators, introduced in [8]. Recall that a locally irregular graph is a graph in which adjacent vertices have different degrees. Now, for a graph G , a set $S \subset V(G)$ of vertices is called an *irregularator* if $G - S$ is locally irregular. In [8], the authors investigated the problem of determining the smallest irregularator of a given graph, which parameter we denote by $\text{Irr}(G)$ for a given graph G . Note that determining $\text{Irr}(G)$

is equivalent to finding the largest locally irregular induced subgraph of G . The authors of [8] mainly investigated the general complexity of determining $\text{Irr}(G)$ for a given graph G , establishing, on the negative side, NP-hardness results and non-approximation results on the problem. On the positive side, they proved that $\text{Irr}(G)$ can be determined in polynomial time provided G is a path, a cycle, a tree, a complete bipartite graph, or a complete graph, and provided FPT results.

Following the aforementioned ideas, we now establish a connection between finding small irregularators and maximising 1's by a proper labelling.

Theorem 19. *If a nice graph G admits an irregularator S , then $\text{maxOne}(G) \geq |E(G - S)|$.*

Proof. Set $H = G - S$. We design a labelling ℓ of G as follows. We start by assigning label 1 to all edges in $E(H)$. At this point, for any two adjacent vertices u and v of H , since H is locally irregular we have $\sigma(u) \neq \sigma(v)$. Along what follows, this will be preserved upon labelling the other edges of G (joining two vertices of S , or one of S and one of H).

We first consider all vertices of S that are not adjacent to any other vertex of S (i.e., isolated vertices of $G[S]$). For every such vertex v , all its incident edges then go to H . We assign labels to these edges so that, whenever labelling any such edge vw (where $w \in V(H)$), we do not introduce any conflict involving w and one of its neighbours. Also, assuming vw_2 and vw_1 are the last two edges incident to v we label (in that order), similarly as in Lemma 14, when labelling vw_2 we also make sure v does not get the same partial sum as w_1 , to guarantee that v and w_1 will not be in conflict when labelling vw_1 afterwards. In case v was actually incident to only one edge, then recall that, whatever label we assign to this edge, v cannot get in conflict with its unique neighbour, since G is nice.

We now consider every edge uv where $u, v \in S$ and $d_S(u) = d_S(v) = 1$. We start by assigning any label to uv , chosen so that if u is incident to only one edge going to H then u does not get the same partial sum as its unique neighbour in H , and similarly regarding v . We then label the edges incident to u and v going to H (such exist since G is nice) similarly as in the previous case, taking into account, when labelling the last edge, that u and v must get distinct sums. In particular, recall that if one of u and v is not incident to any edge going to H (since G is nice, only one of these two vertices can actually satisfy this), then its degree in G is 1, and it cannot be in conflict with its unique neighbour, whatever label be assigned to uv .

It now remains to label the edges of the connected components of $G[S]$ that contain at least two edges. This can be done in a very similar way as what we did in the proof of Lemma 14. Let C be a connected component of $G[S]$ with at least two edges, and let T be a spanning tree of C rooted at some vertex r with degree at least 2 (in T). We first assign label, say, 1 to the edges of $E(C) \setminus E(T)$.

Next we consider the vertices of T in reverse order of their distance to r in T . Whenever considering a new vertex v this way, this means there remains only one incident edge vu in T to be labelled (where u is closer to r than v is), together, possibly, with edges incident to v going to H . We first assign labels to the edges incident to v going to H so that no conflicts involving the neighbours of v arise (which can be done as previously). Last, we label vu , with a label chosen so that the sum of v does not get the same as that of one of its neighbours in H , or of one of its neighbours in T we considered earlier. We are then done with v , and the process can go on.

Just like in the proof of Lemma 14, we also need to pay extra attention when labelling the penultimate edge of C , as the label assigned to that edge must guarantee the two ends of the last edge of C to be labelled do not get in conflict.

Altogether, this whole labelling process results in a proper labelling of G where all edges of H are assigned label 1. The claimed bound then follows. ■

There are cases where the bound in Theorem 19 is rather bad, as illustrated by complete graphs, since, clearly, any nice complete graph K_n satisfies $\text{Irr}(K_n) = n - 1$. Yet, this case is extreme, and, in other cases, such as when local places of a nice graph are locally irregular, it might be that the approach behind Theorem 19 is more promising than that involving special subgraphs and Corollary 15 (which does not take into consideration that large local places can just be assigned 1's right away by a proper labelling).

On a different note, while finding, in a given graph, a largest induced subgraph satisfying some property is a classical problem, [8] did not report concrete applications or problems where finding small irregularities would be useful. To the best of our knowledge, our use of these notions is thus one of its very first applications.

5.3. Modulo methods

We now discuss the use of modulo methods, which have been very classical tools for designing proper labellings with specific properties (see e.g. [2, 4]). The main idea here is to design proper labellings that allow to distinguish adjacent vertices w.r.t. their sums modulo some fixed value (such as the graph's chromatic number). Such labellings can e.g. be attained by switching labels along paths with particular length.

These ideas lead to the following result, for which we provide an abridged proof (as a proof with full details would follow the exact same lines as in [2, 4]).

Theorem 20. *If G is a (nice) connected non-bipartite k -colourable graph for some odd $k \geq 3$, then*

$$\max\text{One}(G) \geq \max\text{One}(G, k) \geq |E(G)| - |V(G)|.$$

Proof. Since G is not bipartite, it contains an odd-length cycle C . We extend C to a particular subgraph H of G , in the following way. We start from $H = C$. Then, as long as there is a vertex v of G that does not belong to H , we add to H the edges of a shortest path from v to some vertex of H . Once all vertices of G are spanned by H , note that H is a unicyclic spanning subgraph of G , where the unique cycle, C , has odd length.

We start building ℓ , a proper labelling of G , by assigning label 1 to all edges of $E(G) \setminus E(H)$. It now remains to label the edges of H . To that end, we consider a proper $\{0, \dots, k-1\}$ -vertex colouring ϕ (where $k \geq 3$ is odd) of G , and label the edges of H so that, eventually, we have $\sigma(v) \equiv \phi(v) \pmod k$ for every vertex $v \in V(G)$. For convenience, and because we are considering sums modulo k , in what follows we assign labels in $\{0, \dots, k-1\}$ to the edges of H , but label 0 can equivalently be regarded as label k .

Start from all edges of H being assigned label, say, 1. Now consider any vertex v of G for which $\sigma(v) \not\equiv \phi(v) \pmod k$ (when taking into account the 1's assigned to the edges of $E(G) \setminus E(H)$), and consider an odd-length closed walk W starting and ending at v , and traversing edges of H only. Such a walk exists because of the presence of C in H . Now, traverse the consecutive edges of W , starting from v and going all the way back to v , and, as going along, apply (modulo k) $+1$ and -1 alternatively to the labels of the traversed edges. That is, we apply $+1$ to the first edge, -1 to the second one, $+1$ to the third one, -1 to the fourth one, and so on. As a result, it can be checked that only the sum of v is altered modulo k , and, due to the length of W , it is actually altered by $+2$. Since k is odd, we can repeat this process, if necessary, until we get $\sigma(v) \equiv \phi(v) \pmod k$ as desired.

Since we can process any other faulty vertex of G this way, eventually the resulting labelling ℓ of G is proper, and it satisfies that all edges of $E(G) \setminus E(H)$ are assigned label 1. Since H is unicyclic, we have $|E(H)| = |V(H)| = |V(G)|$, and the bound thus follows. ■

Note that, in the proof of Theorem 20, it is important that the graph is k -colourable for some $k \geq 3$ odd. As a corollary, we thus derive the following.

Corollary 21. *If G is a (nice) connected k -chromatic graph with $k \geq 3$, then*

- $\maxOne(G) \geq \maxOne(G, k) \geq |E(G)| - |V(G)|$ if k is odd; and
- $\maxOne(G) \geq \maxOne(G, k+1) \geq |E(G)| - |V(G)|$ otherwise.

Corollary 21 might seem weaker than previous Theorem 16. It is worth to emphasise, however, that the proper labellings we design behind Theorem 16 for a nice graph G employ large labels (with maximum value around $2\Delta(G)$ or so), while, behind Corollary 21, the labels used have lower value in general (at

most around $\chi(G)$ or so). Thus, Corollary 21 is somewhat better than Theorem 16 regarding the side question of determining the smallest $k \geq 1$ such that $\maxOne(G) = \maxOne(G, k)$ for a given nice graph G .

5.4. Other methods

Note that using modulo methods, as described in Subsection 5.3, actually stands as a particular application of special subgraphs. It actually turns out that other well known graph objects can be regarded as special subgraphs, so the approach we introduced in Subsection 5.1 can also be related to other classical graph notions.

For instance, in light of Conjecture 10, another approach one could consider, is exploiting maximum cuts to design proper labellings assigning many 1's. Recall that, for a graph G , a *cut* $C = (U, V)$ is a bipartition of $V(G)$, and that the *size* of C is the number of edges of G across the cut (i.e., joining a vertex in U and one in V). It is a well known fact that any *maximum cut* of G , i.e., with maximum size, contains at least half the edges of G . Regarding the parameter \maxOne , one could thus imagine that, perhaps, in some contexts, a proper labelling could be obtained through assigning label 1 to all edges of a maximum cut, and resolving all conflicts through labelling all remaining edges. A problem with this approach, however, is that, through a maximum cut (U, V) of a graph G , it might be that $G[U]$ and $G[V]$ contain isolated vertices (an extreme case being when G is bipartite), while the isolated vertices of $G[U]$ and $G[V]$ induce some structure in G , and, thus, making sure these vertices are not in conflict by some labelling requires to carefully choose the labels assigned to the edges across the cut. In other words, the fact that a labelling of G is proper does not rely solely on the labels assigned in $G[U]$ and $G[V]$.

Another illustration of a case where special subgraphs arise, is for traceable graphs. Recall that a graph G is *traceable* if it contains a Hamiltonian path, i.e., a spanning path. It should be clear that any Hamiltonian path of a nice graph stands as a special subgraph, so, again, Corollary 15 would pop up naturally in such a context. More generally speaking, the same would apply for (nice) graphs having a *2-factor*, i.e., a collection of spanning vertex-disjoint cycles, or more generally a *k-factor* for any $k \geq 3$, being a spanning k -regular subgraph. The same does not necessarily hold for 1-factors, which are exactly *perfect matchings*, since isolated edges in special subgraphs are somewhat tricky to deal with. Still, there are contexts in which we can take advantage of perfect matchings, for instance in (nice) graphs admitting two edge-disjoint perfect matchings, which, together, yield a special subgraph. As an even more special case, we deduce that, for any $k \geq 3$, any (nice) k -regular graph G of class 1 (i.e., with chromatic index k , or, in other words, a partition of the edge set into k perfect matchings) satisfies $\maxOne(G) \geq |E(G)| - |V(G)|$.

6. CONCLUSION

In this work, we have initiated the study of the parameter maxOne , being motivated both by practical concerns (recall the problem of turning a graph into a locally irregular multigraph), and by the fact that this parameter relates to other problems and notions involving proper labellings. The main results we have provided are two-fold, as we both proved that determining $\text{maxOne}(G)$ for a given graph G is a problem on its own (in the sense that it has nothing to do with determining other seemingly close parameters), and came up with several tools and arguments to establish bounds on $\text{maxOne}(G)$. Regarding the latter aspect, we were mainly motivated by Conjecture 10, which we proved true for several classes of graphs. Another interesting aspect is that we established connections between the parameter maxOne and other classical notions and parameters of graph theory.

Many appealing questions and problems remain open at this point, and could be subject to further work. In particular, Conjecture 10 is still open, but recall that Corollary 17 states that only very sparse nice graphs remain to be considered. Maybe this could be achieved by considering the *maximum average degree* (mad for short) parameter, where, recall, for a graph G we denote by $\text{mad}(G)$ the maximum density of a subgraph of G . Note indeed that nice graphs not covered by Corollary 17 have mad at most about 4, while we know Conjecture 10 holds for nice graphs with very small mad (trees and cycles). So one could focus on graphs with small mad . One first step could be to consider planar graphs with large girth (i.e., with long shortest cycles) or graphs with bounded maximum degree, which are sparse in general. It might be that this could be achieved through the tools and arguments we imagined in Section 5, but it might be, too, that other ideas are needed. As pointed out in the previous sections, there are contexts where special subgraphs seem like a viable tool to use, while, in other ones, exploiting irregularities seems more promising. It might thus be that combining all our ideas and tools might be the general way to go.

Apart from Conjecture 10, another general question of interest could be, for a given nice graph G , to wonder about the smallest $k \geq 1$ such that $\text{maxOne}(G) = \text{maxOne}(G, k)$. Recall that some of the tools and approaches we proposed actually deal with that very question, such as Theorem 20. A downside is that this smallest k can be arbitrarily large for a given graph, recall Theorem 5. This question is, thus, probably challenging. Another parameter that is natural to consider for a nice graph G is $\text{maxOne}(G, \chi_S(G))$.

Other appealing questions include algorithmic ones. Recall that, due to Observation 4, determining $\text{maxOne}(G, 2)$ for a given graph G is **NP**-complete in general. Actually, for any given graph G and any fixed $k \geq 2$, we are pretty confident that proofs from [4] could be derived to prove that determining $\text{maxOne}(G, k)$

is NP-complete. However, the situation is not clear about the complexity of determining $\maxOne(G)$ for a given graph G , even when G is a tree, as, even for trees, arbitrarily large labels are sometimes needed in order to maximise 1's by a proper labelling (recall the construction from Theorem 5), which property makes it hard to design NP-hardness reductions. This question could also be considered through the following lens. As highlighted throughout Section 5, in many contexts, given a nice graph, we start from a certain set of edges being assigned label 1, and we hope to be able to label the remaining edges so that a proper labelling results. This converts to a general problem, being, for a given graph G that is already partially labelled (with 1's), to extend the labelling to a proper labelling. In view of the approaches we developed, one could naturally wonder about the general complexity of this problem.

All these directions apart, maximising 1's through proper labellings, as mentioned in Section 2, relates to other close problems and notions, such as proper labellings minimising the label sum, minimising 3's by proper 3-labellings, equitable proper labellings, irregularators, *etc.*, to which are related problems that are all interconnected somehow. For all these problems and notions, many aspects remain open, and, due to these interconnections, it might be that progressing on any of them also provides something regarding the others.

REFERENCES

- [1] O. Baudon, M. Pilśniak, J. Przybyło, M. Senhaji, É. Sopena and M. Woźniak, *Equitable neighbour-sum-distinguishing edge and total colourings*, Discrete Appl. Math. **222** (2017) 40–53.
<https://doi.org/10.1016/j.dam.2017.01.031>
- [2] J. Bensmail, F. Fioravantes and F. Mc Inerney, *On the role of 3s for the 1-2-3 Conjecture*, Theoret. Comput. Sci. **892** (2021) 238–257.
<https://doi.org/10.1016/j.tcs.2021.09.023>
- [3] J. Bensmail, F. Fioravantes, F. Mc Inerney and N. Nisse, *Further results on an equitable 1-2-3 Conjecture*, Discrete Appl. Math. **297** (2021) 1–20.
<https://doi.org/10.1016/j.dam.2021.02.037>
- [4] J. Bensmail, F. Fioravantes and N. Nisse, *On proper labellings of graphs with minimum label sum*, Algorithmica **84** (2022) 1030–1063.
<https://doi.org/10.1007/s00453-021-00903-x>
- [5] J. Bensmail, M. Senhaji and K. Szabo Lyngsie, *On a combination of the 1-2-3 Conjecture and the Antimagic Labelling Conjecture*, Discrete Math. Theor. Comput. Sci. **19**(1) (2017) #21.
<https://doi.org/10.23638/DMTCS-19-1-21>
- [6] G.J. Chang, C. Lu, J. Wu and Q. Yu, *Vertex-coloring edge-weightings of graphs*, Taiwanese J. Math. **15** (2011) 1807–1813.

- [7] G. Chartrand, M.S. Jacobson, J. Lehel, O.R. Oellermann, S. Ruiz and F. Saba, *Irregular networks*, Congr. Numer. **64** (1988) 197–210.
- [8] F. Fioravantes, N. Melissinos and T. Triomatis, *Complexity of finding maximum locally irregular induced subgraphs*, in: Proc. 18th Scandinavian Symposium and Workshops on Algorithm Theory (SWAT 2022), A. Czumaj and Q. Xin (Ed(s)), (Dagstuhl Publishing, Germany, 2022) #24.
- [9] M. Karoński, T. Łuczak and A. Thomason, *Edge weights and vertex colours*, J. Combin. Theory Ser. B **91** (2004) 151–157.
<https://doi.org/10.1016/j.jctb.2003.12.001>
- [10] R. Keusch, *A solution to the 1-2-3 Conjecture* (2023).
arXiv:2303.02611
- [11] B. Seamone, *The 1-2-3 Conjecture and related problems: a survey* (2012).
arXiv:1211.5122

Received 5 June 2023

Revised 15 December 2023

Accepted 15 December 2023

Available online 31 January 2024