Discussiones Mathematicae Graph Theory 45 (2025) 483–507 https://doi.org/10.7151/dmgt.2541

# EXTENDING PARTIAL EDGE COLORINGS OF CARTESIAN PRODUCTS OF GRAPHS

## CARL JOHAN CASSELGREN

Department of Mathematics Linköping University SE-581 83 Linköping, Sweden

e-mail: carl.johan.casselgren@liu.se

FIKRE B. PETROS

AND

SAMUEL A. FUFA

Department of Mathematics Addis Ababa University 1176 Addis Ababa, Ethiopia

e-mail: fikre.bogale@aau.edu.et samuel.fufa@aau.edu.et

### Abstract

We consider the problem of extending partial edge colorings of Cartesian products of graphs. More specifically, we suggest the following Evans-type conjecture. If G is a graph where every precoloring of at most k precolored edges can be extended to a proper  $\chi'(G)$ -edge coloring, then every precoloring of at most k + 1 edges of  $G \square K_2$  is extendable to a proper  $(\chi'(G) + 1)$ edge coloring of  $G \square K_2$ . In this paper we verify that this conjecture holds for trees, complete and complete bipartite graphs, as well as for graphs with small maximum degree. We also prove versions of the conjecture for general regular graphs where the precolored edges are required to be independent.

**Keywords:** precoloring extension, edge coloring, Cartesian product, list coloring.

2020 Mathematics Subject Classification: 05C15.

#### 1. INTRODUCTION

An edge precoloring (or partial edge coloring) of a graph G is a proper edge coloring of some subset  $E' \subseteq E(G)$ ; a *t*-edge precoloring (or just *t*-precoloring) is such a coloring with *t* colors. A *t*-precoloring  $\varphi$  is extendable if there is a proper *t*-edge coloring *f* such that  $f(e) = \varphi(e)$  for any edge *e* that is colored under  $\varphi$ ; *f* is called an extension of  $\varphi$ . In general, the problem of deciding whether a given precoloring is extendable is an  $\mathcal{NP}$ -complete problem, already for 3-regular bipartite graphs [12].

Edge precoloring extension problems seem to have been first considered in connection with the problem of completing partial Latin squares and the wellknown Evans' conjecture that every  $n \times n$  partial Latin square with at most n-1non-empty cells is completable to a partial Latin square [11]. By a well-known correspondence, the problem of completing a partial Latin square is equivalent to asking if a partial edge coloring with  $\Delta(G)$  colors of a balanced complete bipartite graph G is extendable to a proper  $\Delta(G)$ -edge coloring, where  $\Delta(G)$ as usual denotes the maximum degree. Evans' conjecture was proved for large n by Häggkvist [15], and in full generality by Andersen and Hilton [2], and, independently, by Smetaniuk [18].

Another early reference on edge precoloring extension is [16], where the authors study the problem from the viewpoint of polyhedral combinatorics. More recently, the problem of extending a precoloring of a matching has been considered in [9]. In particular, it is conjectured that for every graph G, if  $\varphi$  is a precoloring of a matching M in G using  $\Delta(G) + 1$  colors, and any two edges in M are at distance at least 2 from each other, then  $\varphi$  can be extended to a proper  $(\Delta(G) + 1)$ -edge coloring of G; here, by the *distance* between two edges e and e' we mean the number of edges in a shortest path between an endpoint of e and an endpoint of e'; a *distance-t matching* is a matching where any two edges are at distance at least t from each other. In [9], it is proved that this conjecture holds for e.g. bipartite multigraphs and subcubic multigraphs, and in [14] it is proved that a version of the conjecture with the distance increased to 9 holds for general graphs.

Quite recently, with motivation from results on completing partial Latin squares, questions on extending partial edge colorings of d-dimensional hypercubes  $Q_d$  were studied in [8]. Among other things, a characterization of partial colorings with at most d precolored edges that are extendable to proper d-edge colorings of  $Q_d$  is obtained, thereby establishing an analogue for hypercubes of the characterization by Andersen and Hilton [2] of  $n \times n$  partial Latin squares with at most n non-empty cells that are completable to Latin squares. In particular, every partial coloring with at most d-1 colored edges is extendable to a d-edge coloring of  $Q_d$ . This line of investigation was continued in [6, 7] where similar questions are investigated for trees.

Denote by  $G\Box H$  the Cartesian product of the graphs G and H. Motivated by the result on hypercubes [8], which are iterated Cartesian products of  $K_2$  with itself, in this paper we continue the investigation of edge precoloring extension of graphs with a particular focus on Evans-type questions for Cartesian products of graphs. We are particularly interested in the following conjecture, which would be a far-reaching generalization of a main result of [8].

**Conjecture 1.** If G is a graph where every precoloring of at most k edges can be extended to a proper  $\chi'(G)$ -edge coloring, then every precoloring of at most k + 1 edges of  $G \Box K_2$  is extendable to a proper  $(\chi'(G) + 1)$ -edge coloring of  $G \Box K_2$ .

As we shall see, Conjecture 1 becomes false if we replace  $(\chi'(G) + 1)$  with  $\chi'(G \Box K_2)$  by the example of odd cycles.

It it straightforward that if every precoloring with  $\Delta(G)$  colored edges of a connected Class 1 graph G is extendable to a  $\Delta(G)$ -edge coloring, then G is isomorphic to a star  $K_{1,n}$ . If G, on the other hand, is a connected Class 2 graph where any precoloring with at most  $\Delta(G) + 1$  colored edges is extendable to a  $(\Delta(G) + 1)$ -edge coloring, then G is an odd cycle. Therefore, we shall generally only consider precolorings with at most  $\chi'(G) - 1$  colored edges.

In fact, it is easy to show that odd cycles and stars are the only connected graphs with the property that any partial  $\chi'(G)$ -edge coloring is extendable.

**Proposition 2.** Every partial  $\chi'(G)$ -edge coloring of a connected graph G is extendable if and only if G is isomorphic to a star  $K_{1,n}$  or and odd cycle.

In this paper, we verify that Conjecture 1 holds for trees, complete and complete bipartite graphs. Moreover, we prove a version of Conjecture 1 for regular triangle-free graphs where the precolored edges are required to be independent; a version for graphs with triangles is proved as well. Finally, we prove it for graphs with small maximum degree, namely, graphs with maximum degree two and Class 1 graphs with maximum degree 3.

## 2. Cartesian Products with Trees, Complete Graphs and Complete Bipartite Graphs

In this section, we prove that Conjecture 1 holds for trees, complete bipartite graphs and complete graphs. In the following we shall say that an edge e is  $\varphi$ -colored if  $\varphi$  is a (partial) edge coloring and e is colored under  $\varphi$ . A color c appears at a vertex v if some edge incident with v is colored c; otherwise c is missing at v. Since we only consider edge colorings in this paper, we shall usually omit the word "edge" and just refer to a coloring of a graph.

Before we prove our results, let us briefly outline our general proof idea.

**Proof outline.** We consider a  $\Delta(G)$ -precoloring  $\varphi$  of  $G \times K_2$  using colors  $1, \ldots, \Delta(G)$ , where  $G \times K_2$  consists of two copies  $G_1$  and  $G_2$  of G along with a matching M that joins vertices of  $G_1$  and  $G_2$ . Henceforth, two edges are *corresponding* if their endpoints are joined by two edges of M. Similarly, two vertices are *corresponding* if they are joined by an edge of M. In the proofs we shall distinguish between the case when M contains precolored edges, and the case when it does not.

• When M has no precolored edges, then either all precolored edges are in  $G_1$ , or both  $G_1$  and  $G_2$  contains at least one precolored edge. In the latter case, we can extend the restrictions of  $\varphi$  to  $G_1$  and  $G_2$ , respectively, using  $\Delta(G)$  colors, and then color the edges of M by the color  $\Delta(G) + 1$ .

When all precolored edges are in  $G_1$ , then we remove the color from all edges colored by some fixed color appearing on at least one edge, say 1, and then take an extension of the obtained precoloring of  $G_1$  using colors  $2, \ldots, \Delta(G)+1$ . Next, we recolor the edges  $\varphi$ -colored 1 by the color 1, and thereafter color the edges of  $G_2$  correspondingly, meaning that corresponding edges in  $G_1$  and  $G_2$  get the same color. Finally, we color the edges of M by the color in  $\{1, \ldots, \Delta(G)+1\}$ missing at its endpoints.

• When M contains at least one precolored edge, then we generally aim to select corresponding independent edges of  $G_1$  and  $G_2$  (called *selected edges*), each of which is adjacent to as few precolored edges of  $G_1$  and  $G_2$  as possible, but adjacent to exactly one precolored edge of M. Next, we shall in most proofs color the selected edges by the color of the adjacent edges in M, and consider the resulting colorings of  $G_1$  and  $G_2$  (taken together with the restriction of  $\varphi$  to  $G_1$  and  $G_2$ , respectively). If these colorings are extendable using colors  $1, \ldots, \Delta(G)$ , then we take such extensions, recolor the selected edges by the color  $\Delta(G) + 1$ , and thereafter color the edges of M by its original color, or an arbitrary color not appearing at its endpoints, to obtain an extension of  $\varphi$ .

If the colorings of  $G_1$  and  $G_2$  are not extendable, then we employ some different techniques. The details are given in the proofs below.

Now we turn to the proofs of the main results in this section. Let us first consider trees. In [6], the following was proved.

**Theorem 3** [6]. Every partial coloring of at most  $\Delta(T) - 1$  edges in a tree T is extendable to a proper  $\Delta(T)$ -coloring of T.

Using this result we shall establish Conjecture 1 for the case of trees.

**Theorem 4.** If T is a tree and  $\varphi$  a precoloring of  $\Delta(T)$  edges in  $T \Box K_2$ , then  $\varphi$  can be extended to a proper  $(\Delta(T) + 1)$ -coloring of  $T \Box K_2$ .

486

**Proof.** Let M be a perfect matching in  $T \Box K_2$  such that  $T \Box K_2 - M$  is isomorphic to two copies  $T_1$  and  $T_2$  of T.

If no precolored edges are in M, then we proceed as in the proof outline above, so we assume that M contains at least one precolored edge. By slight abuse of notation, we denote by  $T_i + M$  the graph obtained from  $T_i$  by attaching every edge of M as a pendant edge of  $T_i$ .

If no precolored edges are in  $T_2$ , then since  $T_1 + M$  is a forest with  $\Delta(T)$  precolored edges, by Theorem 3 there is a proper  $(\Delta(T) + 1)$ -coloring of  $T_1 + M$  that agrees with  $\varphi$ . Hence, by coloring every edge of  $T_2$  by the color of its corresponding edge in  $T_1$ , we obtain a proper  $(\Delta(T) + 1)$ -coloring of  $T \Box K_2$ .

Suppose now that both  $T_1$  and  $T_2$  contains at least one precolored edge. Set  $G_i = T_i + M$  and consider the restriction of  $\varphi$  to  $G_1$  and  $G_2$ , respectively. Note that both  $G_1$  and  $G_2$  are forests that each contains at most  $\Delta(T) - 1$  precolored edges. We shall define extensions of these precolorings of  $G_1$  and  $G_2$ , respectively, which agree on M. This yields an extension of  $\varphi$ .

Let  $V_M^1$  be the set containing all vertices of degree  $\Delta(T) + 1$  in  $G_1$  that are incident with some precolored edge from M and  $V_M^2$  be the set containing all vertices of degree  $\Delta + 1$  in  $G_2$  that are incident with some precolored edge from M. Note that  $v_1 \in V_M^1$  if and only if the corresponding vertex  $v_2 \in V_M^2$ . We shall prove that there are matchings  $M_1 \subseteq E(T_1)$  and  $M_2 \subseteq E(T_2)$  such that  $M_1$ covers  $V_M^1$  and  $e_2 \in M_2$  if and only if the corresponding edge  $e_1 \in M_1$ . Moreover we require that if  $u_1v_1 \in M_1$  and the corresponding edge  $u_2v_2 \in M_2$ , then

- (i) exactly one of  $u_1$  and  $v_1$  is in  $V_M^1$ , say  $u_1$ ;
- (ii) the components  $H_1$  of  $G_1 u_1$  and  $H_2$  of  $G_2 u_2$  containing  $v_1$  and  $v_2$ , respectively, contain no precolored edges;
- (iii)  $M_1$  and  $M_2$  do not contain any precolored edges.

Since all vertices of  $V_M^i$  have degree  $\Delta(T)$  in  $T_i$ ,  $T_1 \cup T_2$  contains altogether at most  $\Delta(T) - 1$  precolored edges, and all precolored edges of M are pairwise nonadjacent, we can indeed construct the required matchings  $M_1$  and  $M_2$  by for all precolored edges of M greedily selecting an adjacent edge of  $T_1$  (similarly for  $T_2$ ) so that (i)–(iii) holds.

Next, consider the edges of M that are not precolored but adjacent to some edge of  $M_1$ . We assign some fixed color c, arbitrarily chosen from  $\{1, \ldots, \Delta(T)\}$ , to every uncolored edge of M that is adjacent to an edge of  $M_1$  or  $M_2$ . Taken together with  $\varphi$  this yields a precoloring  $\varphi'$  of the graph  $T \Box K_2$ .

We denote by M' the set containing all the remaining uncolored edges of M. Since all edges of  $M_i$  and M' are pairwise nonadjacent, the set  $M''_i = M_i \cup M'$  is a matching in  $G_i$ . We set  $G''_i = G_i - M''_i$ . Then  $E(G''_1) \cap M = E(G''_2) \cap M$ .

By construction, each of the graphs  $G''_1$  and  $G''_2$  have maximum degree  $\Delta(T)$ , and since both  $G_1$  and  $G_2$  contain at most  $\Delta(T) - 1$  precolored edges under  $\varphi$ , respectively, and (i)–(iii) holds, every connected component of  $G''_1$  and  $G''_2$  contains at most  $\Delta(T) - 1$  precolored edges under the coloring  $\varphi'$ . Hence, it follows from Theorem 3, that there is a proper  $\Delta(T)$ -coloring  $f_i$  of  $G''_i$  that agrees with the restriction of  $\varphi'$  to  $G''_i$ . Note that  $f_1$  and  $f_2$  agree on all edges of M that are in  $G''_1$  (and  $G''_2$ ).

It remains to color the edges of  $M_1'' \cup M_2''$ . We simply assign color  $\Delta(T) + 1$  to all edges of this set. Taken together with  $f_1$  and  $f_2$ , this yields a proper  $(\Delta(T) + 1)$ -edge coloring of  $T \Box K_2$  that is an extension of  $\varphi$ , because  $f_1$  and  $f_2$  agree on  $M \cap E(G_1'')$ . This completes the proof.

Let us now consider complete bipartite graphs. Recall that the precoloring extension problem for the balanced complete bipartite graph  $K_{n,n}$  corresponds to asking whether a partial Latin square can be completed to a Latin square. As mentioned above, motivated by Evans' conjecture [11], Andersen and Hilton [2] completely characterized partial Latin squares of order n with n nonempty cells that cannot be completed to a Latin square of order n. In the language of colorings they proved the following.

**Theorem 5** [3]. Let  $n \ge 2$  be a positive integer. A precoloring  $\varphi$  of at most n edges of  $K_{n,n}$  can be extended to a proper n-coloring of  $K_{n,n}$  if and only if none of the following two conditions holds.

- (a) For some uncolored edge uv there are n differently colored edges with endvertices u or v.
- (b) For some vertex v and some color c, the color c does not appear on any edge incident with v, but every uncolored edge incident with v is adjacent to an edge colored c.

Using this result we now verify that Conjecture 1 holds for the Cartesian product  $K_{n,n} \Box K_2$ .

**Theorem 6.** Let  $n \ge 2$  be a positive integer. If  $\varphi$  is a precoloring of n edges in  $K_{n,n} \Box K_2$ , then  $\varphi$  can be extended to a proper (n + 1)-coloring of  $K_{n,n} \Box K_2$ .

**Proof.** Let M be a perfect matching in  $K_{n,n} \Box K_2$  such that  $K_{n,n} \Box K_2 - M$  is isomorphic to two copies  $K_{n,n}^1$  and  $K_{n,n}^2$  of  $K_{n,n}$ . If no precolored edges are in M, then we proceed as in the proof outline above, so we assume that M contains at least one precolored edge.

Let us first assume that both  $K_{n,n}^1$  and  $K_{n,n}^2$  contains at least one precolored edge, Let  $V_M^i$  be the set containing all vertices in  $K_{n,n}^i$  that are incident with some precolored edge from M. As in the preceding proof,  $v_1 \in V_M^1$  if and only if the corresponding vertex  $v_2 \in V_M^2$ .

Proceeding along the lines in the proof of Theorem 4, we shall construct matchings  $M_1 \subseteq E(K_{n,n}^1)$  and  $M_2 \subseteq E(K_{n,n}^2)$  such that each vertex of  $V_M^1$  is incident with a unique edge of  $M_1$ , and  $e_2 \in M_2$  if and only if the corresponding edge  $e_1$  in  $K_{n,n}^1$  is in  $M_1$ . Furthermore, we shall require that no edge e from  $M_1 \cup M_2$  is adjacent to a precolored edge of  $K_{n,n}^1$  or  $K_{n,n}^2$  of the same color as the precolored edge of M that e is adjacent to, and that no edge of  $M_1 \cup M_2$  is precolored. As above, since the vertex degree in  $K_{n,n}^1$  and  $K_{n,n}^2$  is n,  $K_{n,n}^1$  and  $K_{n,n}^2$  are bipartite, and  $K_{n,n} \Box K_2$  contains altogether n precolored edges, we can simply select the edges of  $M_1$  (and  $M_2$ ) greedily.

Now, from the restriction of  $\varphi$  to  $K_{n,n}^i$ , we define a new precoloring  $\varphi_i$  by coloring every edge of  $M_i$  by the color of the adjacent edge of M under  $\varphi$ . Since  $K_{n,n}^i$  contains at most n-1 precolored edges under  $\varphi_i$ , there is an extension  $f_i$  of  $\varphi_i$  using colors  $1, \ldots, n$ . Now, by recoloring all the edges in  $M_1$  and  $M_2$  by the color n+1 and coloring every uncolored edge of M by the color not appearing at its endpoints, we obtain an extension of  $\varphi$ .

Suppose now that no precolored edges are in  $K_{n,n}^1$  or  $K_{n,n}^2$ , say  $K_{n,n}^2$ . Then  $K_{n,n}^1$  contains at most n-1 precolored edges. Our proof of this case is similar to the proof of the preceding case. As above, let  $V_M^1$  be the set containing all vertices in  $K_{n,n}^1$  that are incident with some precolored edge from M. Since vertices in  $K_{n,n}^1$  have degree n, and  $K_{n,n}^1$  is bipartite, there is a matching  $M_1 \subseteq E(K_{n,n}^1)$  covering  $V_M^1$  satisfying analogous conditions to the matchings constructed in the preceding case.

From the restriction of  $\varphi$  to  $K_{n,n}^1$  we define a new precoloring  $\varphi_1$  of  $K_{n,n}^1$  by coloring every edge of  $M_1$  by the color of its adjacent edge in M. Now, if  $\varphi_1$  is extendable to a proper coloring of  $K_{n,n}^1$  using colors  $1, \ldots, n$ , then we obtain an extension of  $\varphi$  by recoloring all the edges in  $M_1$  by color n + 1, coloring  $K_{n,n}^2$ correspondingly, and then coloring every uncolored edge of M by the unique color from  $\{1, \ldots, n+1\}$  not appearing at its endpoints. So assume that there is no such extension of  $\varphi_1$ . By Theorem 5, this means that the coloring  $\varphi_1$  satisfies condition (a) or (b) of this theorem.

Suppose first that (a) holds. Then all colors  $1, \ldots, n$  appear on some edge under  $\varphi$ , so every color appears on precisely one edge. Since M contains at least one precolored edge, without loss of generality we may assume that one edge  $e_{M_1}$ in  $M_1$  is colored with color 1. Now define a new coloring  $\varphi'_1$  from  $\varphi_1$  by removing color 1 from  $e_{M_1}$  of  $K^1_{n,n}$ . Since  $K^1_{n,n}$  contains exactly n-1  $\varphi'_1$ -colored edges, there is a proper coloring of  $K^1_{n,n}$  using colors  $2, \ldots, n+1$  which is an extension of  $\varphi'_1$ . Now, by recoloring all the edges in  $M_1$  of  $K^1_{n,n}$ , distinct from  $e_{M_1}$ , by color 1, coloring  $K^2_{n,n}$  correspondingly, and then coloring every uncolored edge of Mby the unique color not appearing on any edge incident with one of its endpoints, we obtain an extension of  $\varphi$ .

Suppose now that (b) but not (a) holds. Then there is some color c that appears on at least two edges under  $\varphi$ . If some color c' that is used once by  $\varphi$  appears on some edge in the matching M, then we may remove the color c' from

the edge in  $M_1$  that is colored c', and then proceed as in the preceding paragraph. If, on the other hand, all colors that are used once by  $\varphi$  do not appear on edges of M, then color c is the only color that appears on edges in M; so color c is the only color that appears on edges in  $M_1$ .

Now, since  $K_{n,n}^1$  contains at most  $n-1 \varphi$ -colored edges, there is an extension of the restriction of  $\varphi$  to  $K_{n,n}^1$  using colors  $1, \ldots, n$ . Let  $M_c$  be the set of all edges of  $K_{n,n}^1$  that are colored c and adjacent to a precolored edge of M. By recoloring all the edges of  $M_c$  by n + 1, coloring  $K_{n,n}^2$  correspondingly, and then coloring every uncolored edge of M by the unique color not appearing at its endpoints, we obtain an extension of  $\varphi$ .

Finally, let us consider complete graphs. Again our confirmation of Conjecture 1 is based on a result by Andersen and Hilton [3].

**Theorem 7** [3]. Let  $n \ge 2$  be a positive integer.

- (i) If φ is a precoloring of at most n edges of K<sub>2n</sub>, then φ is extendable to a proper (2n − 1)-coloring unless the precolored edges form a matching, where n − 1 edges are colored by a fixed color c, and one edge is colored by some color c' ≠ c.
- (ii) If φ is a precoloring of at most n + 1 edges of K<sub>2n-1</sub>, then φ is extendable to a proper (2n 1)-coloring unless the precolored edges form a set of n 2 independent edges colored by a fixed color c, and a triangle, disjoint from the independent edges, the edges of which are colored by three different colors that are distinct from c.

In particular, this implies that every partial coloring of at most n-1 edges is extendable to a proper (2n-1)-coloring of  $K_{2n}$ , and similarly, every partial coloring of at most n edges of  $K_{2n-1}$  is extendable to a proper (2n-1)-coloring. The following establishes that Conjecture 1 holds for complete graphs.

**Theorem 8.** Let  $n \ge 2$  be a positive integer.

- (i) If φ is a precoloring of at most n edges of K<sub>2n</sub>□K<sub>2</sub>, then φ is extendable to a proper 2n-coloring of K<sub>2n</sub>□K<sub>2</sub>.
- (ii) If φ is a precoloring of at most n+1 edges of K<sub>2n-1</sub>□K<sub>2</sub>, then φ is extendable to a proper 2n-coloring of K<sub>2n-1</sub>□K<sub>2</sub>.

We note that the number of colors used in part (ii) is best possible, since there are partial colorings of just two edges in  $K_{2n-1} \Box K_2$  that are not extendable to proper (2n-1)-colorings.

Before we prove the general case of Theorem 8, we separately consider the case of  $K_5 \Box K_2$ . We first note the following lemma, which is easily proved using the fact that  $K_{2n-1} \Box K_2$  is Class 1.

**Lemma 9.** If  $\varphi$  is a precoloring of  $K_{2n-1} \Box K_2$  with n+1 precolored edges, where at least n edges have the same color, then  $\varphi$  is extendable to a proper 2n-coloring of  $K_{2n-1} \Box K_2$ .

In the following, we shall say that a matching *covers* a set S of edges, if every edge of S is adjacent to an edge of the matching.

**Lemma 10.** If  $\varphi$  is a partial coloring of at most 4 edges of  $K_5 \Box K_2$ , then  $\varphi$  is extendable to a proper 6-coloring.

**Proof.** Let M be a perfect matching in  $K_5 \Box K_2$  such that  $K_5 \Box K_2 - M$  is isomorphic to two copies  $K_5^1$  and  $K_5^2$  of  $K_5$ . If no precolored edge is in M, then we may proceed as in the proof outline, so we assume that at least one edge of M is precolored.

We shall consider many different cases. By Lemma 9, we may assume that at least two different colors are used in the precoloring  $\varphi$ .

Case 1. All precolored edges are in  $E(K_5^1) \cup M$ . Suppose first that exactly one color appears on the precolored edges of M, say color 1. Then we consider the precoloring of  $K_5^1$  obtained from  $\varphi$  by removing the color 1 from any edge of  $K_5^1$  that is precolored 1. By Theorem 7, this precoloring is extendable to a proper coloring of  $K_5^1$  using colors  $2, \ldots, 6$ . By recoloring any edge of  $K_5^1$  that is  $\varphi$ -precolored 1 by the color 1, coloring  $K_5^2$  correspondingly, and then coloring every uncolored edge of M by an appropriate color missing at its endpoints we obtain an extension of  $\varphi$ .

Suppose now that at least two colors appear on the precolored edges of M. We consider some different subcases.

Case 1.1. Exactly two edges of M are precolored. If there is at least one color c used on the precolored edges of  $K_5^1$  that neither appears on an edge of M, nor is the edge precolored c adjacent to a precolored edge of M, then there is a matching  $M_1 \subseteq E(K_5^1)$  of uncolored edges satisfying the following

- every edge of  $M_1$  is adjacent to exactly one precolored edge of M,
- every precolored edge of M is adjacent to an edge of  $M_1$ ,
- no edge of  $M_1$  is adjacent to two precolored edges  $e_1 \in E(K_5^1)$  and  $e_2 \in M$  that have the same color under  $\varphi$ .

We call such a matching a  $\varphi$ -good matching. Moreover, it is easy to see that we can pick this matching such that if we color the edges of  $M_1$  by the color of the precolored adjacent edges in M, then this coloring along with the restriction of  $\varphi$  to  $K_5^1$  does not satisfy the condition in Theorem 7. Hence, the obtained 5-precoloring of  $K_5^1$  is extendable. By recoloring every edge of  $M_1$  by the color 6, coloring  $K_5^2$  correspondingly, and then coloring the edges of M by an appropriate color not appearing at its endpoints, we obtain an extension of  $\varphi$ . Suppose now that there is no  $\varphi$ -good matching, but there is a color  $c_1$  that appears in  $K_5^1$  but not in M. Since there is no  $\varphi$ -good matching, the edge  $e_1$ colored  $c_1$  is adjacent to an edge e of M that is colored  $c_2 \neq c_1$ , but not to the other precolored edge of M. Moreover, there is another edge  $e'_1$  in  $K_5^1$  colored  $c_2$ . Now, it is easy to see that this implies that there is a proper edge coloring of  $K_5 \Box K_2$  with colors  $\{1, \ldots, 6\} \setminus \{c_2\}$  that agrees with all precolored edges that are not precolored  $c_2$ . Hence,  $\varphi$  is extendable.

Next, assume that the same two colors, say 1 and 2, are used both on the precolored edges of M and on  $K_5^1$ . Then there is at most one vertex of degree 4 in the graph obtained from  $K_5^1$  by removing all precolored edges. We properly color the uncolored edges of  $K_5^1$  using colors 3, 4, 5, 6, color  $K_5^2$  correspondingly, and then proceed as before.

Case 1.2. Exactly three edges of M are precolored. Let  $e_1$  be the precolored edge of  $K_5^1$  and assume first that only two colors, say 1 and 2, are used in the precoloring  $\varphi$ . By Lemma 9, we may assume that  $\varphi(e_1) = 1$  and exactly one edge of M is colored 1. Then we can pick an uncolored edge  $e'_1$  of  $K_5^1$ , that is not adjacent to any edge  $\varphi$ -colored 1. We color  $K_5^1 - \{e_1, e'_1\}$  properly by colors 3, 4, 5, 6, color  $e_1$  and  $e'_1$  by 1 and proceed as before.

Suppose now that three colors appear in the precoloring  $\varphi$ . If the color of the precolored edge in  $K_5^1$  does not appear on an edge of M, then there is some color c that appears on two edges in M, and a color c' that only appears on one edge of M. This implies that there is a matching  $M_1 \subseteq E(K_5^1)$  of uncolored edges satisfying the following

- every edge of  $M_1$  is adjacent to exactly one precolored edge of M,
- every precolored edge of M colored c is adjacent to an edge of  $M_1$ ,
- the edge precolored c' is not adjacent to an edge of  $M_1$ .

Consider the precoloring obtained from the restriction of  $\varphi$  to  $K_5^1$  by in addition coloring every edge of  $M_1$  by the color c. It follows from Theorem 7 that there is an extension of this coloring to  $K_5^1$  using colors  $\{1, 2, 3, 4, 5, 6\} \setminus \{c'\}$ . Now, we obtain an extension of  $\varphi$  by coloring the edges of  $M_1$  by the color c', coloring  $K_5^2$ correspondingly, and finally coloring the edges of M appropriately.

If, on the other hand, there is a color c which appears both in  $K_5^1$  and in M, then we proceed as follows. We pick a proper coloring f of  $K_5^1$  using colors  $\{1, 2, 3, 4, 5, 6\} \setminus \{c\}$ , so that for every vertex  $v \in V(K_5^1)$ , if there is a precolored edge of M incident with v, then the color of the edge of M does not appear at v under f. Next, we recolor the edges of  $K_5^1 \varphi$ -precolored c by the color c, and obtain an extension of  $\varphi$  as before.

The case when four colors appear on edges under  $\varphi$  can be dealt with using a similar argument as in the preceding paragraph.

Case 1.3. Exactly four edges of M are precolored. Suppose now that all four precolored edges are in M. If only two colors appear in  $\varphi$ , then by Lemma 9, we may assume that both colors appear on two edges. Thus there is a matching  $M_1$  in  $K_5^1$  covering all precolored edges of M and such that no edge in  $M_1$  is adjacent to two edges precolored with different colors. Thus,  $\varphi$  is extendable, as before. If, on the other hand, three colors appear in  $\varphi$ , then there is a color cthat appears on at least two edges. We can pick a proper coloring f of  $K_5^1$  using colors  $\{1, \ldots, 6\} \setminus \{c\}$ , such that no precolored edge of M is adjacent to an edge of the same color under f. Hence,  $\varphi$  is extendable. Note that we can use a similar argument if four colors appear on edges under  $\varphi$ .

Note that in all cases above,  $K_5^1$  is first colored, and then  $K_5^2$  is colored correspondingly. We shall use this property when we consider the next case.

Case 2. Both  $K_5^1$  and  $K_5^2$  contains at least one precolored edge. The condition implies that M contains one or two precolored edges. Assume first that Mcontains only one precolored edge, and so we may assume that  $K_5^1$  contains two precolored edges, and  $K_5^2$  one. Let  $e_1$  and  $e'_1$  be the precolored edges of  $K_5^1$ ,  $e_2$ and  $e'_2$  the corresponding edges of  $K_5^2$  respectively, and let  $e''_2$  be the precolored edge of  $K_5^2$ , and  $e''_1$  the corresponding edge of  $K_5^1$ . Furthermore, let e be the precolored edge of M.

Consider the restriction of  $\varphi$  to  $K_5^1$ . If we can assign the color  $\varphi(e_2'')$  to  $e_1''$  so that the resulting coloring  $\varphi_1$  of  $K_5^1$  is proper, then we may proceed as in Case 1 (since in that case  $K_5^2$  is always colored correspondingly). Thus we may assume that either

- (a)  $e_1'' \in \{e_1, e_1'\}$ , or
- (b)  $e_1''$  is adjacent to one of the edges in  $\{e_1, e_1'\}$  and  $e_2''$  has the same color as one adjacent edge in  $\{e_1, e_1'\}$ .

By Lemma 9, we may further assume that no color appears on three edges, and thus at most two edges are precolored by the same color. Then, unless e is adjacent to two precolored edges and there is another edge  $e_1$  precolored  $\varphi(e)$ , that is disjoint from all these three edges, there is an uncolored edge  $e_1^{(3)}$  in  $K_5^1$ that is adjacent to e but not adjacent to a precolored edge of  $K_5^1$  colored  $\varphi(e)$ , and, similarly for the corresponding edge  $e_2^{(3)}$  of  $K_5^2$ . From the restriction of  $\varphi$ to  $K_5^1$  and  $K_5^2$  we obtain new 5-precolorings by coloring the edges  $e_1^{(3)}$  and  $e_2^{(3)}$ by the color  $\varphi(e)$ . Now by Theorem 7 these precolorings are extendable, and we may finish the argument by proceeding as above.

Suppose now that e is adjacent to two precolored edges and there is one additional edge  $e_1$  precolored  $\varphi(e)$  in  $E(K_5^1) \cup E(K_5^2)$ . It is not hard to see that this implies that there are uncolored corresponding edges  $e_1^{(4)} \in E(K_5^1)$ and  $e_2^{(4)} \in E(K_5^2)$  that are not adjacent to any edges precolored  $\varphi(e)$ . We now construct new precolorings from the restrictions of  $\varphi$  to  $K_5^1$  and  $K_5^2$ , respectively, by coloring  $e_1^{(4)}$  and  $e_2^{(4)} \varphi(e)$ , and also  $e_2$  by the color  $\varphi(e)$ . These partial colorings are extendable using colors 1, 2, 3, 4, 5. Moreover, by construction, the extensions of these precolorings satisfy that no edge colored c is adjacent to e. Hence,  $\varphi$  is extendable.

Let us now assume that M contains two precolored edges. So  $K_5^1$  and  $K_5^2$  both contains precisely one precolored edge,  $e_1$  and  $e'_2$ , respectively. Denote the corresponding edges of  $K_5^2$  and  $K_5^1$  by  $e_2$  and  $e'_1$ , respectively. As above, it follows that we may assume that either

- (a)  $e_1 = e'_1$  and  $\varphi(e_1) \neq \varphi(e'_2)$ , or
- (b)  $e_1$  and  $e_2$  are adjacent and  $\varphi(e_1) = \varphi(e'_2)$ .

Suppose first that (a) holds. If both colors in  $\{\varphi(e_1), \varphi(e'_2)\}$ , say 1 and 2, appear on the precolored edges of M, then we pick corresponding uncolored edges  $e''_1 \in E(K_5^1)$  and  $e''_2 \in E(K_5^2)$  that are not adjacent to  $e_1$  or  $e_2$  and color  $e''_1$  and  $e''_2$  by 1 or 2 so that the resulting coloring of  $K_5 \times K_2$  is proper. Thereafter, we color  $K_5^1 - \{e'_1, e''_1\}$  and  $K_5^2 - \{e'_2, e''_2\}$  properly using colors 3, 4, 5, 6, and proceed as above.

If, on the other hand, at least one of the colors in  $\{\varphi(e_1), \varphi(e_2)\}$  do not appear on the precolored edges of M, then there are matchings  $M_1 \subseteq E(K_5^1)$  and  $M_2 \subseteq E(K_5^2)$  of corresponding uncolored edges such that

- every precolored edge of M is adjacent to exactly one edge of  $M_i$ ,
- every edge of  $M_i$  is adjacent to a precolored edge of M,
- no edge of  $M_i$  is adjacent to two precolored edges of the same color.

Now, consider the restriction of  $\varphi$  to  $K_5^1$  and  $K_5^2$ , respectively. By, in addition, coloring the edges of  $M_1$  and  $M_2$  by the color of the adjacent precolored edge of M we obtain extendable 5-precolorings of  $K_5^1$  and  $K_5^2$ , respectively. Given extensions of these precolorings, we may recolor the edges of  $M_1$  and  $M_2$  by color 6, and then color the edges of M appropriately to obtain an extension of  $\varphi$  as before.

Suppose now that (b) holds. Since  $\varphi(e_1) = \varphi(e'_2)$ , and we may by Lemma 9 assume that any color appears on at most two edges under  $\varphi$ , there are matchings  $M_1$  and  $M_2$  as described in the preceding paragraph. Thus, we proceed similarly, and this completes the proof of the lemma.

**Proof of Theorem 8.** We first prove part (i). Denote by M the matching of  $K_{2n} \Box K_2$  such that  $K_{2n} \Box K_2 - M$  is isomorphic to two copies  $K_{2n}^1$  and  $K_{2n}^2$  of  $K_{2n}$ . The cases when no precolored edges are in M can be handled as in the proof of Theorem 6, so we omit the details here.

In the case when M contains at least one precolored edge, then we may select a matching  $M_1$  in  $K_{2n}^1$  as in the proof of Theorem 6 (and possibly also a matching  $M_2$  of corresponding edges in  $K_{2n}^2$ ). The only essential difference in the argument is that since  $K_{2n}$  contains triangles, we can only ensure that n such edges forming the matching  $M_1$  can be selected greedily, although  $K_{2n}$  has vertex degree 2n-1. Nevertheless, since  $K_{2n} \Box K_2$  contains at most n precolored edges, this suffices for our purposes. Apart from this difference, the argument is very similar to the one in the proof of Theorem 6, so we omit the details.

Let us now prove part (ii). Denote by M the matching of  $K_{2n-1} \Box K_2$  such that  $K_{2n-1} \Box K_2 - M$  is isomorphic to two copies  $K_{2n-1}^1$  and  $K_{2n-1}^2$  of  $K_{2n-1}$ .

The case of  $K_3$  follows from the result on odd cycles proved in Section 4, and the case of  $K_5$  is dealt with by the above lemma, so let us assume that  $n \ge 4$ .

We shall consider a number of different cases. In many of these cases we shall use strategies which are similar to the ones used in the proofs of Theorem 6 and/or Lemma 10, so we generally omit many details.

As before, the case when no precolored edge is in M can be dealt with as in the proof outline, so in the following we assume that at least one precolored edge is contained in M. The rest of the proof breaks into the following cases.

- (1) Only one color appear on the precolored edges in M.
- (2) At least two colors appear on the precolored edges in M, but at most one color appears on the edges in  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$ .
- (3) At least two colors appear on the precolored edges of M, and at least two colors appear on the precolored edges in  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$ .

In many of the different subcases below we shall use matchings  $M_1 \subseteq E(K_{2n-1}^1)$ and  $M_2 \subseteq E(K_{2n-1}^2)$  of corresponding uncolored edges that covers all precolored edges of M, and in addition satisfy that

- no edge of  $M_i$  is adjacent to two precolored edges of M of different colors,
- no edge of  $M_i$  is adjacent to an edge of  $K_{2n-1}^i$  and an edge of M that are precolored by the same color.

We simply say that  $(M_1, M_2)$  is a  $\varphi$ -good pair of matchings, and that each of  $M_1, M_2$  is a  $\varphi$ -good matching. When using such matchings, we usually consider the precolorings obtained from the restriction of  $\varphi$  to  $K_{2n-1}^1$  and  $K_{2n-1}^2$ , respectively, by coloring the edges of  $M_1 \cup M_2$  by the color of the adjacent precolored edge of M. The resulting precolorings will be extendable to (2n - 1)-colorings of  $K_{2n-1}^1$  and  $K_{2n-1}^2$ , respectively; indeed, it suffices to verify that none of these precolorings satisfy the condition in part (ii) of Theorem 7. Since such verifications are straightforward, we omit the exact details in the arguments below. Now, from these extensions, we obtain an extension of  $\varphi$  by recoloring the edges of  $M_1 \cup M_2$  by the color 2n, and then coloring the edges of M appropriately.

Case 1. Only one color appears on the precolored edges of M. Assume that color 1 appears on the edges of M. If all precolored edges are in  $M \cup E(K_{2n-1}^1)$ , then consider the precoloring obtained from the restriction of  $\varphi$  to  $K_{2n-1}^1$  by removing the color 1 from all edges  $\varphi$ -colored 1. By Theorem 7, this precoloring is extendable to a proper coloring using colors  $2, \ldots, 2n$ , and we obtain an extension of  $\varphi$  by recoloring the edges of  $K_{2n-1}^1$  that are  $\varphi$ -colored 1 by the color 1, coloring  $K_{2n-1}^2$  correspondingly and then coloring every edge of M by the color 1 or 2n.

Suppose now that both  $K_{2n-1}^1$  and  $K_{2n-1}^2$  contains at least one precolored edge. By Lemma 9, we may assume that there are at least two edges in  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$  precolored by a color distinct from 1. If there is a  $\varphi$ -good pair of matchings  $(M_1, M_2)$ , then we consider the precolorings obtained from the restrictions of  $\varphi$  to  $K_{2n-1}^1$  and  $K_{2n-1}^2$ , respectively, by in addition coloring all edges of  $M_1$ and  $M_2$  by the color 1, and then proceed as outlined above.

Suppose now that there is no  $\varphi$ -good pair of matchings  $(M_1, M_2)$ . Since at most n-2 edges of  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$  are precolored 1, it follows that only one edge  $u_1u_2$  of M is precolored (where  $u_i \in V(K_{2n-1}^i)$ ), there is a matching M' of n-2 edges in  $K_{2n-1}^1$  such that every edge of M' is either precolored 1, or the corresponding edge of  $K_{2n-1}^2$  is precolored 1. Moreover,  $u_1$  is incident with two edges  $e_1$  and  $e_2$  that are independent from M' and satisfy that  $e_i$  or the corresponding edge of  $K_{2n-1}^2$  is precolored by a color distinct from 1. Now, from the restriction of  $\varphi$  to  $K_{2n-1}^1$  we define a new precoloring  $\varphi_1$  by coloring all edges of M' by the color 1, and, in addition, coloring the unique edge of  $K_{2n-1}^1$  that is adjacent to both  $e_1$  and  $e_2$  by the color 1. We define an analogous precoloring  $\varphi_2$  of  $K_{2n-1}^2$ .

By Theorem 7, both  $\varphi_1$  and  $\varphi_2$  are extendable to proper (2n-1)-colorings. Moreover, it is easy to see that neither  $u_1$  nor  $u_2$  is incident with an edge colored 1 in these extensions. Consequently,  $\varphi$  is extendable.

Case 2. At least two colors appear on the precolored edges of M, but at most one color appears on the precolored edges of  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$ .

We first consider the case when all precolored edges are in  $M \cup E(K_{2n-1}^1)$ .

Suppose first that all precolored edges are contained in M. If every color appears on at most one edge in M, then  $\varphi$  is extendable, because we can choose a proper (2n - 1)-coloring f of  $K_{2n-1}^1$  so that for every vertex  $v \in V(K_{2n-1})$ , if the edge of M incident with v is colored  $i \in \{1, \ldots, 2n-1\}$ , then no edge incident with v is colored i under f. A similar argument applies if one color appears on at least two edges in M and all other colors appear on at most one edge.

If, on the other hand, there are two colors  $c_1$  and  $c_2$  that both appear on at least two precolored edges, then there is a  $\varphi$ -good matching  $M_1$  in  $K_{2n-1}^1$ . Thus  $\varphi$  is extendable.

Suppose now that  $E(K_{2n-1}^1)$  contains at least one precolored edge, precolored c say. If all precolored edges of M are colored differently, except that several edges

496

of M may be colored c, then there is a proper coloring f of  $K_{2n-1}^1$  using colors  $\{1, \ldots, 2n\} \setminus \{c\}$ , such that no precolored edge of M is adjacent to an edge of the same color under f. Hence,  $\varphi$  is extendable.

Assume instead that some color  $c_1 \neq c$  appears on at least two edges of Mand every other color appears on at most one edge of M. If at most one edge of  $K_{2n-1}^1$  is precolored c, then we proceed as before and pick a proper coloring fof  $K_{2n-1}^1$  using colors  $\{1, \ldots, 2n\} \setminus \{c_1\}$  that agrees with the restriction of  $\varphi$  to  $K_{2n-1}^1$ , and where no precolored edge of M is adjacent to an edge of the same color under f. If instead color c is used on at least two edges of  $K_{2n-1}^1$ , then it is easy to see that there is a  $\varphi$ -good matching  $M_1$ . Hence,  $\varphi$  is extendable.

Finally, let us assume that there are two colors  $c_1, c_2 \neq c$  that both appear on at least two edges of M. Again, this implies that there is a  $\varphi$ -good matching  $M_1$ , and so,  $\varphi$  is extendable.

Let us now consider the case when both  $K_{2n-1}^1$  and  $K_{2n-1}^2$  contains at least one edge precolored c. If all the colors on edges of M are distinct, except that several edges of M may be colored c, then  $\varphi$  is extendable as in the case when all precolored edges are in  $K_{2n-1}^1$ . Consequently, we assume that there is a color  $c_1 \neq c$ that appears on at least two edges  $u_1u_2, v_1v_2 \in M$ , where  $u_i, v_i \in V(K_{2n-1}^i)$ .

We may assume that at least one precolored edge of  $K_{2n-1}^2$  satisfies that the corresponding edge of  $K_{2n-1}^1$  is adjacent to a precolored edge of  $K_{2n-1}^1$ , since otherwise we can define a precoloring of  $K_{2n-1}^1$  by coloring every edge e that is precolored c, or satisfying that the corresponding edge of  $K_{2n-1}^2$  is precolored c, by the color c, and then proceed as in the case when only edges of  $K_{2n-1}^1$  and M are precolored. This assumption implies that there is a pair of  $\varphi$ -good matchings  $(M_1, M_2)$  (containing  $u_1v_1$  and  $u_2v_2$ , respectively, if these edges are uncolored). As before, this implies that  $\varphi$  is extendable.

Case 3. At least two colors appear on the precolored edges of M and at least two colors appear on the precolored edges of  $E(K_{2n-1}^1) \cup E(K_{2n-1}^2)$ .

Suppose first that all precolored edges lie in  $M \cup E(K_{2n-1}^1)$ . If there is a  $\varphi$ -good matching  $M_1$  in  $K_{2n-1}^1$ , then  $\varphi$  is extendable. Otherwise, if there is no  $\varphi$ -good matching, then at most two edges of M are precolored, because otherwise we could select three edges for  $M_1$ , each of which is adjacent to at least one precolored edge of  $K_{2n-1}^1$ , and thereafter select the rest of the edges of  $M_1$  greedily.

Furthermore, since at least two colors appear on the edges in  $K_{2n-1}^1$ , if there is no  $\varphi$ -good matching, then one precolored edge of M, colored  $c_1$  say, must be adjacent to an edge e' precolored  $c_2 \neq c_1$ , and there is a matching M' of n-2 edges in  $K_{2n-1}^1$ , disjoint from e', all edges of which are precolored  $c_1$ . It is straightforward to verify that this precoloring is extendable, e.g. by first taking an extension of the restriction of  $\varphi$  to the edges precolored by colors distinct from  $c_1$  using colors  $\{1, \ldots, 2n\} \setminus \{c_1\}$ . Let us now consider the case when both  $K_{2n-1}^1$  and  $K_{2n-1}^2$  contains at least one precolored edge. Again, the idea is to find a pair of  $\varphi$ -good matchings  $(M_1, M_2)$ . If there are such matchings  $M_1$  and  $M_2$ , then we can use them for finding an extension of  $\varphi$  as before.

On the other hand, if there are no such matchings, then at most two edges of M are precolored; suppose e.g. that  $e_1, e_2 \in M$  are precolored  $c_1$  and  $c_2$ , respectively. Again it follows that one of  $e_1$  and  $e_2$ , say  $e_1$ , is adjacent to an edge  $e' \in E(K_{2n-1}^1)$  (or  $e' \in E(K_{2n-1}^2)$ ) colored  $c_3 \neq c_1$ , and there is a matching M' in  $K_{2n-1}^1$  ( $K_{2n-1}^2$ ) of n-2 edges, every edge of which is either colored  $c_1$ , or satisfies that the corresponding edge of  $K_{2n-1}^2$  ( $K_{2n-1}^1$ ) is precolored  $c_1$ . Moreover,  $M' \cup \{e', e_2\}$  is independent.

Thus we can color every edge of M' by the color  $c_1$ , retain the color of e' if  $c_3 \neq c_2$ , and also color the unique edge adjacent to both e' and  $e_2$ , but not  $e_1$ , by the color  $c_1$  to obtain a precoloring of  $K_{2n-1}^1$  that is extendable using colorings  $\{1, \ldots, 2n\} \setminus \{c_2\}$ . Moreover, every extension of this precoloring satisfies that no edge adjacent to  $e_1$  is colored  $c_1$ . Hence,  $\varphi$  is extendable. This completes the proof of the theorem.

### 3. CARTESIAN PRODUCTS WITH GENERAL GRAPHS

We have not been able to confirm Conjecture 1 in the general case, but we can prove it for the case of regular triangle-free graphs when the precolored edges are independent.

**Theorem 11.** If G is a triangle-free regular graph where every precoloring of at most  $k < \Delta(G)$  independent edges are extendable to a  $\chi'(G)$ -coloring, then every precoloring of at most k + 1 independent edges in  $G \square K_2$  is extendable to a  $(\chi'(G) + 1)$ -coloring.

**Proof.** Without loss of generality, we assume that k + 1 edges of  $G \Box K_2$  are precolored. We denote this precoloring by  $\varphi$ , by  $G_1$  and  $G_2$  the copies of G in  $G \Box K_2$ , respectively, and by M the perfect matching between  $G_1$  and  $G_2$ .

If no precolored edges are in M, then we proceed as in the proof outline above, so we assume that at least one precolored edge is contained in M, and consider two different cases.

Case 1. All precolored edges are in  $E(G_1) \cup M$ . We assume that at least one precolored edge is in M. Let  $E_M$  be the set of all precolored edges in M. As in the proof outline, we shall select a matching  $M_1$  of  $|E_M|$  uncolored edges in  $G_1$  satisfying the following:

- (i) every edge of  $E_M$  is adjacent to exactly one edge of  $M_1$ ;
- (ii) every edge of  $M_1$  is adjacent to exactly one precolored edge of G.

498

Since  $G_1$  is regular and triangle-free, at most one endpoint of a precolored edge of  $G_1$  is adjacent to an endpoint of a precolored edge in M. Thus, since  $G \Box K_2$  contains at most  $\Delta(G_1)$  precolored edges and all those precolored edges are independent, it is straightforward to verify that there is a set  $M_1 \subseteq E(G_1)$ satisfying (i)–(ii); indeed, since every vertex of  $G_1$  has degree  $\Delta(G_1)$ , we can simply select edges adjacent to the precolored edges of M greedily.

Now, by coloring all edges of  $M_1$  by the color of the adjacent edge of M, and taking this coloring together with the restriction of  $\varphi$  to  $G_1$ , we obtain a precoloring  $\varphi_1$  of  $G_1$  with k + 1 precolored independent edges. Without loss of generality we assume that some edge is colored 1 under  $\varphi_1$ . By removing the color 1 from every such edge  $\varphi_1$ -precolored 1, we obtain a precoloring that is extendable to a proper coloring of  $G_1$  using colors  $2, \ldots, \chi'(G) + 1$ .

Next, for every edge of  $G_1$  that is  $\varphi$ -precolored 1, we recolor this edge by 1. Similarly, for every  $\varphi_1$ -precolored edge e of  $M_1$  such that  $\varphi_1(e) \neq 1$ , we recolor e by the color 1 and the adjacent edge of M by the color  $\varphi_1(e)$ . Finally, we recolor any edge of M that is  $\varphi$ -precolored 1 by the color 1. Since all  $\varphi_1$ -precolored edges are independent, the resulting partial coloring of G is proper. By coloring  $G_2$  correspondingly and coloring all uncolored edges of M by a color missing at its endpoints, we obtain a proper coloring that is an extension of  $\varphi$ .

Case 2.  $E(G_1), E(G_2)$  and M contains at least one precolored edge each. Let  $\varphi_1$  and  $\varphi_2$  be the restrictions of  $\varphi$  to  $G_1$  and  $G_2$ , respectively. As in the preceding case we shall select a matching of  $|E_M|$  uncolored edges  $M_1 \subseteq E(G_1)$  and a matching  $M_2 \subseteq E(G_2)$  of uncolored corresponding edges satisfying the following:

- (i) every edge of  $E_M$  is adjacent to exactly one edge of  $M_i$ ;
- (ii) every edge of  $M_1 \cup M_2$  is adjacent to exactly one precolored edge of G.

The existence of such sets  $M_1$  and  $M_2$  follows as in Case 2, since  $G_1$  and  $G_2$  are  $\Delta(G)$ -regular and  $G \square K_2$  contains altogether at most  $\Delta(G)$  precolored edges.

Now, consider the precolorings obtained from  $\varphi_1$  and  $\varphi_2$ , respectively, by coloring every edge of  $M_1$  and  $M_2$  by the color of the adjacent precolored edge of M. Since  $G_1$  and  $G_2$  both contains at least one  $\varphi$ -precolored edge, the obtained precolorings  $\varphi'_1$  and  $\varphi'_2$ , respectively, are by assumption extendable to proper  $\chi'(G)$ -colorings. Now, we recolor every edge of  $M_1 \cup M_2$  by the color  $\chi'(G) + 1$ , and then color every edge of M that is adjacent to an edge of  $M_1 \cup M_2$  by the color of the adjacent edge of  $M_1 \cup M_2$  (i.e., the color under  $\varphi$  if the edge of M is precolored). We color every edge of M that is not adjacent to an edge of  $M_1 \cup M_2$  by the color  $\chi'(G) + 1$ . Since  $M_i$  is a matching, the resulting coloring is proper, and thus also an extension of  $\varphi$ .

For graphs with triangles we have the following variant of Theorem 11.

**Theorem 12.** If G is a regular graph where every precoloring of at most  $k < \Delta(G)/2$  independent edges is extendable to a  $\chi'(G)$ -coloring, then every precoloring of at most k + 1 independent edges in  $G \Box K_2$  is extendable to a  $(\chi'(G) + 1)$ -coloring.

The proof of this theorem is almost identical to the proof of the preceding one. The only essential difference is that when G is not triangle-free we have to assume that at most  $\Delta(G)/2$  edges are precolored to be able to ensure that we can select independent edges in  $G_1$  and  $G_2$  that are adjacent to precolored edges of M and also not adjacent to any precolored edges in  $G_1$  or  $G_2$ . We omit the details.

### 4. CARTESIAN PRODUCTS WITH SUBCUBIC GRAPHS

In this section, we consider Conjecture 1 for subcubic graphs, that is, graphs with maximum degree at most 3. First we prove that it holds for graphs with maximum degree two. The case of paths was considered above, so it suffices to consider cycles. We shall need some well-known auxiliary results on list coloring.

**Lemma 13.** For every path P, if one edge  $e \in E(P)$  has a list of size at least 1 and all other edges have lists of at least 2 colors, then P has a proper coloring using colors from the lists.

**Lemma 14.** If L is a list assignment for the edges of a cycle C, where every list has size at least two and not all edges have the same list, then C has a proper coloring using colors from the lists.

**Proposition 15.** Let  $n \ge 2$  be a positive integer. If  $\varphi$  is a precoloring of two edges in  $C_{2n} \Box K_2$ , then  $\varphi$  can be extended to a proper 3-coloring of  $C_{2n} \Box K_2$ .

**Proof.** Let M be matching of  $C_{2n} \Box K_2$ , so that  $C_{2n} \Box K_2 - M$  consists of two copies  $C_{2n}^1$  and  $C_{2n}^2$  of  $C_{2n}$ . The cases when no precolored edge is contained in the matching M can be dealt with as above. If M contains exactly one precolored edge, then if the two precolored edges have distinct colors, then  $\varphi$  is trivially extendable. If, on the other hand, the two precolored have the same color, say 1, then we can properly color the uncolored edges in the copy of  $C_{2n}$  containing a precolored edge by colors 2 and 3; thus  $\varphi$  is extendable.

It remains to consider the case when no precolored edges are in  $C_{2n}^1$  or  $C_{2n}^2$ . If the two precolored edges have the same color, then  $\varphi$  is trivially extendable. If they have different colors and have distance at least 2, then  $\varphi$  is extendable by Lemma 14. The case when the precolored edges are at distance 1 is straightforward.

500

Note that Proposition 15 does not hold for odd cycles. For instance, consider the Cartesian product  $C_{2n+1} \Box K_2$  where two corresponding edges of the two copies of  $C_{2n+1}$  are colored by 1 and 2, respectively. If this precoloring is extendable to a proper 3-coloring of  $C_{2n+1} \Box K_2$ , then every edge in the matching M joining vertices of the copies of  $C_{2n+1}$  must be colored 3. Hence, the precoloring is not extendable.

Nevertheless, for odd cycles, we have the following analogue of Proposition 15.

**Proposition 16.** Let  $n \ge 1$  be a positive integer. If  $\varphi$  is a precoloring of three edges in  $C_{2n+1} \Box K_2$ , then  $\varphi$  can be extended to a proper 4-coloring of G.

**Proof.** Let M be a perfect matching in  $C_{2n+1} \Box K_2$  such that  $C_{2n+1} \Box K_2 - M$  is isomorphic to two copies  $C_{2n+1}^1$  and  $C_{2n+1}^2$  of  $C_{2n+1}$ . Without loss of generality we assume that the precoloring of G uses colors 1, 2, 3. We shall consider some different cases. Again, we omit the details in the case when M contains no precolored edges.

Case 1. M contains exactly one precolored edge. For any uncolored edge  $e \in E(C_{2n+1} \Box K_2)$ , we define a color list  $L(e) \subseteq \{1, 2, 3, 4\}$  by setting

$$L(e) = \{1, 2, 3, 4\} \setminus \{\varphi(e') : e' \text{ is adjacent to } e\}.$$

If no precolored edges are in  $C_{2n+1}^2$ , then at most one uncolored edge of  $C_{2n+1}^1$ is adjacent to precolored edges of three distinct colors, which implies that at most one edge e has a list L(e) of size 1 and all other edges of  $C_{2n+1}^1$  have lists of size at least 2. By Lemma 13, there is a proper coloring  $\varphi_1$  of  $C_{2n+1}^1$ , which is an extension of the restriction of  $\varphi$  to  $C_{2n+1}^1$ . Hence, we obtain an extension of  $\varphi$ by coloring every edge of  $C_{2n+1}^2$  by the color of its corresponding edge in  $C_{2n+1}^1$ , and then coloring every edge of M with some color in  $\{1, 2, 3, 4\}$  that is missing at its endpoints.

Suppose now that both  $C_{2n+1}^1$  and  $C_{2n+1}^2$  each contains exactly one precolored edge. Let  $e_1 \in E(C_{2n+1}^1)$ ,  $e_2 \in M$ , and  $e_3 \in E(C_{2n+1}^2)$  be the precolored edges of  $C_{2n+1}^1$ , M and  $C_{2n+1}^2$ , respectively.

First we treat the case when n = 1; it needs to be considered separately. If  $e_1$  and  $e_2$ , and  $e_2$  and  $e_3$  are adjacent, then  $\varphi$  is extendable since  $\varphi(e_2) \notin \{\varphi(e_1), \varphi(e_3)\}$ ; we can e.g. first properly color  $C_3^1$  and  $C_3^2$  using colors from  $\{1, 2, 3, 4\} \setminus \{\varphi(e_2)\}$ , and then color the remaining uncolored edges of M. If exactly two of the edges  $e_1, e_2, e_3$  are pairwise adjacent, say  $e_1$  and  $e_2$ , and  $\varphi(e_3) = \varphi(e_2)$ , then we can instead use colors  $\varphi(e_1), \varphi(e_2)$  and one additional color from  $\{1, 2, 3, 4\}$  for coloring  $C_3^1$  and  $C_3^2$ ; if  $\varphi(e_3) \neq \varphi(e_2)$ , then we proceed similarly using colors  $\{1, 2, 3, 4\} \setminus \{\varphi(e_2)\}$ .

Suppose now that all the edges  $e_1, e_2, e_3$  are pairwise nonadjacent. If at most two colors  $c_1$  and  $c_2$  appear on the precolored edges, then we may assume that

 $c_1$  appears on at most one edge under  $\varphi$ . We color  $C_3 \Box K_2$  properly using the colors in  $\{1, 2, 3, 4\} \setminus \{c_1\}$ , so that the edges precolored  $c_2$  get the color  $c_2$ . Then we recolor the edge precolored  $c_1$  by the color  $c_1$ . If, on the other hand, three colors appear on the precolored edges, then we properly color  $C_3^1$  and  $C_3^2$  using colors  $\{1, 2, 3, 4\} \setminus \{\varphi(e_2)\}$ .

Now we treat the case when  $n \ge 2$ . We define a list assignment L for  $C_{2n+1}^1$  by setting

$$L(e) = \{1, 2, 3\} \setminus \{\varphi(e') : e' \in E(C_{2n+1} \Box K_2) \text{ is adjacent to } e\}.$$

Since at most one edge is adjacent to both  $e_1$  and  $e_2$ , at most one uncolored edge e of  $C_{2n+1}^1$  satisfies that |L(e)| = 1, and all other edges of  $C_{2n+1}^1$  have lists of size at least two. By Lemma 13, there is a proper coloring  $\varphi_1$  of  $C_{2n+1}^1$  using colors 1, 2, 3 which is an extension of the restriction of  $\varphi$  to  $C_{2n+1}^1$ . Arguing similarly, we can define a proper coloring  $\varphi_2$  of  $C_{2n+1}^2$  using colors 1, 2, 3 which is an extension of the restriction of  $\varphi$  to  $C_{2n+1}^2$ . Finally, we obtain an extension of  $\varphi$  by coloring every uncolored edge of M with color 4.

Case 2. M contains at least two precolored edges. For the uncolored edges of  $C_{2n+1} \Box K_2$ , we define a list assignment L by setting

$$L(e) = \{1, 2, 3, 4\} \setminus \{\varphi(e') : e' \in E(C_{2n+1} \Box K_2) \text{ is adjacent to } e\}.$$

If M contains exactly two precolored edges, then exactly one precolored edge is in  $C_{2n+1}^1$  or  $C_{2n+1}^2$ , say  $C_{2n+1}^1$ . Then at most one edge of  $C_{2n+1}^1$  is adjacent to precolored edges of three distinct colors, so at most one edge e satisfies that |L(e)| = 1, and all other edges of  $C_{2n+1}^1$  have lists of size at least 2. Hence by Lemma 13, there is a proper coloring  $\varphi_1$  of  $C_{2n+1}^1$  using colors from the lists. By coloring  $C_{2n+1}^2$  correspondingly, and then coloring the uncolored edges of M, we obtain an extension of  $\varphi$ . If, on the other hand, M contains all three precolored edges, then we can obtain an extension of  $\varphi$  by first coloring the uncolored edges of M, and then apply Lemma 14.

Note that even though any partial 3-coloring of  $C_{2n+1}$  is extendable, the upper bound of three precolored edges in Proposition 16 is best possible. For instance, consider a precoloring where two adjacent edges  $e_1$  and  $e_2$  of  $C_{2n+1}^1$  are precolored 1 and 2, respectively, and the corresponding edges  $e'_1$  and  $e'_2$  of  $C_{2n+1}^2$  are precolored 3 and 4, respectively (using the same notation as in the preceding proof).

Next, we shall verify that Conjecture 1 holds for Class 1 graphs of maximum degree 3.

**Theorem 17.** If G is a Class 1 graph with  $\Delta(G) = 3$  and every partial 3-coloring of at most k < 3 edges in G is extendable, then every partial 4-coloring of at most k + 1 edges in  $G \square K_2$  is extendable.

**Proof.** The case when k = 1 is trivial, so let us assume that k = 2. Let  $G_1$  and  $G_2$  be copies of G in  $G \square K_2$  and M the perfect matching joining vertices of  $G_1$  with corresponding vertices of  $G_2$ .

Consider a partial 3-coloring  $\varphi$  of  $G \Box K_2$  with three precolored edges. Note that we may assume that G is connected, since otherwise we just consider every component of G separately. Next, we prove that G contains no triangle.

If G is a triangle, then we can apply Proposition 16, so we may assume that this is not the case. Thus if G contains a triangle xyzx, then at least one vertex, say x has degree 3. Then we can color an edge incident with x, which is not contained in the triangle, by the color 2, and yz by the color 1. The resulting partial coloring is not extendable to a proper 3-coloring, contradicting that any partial coloring with 2 precolored edges is extendable. Thus G is triangle-free.

In the remaining part of the proof of Theorem 17 we shall consider a large number of different cases. In many of these cases, we shall employ ideas that have been used above. Thus, in several places we just sketch the arguments, rather than giving all the details. As usual, we shall omit the details in the case when no precolored edges are contained in M, since we may proceed as in the proof outline above.

Case 1. M contains exactly one precolored edge. If the color of the precolored edge e of M only appears on e under  $\varphi$ , then the result is trivial. So assume that  $e = u_1 u_2$  is a precolored edge of M, that  $\varphi(e) = 1$ , where  $u_i \in V(G_i)$ , and that at least one other edge is precolored 1 under  $\varphi$ .

If the remaining two precolored edges are in  $G_1$ , then we consider the precoloring  $\varphi'$  of  $G_1$  obtained from the restriction of  $\varphi$  to  $G_1$  by removing color 1 from all edges precolored 1 in  $G_1$ . Then  $\varphi'$  is extendable to a proper 3-coloring using colors 2, 3, 4. Next, we recolor the edges precolored 1 by the color 1, color  $G_2$ correspondingly, and color every edge of M by a color missing at its endpoints to obtain an extension of  $\varphi$ .

Suppose now that both  $G_1$  and  $G_2$  contains precolored edges. If  $d_{G_1}(u_1) = 3$ , then  $d_{G_2}(u_2) = 3$ , and since G - M contains at most two precolored edges and G is triangle-free, there are uncolored corresponding edges  $e_1 \in E(G_1)$  and  $e_2 \in E(G_2)$ , adjacent to e, such that neither  $e_1$ , nor  $e_2$ , is adjacent to an edge of G - M precolored 1. We define a new precoloring  $\varphi'$  from the restriction of  $\varphi$ to G - M by coloring  $e_1$  and  $e_2$  by the color 1. The restrictions of  $\varphi'$  to  $G_1$  and  $G_2$ , respectively, are extendable to proper 3-colorings of  $G_1$  and  $G_2$ , respectively. Next we color  $e_1$  and  $e_2$  by the color 4, e by the color 1, and every uncolored edge of M by a color in  $\{1, 2, 3, 4\}$  missing at its endpoints.

Now assume that  $d_{G_1}(u_1) = 2$ . If there is an uncolored edge  $e_1 \in E(G_1)$ adjacent to e such that neither  $e_1$ , nor the corresponding edge  $e_2 \in E(G_2)$ , is adjacent to an edge of G - M precolored 1, then we proceed as in the preceding paragraph. Otherwise, there are corresponding edges  $e_1 \in E(G_1)$  and  $e_2 \in E(G_2)$  that are adjacent to e and satisfies that  $e_1$  is precolored or adjacent to an edge precolored 1, and  $e_2$  is neither precolored, nor adjacent to a precolored edge of  $G_2$ ; moreover, there are corresponding edges  $e'_1 \in E(G_1)$  and  $e'_2 \in E(G_2)$  adjacent to e that satisfy analogous conditions with the roles of  $G_1$  and  $G_2$  interchanged. Hence, from the restriction of  $\varphi$  to  $G_1$  and  $G_2$ , respectively, we obtain extendable partial 3-colorings by coloring  $e'_1$  and  $e_2$  with colors from  $\{2,3\}$ . Note that in these extensions no edge colored 1 is adjacent to e. Hence,  $\varphi$  is extendable. A similar argument applies when  $d_{G_1}(u_1) = 1$ .

Case 2. M contains exactly two precolored edges. We assume that  $G_1$  contains the third precolored edge  $e_3$ . If all precolored edges of M have the same color, then we proceed as in Case 2 when  $E(G_1) \cup M$  contains all precolored edges. Thus, we assume that two different colors 1 and 2 appear on the precolored edges of M; let  $u_1$  and  $v_1$  be the endpoints of these edges in  $G_1$ , respectively, where  $u_1$ is incident with an edge of M precolored 1.

Suppose first that  $e_3$  is colored by some color appearing on M, say 1. If there is an uncolored edge e' incident with  $u_1$  that is neither incident with  $v_1$ , nor adjacent to  $e_3$ , then we color e' by the color 1 and take an extension of the coloring of e' and  $e_3$  using colors 1, 3, 4. We now proceed as before to obtain an extension of  $\varphi$ . Otherwise, if there is no such edge e', then

- (a)  $d_{G_1}(u_1) = 1$  and  $u_1$  is adjacent to an endpoint of  $e_3$ , or
- (b)  $d_{G_1}(u_1) = 1$  and  $u_1$  and  $v_1$  are adjacent, or
- (c)  $d_{G_1}(u_1) = 2$  and  $u_1$  is adjacent both to  $v_1$  and an endpoint of  $e_3$  (and these vertices are distinct).

If (a) holds, then we may simply take an extension of the restriction of  $\varphi$  to  $G_1$  using colors 1, 3, 4; if (b) or (c) holds, then we color  $u_1v_1$  by color 3 and take an extension of the obtained coloring of  $G_1$  using colors 1, 3, 4. In all cases, it is straightforward that  $\varphi$  is extendable.

Suppose now that  $\varphi(e_3) = 3$ . If there is no uncolored edge incident with  $u_1$  or  $v_1$ , then the result is trivial. Otherwise, if there is such an edge, then we proceed as in the preceding paragraph.

Case 3. M contains exactly three precolored edges. If all three precolored edges of M have the same color, then the result is trivial.

Suppose now that two colors appear on the precolored edges in M, say 1 and 2, and that color 2 appears on only one edge. Denote this edge by  $e = u_1u_2$ , where  $u_1 \in V(G_1)$ . If  $d_{G_1}(u_1) \leq 2$ , then properly color the edges incident with  $u_1$  by 3 and 4. This precoloring of  $G_1$  is extendable to a proper coloring of  $G_1$  using colors 2, 3, 4. We obtain an extension of  $\varphi$  by coloring  $G_2$  correspondingly, coloring  $u_1u_2$  by the color 2 and all other edges of M by the color 1. If  $d_{G_1}(u_1) = 3$ , then there is an edge  $u_1x$  of  $G_1$  that is adjacent to only one precolored edge of M. We

define a precoloring of  $G_1$  by coloring  $u_1x$  by the color 2. This precoloring is extendable to a precoloring of  $G_1$  by colors 2, 3, 4. We now obtain an extension of  $\varphi$  by recoloring  $u_1x$  and proceeding as before.

Suppose now that three colors appear on the edges of M, i.e., that  $u_1u_2$  is colored 1,  $v_1v_2$  is colored 2, and  $w_1w_2$  is colored 3, where  $u_i, v_i, w_i \in V(G_i)$ .

If there exist two distinct vertices  $x, y \in V(G_1) \setminus \{u_1, v_1, w_1\}$  such that x and y can be matched to distinct vertices in  $\{u_1, v_1, w_1\}$  by two independent edges, say  $u_1x$  and  $v_1y$ , then we color these edges by 1 and 2, respectively and take an extension of this precoloring of  $G_1$  using colors 1, 2, 4. Next, we recolor both  $u_1x$  and  $v_1y$  by the color 3, color  $u_1u_2$  by the color 1,  $v_1v_2$  by the color 2 and all other edges of M by the color 3; this yields an extension of  $\varphi$ . Otherwise, if no such edges exist, then  $\{u_1, v_1, w_1\}$  has at most two neighbors outside  $\{u_1, v_1, w_1\}$  in  $G_1$ . Moreover, since  $G_1$  is connected, there must be at least one neighbor of  $\{u_1, v_1, w_1\}$  in  $V(G_1) \setminus \{u_1, v_1, w_1\}$  in  $G_1$ .

Suppose first that  $\{u_1, v_1, w_1\}$  has exactly one neighbor  $x \notin \{u_1, v_1, w_1\}$  in  $G_1$ . If all vertices in  $\{u_1, v_1, w_1\}$  are adjacent to x, then we color  $xu_1$  by 1 and  $xv_1$  by color 4. By assumption, this coloring of  $G_1$  is extendable to a proper coloring of  $G_1$  using colors 1, 2, 4. Since G is triangle-free, it follows that  $\varphi$  is extendable (by recoloring  $xu_1$  by 3).

If only one vertex in  $\{u_1, v_1, w_1\}$  is adjacent to x, say  $u_1$ , then either both  $v_1$  and  $w_1$  are adjacent to  $u_1$ , or both  $u_1$  and  $w_1$  are adjacent to  $v_1$ . In the first case we color  $u_1v_1$  by 4 and  $u_1w_1$  by 2; in the latter case we color  $v_1w_1$  by 4 and  $u_1w_1$  by 2; in the latter case we color  $v_1w_1$  by 4 and  $u_1v_1$  by 3. Both these partial colorings of  $G_1$  are extendable to proper colorings of  $G_1$  using colors 2, 3, 4. Hence,  $\varphi$  is extendable.

Suppose now that two vertices in  $\{u_1, v_1, w_1\}$  are adjacent to x, say  $u_1$  and  $v_1$ . If  $w_1$  is adjacent to both  $u_1$  and  $v_1$ , then we color  $xv_1$  by the color 1, and  $u_1w_1$  by the color 2. This precoloring of  $G_1$  is extendable (using colors 1, 2, 4), and so,  $\varphi$  is extendable. If  $w_1$  is only adjacent to one of  $u_1$  and  $v_1$ , then we may proceed similarly.

Let us finally consider the case when  $\{u_1, v_1, w_1\}$  has exactly two neighbors  $x, y \notin \{u_1, v_1, w_1\}$  in  $G_1$ . Then x and y has only one common neighbor in  $\{u_1, v_1, w_1\}$ , say  $u_1$ . Thus  $v_1$  and  $w_1$  are only adjacent to vertices in  $\{u_1, v_1, w_1\}$ . Since G is triangle-free, we can properly color the edges incident with  $w_1$  and  $v_1$  by two colors from  $\{2, 3, 4\}$  so that no vertex is incident with two edges of the same color. By assumption, there is an extension of the obtained coloring of  $G_1$  using colors 2, 3, 4. Hence,  $\varphi$  is extendable. This completes the proof of the theorem.

Unfortunately, we are not able to prove a corresponding result for Class 2 graphs with maximum degree 3, since we cannot handle the presence of more precolored edges using our method. Nevertheless, we note that since  $\Delta(L(G)) \leq 4$  if G is 3-regular, where L(G) denotes the line graph of G, it follows from

the characterization of non-degree-choosable graphs<sup>1</sup> proved in [5, 10] that one can decide in a polynomial time whether a given partial 4-coloring of a graph with maximum degree 3 is extendable to a proper 4-coloring. In particular, any partial coloring with at most three precolored edges of a subcubic Class 2 graph is extendable (while a precoloring of four edges is obviously not always possible to extend). Thus, subcubic Class 2 graphs constitutes another large family of graphs which admits an Evans-type result. So while there are well-known examples of subcubic Class 1 graphs that do not admit an Evans-type result (see e.g. [8]), such examples do not exist for Class 2 graphs.

Furthermore, let us note that the condition on degree here is best possible since there are 4-regular Class 2 graphs where not every partial coloring of at most 4 edges is extendable to a proper 5-coloring [3]; indeed  $K_5$  is such a graph.

### Acknowledgements

We thank the referees for helpful comments which lead to improvements of the paper.

Casselgren was supported by a grant from the Swedish Research council VR (2017-05077). Petros and Fufa thank the International Science Program in Uppsala, Sweden, for financial support.

#### References

- M.O. Albertson and E.H. Moore, *Extending graph colorings using no extra colors*, Discrete Math. 234 (2001) 125–132. https://doi.org/10.1016/S0012-365X(00)00376-9
- [2] L.D. Andersen and A.J.W. Hilton, *Thank Evans!*, Proc. Lond. Math. Soc. 47 (1983) 507–522. https://doi.org/10.1112/plms/s3-47.3.507
- [3] L.D. Andersen and A.J.W. Hilton, Symmetric latin square and complete graph analogues of the Evans conjecture, J. Combin. Des. 4 (1994) 197–252. https://doi.org/10.1002/jcd.3180020404
- [4] A.S. Asratian, T.M.J. Denley and R. Häggkvist, Bipartite Graphs and their Applications (Cambridge University Press, 1998). https://doi.org/10.1017/CBO9780511984068
- [5] O.V. Borodin, Criterion of chromaticity of a degree prescription, Abstracts of IV All-Union Conference on Theoretical Cybernetics, Novosibirsk (1977) 127–128, in Russian.
- [6] C.J. Casselgren and F.B. Petros, *Edge precoloring extension of trees*, Australas. J. Combin. 81 (2021) 233–244.

<sup>&</sup>lt;sup>1</sup>A graph G is *degree-choosable* if it has an L-coloring whenever L is a list assignment such that  $|L(v)| \ge d_G(v)$  for all  $v \in V(G)$ .

- [7] C.J. Casselgren and F.B. Petros, *Edge precoloring extension of tree* II, Discuss. Math. Graph Theory 44 (2024) 613–637. https://doi.org/10.7151/dmgt.2461
- C.J. Casselgren, K. Markström and L.A. Pham, Edge precoloring extension of hypercubes, J. Graph Theory 95 (2020) 410–444. https://doi.org/10.1002/jgt.22561
- [9] K. Edwards, A. Girão, J. van den Heuvel, R.J. Kang, G.J. Puleo and J.-S. Sereni, Extension from precoloured sets of edges, Electron. J. Combin. 25 (2018) #P3.1. https://doi.org/10.37236/6303
- [10] P. Erdős, A.L. Rubin and H. Taylor, *Choosability in graphs*, in: Proc. West Coast Conference on Combinatorics, Graph Theory and Computing, Congr. Numer. XXVI (1979) 125–157.
- [11] T. Evans, *Embedding incomplete latin squares*, Amer. Math. Monthly **67** (1960) 958–961. https://doi.org/10.2307/2309221
- [12] J. Fiala, NP-completeness of the edge precoloring extension problem on bipartite graphs, J. Graph Theory 43 (2003) 156–160. https://doi.org/10.1002/jgt.10088
- [13] F. Galvin, The list chromatic index of a bipartite multigraph, J. Combin. Theory Ser. B 63 (1995) 153–158. https://doi.org/10.1006/jctb.1995.1011
- [14] A. Girão and R.J. Kang, A precolouring extension of Vizing's theorem, J. Graph Theory 92 (2019) 255–260. https://doi.org/10.1002/jgt.22451
- [15] R. Häggkvist, A solution of the Evans conjecture for latin squares of large size, Colloq. Math. Soc. János Bolyai 18 (1978) 495–513.
- [16] O. Marcotte and P.D. Seymour, *Extending an edge-coloring*, J. Graph Theory 14 (1990) 565–573. https://doi.org/10.1002/jgt.3190140508
- [17] H.J. Ryser, A combinatorial theorem with an application to Latin rectangles, Proc. Amer. Math. Soc. 2 (1951) 550–552.
- [18] B. Smetaniuk, A new construction for Latin squares I. Proof of the Evans conjecture, Ars Combin. 11 (1981) 155–172.

Received 18 April 2023 Revised 29 January 2024 Accepted 29 January 2024 Available online 28 February 2024

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License https://creativecommons.org/licenses/by-nc-nd/4.0/