

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES

BRYAN FREYBERG AND RYAN PETERS

University of Minnesota Duluth
Duluth, MN, USA

e-mail: frey0031@d.umn.edu
pet04060@d.umn.edu

Abstract

Let G be a forest with six edges. We prove that G decomposes the complete graph K_n if and only if $n \equiv 0, 1, 4, \text{ or } 9 \pmod{12}$, unless $n = 9$ and G is one of nine exceptional forests.

Keywords: graph decomposition, forests, ρ -labeling.

2020 Mathematics Subject Classification: 05C51.

1. INTRODUCTION

For some graph G , a G -decomposition of a graph H is a set $\mathcal{G} = \{G_1, G_2, \dots, G_t\}$ of pairwise edge-disjoint subgraphs of H , each of which is isomorphic to G , such that $E(H) = \bigcup_{i=1}^t E(G_i)$. The elements of \mathcal{G} are called G -blocks.

Finding G -decompositions is a classical combinatorial problem that spans the related fields of graph theory, design theory, and error-correcting codes. The problem was given new life in the 1960s by Kotzig and Rosa who produced a new method for decomposing graphs now known as graph labeling.

We only consider G -decompositions of the complete graph K_n , so from here on, we refer to G -decompositions of K_n simply as G -decompositions. Determining the set of all integers n such that a G -decomposition exists is known as the decomposition spectrum for G . We are particularly interested in solving this spectrum problem for all small graphs. The decomposition spectrum for graphs with less than six edges is completely determined. In this article, we complete the decomposition spectrum for graphs with six edges. An overview of the results for graphs with a small number of edges is given in [9], and we reproduce an updated version of it here.

1.1. Graphs with at most four edges

For graphs with one or two edges, the cases are trivial. Graphs with three edges are only K_3 , $K_{1,3}$, P_4 , $P_2 \cup P_3$, or $3K_2$. For K_3 we have a Steiner Triple System, and it has been solved by Kirkman [14]. $K_{1,3}$ was classified by Cain [5], P_4 was settled by Bermond [2], $P_2 \cup P_3$ was solved by Bermond, Huang, Rosa and Sotteau [3], and $3K_2$ was classified by de Werra [6].

There are five types of connected graphs with four edges. Kotzig covered the C_4 case [15], and Bermond and Schönheim the cases containing K_3 (K_3 with a pendant edge and $K_3 \cup P_2$ [4]). The path P_5 and the tree with a unique vertex of degree three were settled by Huang and Rosa [12], and the star $K_{1,4}$ by Yamamoto *et al.* [18].

The forests with six vertices, that is, $2P_3$, $P_4 \cup P_2$ and $K_{1,3} \cup K_2$ were classified by Yin and Gong [19].

The case $P_3 \cup 2P_2$ appears to be folklore, and follows from its σ^+ - or 1-rotational σ^+ -labeling. The matching $4K_2$ was classified by de Werra [6].

1.2. Graphs with five edges

Graphs with five vertices were studied by Bermond, Huang, Rosa and Sotteau [3]. Huang and Rosa [12] covered the cases for all trees with up to nine vertices. Disconnected graphs containing a cycle were settled by Yin and Gong except for $K_3 \cup 2K_2$ [19]. The matching with five edges was solved by de Werra [6], while the remaining forests are treated in [10].

1.3. Graphs with six edges

The only graph on four vertices with six edges is K_4 , which has been covered by Hanani [11]. The graphs on five vertices have been completely settled by Bermond, Huang, Rosa and Sotteau [3] except for $\overline{P_5}$. This case was solved by Kang and Wang [13].

All graphs with six edges and six vertices decompose were settled by Yin and Gong [19].

Graphs with six edges and more than six vertices are either trees (treated in [12]) or disconnected. This leaves only disconnected unicyclic graphs with six edges, which were settled in [1], and forests with six edges. As previously noted, the matching with six edges was solved by de Werra in [6]. We settle the remaining forests with six edges in this article by proving the following theorem.

Theorem 1. *Let G be a forest with exactly six edges. There exists a G -decomposition of K_n if and only if $n > 4$ and $n \equiv 0, 1, 4, \text{ or } 9 \pmod{12}$, unless $n = 9$ and G is one of the nine exceptional forests listed below.*

- $K_{1,5} \cup K_2$
- $K_{1,4} \cup 2K_2$
- $K_{1,4} \cup P_3$
- $K_{1,3} \cup 3K_2$
- $P_4 \cup 3K_2$
- $2P_3 \cup 2K_2$
- $P_3 \cup 4K_2$
- $6K_2$
- $2K_{1,3}$

1.4. Catalog and necessary conditions

Suppose G is a graph with six edges and a G -decomposition exists. Then 6 must divide $|E(K_n)| = \binom{n}{2}$. This implies $n \equiv 0, 1, 4, \text{ or } 9 \pmod{12}$. We will prove these conditions are sufficient, up to a few small exceptions, when G is a forest by using labeling techniques based on Rosa’s in the sections that follow [17].

Rosa’s revolutionary approach to decomposing complete graphs is based on assigning lengths to each edge of the complete graph K_{2m+1} as follows. Let $V(K_{2m+1}) = \{0, 1, \dots, 2m\}$ and define the *length* of edge $uv \in E(K_{2m+1})$ as $\ell(uv) = \min\{|u-v|, 2m+1-|u-v|\}$. This length function partitions the $m(2m+1)$ edges of K_{2m+1} into $2m+1$ edges of each length in $\{1, 2, \dots, m\}$. If G is a graph with m edges, the problem of finding a G -decomposition of K_{2m+1} now reduces to finding an injective assignment of labels $f : V(G) \rightarrow \{0, 1, \dots, 2m\}$ such that $\{\ell(uv) : uv \in E(G)\} = \{1, 2, \dots, m\}$. This is because once such an assignment of labels is found, it may be cyclically increased by 1 modulo $2m+1$, to produce $2m+1$ edge-disjoint copies of G which uses $m(2m+1)$ edges of K_{2m+1} . Since this operation we call *clicking* preserves edge length, we see that no edge has been repeated, and we have therefore constructed a G -design of K_{2m+1} .

The forests with six edges and at least two components are cataloged in Figures 1 and 2. We use the naming convention $G(k; e_1, e_2, \dots, e_k)_t$ to denote the t^{th} forest with k connected components and e_i edges in the i^{th} connected component for $1 \leq i \leq k$. To simplify the names, we will use exponential notation for components of the same size.

2. DECOMPOSITIONS OF K_9

In this section, we characterize all forests with six edges that decompose K_9 . There are nine such forests that do not decompose K_9 .

Lemma 2. *Let G be a forest with exactly six edges. If G is isomorphic to one of the graphs below, then G does not decompose K_9 .*




Figure 1. Forests with six edges and two connected components.

- $G(2; 1, 5)_1$
- $G(2; 2, 4)_1$
- $G(2; 3, 3)_1$
- $G(3; 1^2, 4)_1$
- $G(4; 1^3, 3)_1$
- $G(4; 1^3, 3)_2$
- $G(4; 1^2, 2^2)_1$
- $G(5; 1^4, 2)_1$
- $G(6; 1^6)_1$

Proof. Suppose $\mathcal{G} = \{G_1, G_2, \dots, G_6\}$ is a $G(2, 1, 5)_1$ -decomposition of K_9 . By the pigeonhole principle, there exists a vertex $v \in V(K_9)$ such that v does not appear as the center of the star $K_{1,5}$ in any G_i for $1 \leq i \leq 6$. But then, $\deg(v) \leq 6$ since all other vertices have degree 1. But of course every vertex in $V(K_9)$ has degree 8, so we have a contradiction. Essentially the same argument proves that $G(3; 1^2, 4)_1$ cannot decompose K_9 . If $G \cong G(2; 2, 4)_1$ or $G(2; 3, 3)_1$, the




Figure 2. Forests with six edges and more than two connected components.

decomposition was shown not to exist by Meszka [16].

If G is isomorphic to one of the five remaining forests on the list, then G contains 10 or more vertices, so of course a G -decomposition of any graph with 9 vertices cannot exist. ■

The next theorem shows that the nine forests from Lemma 2 are the only ones that do not decompose K_9 .

Theorem 3. *A forest with exactly six edges decomposes K_9 if and only if it is not isomorphic to one of the graphs below.*

- $G(2; 1, 5)_1$
- $G(2; 2, 4)_1$
- $G(2; 3, 3)_1$
- $G(3; 1^2, 4)_1$
- $G(4; 1^3, 3)_1$
- $G(4; 1^3, 3)_2$

- $G(4; 1^2, 2^2)_1$
- $G(5; 1^4, 2)_1$
- $G(6; 1^6)_1$

Proof. Let G be a forest with six edges. If G is isomorphic to one of the nine graphs listed in the theorem statement, the decomposition does not exist by Lemma 2. For the remaining forests, we construct a G -decomposition as follows.

Let $V(K_9) = \{0, 1, \dots, 8\}$ and assign each edge $uv \in E(K_9)$ a length

$$\ell(uv) = \min\{|u - v|, 9 - |u - v|\}$$

and a color

$$c(uv) = \begin{cases} \text{blue} & \text{if } u + v \equiv 0 \pmod{3}, \\ \text{red} & \text{if } u + v \equiv 1 \pmod{3}, \\ \text{green} & \text{if } u + v \equiv 2 \pmod{3}. \end{cases}$$

Observe that K_9 contains exactly nine edges of each of the lengths $\{1, 2, 3, 4\}$. Furthermore, among the nine edges of each length, there are exactly three of each color.

Let G_1 and G_2 be the blocks in the appropriate subfigure of Figure 3. Click G_1 by 3 to form blocks G_3 and G_5 . Repeat this procedure with G_2 , forming blocks G_4 and G_6 . Notice that clicking by 3 preserves both length and color. Therefore, it suffices for the reader to check that the edges in blocks G_1 together with G_2 fill the color spectrum for each edge length 1, 2, 3 and 4. ■

3. DECOMPOSITIONS OF K_n FOR $n \equiv 0$ OR $1 \pmod{12}$

Freyberg and Tran introduced the following labeling in [8]. Let G be a bipartite graph with m edges and bipartition $V(G) = A \cup B$. A σ^{+-} -labeling of G is an injection $f : V(G) \rightarrow \{0, 1, \dots, 2m - 2\}$ which induces a bijective length function $\ell : E(G) \rightarrow \{1, 2, \dots, m\}$ where $\ell(uv) = f(v) - f(u)$ for every edge $uv \in E(G)$ with $u \in A$ and $v \in B$, and f has the additional property that $f(u) - f(v) \neq m$ for all $u \in A$ and $v \in B$.

Readers familiar with the Rosa-type labelings may recognize that a σ^{+-} -labeling is a restricted ρ -labeling (see [7] for a nice overview). The usefulness of a σ^{+-} -labeling is punctuated by the following.

Theorem 4 (Freyberg and Tran, [8]). *Let G be a bipartite graph with m edges and a σ^{+-} -labeling such that the edge of length m is a pendant edge e . Then there exists a G -decomposition of K_{2mx} and K_{2mx+1} for every positive integer x .*

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 683






Figure 3. Blocks for G -decompositions of K_9 .




(h) $G(2; 3, 3)_2$

(i) $G(2; 3, 3)_3$




(j) $G(3; 1, 1, 4)_2$

(k) $G(3; 1, 1, 4)_3$



(l) $G(3; 1, 2, 3)_1$

(m) $G(3; 1, 2, 3)_2$



(n) $G(3; 2, 2, 2)_3$

Figure 3. (Cont.) Blocks for G -decompositions of K_9 .

This leads to the main result of this section.

Theorem 5. *Let G be a forest with exactly six edges. Then G decomposes K_n whenever $n \equiv 0$ or $1 \pmod{12}$.*

Proof. If G is a matching, then the proof was given in [6]. If not, then the proof is by Theorem 4 and the σ^{+-} labeling of G given in Figures 4 and 5. Red-labeled vertices belong to A while black-labeled vertices belong to B . The length of each edge is displayed in blue. ■

4. DECOMPOSITIONS OF K_n FOR $n \equiv 4$ OR $9 \pmod{12}$ AND $n > 9$

If n is odd, we let $V(K_n) = \{0, 1, \dots, n - 1\}$ and define the length of an edge $uv \in E(K_n)$ as $\ell(uv) = \min\{|u - v|, n - |u - v|\}$, just as in the proof of Theorem 3. If $\ell(uv) = n - |u - v|$, we call uv a *wrap-around edge*. It follows that K_n contains exactly n edges of each length from the set $\{1, 2, \dots, \frac{n-1}{2}\}$. On the other hand, if n is even we let $V(K_n) = \{0, 1, \dots, n - 2\} \cup \{\infty\}$ and for each $uv \in E(K_n)$, define $\ell(uv) = \min\{|u - v|, n - 1 - |u - v|\}$ whenever $u, v \in \{0, 1, \dots, n - 2\}$, and $\ell(uv) = \infty$ otherwise. In this case, K_n has exactly $n - 1$ edges of each length from $\{1, 2, \dots, \frac{n-2}{2}, \infty\}$. We prove the following theorems using this observation of edge length distribution.

Theorem 6. *Let G be a forest with six edges and an integer $n = 12k + 9$ be given. If $k \geq 1$, then there exists a G -decomposition of K_n .*

Proof. Let $G \cong G(2; 1, 5)_1$. Label blocks G_1, G_2, \dots, G_{k+2} as in Figure 29. Notice the blocks have the following properties:

1. The blocks contain no wrap-around edges.
2. G_1 contains exactly three edges of length 1.
3. G_1 contains exactly three edges of length 7.

Let x_0y_0, x_1y_1 , and x_2y_2 be any three edges of the same length in G_1 . Then $x_i + y_i \equiv i \pmod{3}$ (see subscripts of the vertex labels in G_1).

4. $\bigcup_{i=2}^{k+1} G_i$ contains exactly one edge of each length $\{\infty, 2, 3, \dots, 6k + 1\} \setminus \{7\}$.

Clicking $G_1 \cup G_2$ by 3 and G_i by 1 for $3 \leq i \leq k + 2$ yields n edges of each length. Property 1 preserves the length of the edges in $G_1 \cup G_2$, for all i . Property 3 above ensures no edge repeats when clicking by 3. Hence, we have described a G -decomposition of K_n .

The proof follows in the same way for the remaining forests. See Figures 30 through 51 for the labelings. ■




Figure 4. σ^{+-} -labeling of forests with two components.

The proof of the next theorem is very similar.

Theorem 7. *Let G be a forest with six edges and an integer $n = 12k + 4$ be given. If $k \geq 1$, then there exists a G -decomposition of K_n .*

Proof. Let G be a forest with six edges and $G \not\cong G(2; 1, 5)_1$. Label blocks G_1, G_2, \dots, G_{k+1} as in the appropriate figure (see Figures 7 through 28). Notice the blocks have the following properties.




Figure 5. σ^{+-} -labeling of forests with more than two components.

Clicking G_1 by 3 and G_i by 1 for $2 \leq i \leq k + 1$ yields $n - 1$ edges of each length. Property 4 above ensures no edge has been repeated in clicking G_1 by 3. Property 1 preserves the length of the edges G_1 for all k . Hence, we have described a G -decomposition of K_n .

The construction for the remaining forest $G(2; 1, 5)_1$, follows in the same way except that there are four starter blocks instead of two (see Figure 6). Click block G_i by 3 for $1 \leq i \leq 4$, and by 1 for $5 \leq i \leq k + 3$ to obtain the decomposition. ■

5. CONCLUSION

The proof of Theorem 1 follows directly from Theorems 3, 5, 6, and 7. Therefore, we have solved the K_n decomposition spectrum problem for forests with six edges. As a result, we now have a complete answer for which n and G there exists a G -decomposition of K_n where G is any graph with six or less edges.

REFERENCES

- [1] J. Ahern, B. Freyberg, D. Froncek and M. Keranen, *Decompositions of complete graphs into disconnected unicyclic graphs with six edges* (2024), manuscript.
- [2] J.C. Bermond, *Cycles dans les graphes et G -configurations*, PhD Thesis, University of Paris XI (Orsay, 1975).
- [3] J.C. Bermond, C. Huang, A. Rosa and D. Sotteau, *Decomposition of complete graphs into isomorphic subgraphs with five vertices*, *Ars Combin.* **10** (1980) 211–254.
- [4] J.C. Bermond and J. Schönheim, *G -decomposition of K_n , where G has four vertices or less*, *Discrete Math.* **19** (1977) 113–120.
[https://doi.org/10.1016/0012-365X\(77\)90027-9](https://doi.org/10.1016/0012-365X(77)90027-9)
- [5] P. Cain, *Decomposition of complete graphs into stars*, *Bull. Aust. Math. Soc.* **10** (1974) 23–30.
<https://doi.org/10.1017/S0004972700040582>
- [6] D. de Werra, *On some combinatorial problems arising in scheduling*, *Canad. Operational Res. Soc. J.* **8** (1970) 165–175.
- [7] S.I. El-Zanati and C. Vanden Eynden, *On Rosa-type labelings and cyclic graph decompositions*, *Math. Slovaca* **59** (2009) 1–18.
<https://doi.org/10.2478/s12175-008-0108-x>
- [8] B. Freyberg and N. Tran, *Decomposition of complete graphs into bipartite unicyclic graphs with eight edges*, *J. Combin. Math. Combin. Comput.* **114** (2020) 133–142.
- [9] D. Froncek and B. Kubesa, *Decomposition of complete graphs into connected unicyclic bipartite graphs with seven edges*, *Bull. Inst. Combin. Appl.* **93** (2021) 52–80.
- [10] D. Froncek and A. Sailstad, *Decompositions of complete graphs into forests with five edges* (2024), manuscript.
- [11] H. Hanani, *The existence and construction of balanced incomplete block designs*, *Ann. Math. Statist.* **32** (1961) 361–386.
<https://doi.org/10.1214/aoms/1177705047>
- [12] C. Huang and A. Rosa, *Decomposition of complete graphs into trees*, *Ars Combin.* **5** (1978) 23–63.
- [13] Q. Kang and Z. Wang, *Optimal packings and coverings of λK_v with graphs $K_5 - P_5$* , *Bull. Inst. Combin. Appl.* **41** (2004) 22–41.

- [14] T.P. Kirkman, *On a problem in combinations*, Cambridge Dublin Math. J. **2** (1847) 191–204.
- [15] A. Kotzig, *Decomposition of a complete graph into $4k$ -gons*, Mat. Čas. **15** (1965) 229–233, in Russian.
- [16] M. Meszka (2024), personal communication.
- [17] A. Rosa, *On certain valuations of the vertices of a graph*, in: Theory of Graphs, International Symp. Rome 1966 (Dunod Gordon and Breach, Science Publishers, Inc. New York and Dunod Paris, 1967) 349–355.
- [18] S. Yamamoto, H. Ikeda, S. Shige-eda, K. Ushio and N. Hamada, *On claw-decomposition of complete graphs and complete bigraphs*, Hiroshima Math. J. **5** (1975) 33–42.
- [19] J. Yin and B. Gong, *Existence of G -designs with $|V(G)| = 6$* , in: Combinatorial Designs and Applications, W.D. Wallis *et al.* (Ed(s)) (Marcel Dekker, New York, 1990) 201–218.

6. APPENDIX

6.1. Labelings for $n \equiv 4 \pmod{12}$




Figure 6. $G(2; 1, 5)_1$




Figure 7. $G(2; 1, 5)_2$




Figure 8. $G(2; 1, 5)_3$




Figure 9. $G(2; 1, 5)_4$




Figure 10. $G(2; 1, 5)_5$




Figure 11. $G(2; 1, 5)_6$




Figure 12. $G(2; 2, 4)_1$




Figure 13. $G(2; 2, 4)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 693




Figure 14. $G(2; 2, 4)_3$




Figure 15. $G(2; 3^2)_1$




Figure 16. $G(2; 3^2)_2$




Figure 17. $G(2; 3^2)_3$




Figure 18. $G(3; 1^2, 4)_1$




Figure 19. $G(3; 1^2, 4)_2$




Figure 20. $G(3; 1^2, 4)_3$




Figure 21. $G(3; 1, 2, 3)_1$




Figure 22. $G(3; 1, 2, 3)_2$




Figure 23. $G(3; 2^3)_1$




Figure 24. $G(4; 1^3, 3)_1$




Figure 25. $G(4; 1^3, 3)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 697




Figure 26. $G(4; 1^2, 2^2)_1$




Figure 27. $G(5; 1^4, 2)_1$




Figure 28. $G(6; 1^6)_1$

6.2. Labelings for $n \equiv 9 \pmod{12}$

The vertex labels for some of the blocks in the figures in the next two subsections are of the form i_j . The subscript j indicates the modulo 3 equivalence class of i . For example, if $i = 7$, then $j = 1$ since $7 \equiv 1 \pmod{3}$.




Figure 29. $G(2; 1, 5)_1$




Figure 30. $G(2; 1, 5)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 699




Figure 31. $G(2; 1, 5)_3$




Figure 32. $G(2; 1, 5)_4$




Figure 33. $G(2; 1, 5)_5$




Figure 34. $G(2; 1, 5)_6$




Figure 35. $G(2; 2, 4)_1$




Figure 36. $G(2; 2, 4)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 701




Figure 37. $G(2; 2, 4)_3$




Figure 38. $G(2; 3^2)_1$




Figure 39. $G(2; 3^2)_2$




Figure 40. $G(2; 3^2)_3$




Figure 41. $G(3; 1^2, 4)_1$




Figure 42. $G(3; 1^2, 4)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 703




Figure 43. $G(3; 1^2, 4)_3$




Figure 44. $G(3; 1, 2, 3)_1$




Figure 45. $G(3; 1, 2, 3)_2$




Figure 46. $G(3; 2^3)$




Figure 47. $G(4; 1^3, 3)_1$




Figure 48. $G(4; 1^3, 3)_2$

DECOMPOSITION OF COMPLETE GRAPHS INTO FORESTS WITH SIX EDGES 705




Figure 49. $G(4; 1^2, 2^2)_1$




Figure 50. $G(5; 1^4, 2)_1$




Figure 51. $G(6; 1^6)_1$

Received 22 February 2024

Revised 6 June 2024

Accepted 7 June 2024

Available online 18 July 2024