

## CRITICALITY OF SWITCHING CLASSES OF REVERSIBLE 2-STRUCTURES LABELED BY AN ABELIAN GROUP

HOUMEM BELKHECHINE

*University of Carthage  
Institut Préparatoire aux Études d'Ingénieurs de Bizerte, BP 64  
7021 Bizerte, Tunisie*

**e-mail:** houmem@gmail.com

PIERRE ILLE

*Institut de Mathématiques de Marseille  
CNRS UMR 7373, 13453 Marseille, France*

**e-mail:** pierre.ille@univ-amu.fr

AND

ROBERT E. WOODROW

*Department of Mathematics and Statistics  
University of Calgary, 2500 University Drive  
Calgary, Alberta, Canada T2N 1N4*

**e-mail:** woodrow@ucalgary.ca

### Abstract

Let  $V$  be a finite vertex set and let  $(\mathbb{A}, +)$  be a finite abelian group. An  $\mathbb{A}$ -labeled and reversible 2-structure defined on  $V$  is a function  $g : (V \times V) \setminus \{(v, v) : v \in V\} \rightarrow \mathbb{A}$  such that for distinct  $u, v \in V$ ,  $g(u, v) = -g(v, u)$ . The set of  $\mathbb{A}$ -labeled and reversible 2-structures defined on  $V$  is denoted by  $\mathcal{L}(V, \mathbb{A})$ . Given  $g \in \mathcal{L}(V, \mathbb{A})$ , a subset  $X$  of  $V$  is a *clan* of  $g$  if for any  $x, y \in X$  and  $v \in V \setminus X$ ,  $g(x, v) = g(y, v)$ . For example,  $\emptyset$ ,  $V$  and  $\{v\}$  (for  $v \in V$ ) are clans of  $g$ , called *trivial*. An element  $g$  of  $\mathcal{L}(V, \mathbb{A})$  is *primitive* if  $|V| \geq 3$  and all the clans of  $g$  are trivial.

The set of the functions from  $V$  to  $\mathbb{A}$  is denoted by  $\mathcal{S}(V, \mathbb{A})$ . Given  $g \in \mathcal{L}(V, \mathbb{A})$ , with each  $s \in \mathcal{S}(V, \mathbb{A})$  is associated the *switch*  $g^s$  of  $g$  by  $s$  defined as follows: given distinct  $x, y \in V$ ,  $g^s(x, y) = s(x) + g(x, y) - s(y)$ . The *switching class* of  $g$  is  $\{g^s : s \in \mathcal{S}(V, \mathbb{A})\}$ . Given a switching class  $\mathfrak{S} \subseteq \mathcal{L}(V, \mathbb{A})$  and  $X \subseteq V$ ,  $\{g|_{(X \times X) \setminus \{(x, x) : x \in X\}} : g \in \mathfrak{S}\}$  is a switching class, denoted by  $\mathfrak{S}[X]$ .

Given a switching class  $\mathfrak{S} \subseteq \mathcal{L}(V, \mathbb{A})$ , a subset  $X$  of  $V$  is a *clan* of  $\mathfrak{S}$  if  $X$  is a clan of some  $g \in \mathfrak{S}$ . For instance, every  $X \subseteq V$  such that  $\min(|X|, |V \setminus X|) \leq 1$  is a clan of  $\mathfrak{S}$ , called *trivial*. A switching class  $\mathfrak{S} \subseteq \mathcal{L}(V, \mathbb{A})$  is *primitive* if  $|V| \geq 4$  and all the clans of  $\mathfrak{S}$  are trivial. Given a primitive switching class  $\mathfrak{S} \subseteq \mathcal{L}(V, \mathbb{A})$ ,  $\mathfrak{S}$  is *critical* if for each  $v \in V$ ,  $\mathfrak{S} - v$  is not primitive. First, we translate the main results on the primitivity of  $\mathbb{A}$ -labeled and reversible 2-structures in terms of switching classes. For instance, we prove the following. For a primitive switching class  $\mathfrak{S} \subseteq \mathcal{L}(V, \mathbb{A})$  such that  $|V| \geq 8$ , there exist  $u, v \in V$  such that  $u \neq v$  and  $\mathfrak{S}[V \setminus \{u, v\}]$  is primitive. Second, we characterize the critical switching classes by using some of the critical digraphs described in [Y. Boudabous and P. Ille, *Indecomposability graph and critical vertices of an indecomposable graph*, Discrete Math. **309** (2009) 2839–2846].

**Keywords:** labeled and reversible 2-structure, switching class, clan, primitivity, criticality.

**2010 Mathematics Subject Classification:** 05C75.

#### REFERENCES

- [1] P. Bonizzoni, *Primitive 2-structures with the  $(n - 2)$ -property*, Theoret. Comput. Sci. **132** (1994) 151–178.  
doi:10.1016/0304-3975(94)90231-3
- [2] Y. Boudabous and P. Ille, *Indecomposability graph and critical vertices of an indecomposable graph*, Discrete Math. **309** (2009) 2839–2846.  
doi:10.1016/j.disc.2008.07.015
- [3] Y. Cheng and A.L. Wells, *Switching classes of directed graphs*, J. Combin. Theory Ser. B **40** (1986) 169–186.  
doi:10.1016/0095-8956(86)90075-4
- [4] A. Cournier and P. Ille, *Minimal indecomposable graph*, Discrete Math. **183** (1998) 61–80.  
doi:10.1016/S0012-365X(97)00077-0
- [5] A. Ehrenfeucht, T. Harju and G. Rozenberg, *The Theory of 2-Structures, A Framework for Decomposition and Transformation of Graphs* (World Scientific, 1999).  
doi:10.1142/4197
- [6] A. Ehrenfeucht and G. Rozenberg, *Theory of 2-structures, Part I: clans, basic subclasses, and morphisms*, Theoret. Comput. Sci. **70** (1990) 277–303.  
doi:10.1016/0304-3975(90)90129-6
- [7] A. Ehrenfeucht and G. Rozenberg, *Primitivity is hereditary for 2-structures*, Theoret. Comput. Sci. **70** (1990) 343–358.
- [8] P. Ille, *Recognition problem in reconstruction for decomposable relations*, in: Finite and Infinite Combinatorics in Sets and Logic, B. Sands, N. Sauer and R. Woodrow (Ed(s)), (Kluwer Academic Publishers, 1993) 189–198.  
doi:10.1007/978-94-011-2080-7\_13

- [9] P. Ille, *Indecomposable graphs*, Discrete Math. **173** (1997) 71–78.  
doi:10.1016/S0012-365X(96)00097-0
- [10] P. Ille and R. Woodrow, *Decomposition tree of a lexicographic product of binary structures*, Discrete Math. **311** (2011) 2346–2358.  
doi:10.1016/j.disc.2011.05.037
- [11] J.H. Schmerl and W.T. Trotter, *Critically indecomposable partially ordered sets, graphs, tournaments and other binary relational structures*, Discrete Math. **113** (1993) 191–205.  
doi:10.1016/0012-365X(93)90516-V
- [12] J.J. Seidel, *Strongly regular graphs of L2 type and of triangular type*, in: Proc. Kon. Nederl. Akad. Wetensch. Ser. A (Indag. Math. **29**, 1967) 188–196.  
doi:10.1016/S1385-7258(67)50031-8

Received 27 September 2015

Revised 29 March 2016

Accepted 29 March 2016