

SOME REMARKS ON α -DOMINATION

FRANZ DAHME

Lehrstuhl II für Mathematik, RWTH-Aachen
52056 Aachen, Germany

e-mail: franz.dahme@web.de

DIETER RAUTENBACH

Forschungsinstitut für Diskrete Mathematik
Lennéstr. 2, D-53113 Bonn, Germany

e-mail: rauten@or.uni-bonn.de

AND

LUTZ VOLKMANN

Lehrstuhl II für Mathematik, RWTH-Aachen
52056 Aachen, Germany

e-mail: volkm@math2.rwth-aachen.de

Abstract

Let $\alpha \in (0, 1)$ and let $G = (V_G, E_G)$ be a graph. According to Dunbar, Hoffman, Laskar and Markus [3] a set $D \subseteq V_G$ is called an α -dominating set of G , if $|N_G(u) \cap D| \geq \alpha d_G(u)$ for all $u \in V_G \setminus D$. We prove a series of upper bounds on the α -domination number of a graph G defined as the minimum cardinality of an α -dominating set of G .

Keywords: α -domination; domination.

2000 Mathematics Subject Classification: 05C69.

1. Introduction

Let $\alpha \in (0, 1)$ and let $G = (V_G, E_G)$ be a simple and finite graph. The neighbourhood and the degree of a vertex $u \in V_G$ in the graph G are denoted by $N_G(u)$ and $d_G(u)$, respectively. For all further notation and terminology we refer the reader to [5].

In [3] Dunbar, Hoffman, Laskar and Markus introduce the concept of an α -dominating set. They define a set $D \subseteq V_G$ to be an α -dominating set of G , if $|N_G(u) \cap D| \geq \alpha d_G(u)$ for all $u \in V_G \setminus D$, i.e., for every vertex that does not belong to D at least the α -fraction of its neighbours belongs to D .

In the present paper we prove a series of upper bounds on the α -domination number $\gamma_\alpha(G)$ of a graph G defined as the minimum cardinality of an α -dominating set of G . For further results on this parameter we refer the reader to [3] and [6].

2. Results

Our first bound follows easily from a colouring result of Cowen and Emerson (cf. Theorem 5.1 in [9]). In order to make our presentation self-contained we include a short proof using a classical Erdős-type exchange argument.

Theorem 2.1. *If $\alpha \in (0, 1)$ and G is a graph of order n , then $\gamma_\alpha(G) \leq (1 - \frac{1}{\lceil \frac{1}{1-\alpha} \rceil})n$.*

Proof. Let $r = \lceil \frac{1}{1-\alpha} \rceil$ and let $V_G = V_1 \cup V_2 \cup \dots \cup V_r$ be a partition of V_G such that $\sum_{i=1}^r \sum_{u \in V_i} |N_G(u) \cap V_i|$ is minimum. Note that $r(1-\alpha) \geq 1$.

If $|N_G(u_0) \cap V_{i_0}| > (1-\alpha)d_G(u_0)$ for some $i_0 \in \{1, 2, \dots, r\}$ and some $u_0 \in V_{i_0}$, then there is some index i_1 with $1 \leq i_1 \leq r$ such that $|N_G(u_0) \cap V_{i_1}| < (1-\alpha)d_G(u_0)$. If $V'_{i_0} = V_{i_0} \setminus \{u_0\}$, $V'_{i_1} = V_{i_1} \cup \{u_0\}$ and $V'_l = V_l$ for $1 \leq l \leq r$ with $l \notin \{i_0, i_1\}$, then

$$\begin{aligned} \sum_{i=1}^r \sum_{u \in V'_i} |N_G(u) \cap V'_i| &= \sum_{i=1}^r \sum_{u \in V_i} |N_G(u) \cap V_i| - 2|N_G(u_0) \cap V_{i_0}| + 2|N_G(u_0) \cap V_{i_1}| \\ &< \sum_{i=1}^r \sum_{u \in V_i} |N_G(u) \cap V_i| \end{aligned}$$

which is a contradiction.

For each $i \in \{1, 2, \dots, r\}$ we therefore have $|N_G(u) \cap V_i| \leq (1 - \alpha)d_G(u)$ for all $u \in V_i$, which implies that $V_G \setminus V_i$ is an α -dominating set of G . Since $\max\{|V_i| \mid 1 \leq i \leq r\} \geq \frac{n}{r}$, the desired bound follows. ■

The following corollary is actually equivalent to Theorem 2.1.

Corollary 2.2. *Let $i \in \mathbf{N}_0 = \{0, 1, 2, \dots\}$ and let $\alpha \in \mathbf{R}$ be such that $\frac{i}{i+1} < \alpha \leq \frac{i+1}{i+2}$. If G is a graph of order n , then $\gamma_\alpha(G) \leq \frac{i+1}{i+2}n < \frac{n}{2-\alpha}$.*

Proof. If $\frac{i}{i+1} < \alpha \leq \frac{i+1}{i+2}$ for some $i \in \mathbf{N}_0$, then $i + 1 < \frac{1}{1-\alpha} \leq i + 2$ and $\frac{i+1}{i+2} < \frac{1}{2-\alpha}$. These inequalities easily imply the desired result. ■

Corollary 2.3 (Corollary 10 in [3]). *If $\alpha \in (0, \frac{1}{2}]$ and G is a graph of order n , then $\gamma_\alpha(G) \leq \frac{n}{2}$.*

The following observation shows that the bound given in Theorem 2.1 and Corollary 2.2 is essentially best-possible.

Observation 2.4. *Let $i \in \mathbf{N}_0 = \{0, 1, 2, \dots\}$ and let $\alpha \in \mathbf{R}$ be such that $\frac{i}{i+1} < \alpha \leq \frac{i+1}{i+2}$. For every $\epsilon > 0$, there is a connected graph G of order n such that $\gamma_\alpha(G) \geq (\frac{i+1}{i+2} - \epsilon)n$.*

Proof. For $l \geq 1$ let the connected graph $G(l)$ arise from the disjoint union of l cliques H_1, H_2, \dots, H_l of order $i + 2$ and one isolated vertex v by joining v and one vertex u_j of H_j by a new edge for $1 \leq j \leq l$.

Let D be a minimum α -dominating set of $G(l)$. If $V_{H_j} \setminus \{u_j\} \subseteq D$ for some $1 \leq j \leq l$, then $|D \cap V_{H_j}| \geq i + 1$. If $V_{H_j} \setminus \{u_j\} \not\subseteq D$ for some $1 \leq j \leq l$, then there is a vertex $u'_j \in V_{H_j} \setminus (\{u_j\} \cup D)$ with $d_{G(l)}(u'_j) = i + 1$ and thus $|D \cap V_{H_j}| \geq \lceil \alpha d_{G(l)}(u'_j) \rceil = \lceil \alpha(i + 1) \rceil = i + 1$. Altogether, we obtain $\gamma_\alpha(G(l)) \geq l(i + 1)$ and thus $\frac{\gamma_\alpha(G(l))}{n_{G(l)}} \geq \frac{l(i+1)}{l(i+2)+1}$. Since $\lim_{l \rightarrow \infty} \frac{l(i+1)}{l(i+2)+1} = \frac{i+1}{i+2}$, the desired result follows. ■

Note that the graphs we considered for the proof of Observation 2.4 were connected but not 2-connected. In fact, we believe the following.

Conjecture 2.5. *If $\alpha \in (0, 1)$ and G is a 2-(vertex-)connected graph of order n , then $\gamma_\alpha(G) \leq \lceil \alpha(n - 1) \rceil$.*

Using a probabilistic argument, our next result shows that the bound in Conjecture 2.5, i.e., $\approx \alpha n$, is essentially the right bound for graphs with a sufficiently large minimum degree.

Theorem 2.6. *For $\alpha \in (0, 1)$ and $\epsilon > 0$ there is some constant $\delta_{\alpha, \epsilon}$ depending only on α and ϵ such that $\gamma_\alpha(G) \leq (\alpha + \epsilon)n$ for all graphs G of order n with minimum degree at least $\delta_{\alpha, \epsilon}$.*

Proof. Let G be a graph of order n and minimum degree δ . Without loss of generality, we may assume that $\alpha + \frac{\epsilon}{2} \leq 1$. Independently for each vertex $u \in V_G$ we introduce a random indicator variable Y_u such that $\Pr[Y_u = 1] = \alpha + \frac{\epsilon}{2}$ and $\Pr[Y_u = 0] = 1 - (\alpha + \frac{\epsilon}{2})$. Let $D_0 = \{u \in V_G \mid Y_u = 1\}$ and

$$D_1 = \{u \in V_G \mid Y_u = 0 \text{ and } |N_G(u) \cap D_0| < \alpha d_G(u)\}.$$

Clearly, $D_0 \cup D_1$ is an α -dominating set of G . The expected value of $|D_0|$ is $\mathbb{E}[|D_0|] = (\alpha + \frac{\epsilon}{2})n$. It remains to estimate the expected value $\mathbb{E}[|D_1|]$.

For $u \in V_G$ let $Z_u = |N_G(u) \cap D_0| = \sum_{v \in N_G(u)} Y_v$, i.e., Z_u is the sum of mutually independent indicator variables. Clearly, $\mathbb{E}[Z_u] = (\alpha + \frac{\epsilon}{2})d_G(u)$. Using Corollary A.1.14 in [1] – a well-known bound of large deviation whose origin goes back to Chernoff [2] – we obtain that there is some constant $c(\alpha, \epsilon) > 0$ depending only on α and ϵ such that

$$\begin{aligned} \Pr[|N_G(u) \cap D_0| < \alpha d_G(u)] &= \Pr[Z_u < \alpha d_G(u)] \\ &= \Pr\left[\left(\alpha + \frac{\epsilon}{2}\right) d_G(u) - Z_u > \frac{\epsilon}{2} d_G(u)\right] \\ &\leq \Pr\left[\left|Z_u - \left(\alpha + \frac{\epsilon}{2}\right) d_G(u)\right| > \frac{\epsilon}{2} d_G(u)\right] \\ &= \Pr\left[|Z_u - \mathbb{E}[Z_u]| > \frac{\epsilon}{2\alpha + \epsilon} \mathbb{E}[Z_u]\right] \\ &\leq 2e^{-c(\alpha, \epsilon)\mathbb{E}[Z_u]} \\ &= 2e^{-c(\alpha, \epsilon)(\alpha + \frac{\epsilon}{2})d_G(u)}. \end{aligned}$$

Therefore,

$$\begin{aligned} E[|D_1|] &= \sum_{u \in V_G} \Pr[u \in D_1] \\ &= \sum_{u \in V_G} \Pr[Y_u = 0] \cdot \Pr[|N_G(u) \cap D_0| < \alpha d_G(u)] \\ &\leq \sum_{u \in V_G} \left(1 - \left(\alpha + \frac{\epsilon}{2}\right)\right) 2e^{-c(\alpha, \epsilon)(\alpha + \frac{\epsilon}{2})d_G(u)} \\ &\leq n \left(1 - \left(\alpha + \frac{\epsilon}{2}\right)\right) 2e^{-c(\alpha, \epsilon)(\alpha + \frac{\epsilon}{2})\delta}. \end{aligned}$$

Since (for fixed α and ϵ and hence also fixed $c(\alpha, \epsilon)$)

$$\lim_{\delta \rightarrow \infty} \left(1 - \left(\alpha + \frac{\epsilon}{2}\right)\right) 2e^{-c(\alpha, \epsilon)(\alpha + \frac{\epsilon}{2})\delta} = 0,$$

the desired result follows. ■

We proceed to some best-possible bounds for bipartite graphs and for cacti, i.e., graphs all cycles of which are edge-disjoint.

Proposition 2.7. *Let $\alpha \in (0, 1)$.*

- (i) *If G is a bipartite graph of order n , then $\gamma_\alpha(G) \leq \frac{1}{2}n$.*
- (ii) *If G is a non-bipartite cactus of order n and the length of a shortest odd cycle in G is at least g_{odd} , then $\gamma_\alpha(G) \leq \frac{1}{2}\left(1 + \frac{1}{g_{\text{odd}}}\right)n$.*

Proof. (i) Since the smaller partite set of G is an α -dominating set of G , the result is obvious.

(ii) Let G be a non-bipartite cactus of order $n \geq 2$ such that the length of a shortest odd cycle in G is at least g_{odd} . We may assume, without loss of generality, that G is connected. We prove the bound by induction on the number b of blocks of G .

First we assume that $b = 1$, i.e., G is an odd cycle of length at least g_{odd} . This implies

$$\gamma_\alpha(G) = \left\lceil \frac{n}{2} \right\rceil = \frac{n+1}{2} \leq \frac{1}{2} \left(1 + \frac{1}{g_{\text{odd}}}\right) n.$$

Now let $b > 1$ and let $B = (V_B, E_B)$ be an endblock of G , i.e. G has exactly one cutvertex v . As above, we see that there is an α -dominating set D_B of B such that $|D_B| \leq \frac{1}{2}(1 + \frac{1}{g_{\text{odd}}})|V_B|$ and $v \in D_B$.

Let H_1, H_2, \dots, H_l be the components of $G - V_B$. Clearly, H_i is a cactus having less blocks than G such that H_i is either bipartite or the length of a shortest odd cycle in H_i is at least g_{odd} for $1 \leq i \leq l$. By (i) and the induction hypothesis, there is an α -dominating set D_i of H_i such that $|D_i| \leq \frac{1}{2}(1 + \frac{1}{g_{\text{odd}}})|V_{H_i}|$ for $1 \leq i \leq l$.

Since $v \in D_B$, it follows that $D = D_B \cup D_1 \cup D_2 \cup \dots \cup D_l$ is an α -dominating set of G such that $|D| \leq \frac{1}{2}(1 + \frac{1}{g_{\text{odd}}})(|V_B| + |V_{H_1}| + |V_{H_2}| + \dots + |V_{H_l}|) = \frac{1}{2}(1 + \frac{1}{g_{\text{odd}}})n$ and the proof is complete. \blacksquare

The term $\frac{n}{2}$ in Corollary 2.3 and Proposition 2.7 is a well-known upper bound for the (ordinary) domination number $\gamma(G)$ of a graph G of order n without isolated vertices [7].

The graphs for which the equality $\gamma(G) = \frac{n}{2}$ holds have been well-studied and characterized in [4] and [8]. The main obstacle for proving an analogous characterization for the α -domination number is the fact that — in contrast to the (ordinary) domination number — this parameter has no monotonicity with respect to spanning subgraphs, i.e., if $G = (V_G, E_G)$ and $H = (V_H, E_H)$ are graphs such that $V_G = V_H$ and $E_H \subseteq E_G$, then $\gamma(H) \geq \gamma(G)$ but no such estimate holds for γ_α . As an example consider the star $K_{1,5}$ of order 6, the complete graph K_6 of order 6 and the tree E that arises by subdividing two edges of $K_{1,3}$. Clearly, $K_{1,5}$ and E are two spanning subgraphs of K_6 but $1 = \gamma_{\frac{2}{5}}(K_{1,5}) < 2 = \gamma_{\frac{2}{5}}(K_6) < 3 = \gamma_{\frac{2}{5}}(E)$.

We close with a corresponding characterization for trees.

Theorem 2.8. *If $\alpha \in (0, 1)$ and T is a tree of order n , then $\gamma_\alpha(T) = \frac{n}{2}$ if and only if T has a perfect matching M such that $\min\{d_T(u), d_T(v)\} < \lceil \frac{1}{1-\alpha} \rceil$ for all $uv \in M$.*

Proof. We first prove the ‘if’-part. Let T be a tree that has a perfect matching M such that $\min\{d_T(u), d_T(v)\} < \lceil \frac{1}{1-\alpha} \rceil$ for all $uv \in M$. Let D be a minimum α -dominating set of T . If there is some edge $u_0v_0 \in M$ such that $u_0, v_0 \notin D$, then $d_T(u_0) - 1 \geq |N_G(u_0) \cap D| \geq \alpha d_T(u_0)$ and $d_T(v_0) - 1 \geq |N_G(v_0) \cap D| \geq \alpha d_T(v_0)$. This implies the contradiction $\min\{d_T(u_0), d_T(v_0)\} \geq \lceil \frac{1}{1-\alpha} \rceil$. Therefore, $|\{u, v\} \cap D| \geq 1$ for all $uv \in M$.

In view of Proposition 2.7, this implies

$$\frac{n}{2} = |M| \leq \sum_{uv \in M} |\{u, v\} \cap D| = |D| = \gamma_\alpha(T) \leq \frac{n}{2}$$

and the proof of the ‘if’-part is complete.

Now we prove the ‘only if’-part by induction on the order n of T . If $n \leq 2$, then the result is immediate. Now let T be a tree of order $n \geq 3$ such that $\gamma_\alpha(T) = \frac{n}{2}$. Let u be an endvertex of T and let v be the unique neighbour of u in T .

If v is adjacent to an endvertex u' of T different from u , then let $T' = T - \{u\}$. It is obvious that there is a minimum α -dominating set D' of T' that contains v . Therefore, D' is also an α -dominating set of T and we obtain the contradiction $\gamma_\alpha(T) \leq \gamma_\alpha(T') \leq \frac{n-1}{2}$. Hence u is the unique endvertex of T adjacent to v and all components T_1, T_2, \dots, T_l of $T - \{u, v\}$ are trees of order at least two. For $1 \leq i \leq l$ let D_i be a minimum α -dominating set of T_i . Since the set $\{v\} \cup D_1 \cup D_2 \cup \dots \cup D_l$ is an α -dominating set of T , we obtain

$$\frac{n-2}{2} = \gamma_\alpha(T) - 1 \leq \sum_{i=1}^l |D_i| = \sum_{i=1}^l \gamma_\alpha(T_i) \leq \sum_{i=1}^l \frac{|V_{T_i}|}{2} = \frac{n-2}{2}.$$

This easily implies that $\gamma_\alpha(T_i) = \frac{|V_{T_i}|}{2}$ and, by induction, the tree T_i has a perfect matching M_i for $1 \leq i \leq l$. Clearly, $M = \{uv\} \cup M_1 \cup M_2 \cup \dots \cup M_l$ is a perfect matching of T .

For contradiction we assume that $\min\{d_T(u_0), d_T(v_0)\} \geq \lceil \frac{1}{1-\alpha} \rceil$ for some edge $u_0v_0 \in M$. Since T is a tree, it is straightforward to verify that the set

$$D = \{w \in V_T \mid \text{dist}_G(w, u_0) \text{ is odd and } \text{dist}_G(w, u_0) < \text{dist}_G(w, v_0)\} \\ \cup \{w \in V_T \mid \text{dist}_G(w, v_0) \text{ is odd and } \text{dist}_G(w, v_0) < \text{dist}_G(w, u_0)\}$$

is an α -dominating set of T such that $u_0, v_0 \notin D$ and $|\{u, v\} \cap D| = 1$ for all $uv \in M \setminus \{u_0v_0\}$. This leads to the contradiction $\gamma_\alpha(T) \leq |D| = \frac{n-2}{2}$ and the proof is complete. ■

Note that $\lceil \frac{1}{1-\alpha} \rceil = 2$ for $\alpha \in (0, \frac{1}{2}]$. Thus for these values of α the trees described by the condition given in Theorem 2.8 correspond exactly to the trees T of order n that satisfy $\gamma(T) = \frac{n}{2}$ (every non-endvertex of these trees is adjacent to a unique endvertex).

References

- [1] N. Alon and J. Spencer, *The probabilistic method*, 2nd ed., (Wiley-Interscience Series in Discrete Math. and Optimization, 2000).
- [2] H. Chernoff, *A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations*, *Ann. Math. Stat.* **23** (1952) 493–507.
- [3] J.E. Dunbar, D.G. Hoffman, R.C. Laskar and L.R. Markus, *α -domination*, *Discrete Math.* **211** (2000) 11–26.
- [4] J.F. Fink, M.S. Jacobson, L.F. Kinch and J. Roberts, *On graphs having domination number half their order*, *Period. Math. Hungar.* **16** (1985) 287–293.
- [5] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of domination in graphs* (Marcel Dekker, New York, 1998).
- [6] F. Dahme, D. Rautenbach and L. Volkmann, *α -domination perfect trees*, manuscript (2002).
- [7] O. Ore, *Theory of Graphs*, Amer. Math. Soc. Colloq. Publ., 38 (Amer. Math. Soc., Providence, RI, 1962).
- [8] C. Payan and N.H. Xuong, *Domination-balanced graphs*, *J. Graph Theory* **6** (1982) 23–32.
- [9] D.R. Woodall, *Improper colourings of graphs*, *Pitman Res. Notes Math. Ser.* **218** (1988) 45–63.

Received 31 March 2003
Revised 12 December 2003