

## PACKING OF THREE COPIES OF A DIGRAPH INTO THE TRANSITIVE TOURNAMENT

MONIKA PILŚNIAK\*

*Faculty of Applied Mathematics AGH*  
*Department of Discrete Mathematics*  
*al. Mickiewicza 30, 30-059 Kraków, Poland*  
**e-mail:** pilsniak@uci.agh.edu.pl

### Abstract

In this paper, we show that if the number of arcs in an oriented graph  $\vec{G}$  (of order  $n$ ) without directed cycles is sufficiently small (not greater than  $\frac{2}{3}n - 1$ ), then there exist arc disjoint embeddings of three copies of  $\vec{G}$  into the transitive tournament  $TT_n$ . It is the best possible bound.

**Keywords:** packing of digraphs, transitive tournament.

**2000 Mathematics Subject Classification:** 05C70, 05C35.

### 1. Introduction. Results

Let  $\vec{G}$  be a digraph of order  $n$  with the vertex set  $V(\vec{G})$  and the arc set  $E(\vec{G})$ . A digraph  $\vec{G}$  is called *transitive* when it satisfies the condition of transitivity: if  $(u, v)$  and  $(v, w)$  are two arcs of  $\vec{G}$  then  $(u, w)$  is the arc, too. For any vertex  $v \in V(\vec{G})$  let us denote by  $d^+(v)$  the *outdegree* of  $v$ , i.e., the number of vertices of  $\vec{G}$  that are adjacent from  $v$ . By  $d^-(v)$  we denote the *indegree* of  $v$ , i.e., the number of vertices adjacent to  $v$ . The *degree* of a vertex  $v$ , denoted by  $d(v)$ , is the sum  $d(v) = d^-(v) + d^+(v)$ . A digraph without directed cycles of length two is called an *oriented graph*. Replacing every arc  $(u, v)$  in an oriented graph  $\vec{G}$  by an edge  $uv$  yields its *underlying graph*.

---

\*The research partly supported by KBN grant 2 P03A 016 18.

A *tournament* is an oriented graph such that its underlying graph is complete. A transitive tournament of order  $n$  will be denoted by  $TT_n$ . As it is unique up to isomorphism, throughout the paper, we will view  $TT_n$  as shown in Figure 1. And we can denote the vertices in  $TT_n$  by consecutive integers in such way that if  $i < j$ , then  $(i, j)$  is an arc of  $TT_n$ . The vertices 1, 2 and  $n$  will be called the *first*, the *second* and the *last vertex* of  $TT_n$ , respectively. We define the *length of an arc*  $(i, j)$  as the difference  $j - i$ .




Figure 1: Transitive tournament  $TT_n$

An (*oriented*) *path* between two distinct vertices  $u$  and  $v$  in an oriented graph  $\vec{G}$  is a finite sequence

$$u = v_0, v_1, \dots, v_{k-1}, v_k = v$$

of vertices, beginning with  $u$  and ending with  $v$  and edges  $v_{i-1}v_i \in E(\vec{G})$  for  $i \in \{1, \dots, k\}$ . A *semipath* between two distinct vertices  $u$  and  $v$  is a path between  $u$  and  $v$  in the underlying graph  $G$ .

A vertex  $x \in V(\vec{G})$  is an *end-vertex* if its degree  $d(x) = 1$ . An arc beginning or ending in  $x$  we call an *end-arc*.

Let  $u$  and  $v$  be end-vertices. The arcs  $u'u, v'v$  (or  $uu', vv'$ ) are called *independent* when  $u' \neq v'$ .

Let  $\vec{G}(V, E)$  be an oriented graph of order  $n$ . An *embedding of  $\vec{G}$  into  $TT_n$*  is a couple  $(\sigma, \sigma')$  in which  $\sigma$  is a bijection  $V \rightarrow \{1, \dots, n\} = V(TT_n)$  and  $\sigma'$  is an injection  $E \rightarrow E(TT_n)$  induced by  $\sigma$  (i.e., for any edge  $ij \in E$ ,  $\sigma'(ij) = \sigma(i)\sigma(j)$ ). We will speak more simply of the embedding  $\sigma$  of  $\vec{G}$ . If  $V(\vec{G}) = k < n$  we can also speak about an embedding of  $\vec{G}$  by adding  $(n - k)$  isolated points to  $\vec{G}$  and we say that  $\vec{G}$  is embeddable into  $TT_n$  if  $\vec{G}' := \vec{G} \cup \{\text{isolated vertices}\}$  is embeddable.

A *k-packing* of  $k$  oriented graphs  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_k$  of order  $n$  into  $TT_n$  is a  $k$ -tuple  $(\sigma_1, \dots, \sigma_k)$  in which  $\sigma_i$  is an embedding of  $\vec{G}_i$  for  $1 \leq i \leq k$  such that the  $k$  sets  $\sigma'_i(E_i)$  are disjoint.

We say that  $\vec{G}$  is  $k$ -packable into  $TT_n$  if a packing of  $k$  copies of  $\vec{G}$  into  $TT_n$  exists.

There are many results concerning packing of graphs. The basic result was proved, independently, in [2], [3] and [6].

**Theorem 1.** *Let  $G, H$  be graphs of order  $n$ . If  $|E(G)| \leq n-2$  and  $|E(H)| \leq n-2$  then  $G$  and  $H$  are packable into  $K_n$ .*

B. Bollobás and S.E. Eldridge made the following conjecture.

**Conjecture 2.** *Let  $G_1, G_2, \dots, G_k$  be  $k$  graphs of order  $n$ . If  $|E(G_i)| \leq n-k$ ,  $i = 1, \dots, k$ , then  $G_1, G_2, \dots, G_k$  are packable into  $K_n$ .*

The case  $k = 3$  of Conjecture 2 was proved by H. Kheddouci, S. Marshall, J.F. Saclé and M. Woźniak in [5].

If one restrains the study to the packing of three copies of the same graph, the hypothesis on size can slightly improved. The following theorem was proved in [7].

**Theorem 3.** *Let  $G$  be a graph of order  $n$ ,  $G \neq K_3 \cup 2K_1$ ,  $G \neq K_4 \cup 4K_1$ . If  $|E(G)| \leq n-2$ , then a 3-packing of  $G$  into  $K_n$  exists.*

The main result of this paper is similar to the basic result of Conjecture 2 for case  $k = 3$  but for an acyclic digraph and its 3-packing into  $TT_n$ .

The motivation for us is the paper by A. Görlich, M. Pilśniak, M. Woźniak [4] where the existence of a 2-packing of  $\vec{G}$  into  $TT_n$  was shown. More precisely, the following result was proved therein.

**Theorem 4.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$  such that  $|E(\vec{G})| \leq \frac{3(n-1)}{4}$ . Then  $\vec{G}$  is 2-packable into  $TT_n$ .*

The basic references of studies addressing packing problems can be found in [1, 8, 9, 10].

## 2. Some Lemmas

Before starting the proof of the main theorem we need some preliminary lemmas.

**Lemma 5.** *Let  $\vec{G}$  be a digraph isomorphic to a path of length  $k$ . If  $k = \lfloor \frac{2}{3}n - 1 \rfloor$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .*

**Proof.** Notice that for  $n \leq 3$  the length of a path is zero or one and it is clear that it is 3-packable into  $TT_n$ .

We use induction on the order of the transitive tournament. For  $n = 4$  the length of a path  $\vec{P}$  is one, let  $\vec{P} = v_0, v_1$ . We can define its embedding  $\sigma_1(v_0) = 1$  and  $\sigma_1(v_1) = 4$  in  $TT_4$  and the embeddings  $\sigma_2$  and  $\sigma_3$  as follows:  $\sigma_2(v_0) = 2$ ,  $\sigma_3(v_0) = 3$  and  $\sigma_2(v_1) = 3$ ,  $\sigma_3(v_1) = 4$ .

For  $n = 5$  the length of a path  $\vec{P}$  is two, let  $\vec{P} = v_0, v_1, v_2$ . We can define its embedding  $\sigma_1(v_0) = 3$ ,  $\sigma_1(v_1) = 4$  and  $\sigma_1(v_2) = 5$  in  $TT_5$  and the embeddings  $\sigma_2$  and  $\sigma_3$  as follows:  $\sigma_2(v_0) = \sigma_3(v_0) = 1$ , and  $\sigma_2(v_1) = 2$ ,  $\sigma_3(v_1) = 3$ , and  $\sigma_2(v_2) = 4$ ,  $\sigma_3(v_2) = 5$ .

For  $n = 6$  the length of a path  $\vec{P}$  is three, let  $\vec{P} = v_0, v_1, v_2, v_3$ . We can define its embedding  $\sigma_1(v_0) = 1$ ,  $\sigma_1(v_1) = 4$ ,  $\sigma_1(v_2) = 5$  and  $\sigma_1(v_3) = 6$  in  $TT_6$  and the embeddings  $\sigma_2$  and  $\sigma_3$  as follows:  $\sigma_2(v_0) = \sigma_3(v_0) = 1$   $\sigma_2(v_1) = 2$ ,  $\sigma_3(v_1) = 3$ , and  $\sigma_2(v_2) = 3$ ,  $\sigma_3(v_2) = 4$ , and  $\sigma_2(v_3) = 5$ ,  $\sigma_3(v_3) = 6$ .

Now, let  $n \geq 7$  and we assume that our result is true for all  $n' < n$ . Let  $v_0, \dots, v_k$  be the path  $\vec{P}$  of length  $k = \lfloor \frac{2}{3}n - 1 \rfloor$  in  $TT_n$ . By induction, there exist the embeddings  $\sigma'_1$ ,  $\sigma'_2$  and  $\sigma'_3$  of path  $v_0, \dots, v_{k-2}$  into  $TT_{n-3}$ . Moreover, we can assume that vertices  $\sigma'_1(v_{k-2}) = \sigma'_3(v_{k-2}) = n - 3$  and the number of  $\sigma'_2(v_{k-2})$  in  $TT_{n-3}$  is less than  $n - 3$ . Now we add three vertices to  $TT_{n-3}$  at the end. Two vertices  $v_{k-1}, v_k$  of the path obtain the numbers:  $n - 1$  and  $n$ , so  $\sigma_1(v_{k-1}) = n - 1$ ,  $\sigma_1(v_k) = n$ . We define the embeddings  $\sigma_2$  and  $\sigma_3$  in  $TT_n$  as follows:  $\sigma_2(v_{k-1}) = \sigma_3(v_{k-1}) = n - 2$  and  $\sigma_2(v_k) = n - 1$ ,  $\sigma_3(v_k) = n$ , and  $\sigma_1(v_i) = \sigma'_1(v_i)$ ,  $\sigma_2(v_i) = \sigma'_2(v_i)$ ,  $\sigma_3(v_i) = \sigma'_3(v_i)$  for  $i \in \{0, \dots, k - 2\}$ .

Thus, by induction, the proof is complete. ■

The following result may be proved in a similar way as Lemma 4.15 in [8].

**Lemma 6.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that*

- (a)  $x'x, y'y, z'z$ , or
- (b)  $xx', yy', zz'$

*are three independent end-arcs in  $E(\vec{G})$ . If  $\vec{H} := \vec{G} - \{x, y, z\}$  is 3-packable into  $TT_{n-3}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .*

**Lemma 7.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that  $z$  is an isolated vertex and*

- (c)  $x'x, y'y$ , or
- (d)  $xx', yy'$

are two independent end-arcs in  $E(\vec{G})$ . If  $\vec{H} := \vec{G} - \{x, y, z\}$  is 3-packable into  $TT_{n-3}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .




Figure 2. Two cases from Lemma 6




Figure 3. Two cases from Lemma 7

**Proof.** This lemma follows immediately from Lemma 6, (see Figure 2 and Figure 3). ■

**Lemma 8.** Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that  $y, z$  are two isolated vertices and  $x$  is a vertex such that

- (e)  $d^-(x) \geq 2, d^+(x) = 0$ , or
- (f)  $d^+(x) \geq 2, d^-(x) = 0$ .

If  $\vec{H} := \vec{G} - \{x, y, z\}$  is 3-packable into  $TT_{n-3}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .




Figure 4. Two cases from Lemma 8

**Proof.** Without loss of generality we can consider only the case (e). By assumption there exist arc disjoint embeddings  $\sigma'_1, \sigma'_2$  and  $\sigma'_3$  of  $\vec{H}$  into  $TT_{n-3}$ . Add three vertices to  $TT_{n-3}$  at the end and we obtain the transitive tournament  $TT_n$ .

Now, we define the embeddings of  $\vec{G}$ :  $\sigma_1(v) = \sigma'_1(v), \sigma_2(v) = \sigma'_2(v), \sigma_3(v) = \sigma'_3(v)$  for all vertices of  $\vec{H}$ , and  $\sigma_1(x) = n - 2, \sigma_2(x) = n - 1, \sigma_3(x) = n$ . This is the correct 3-packing of  $\vec{G}$  into  $TT_n$ , which completes the proof. ■

**Lemma 9.** Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that  $y, z$  are two isolated vertices in  $\vec{G}$ , the end-vertices  $x_1, \dots, x_k$  are adjacent to a vertex  $x$ , which is such that  $d^+(x) = t \geq 1, d^-(x) = k \geq 2$  and  $k + t \geq 4$ .

If  $\vec{H} := \vec{G} - \{x, y, z, x_1, \dots, x_k\}$  is 3-packable into  $TT_{n-3-k}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .




Figure 5. The case from Lemma 9

**Proof.** Let us imagine a transitive tournament  $TT_{n-3-k}$  with the vertices numbered from  $k + 4$  to  $n$ . Let us assume that embeddings  $\sigma'_1, \sigma'_2$  and  $\sigma'_3$  of

$\vec{H}$  exist in  $TT_{n-3-k}$ . Let us add  $k + 3$  vertices to  $TT_{n-3-k}$  at the beginning and we obtain the transitive tournament  $TT_n$ .

Now, we define the embeddings  $\sigma_1, \sigma_2$  and  $\sigma_3$  of  $\vec{G}$  into  $TT_n$  as follows:  $\sigma_1(x_i) = \sigma_2(x_i) = \sigma_3(x_i) = i$  for  $i \in \{1, \dots, k\}$ ,  $\sigma_1(x) = k + 1$ ,  $\sigma_2(x) = k + 2$ ,  $\sigma_3(x) = k + 3$ , and  $\sigma_1(v) = \sigma'_1(v)$ ,  $\sigma_2(v) = \sigma'_2(v)$ ,  $\sigma_3(v) = \sigma'_3(v)$  for all the remaining vertices. We obtain a 3-packing of  $\vec{G}$ . ■

**Lemma 10.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that  $x, y$  are two isolated vertices in  $\vec{G}$ ,  $a, b$  are two end-vertices adjacent to a vertex  $c$ . Let  $d$  be a vertex adjacent from  $c$  such that  $d^-(c) = 2$ ,  $d^+(c) = 1$ ,  $d^-(d) = 1$ ,  $d^+(d) \geq 1$ . If  $\vec{H} := \vec{G} - \{x, y, a, b, c, d\}$  is 3-packable into  $TT_{n-6}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .*




Figure 6. The case from Lemma 10

**Proof.** Let us imagine a transitive tournament  $TT_{n-6}$  with the vertices numbered from 7 to  $n$ . Let us assume that embeddings  $\sigma'_1, \sigma'_2$  and  $\sigma'_3$  of  $\vec{H}$  exist in  $TT_{n-6}$ . Let us add the vertices  $a, b, c, d, x, y$  to  $TT_{n-6}$  at the beginning and we obtain a transitive tournament  $TT_n$ .

We can define the embedding  $\sigma_1$  of  $\vec{G}$  into  $TT_n$  as follows:  $\sigma_1(a) = 1$ ,  $\sigma_1(b) = 2$ ,  $\sigma_1(c) = 3$ ,  $\sigma_1(d) = 4$ ,  $\sigma_1(x) = 5$ ,  $\sigma_1(y) = 6$  and  $\sigma_1(v) = \sigma'_1(v)$  for all the remaining vertices. Now, we define the embeddings  $\sigma_2$  and  $\sigma_3$  of  $\vec{G}$  into  $TT_n$  as follows:  $\sigma_2(a) = \sigma_3(a) = 1$ ,  $\sigma_2(b) = \sigma_3(b) = 2$ ,  $\sigma_2(c) = 4$  and  $\sigma_3(c) = 5$ ,  $\sigma_2(d) = 5$  and  $\sigma_3(d) = 6$  and  $\sigma_2(v) = \sigma'_2(v)$ ,  $\sigma_3(v) = \sigma'_3(v)$  for all the remaining vertices. So a 3-packing of  $\vec{G}$  into  $TT_n$  exists. ■

**Lemma 11.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$ . Suppose that  $a_k$  ( $k > 1$ ) is a vertex in  $\vec{G}$  such that a path of length  $k - 1$  from  $a_1$  to  $a_k$  exists and  $d^+(a_k) \geq 2$ . Moreover, suppose that  $y_1, \dots, y_{k'}$  ( $k' = \lfloor \frac{k+3}{2} \rfloor$ ) are isolated vertices in  $\vec{G}$ .*

If  $\vec{H} := \vec{G} - \{y_1, \dots, y_{k'}, a_1, \dots, a_k\}$  is 3-packable into  $TT_{n-k-k'}$ , then  $\vec{G}$  is 3-packable into  $TT_n$ .




Figure 7. The case from Lemma 11

**Proof.** Let us imagine a transitive tournament  $TT_{n-k-k'}$  with the vertices numbered from  $k + k' + 1$  to  $n$ . Let us assume that there are embeddings  $\sigma'_1, \sigma'_2$  and  $\sigma'_3$  of  $\vec{H}$  into  $TT_{n-k-k'}$ . Let us add  $k + k'$  vertices to  $TT_{n-k-k'}$  at the beginning and we obtain the transitive tournament  $TT_n$ .

In Lemma 5 we show that the path of length  $k - 1$  is 3-packable into  $TT_{\lfloor \frac{3}{2}k + \frac{1}{2} \rfloor}$ . So there are embeddings  $\sigma''_1, \sigma''_2$  and  $\sigma''_3$  of this path into  $TT_{k+k'-1}$ . Now we extend the embeddings  $\sigma''_1, \sigma''_2$  and  $\sigma''_3$  to embeddings  $\sigma^*_1, \sigma^*_2$  and  $\sigma^*_3$  into  $TT_{k+k'}$  with the last isolated vertex added. We will modify these embeddings if necessary so that  $\sigma^*_1(a_k) \neq \sigma^*_2(a_k) \neq \sigma^*_3(a_k)$ .

We consider three cases:

1. In the case of two embeddings of a path, the vertex  $a_k$  is embedded in the same vertex of  $TT_{k+k'-1}$ , for example  $\sigma''_1(a_k) \neq \sigma''_2(a_k) = \sigma''_3(a_k)$ ,
2. In the case of three embeddings of a path, the vertex  $a_k$  is embedded in the same vertex of  $TT_{k+k'-1}$  but not in the last, say  $\sigma''_1(a_k) = \sigma''_2(a_k) = \sigma''_3(a_k) = i < k + k' - 1$ ,
3. In the case of three embeddings of a path, the vertex  $a_k$  is embedded in the last vertex of  $TT_{k+k'-1}$ .

In the first case we may choose for  $\sigma^*_2(a_k)$  the last vertex of  $TT_{k+k'}$ .

In the second case we may choose  $\sigma^*_2(a_k) = k + k' - 1$  and  $\sigma^*_3(a_k) = k + k'$ .

In the third case we must have  $\sigma''_1(a_{k-1}) > \sigma''_2(a_{k-1}) > \sigma''_3(a_{k-1})$ . If  $\sigma''_1(a_k) - \sigma''_1(a_{k-1}) > 1$ , then we may assume  $\sigma^*_1(a_k)$  is in the  $k + k' - 2$  vertex, and  $\sigma^*_2(a_k) = k + k'$ . If  $\sigma''_1(a_k) - \sigma''_1(a_{k-1}) = 1$  (in  $TT_{k+k'-1}$ ), then either we may assume  $\sigma^*_2(a_k)$  is in the  $k + k' - 2$  vertex or we may assume  $\sigma^*_3(a_k)$  is in the  $k + k' - 2$  vertex and the other one in  $k + k'$  vertex.



Now  $\sigma_1^*(a_k) \neq \sigma_2^*(a_k) \neq \sigma_3^*(a_k)$  and we can define the embeddings  $\sigma_1, \sigma_2$  and  $\sigma_3$  of  $\vec{G}$  into  $TT_n$  as follows:  $\sigma_1(a_i) = \sigma_1^*(a_i), \sigma_2(a_i) = \sigma_2^*(a_i), \sigma_3(a_i) = \sigma_3^*(a_i)$  for all  $i \in \{1, \dots, k\}, \sigma_1(y_j) = \sigma_1^*(y_j), \sigma_2(y_j) = \sigma_2^*(y_j), \sigma_3(y_j) = \sigma_3^*(y_j)$  for all  $j \in \{1, \dots, k'\}$  and  $\sigma_1(v) = \sigma_1'(v), \sigma_2(v) = \sigma_2'(v), \sigma_3(v) = \sigma_3'(v)$  for all the remaining vertices. ■

### 3. The Main Result

In this section, we consider the existence of a 3-packing of  $\vec{G}$  into  $TT_n$  and we prove the following theorem.

**Theorem 12.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$  such that  $|E(\vec{G})| \leq \frac{2}{3}n - 1$ . Then  $\vec{G}$  is 3-packable into  $TT_n$ .*

#### 3.1 The bound in Theorem 12 is the best possible

First, we show that the size condition in Theorem 12 cannot be weakened.

Let us consider a path of length  $k$  and suppose that a 3-packing of such a path into  $TT_n$  exists, where  $n > k$ . It means that  $\vec{G}, \vec{G}'$  and  $\vec{G}''$  are three arc disjoint subgraphs of the transitive tournament  $TT_n$  isomorphic to such a path. Let  $k_1, k'_1$  and  $k''_1$  denote the number of arcs of length one in  $\vec{G}, \vec{G}'$  and  $\vec{G}''$ ,  $k_2, k'_2$  and  $k''_2$  denote the number of arcs of length two and  $k_3, k'_3$  and  $k''_3$  denote the number of arcs of length greater than two, respectively. Thus

$$(*) \quad \left. \begin{aligned} k_1 + k_2 + k_3 &= k, \\ k'_1 + k'_2 + k'_3 &= k, \\ k''_1 + k''_2 + k''_3 &= k. \end{aligned} \right\}$$

Since  $\vec{G}, \vec{G}'$  and  $\vec{G}''$  are subgraphs of  $TT_n$ , we have

$$k_1 + 2k_2 + 3k_3 \leq n - 1,$$

$$k'_1 + 2k'_2 + 3k'_3 \leq n - 1,$$

$$k''_1 + 2k''_2 + 3k''_3 \leq n - 1.$$

By adding the last three inequalities we get

$$k_1 + k'_1 + k''_1 + 2k_2 + 2k'_2 + 2k''_2 + 3k_3 + 3k'_3 + 3k''_3 \leq 3n - 3.$$

But on the other hand, since  $\vec{G}$ ,  $\vec{G}'$  and  $\vec{G}''$  are arc disjoint and the total number of arcs of length 1 in  $TT_n$  is equal to  $(n - 1)$ , we have:

$$2(k_1 + k'_1 + k''_1) \leq 2(n - 1)$$

and since the total number of arcs of length 2 in  $TT_n$  is equal to  $(n - 2)$ , we have:

$$k_2 + k'_2 + k''_2 \leq n - 2.$$

By adding these three inequalities and using (\*) we get

$$9k \leq 6n - 7.$$

Finally, we obtain

$$k \leq \frac{2}{3}n - 1.$$

### 3.2 Proof of Theorem 12

At the beginning, we can notice that for  $n \leq 4$  an oriented graph satisfying the assumption of Theorem 12 has zero or one arc and, obviously, is 3-packable into  $TT_n$ . For  $n = 5$  an oriented graph satisfying the assumption of Theorem 12 has at most two arcs and it is also easily seen that it is 3-packable.

Now, let us assume that  $\vec{G}$  is a counterexample of Theorem 12 for minimum possible  $n \geq 6$ .

Let us notice that for  $6 \leq n \leq 9$ , if  $\vec{G}$  does not have any isolated vertex and has, of course, at most  $\frac{2}{3}n - 1$  edges, then  $\vec{G}$  has only tree-components and at least three of them are isolated arcs. So by Lemma 6, we get a contradiction with the minimality of  $\vec{G}$ .

As above, if  $\vec{G}$  (for  $6 \leq n \leq 9$ ) has only one isolated vertex, then  $\vec{G}$  has at least two isolated arcs (for  $7 \leq n \leq 9$ ) or one isolated arc and one end-arc ( $n = 6$ ). So by Lemma 7, we get a contradiction with the minimality of  $\vec{G}$ . Hence in the next part of the proof we can assume that for  $n \leq 9$   $\vec{G}$  has at least two isolated vertices.

It is obvious that every oriented graph  $\vec{G}$ , for  $n \geq 10$  which satisfies the conditions of Theorem 12 is not connected and at least  $\lceil \frac{n}{3} + \frac{7}{9} \rceil$  of its components are oriented trees (including, the isolated points as trivial oriented trees). If in  $\vec{G}$  there are more than four non-trivial oriented trees as its components, then  $\vec{G}$  has at least five independent end-vertices. So three

of them have to be such as in case (a) or (b) in Lemma 6. We get a contradiction with the minimality of  $\vec{G}$ . Hence  $\vec{G}$  has at most four components being non-trivial oriented trees and at least  $\lceil \frac{n}{3} + \frac{7}{9} \rceil$  of its components are oriented trees. For order  $n \geq 10$  we obtain an isolated point in  $\vec{G}$ .

Now, if in  $\vec{G}$  there are more than two non-trivial oriented trees as its components, then  $\vec{G}$  has at least three independent end-vertices. So two of them have to be such as in case (c) or (d) in Lemma 7 and since in  $\vec{G}$  there is an isolated vertex, we get a contradiction with the minimality of  $\vec{G}$ .

Hence from this moment in the proof (for order  $n \geq 6$ )  $\vec{G}$  has at most two components being non-trivial oriented trees and at least  $\max\{2, \lceil \frac{n}{3} - \frac{11}{9} \rceil\}$  of its components are isolated vertices.

Let  $\vec{H}$  be a non-trivial connected component of  $\vec{G}$  of the greatest order. Let a vertex  $x \in V(\vec{H})$  be such that  $d^-(x) = 0$ . It is easily seen that there is not more than one vertex adjacent from  $x$ , since if there is more than one, then  $\vec{G}$  satisfies the assumptions of Lemma 8 and it leads to a contradiction with the minimality of  $\vec{G}$ .

It means that  $d^+(x) = 1$ . If  $y$  is a neighbour of  $x$ ,  $\vec{G}$  satisfies one of the following properties:

1.  $d^-(y) \geq 3$ ;
2.  $d^-(y) = 2$  and  $d^+(y) \geq 2$ ;
3.  $d^-(y) = 2$  and  $d^+(y) \leq 1$ ;
4. there is a path  $(a_1 = x, a_2 = y, \dots, a_k)$ ,  $k \geq 2$  and  $d^+(a_k) \geq 2$ ;
5.  $\vec{G}$  is an oriented path.

It is easily seen that in the first, the second and the third case we may assume that all vertices adjacent to  $y$  are end-vertices. If not, in the graph  $\vec{G}$  either there are two end-vertices like in Lemma 7 or there is a vertex with indegree zero and outdegree greater than or equal to 2, hence it satisfies the assumptions of Lemma 8. In both the cases we obtain a contradiction with the minimality of  $\vec{G}$ .

*Case 1.* It is obvious that in this case such a graph is 3-packable since either  $d^+(y) = 0$  and it satisfies the assumptions of Lemma 8 or  $d^+(y) > 0$  and the assumptions of Lemma 9.

*Case 2.* Such a graph is 3-packable since it satisfies the assumptions of Lemma 9.

*Case 3.* As in the first case, if  $d^+(y) = 0$ , it satisfies the assumptions of Lemma 8.

Let  $d^+(y) = 1$  and  $z$  be a vertex adjacent from  $y$ . If  $d(z) = 1$ , assume first that  $\vec{H}$  is a not unique non-trivial component of  $\vec{G}$ . In the second non-trivial component  $\vec{K}$  of  $\vec{G}$  there is a vertex  $v \in V(\vec{K})$  such that  $d^-(v) = 0$ . For the same reason as before the outdegree of  $v$  must be equal to 1. And then there are two end-arcs: one ending in  $x$  and the other ending in  $v$ , so by Lemma 7  $\vec{G}$  is 3-packable, which contradicts the minimality of  $\vec{G}$ . Hence in this case  $\vec{G}$  has a unique non-trivial component  $\vec{H}$ . So  $\vec{H}$  has three arcs and in  $\vec{G}$ , which satisfies the assumption of Theorem 12, there are two isolated vertices. Three copies of such a graph can be packed in the same way as in the proof of Lemma 10, but  $\vec{G}$  is not 3-packable, so  $d^-(z) > 1$ .

If  $d^-(z) > 1$ , then two end-vertices, like in Lemma 7, exist in the graph  $\vec{G}$  and  $\vec{G}$  is 3-packable. If  $d^-(z) = 1$  and  $d^+(z) \geq 1$  such a graph is 3-packable since it satisfies the assumptions of Lemma 10.

*Case 4.* We may observe that if  $d^-(a_i) > 1$ , for any  $i > 2$ , in the graph  $\vec{G}$  either there are two end-vertices like in Lemma 7 or there is a vertex with indegree zero and outdegree greater than or equal to 2, hence it satisfies the assumptions of Lemma 8. In both the cases we obtain a contradiction with the minimality of  $\vec{G}$ .

It is obvious that in the fourth case such a graph is 3-packable since it satisfies the assumptions of Lemma 11.

*Case 5.* Such a graph is 3-packable since it satisfies the assumptions of Lemma 5.

Therefore the set of counterexamples is empty and the proof of Theorem 12 is complete. ■

#### 4. A Conjecture — $m$ -Packable into $TT_n$

Finally we can make a general conjecture.

**Conjecture 13.** *Let  $\vec{G}$  be an acyclic digraph of order  $n$  such that  $|E(\vec{G})| \leq \frac{m+1}{2m}n - \frac{m^2+5}{6m}$ . Then  $\vec{G}$  is  $m$ -packable into  $TT_n$ .*

We show only that the size condition in Theorem 13 cannot be weakened. Let us consider a path of length  $k$ . Then we suppose that there is an

$m$ -embedding of such a path into  $TT_n$ , where  $n > k$ . It means that  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_m$  are  $m$  arc disjoint subgraphs of the transitive tournament  $TT_n$  isomorphic to such a path. Let for  $1 \leq i \leq m - 1$ ,  $k_1^i, k_2^i, \dots, k_m^i$  denote the numbers of arcs in  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_m$  of length  $i$  in  $TT_n$  and  $k_1^m, k_2^m, \dots, k_m^m$  denote the number of arcs in  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_m$  of length greater than  $m - 1$ , respectively. Thus

$$(*) \left. \begin{aligned} k_1^1 + k_1^2 + \dots + k_1^m &= k, \\ k_2^1 + k_2^2 + \dots + k_2^m &= k, \\ &\dots \\ k_m^1 + k_m^2 + \dots + k_m^m &= k. \end{aligned} \right\}$$

Since  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_m$  are subgraphs of  $TT_n$  we have for each  $\vec{G}_m$

$$k_i^1 + 2k_i^2 + \dots + mk_i^m \leq n - 1.$$

By adding those inequalities we get

$$\sum_{i=1}^m k_i^1 + 2 \sum_{i=1}^m k_i^2 + \dots + m \sum_{i=1}^m k_i^m \leq mn - m.$$

But on the other hand, since  $\vec{G}_1, \vec{G}_2, \dots, \vec{G}_m$  are arc disjoint and the total number of arcs of length 1 is equal to  $n - 1$  we have:

$$(m - 1) \sum_{i=1}^m k_i^1 \leq (m - 1)(n - 1),$$

Since the total number of arcs of length 2 is equal to  $n - 2$  we have:

$$(m - 2) \sum_{i=1}^m k_i^2 \leq (m - 2)(n - 2)$$

and similar inequalities, up to

...

$$\sum_{i=1}^m k_i^{m-1} \leq (n - m + 1).$$

By adding these inequalities and using (\*) we obtain

$$m^2k \leq (m+m-1+m-2+\dots+1)n - (m+1)(m-1) + 2(m-2) + \dots + (m-1)1$$

hence finally

$$k \leq \frac{m+1}{2m}n - \frac{m^2+5}{6m}.$$

### Acknowledgement

The author is indebted to Mariusz Woźniak for his helpful comments and remarks.

### References

- [1] B. Bollobás, *Extremal Graph Theory* (Academic Press, London, 1978).
- [2] B. Bollobás and S.E. Eldridge, *Packings of graphs and applications to computational complexity*, J. Combin. Theory **25** (B) (1978) 105–124.
- [3] D. Burns and S. Schuster, *Every  $(n, n-2)$  graph is contained in its complement*, J. Graph Theory **1** (1977) 277–279.
- [4] A. Görlich, M. Pilśniak and M. Woźniak, *A note on a packing problem in transitive tournaments*, preprint Faculty of Applied Mathematics, University of Mining and Metallurgy, No.37/2002.
- [5] H. Kheddouci, S. Marshall, J.F. Saclé and M. Woźniak, *On the packing of three graphs*, Discrete Math. **236** (2001) 197–225.
- [6] N. Sauer and J. Spencer, *Edge disjoint placement of graphs*, J. Combin. Theory **25** (B) (1978) 295–302.
- [7] M. Woźniak and A.P. Wojda, *Triple placement of graphs*, Graphs and Combin. **9** (1993) 85–91.
- [8] M. Woźniak, *Packing of graphs*, Dissertationes Math. **362** (1997).
- [9] H.P. Yap, *Some Topics in Graph Theory*, London Math. Society, Lectures Notes Series, Vol. 108 (Cambridge University Press, Cambridge, 1986).
- [10] H.P. Yap, *Packing of graphs – a survey*, Discrete Math. **72** (1988) 395–404.

Received 29 April 2003

Revised 8 March 2004